

RESEARCH

Open Access

Thermal adaptability of *Kluyveromyces marxianus* in recombinant protein production

Stefano Raimondi¹, Elena Zanni², Alberto Amaretti¹, Claudio Palleschi², Daniela Uccelletti^{2*} and Maddalena Rossi^{1*}

Abstract

Background: *Kluyveromyces marxianus* combines the ease of genetic manipulation and fermentation with the ability to efficiently secrete high molecular weight proteins, performing eukaryotic post-translational modifications. It is able to grow efficiently in a wide range of temperatures. The secretion performances were analyzed in the host *K. marxianus* L3 in the range between 5°C and 40°C by means of 3 different reporter proteins, since temperature appears a key parameter for production and secretion of recombinant proteins.

Results: The recombinant strains were able to grow up to 40°C and, along the tested temperature interval (5-40°C), the specific growth rates (μ) were generally lower as compared to those of the untransformed strain. Biomass yields were slightly affected by temperature, with the highest values reached at 15°C and 30°C. The secretion of the endogenous β -fructofuranosidase, used as an internal control, was efficient in the range of the tested temperature, as evaluated by assaying the enzyme activity in the culture supernatants. The endogenous β -fructofuranosidase production was temperature dependent, with the highest yield at 30°C. The heterologous proteins HSA, GAA and Sod1p were all successfully produced and secreted between 5°C and 40°C, albeit each one presented a different optimal production temperature (15, 40, 5-30°C for HSA, GAA and Sod1p, respectively).

Conclusions: *K. marxianus* L3 has been identified as a promising and flexible cell factory. In a sole host, the optimization of growth temperatures for the efficient secretion of each individual protein can be carried out over a wide range of temperatures.

Background

Kluyveromyces marxianus is one of the alternative yeasts for which an efficient transformation system was developed [1]. It has a long history of usage in food fermentation, and holds the GRAS status (Generally Regarded As Safe) that allows straightforward food and pharmaceutical applications of this microorganism and its derivatives [2]. *K. marxianus* grows on cheap lactose based media, such as whey and inulin, ferments xylose to ethanol, and does not require expensive explosion-proof plants which are necessary for methylotrophic yeasts [3,4].

Ethanol production represents one of the pivotal field of *K. marxianus* utilization. In the fermentation of renewable biomass residues, its capability to simultaneously consume glucose and xylose can be exploited in order to maximize

ethanol productivity [4]. Recently, *K. marxianus* was also employed for tequila production at industrial level: the strain showed a higher osmotic (22% w/v) and ethanol (10% v/v) tolerance, producing high levels of volatiles [5].

However, the intrinsic advantages of *K. marxianus* over other yeasts will be better utilized for a variety of biotechnological applications as more and well-defined genetic engineering tools will become available. Recently, four *S. cerevisiae* promoters were characterized in *K. marxianus* for promoter strength, time-dependent changes, and stochastic gene expression pattern [6]. In addition, the genome of the thermotolerant *K. marxianus* KCTC 17555 that can convert inulin-rich plant biomass into ethanol and/or platform biochemicals was sequenced and analysed [7].

For heterologous gene expression, *K. marxianus* represents a valid alternative because of its outstanding secretory capabilities [8-10]. *K. marxianus* can also produce eukaryotic proteins since it modifies foreign proteins according to a general eukaryotic scheme. Moreover, *K. marxianus* has been exploited as a cell factory to obtain

* Correspondence: daniela.uccelletti@uniroma1.it; maddalena.rossi@unimore.it

¹Department of Life Sciences, University of Modena and Reggio Emilia, Via Campi 183, 41125, Modena, Italy

²Department of Biology and Biotechnology Charles Darwin, University of Rome "La Sapienza", Piazzale Aldo Moro 5, 00185, Rome, Italy

valuable enzymes, showing retention of activity in a large temperature interval [11].

A peculiar trait of this yeast is the capacity to grow within a wide interval of temperatures, ranging from 4°C to 45°C. This feature has been investigated in a study where the adaptive responses at different temperatures of psychrophilic and mesophilic yeasts were compared [12]. The main objective of the current study was to explore the capability of *K. marxianus* to secrete various heterologous proteins at different temperatures. In fact, for both eukaryotic and prokaryotic hosts, it has been reported that growth temperature affects the yield of properly folded recombinant proteins [13,14]. In *E. coli* a common strategy to express the target protein in a soluble state consists in evaluating different growth temperatures. Temperature lowering improves the solubility of the proteins by diminishing aggregation as a result of a decrease in the production rate, allowing the newly synthesized chain to fold properly [15,16]. Several studies explored the effect of the growth temperature on the expression and secretion of heterologous proteins also in yeasts, albeit, in most cases, the authors investigated limited arrays of temperatures, generally ranging between 15 and 30°C [17-19].

Herein, the impact of different temperatures on endogenous and recombinant protein production and secretion was assessed in *K. marxianus* within a wider range of temperatures, ranging between 5 and 40°C. The reporter genes encoding the glucoamylase (GAA) from the yeast *Arxula adenivorans* [20], the recombinant human serum albumin (HSA) [21], and the *Kluyveromyces lactis* Cu²⁺/Zn²⁺ superoxide dismutase (SOD1) [22] were used.

Results and discussion

To the best of our knowledge, the capability to grow both at mesophilic and psychrophilic temperatures has never been investigated for a unique eukaryotic host of recombinant proteins. *Kluyveromyces marxianus* can grow in a wide range of temperatures, it is still an under-exploited biotechnological system, and few studies have so far explored its potential for heterologous protein production [8,9]. In the present work, *K. marxianus* has been challenged as a cell factory model to investigate the impact of temperature on the secretion efficiency of endogenous and recombinant proteins.

Specific growth rate and biomass yield of *Kluyveromyces marxianus* L3

Growth rate and biomass yield are physiological features of major importance for a “cell factory” organism in order to reach high volumetric productivities of recombinant proteins. It is generally assumed that growth at temperatures below the optimal value positively affects production and secretion of recombinant proteins. In this perspective,

we investigated *K. marxianus* L3 specific growth rates (μ) and biomass yield (g L⁻¹) at different temperatures (5, 15, 30, 40, and 45°C; Table 1), exploiting the advantage of this heat-tolerant yeast, which optimum temperature for growth falls generally around 42°C.

As expected, the temperature markedly affected the specific growth rate (μ). The μ increased if the temperature raised, and it was highest at 45°C (0.464 h⁻¹). This behavior was in accordance with data reported in literature indicating that all the *K. marxianus* strains are able to grow at 42°C, and only few strains can grow up to 48°C [23].

Biomass yields were slightly affected by temperature, with the highest values reached at 15 and 30°C (12.8 and 12.9 g L⁻¹, respectively). These high yields confirmed the excellent capability of *K. marxianus* to convert substrate into biomass [24]. The lower values at the highest temperatures (10.6 and 9.8 g L⁻¹ at 40 and 45°C, respectively) suggested that, even if the growth rate was high, cell reproduction at the highest temperature required a metabolic cost that affected the biomass yield.


Secretion of endogenous β -fructofuranosidase

In order to assess the secretory capability of the host strain *K. marxianus* L3, the activity of endogenous β -fructofuranosidase (EC 3.2.1.26) was analyzed in the supernatant of cultures grown between 5 and 45°C. At all the tested temperatures, enzyme activity increased during both the growth and the stationary phases (Figure 1). The extracellular β -fructofuranosidase activity was temperature dependent, with the highest yield at 30°C (Figure 1). With the exception of 5°C, the enzyme production was abundant also at lower and higher temperatures. This result was consistent with the data reported in literature [25-27], and confirmed that *K. marxianus* can be regarded as a good secretor, acting as a thermally flexible cell factory able to secrete active forms of an homologous enzyme in a wide range of temperatures.

Table 1 Effects of the temperature on growth of *K. marxianus* L3 wild-type and recombinant strains

T (°C)	Wild-type DW (g L ⁻¹)	Wild-type Q _x (g L ⁻¹ h ⁻¹)	Wild-type μ (h ⁻¹)	rHSA μ (h ⁻¹)	GAA μ (h ⁻¹)	KISod1p μ (h ⁻¹)
5	10.5 [#]	0.096	0.041	0.034 ^a	0.031 ^a	0.034 ^a
15	12.8 [‡]	0.355	0.171	0.138 ^a	0.160	0.144 ^a
30	12.9 [‡]	0.479	0.377	0.321 ^a	0.341 ^a	0.271
40	10.6 [#]	0.443 [‡]	0.439	0.293 ^a	0.369	0.313 ^a
45	9.8	0.426 [‡]	0.464	-	-	-

The strains were cultured at different temperatures in YPD medium containing 20 g L⁻¹ glucose. The specific growth rates (μ), the biomass yield (expressed as dry weight grams per litre of culture) and the biomass volumetric productivities (Q_x) are reported as a function of growth temperature. The values are means of three independent experiments, with standard deviations always lower than 5%. Superscripts within a column (symbols) and within a row (letters) indicate statistically similar means, P > 0.05.



Secretion of heterologous proteins

In a previous study the comparison of expression and secretion of recombinant proteins at different temperatures has been carried out by using different hosts [28]. Herein, *K. marxianus* L3 was challenged for the secretion of three recombinant proteins in a wide range of temperatures. Specific growth rates (μ) were determined for each recombinant strain expressing the heterologous proteins HSA, GAA, or KISod1p at 5, 15, 30, 40, and 45°C (Table 1). The recombinant strains were not able to grow at 45°C, possibly because of a low segregational stability of recombinant plasmids at the highest growth temperature, as reported in both prokaryotic and eukaryotic hosts [29,30].

In the explored range of temperature, the μ increased by raising the temperature, with the exception of HSA that presented $\mu_{40^\circ\text{C}} < \mu_{30^\circ\text{C}}$. When heterologous proteins were expressed, generally μ were lower respect to the host (Table 1), reflecting a higher metabolic load [31,32].

HSA production

HSA was successfully produced and secreted in the range between 5 and 40°C, and the growth phase affected specific HSA production (Figure 2A). The yields, expressed as relative abundances, showed a peak at 15°C both in exponential and stationary phases. A notably secretion efficiency was observed at all tested temperatures, when compared to the highest reference value (100% at 15°C). Compared to secretion levels, intracellular HSA production in both the




Figure 2 Effects of the growth temperature on HSA secretion.

Culture samples, grown at 5 - 40°C, have been analyzed both at the growth and stationary phases. **A)** Western blot analysis of secreted rHSA; **B)** Western blot analysis of intracellular rHSA; **C)** Q-PCR analysis of rHSA mRNA. Data are reported as a function of growth phase and temperature. The values, means of three independent experiments \pm standard deviations, have been normalized setting at 1 the highest value. Superscripts indicate statistically similar means ($P > 0.05$). Symbols are used to compare sample grown at the same temperature, collected at different growth phases (exponential or stationary). Letters are used to compare samples collected at the same growth phase, cultured at diverse temperatures.

exponential and stationary phases presented the opposite behavior, being the lowest at 15°C and the highest at 40°C (Figure 2B). Conceivably, the lower intracellular HSA at 15°C was associated to a more efficient secretion, and *vice versa* for the higher temperatures. A RT-PCR analysis was

performed to determine transcription levels at the different temperatures (Figure 2C). At 5°C and 15°C very low amounts of HSA mRNA were observed during both exponential and stationary phases. At the higher temperatures, the relative amount of the messenger was always greater in stationary phase, and increased by rising the temperature. Although high temperatures seem not to drive the *ADH4* promoter activity in *S. cerevisiae* or in *K. lactis* [33], we could not exclude this possibility in the *K. marxianus* background. This aspect will deserve further investigations.

GAA production

Heterologous GAA was successfully produced by *K. marxianus* at all the tested temperatures and the active enzyme accumulated in the medium over the time (Figure 3A). During the exponential phase production increased by rising the growth temperature up to 40°C. In the stationary phase, the total amount of active GAA secreted in the medium was similar over the range of temperatures between 5 and 30°C, whereas it was higher at 40°C. Transcript analysis showed that the relative amount of GAA mRNA had a slight variation between 5°C and 30°C in both growth phases; a significant increase was instead observed at 40°C ($p < 0.05$) (Figure 3B).

KISod1p production

The secretion efficiency of active Sod1p was compared at different temperatures by quantitative analysis of zymogram bands (Figure 4A). During the exponential phase, the capability to secrete active Sod1p increased by lowering the temperature. In stationary phase the production of active enzyme was similar between 5°C and 30°C and was significantly reduced at 40°C.

Transcriptional analysis showed that *SOD1* expression was almost not responsive to temperature up to 30°C; the level of transcript increased significantly at 40°C (Figure 4B). Actually, the genes *KmSOD1* of *K. marxianus* and *KISOD1* of *K. lactis* present a 83% identity, it is then conceivable that the employed probe hybridized to both transcripts. Furthermore, endogenous Sod1p should be upregulated as a key part of a heat shock response. In *S. cerevisiae* oxidative stress and antioxidant enzymes play a pivotal role in heat-induced cell death and it has been observed that overexpression of catalase and superoxide dismutase genes caused increased thermotolerance [34].

Taken together, our results support *K. marxianus* L3 as a thermally flexible cell factory. This strain can be adopted in a process optimized for the 'most productive temperature' whenever no particular constraints were placed by the produced protein. Conversely, the strain L3 can be employed at the 'most suitable temperature' in case of specific protein requirements (e.g. stability at low temperature).





Figure 3 Effects of the growth temperature on GAA secretion.

Culture samples, grown at 5 - 40°C, have been analyzed both at the growth and stationary phases. **A)** extracellular GAA activity; **B)** northern blot analysis of GAA mRNA. Data are means of three independent experiments \pm standard deviations. The values, means of three independent experiments \pm standard deviations, have been normalized setting at 1 the highest value. Superscripts indicate statistically similar means ($P > 0.05$). Symbols are used to compare samples grown at the same temperature, collected at different phases (exponential or stationary). Letters are used to compare samples collected at the same growth phase, cultured at different temperatures.

Conclusions

In this study, the yeast *K. marxianus* has proved to be a handy model system to investigate the impact of temperature on cell factory efficiency. It resulted in a thermally flexible host able to produce and secrete three different recombinant proteins between 5 and 40°C. Notably, for the first time, the impact of a wide range of temperatures on the production of diverse heterologous proteins has been explored by using a single biotechnological platform. A peculiar behavior of each recombinant protein with respect to the temperature was observed, strengthening the advantage of using *K. marxianus* as unique host. Because of its ability to grow between 5°C and 40°C, and efficiently secrete heterologous proteins, it can be exploited as a first-strike model system to study the temperature tuning and optimization of the production. Moreover *K. marxianus*



meets the criteria of process safety and the demand for fast growth and high yield biomass.

Methods

Strains and media

Kluyveromyces marxianus L3 [35] was used as expression host. Yeast cells were grown in YPD medium (10 g l⁻¹ yeast extract, 10 g l⁻¹ peptone, and 20 g l⁻¹ glucose). Supplement of antibiotic G418 (0.2 g l⁻¹) was added for the maintenance of recombinant plasmids. Standard batch cultures were carried out in 500 ml baffled Erlenmeyer flasks with 50 ml YPD medium. Flasks were inoculated (10% v/v) with exponential phase cultures, grown at the same temperature, to an OD₆₀₀ of 1.0, then were incubated at 5, 15, 30, 40, and 45°C for 7, 4, 3, 2 and 2 days, respectively. For each cultures, samples corresponding to the exponential growth phase and to the stationary phase were analyzed.

For strains harboring the plasmids pYG-K1KISOD1 [22] and pGAM-GAM [20], heterologous protein expression was induced after 24 h of growth by the addition of ethanol to a final concentration of 2% v/v. All the experiments were performed in triplicate.

Escherichia coli DH5α (φ80lacZΔM15, recA1, end A1, gyr A96, thi-1, hsd R17, rel A1) was used for general cloning purposes.

Plasmids and DNA manipulation

The pGM-GAM, pYG132, and pYG-K1KISOD1 plasmids were utilized for the expression of the reporter proteins. pGM-GAA contains the glucoamylase (GAA) gene from the yeast *Arxula adenivorans* under the control of the *S. cerevisiae* GAPDH promoter and of the *S. cerevisiae* PHO5 terminator [20]. pYG132 harbors an expression cassette for the secretion of the recombinant human serum albumin (HSA), driven by the native signal sequence, under the control of the ethanol inducible *K. lactis* ADH4 promoter and of the *K. lactis* PGK terminator [21]. pYG-K1KISOD1 carries the *K. lactis* KISOD1 gene encoding the the Cu²⁺/Zn²⁺ superoxide dismutase fused with the K1 secretion signal, under the control of the *K. lactis* ADH4 promoter and PGK terminator [22]. *K. marxianus* L3 was transformed by electroporation with a Biorad Gene-Pulser apparatus, as reported in Raimondi et al. [22].

Calculation of the specific growth rate, biomass yield and volumetric productivity

For biomass determination, an appropriate volume of the culture was centrifuged. The pellet was washed in d. d. water and was desiccated under an infrared heat lamp. The dry weight was determined gravimetrically and the biomass yield was expressed as grams of dry weight per litre of culture broth. The specific growth rate (μ) was calculated using dry weight values in the exponential phase of the growth curve. Glucose consumption was analyzed by an HPLC equipped with refractive index detector (HPLC System, 1200 Series, Agilent Technologies, Santa Clara, CA). The analysis was performed with an Aminex HPX-87H ion exclusion column and 0.005 M H₂SO₄ (0.6 ml min⁻¹) as the mobile phase. The volumetric productivities of biomass Q_X were calculated by dividing the biomass yield with the corresponding culture time.

β-fructofuronidase activity

β-fructofuronidase was determined as inulinase activity according to Burkert et al. [36]. Culture was centrifuged for 5 minutes at 18000 g and 1 ml of supernatant, properly diluted in acetate buffer 0.1 M at pH 5, was mixed with 9 ml of 20 g l⁻¹ sucrose dissolved in acetate buffer. The sample was incubated at 60°C and the rate of glucose and fructose production was determined by the 3,5-dinitrosalicylic acid method [37]. One unit of invertase is defined as the amount

of enzyme catalyzing the liberation of 1 μmol of reducing sugars per minute under the conditions above mentioned.

Western-blot analysis of recombinant HSA

For HSA Western-blot analysis, proteins from the supernatant (10 μl of 1:10 dilution) were separated with SDS-PAGE using the buffer system of Laemmli [38] and 12% acrylamide gels. After electroblotting onto a polyvinylidene difluoride membrane (Biorad), target proteins were detected with specific polyclonal antibodies of rabbit. Anti-HSA primary antibodies (Sigma) were used in 1:10,000 dilution. Monoclonal anti-rabbit IgG conjugated with peroxidase (Promega) were used as secondary antibodies. Immunologically active proteins were visualized with enhanced chemiluminescence detection system (GE Healthcare), according to the manufacturer's instructions. The densitometric analysis was performed with an image analyzer (Phoretix 1D; Non Linear Dynamics Ltd.).

RT-PCR analysis

Total RNA of *K. marxianus* cells was extracted by the hot phenol method [39]. RNA was quantified by absorbance (A_{260}) and subjected to TURBO DNase treatment according to the manufacturer's instructions (Ambion). Reverse transcription was performed using a Promega Reverse Transcription System with 1 μg total RNA to yield 20 μl cDNA. After cDNA generation, samples were purified, quantified with Nanodrop™ 3300 (Thermo Scientific), diluted with DEPC water (Ambion) to a final concentration of 10 ng μl^{-1} , and 2 μl were used for SYBR Green SensiMix (Bioline). Oligonucleotides (purchased from SIGMA) were designed with Primer3 (<http://frodo.wi.mit.edu/primer3/> website), considering an amplicon size of 95–220 bp and a T_m of approximately 55°C. For the HSA the primers used were 5'- GTTGAACCTTCGTGAAAC-3' and 5'- AAGTAAGGATGTCTTC TGGC-3'; for the *KmACT1* they were 5'- CTCCTT GCCTCATGCTATC-3' and 5'-GAAGGAGTAACCACG TTCAG-3'.

Glucoamylase activity

GAA activity was determined at 37°C as starch-hydrolyzing activity of the cell-free culture broths, according to Morlino et al. [20] One unit of GAA activity was defined as the quantity of enzyme needed to decrease the absorbance at 580 nm by one absorbance unit per minute. GAA activity was expressed as volumetric activity (U l^{-1}).

Superoxide dismutase activity

SOD activity was determined by on native gel by the nitroblue tetrazolium method [40]. During illumination with light, the gel became uniformly blue-purple except at positions containing active SOD protein. The densitometric

analysis was carried out with the image analyzer Phoretix 1D (Non Linear Dynamics Ltd).

Northern blot analysis

Total RNA was prepared by extraction with hot acidic phenol [41]. Northern blot analysis was performed as previously described according to Sambrook et al. [42]. The *A. adenivorans* GAA probe corresponded to the 2.1-kb *HindIII* region of the pGM-GAM plasmid. The 0.7 kb fragment of the *KlKISOD1* gene, amplified using the set of primers K1-KISOD1_f (5'-GC AAG CTT ATG AAT ATA TTT TAC ATA TTT TTG TTT TTG CTG TCA TTC GTT CAA GGT ACC CGG GGA GTT AAT GCA GTT GCA G-3') and KISOD1_r (5'-GC AAG CTT TTA AGC GTT AGA GAT ACC-3'), was used to detect *KISOD1* mRNA. The probes were labeled with [α - ^{32}P]dATP by use of the Ready Prime DNA labeling system (GE Healthcare), according to the manufacturer's instructions.

Statistical analysis

All values are means of three separate experiments. Differences in means were analyzed using Student's *t* test with independent measures. Differences were considered statistically significant if $P < 0.05$.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SR, DU and MR conceived the study, participated in its design and wrote the manuscript. SR, EZ, and AA performed the experimental work and drafted the manuscript. CP has been involved in drafting the manuscript and revising it critically. All authors read and approved the final manuscript.

Received: 21 December 2012 Accepted: 26 March 2013

Published: 15 April 2013

References

1. Iborra F: High efficiency transformation of *Kluyveromyces marxianus* by a replicative plasmid. *Curr Genet* 1993, **24**:181–183.
2. Fonseca GG, Heinze E, Wittmann C, Gombert AK: The yeast *Kluyveromyces marxianus* and its biotechnological potential. *Appl Microbiol Biotechnol* 2008, **79**:339–354.
3. Diniz RH, Silveira WB, Fietto LG, Passos FM: The high fermentative metabolism of *Kluyveromyces marxianus* UFV-3 relies on the increased expression of key lactose metabolic enzymes. *Antonie Van Leeuwenhoek* 2012, **101**:541–550.
4. Dos Santos VC, Bragança CR, Passos FJ, Passos FM: Kinetics of growth and ethanol formation from a mix of glucose/xylose substrate by *Kluyveromyces marxianus* UFV-3. *Antonie Van Leeuwenhoek* 2013, **103**:153–161.
5. López-Alvarez A, Díaz-Pérez AL, Sosa-Aguirre C, Macías-Rodríguez L, Campos-García J: Ethanol yield and volatile compound content in fermentation of agave must by *Kluyveromyces marxianus* UMPe-1 comparing with *Saccharomyces cerevisiae* baker's yeast used in tequila production. *J Biosci Bioeng* 2012, **113**(5):614–618.
6. Lee KS, Kim JS, Heo P, Yang TJ, Sung YJ, Cheon Y, Koo HM, Yu BJ, Seo JH, Jin YS, Park JC, Kweon DH: Characterization of *Saccharomyces cerevisiae* promoters for heterologous gene expression in *Kluyveromyces marxianus*. *Appl Microbiol Biotechnol* 2012. Epub ahead of print.
7. Jeong H, Lee DH, Kim SH, Kim HJ, Lee K, Song JY, Kim BK, Sung BH, Park JC, Sohn JH, Koo HM, Kim JF: Genome sequence of the thermotolerant yeast *Kluyveromyces marxianus* var. *marxianus* KCTC 17555. *Eukaryot Cell* 2012, **11**(12):1584–1585.

8. Rocha SN, Abrahão-Neto J, Cerdán ME, Gombert AK, González-Siso MI: **Heterologous expression of a thermophilic esterase in *Kluyveromyces* yeasts.** *Appl Microbiol Biotechnol* 2011, **89**:375–385.
9. Rocha SN, Abrahão-Neto J, Cerdán ME, González-Siso MI, Gombert AK: **Heterologous expression of glucose oxidase in the yeast *Kluyveromyces marxianus*.** *Microb Cell Fact* 2010, **9**:4.
10. Raimondi S, Uccelletti D, Amaretti A, Leonardi A, Palleschi C, Rossi M: **Secretion of *Kluyveromyces lactis* Cu/Zn SOD: strategies for enhanced production.** *Appl Microbiol Biotechnol* 2010, **86**:871–878.
11. Foukis A, Stergiou PY, Theodorou LG, Papagianni M, Papamichael EM: **Purification, kinetic characterization and properties of a novel thermo-tolerant extracellular protease from *Kluyveromyces marxianus* IFO 0288 with potential biotechnological interest.** *Bioresour Technol* 2012, **123**:214–220.
12. Rossi M, Buzzini P, Cordisco L, Amaretti A, Sala M, Raimondi S, Ponzoni C, Pagnoni UM, Matteuzzi D: **Growth, lipid accumulation, and fatty acid composition in obligate psychrophilic, facultative psychrophilic, and mesophilic yeasts.** *FEMS Microbiol Ecol* 2009, **69**:363–372.
13. García-Fruitós E, Vazquez E, Gonzalez-Montalbán N, Ferrer-Miralles N, Villaverde A: **Analytical approaches for assessing aggregation of protein biopharmaceuticals.** *Curr Pharm Biotechnol* 2011, **12**:1530–1536.
14. Maldonado LMTP, Hernández VEB, Rivero EM, de la Rosa AP B, Flores JLF, Acevedo LGO, De León Rodríguez A: **Optimization of culture conditions for a synthetic gene expression in *Escherichia coli* using response surface methodology: the case of human interferon beta.** *Biomol Eng* 2007, **24**:217–222.
15. Noguère C, Larsson AM, Guyot JC, Bignon C: **Fractional factorial approach combining 4 *Escherichia coli* strains, 3 culture media, 3 expression temperatures and 5 N-terminal fusion tags for screening the soluble expression of recombinant proteins.** *Protein Expr Purif* 2012, **84**:204–213.
16. Baneyx F, Mujacic M: **Recombinant protein folding and misfolding in *Escherichia coli*.** *Nat Biotechnol* 2004, **22**:1399–1408.
17. Dragosits M, Stadlmann J, Albiol J, Baumann K, Maurer M, Gasser B, Sauer M, Altmann F, Ferrer P, Mattanovich D: **The effect of temperature on the proteome of recombinant *Pichia pastoris*.** *J Proteome Res* 2009, **8**:1380–1392.
18. Li Z, Xiong F, Lin Q, D'Anjou M, Daugulis AJ, Yang DS, Hew CL: **Low-temperature increases the yield of biologically active herring antifreeze protein in *Pichia pastoris*.** *Protein Expr Purif* 2001, **21**:438–445.
19. Gasser B, Maurer M, Rautio J, Sauer M, Bhattacharyya A, Saloheimo M, Penttilä M, Mattanovich D: **Monitoring of transcriptional regulation in *Pichia pastoris* under protein production conditions.** *BMC Genomics* 2007, **8**:179.
20. Morlino GB, Tizzani L, Bianchi MM, Frontali L: **Inducible amplification of gene copy number and heterologous protein production in the yeast *Kluyveromyces lactis*.** *Appl Environ Microbiol* 1999, **65**:4808–4813.
21. Saliola M, Mazzoni C, Solimando N, Crisà A, Falcone C, Jung G: **FleerR: Use of the *KIADH4* promoter for ethanol-dependent production of recombinant Human Serum Albumin in *Kluyveromyces lactis*.** *Appl Environ Microbiol* 1999, **65**:53–60.
22. Raimondi S, Uccelletti D, Matteuzzi D, Pagnoni UM, Rossi M, Palleschi C: **Characterization of the superoxide dismutase *SOD1* gene of *Kluyveromyces marxianus* L3 and improved production of SOD activity.** *Appl Microbiol Biotechnol* 2008, **77**:1269–1277.
23. Lane MM, Burke N, Karreman R, Wolfe KH, O'Byrne CP, Morrissey JP: **Physiological and metabolic diversity in the yeast *Kluyveromyces marxianus*.** *Antonie Van Leeuwenhoek* 2011, **100**:507–519.
24. Fonseca GG, Gombert AK, Heinze E, Wittmann C: **Physiology of the yeast *Kluyveromyces marxianus* during batch and chemostat cultures with glucose as the sole carbon source.** *FEMS Yeast Res* 2007, **7**:422–435.
25. Masoud W, Jespersen L: **Pectin degrading enzymes in yeasts involved in fermentation of *Coffea arabica* in East Africa.** *Int J Food Microbiol* 2006, **110**:291–296.
26. Arizon J, Morel S, Gschaedler A, Monsan P: **Purification and substrate specificities of a fructanase from *Kluyveromyces marxianus* isolated from the fermentation process of Mezzal.** *Bioresour Technol* 2011, **102**:3298–3303.
27. Rouwenhorst RJ, Ritmeester WS, Scheffers WA, Van Dijken JP: **Localization of inulinase and invertase in *Kluyveromyces* species.** *Appl Environ Microbiol* 1990, **56**:3329–3336.
28. Dragosits M, Frascotti G, Bernard-Granger L, Vázquez F, Giuliani M, Baumann K, Rodríguez-Carmona E, Tokkanen J, Parrilli E, Wiebe MG, Kunert R, Maurer M, Gasser B, Sauer M, Branduardi P, Pakula T, Saloheimo M, Penttilä M, Ferrer P, Luisa Tutino M, Villaverde A, Porro D, Mattanovich D: **Influence of growth temperature on the production of antibody Fab fragments in different microbes: a host comparative analysis.** *Biotechnol Prog* 2011, **27**:38–46.
29. Zhang Y, Li T, Liu J: **Low temperature and glucose enhanced T7 RNA polymerase-based plasmid stability for increasing expression of glucagons-like peptide-2 in *Escherichia coli*.** *Protein Expr Purif* 2003, **29**:132–139.
30. Zhang Z, Moo-Young M, Chisti Y: **Plasmid stability in recombinant *Saccharomyces cerevisiae*.** *Biotechnol Adv* 1996, **14**:401–435.
31. Glick BR: **Metabolic load and heterologous gene expression.** *Biotechnol Adv* 1995, **13**:247–261.
32. Neubauer P, Lin HY, Mathiszk B: **Metabolic load of recombinant protein production: inhibition of cellular capacities for glucose uptake and respiration after induction of a heterologous gene in *Escherichia coli*.** *Biotechnol Bioeng* 2003, **83**:53–64.
33. Mazzoni C, Santori F, Saliola M, Falcone C: **Molecular analysis of UAS(E), a cis element containing stress response elements responsible for ethanol induction of the *KIADH4* gene of *Kluyveromyces lactis*.** *Res Microbiol* 2000, **151**:19–28.
34. Davidson JF, Whyte B, Bissinger PH, Schiestl RH: **Oxidative stress is involved in heat-induced cell death in *Saccharomyces cerevisiae*.** *PNAS* 1996, **93**:5116–5121.
35. Dellomonaco C, Amaretti A, Zanon S, Pompei A, Matteuzzi D, Rossi M: **Fermentative production of superoxide dismutase with *Kluyveromyces marxianus*.** *J Ind Microbiol Biotechnol* 2007, **34**:27–34.
36. Burkert JFM, Kalil SJ, Mauerer Filho F, Rodrigues MI: **Parameters optimization for enzymatic assays using experimental design.** *Braz J ChemEng* 2006, **23**:163–170.
37. Miller GL: **Use of dinitrosalicylic acid reagent for determination of reducing sugar.** *Anal Chem* 1959, **31**:426–428.
38. Laemmli UK: **Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4.** *Nature* 1970, **227**:680–685.
39. Schmitt ME, Brown TA, Trumpower BL: **A rapid and simple method for preparation of RNA from *Saccharomyces cerevisiae*.** *Nucleic Acids Res* 1990, **18**:3091–3092.
40. McCord JM, Keele BB, Fridovich I: **An enzyme-based theory of obligate anaerobiosis: the physiological function of superoxide dismutase.** *Proc Natl Acad Sci* 1971, **68**:1024–1027.
41. Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K: *Current protocols in molecular biology.* New York: Greene Publishing Associates and Wiley-Interscience; 1994.
42. Sambrook J, Fritsch EF, Maniatis T: *Molecular cloning: a laboratory manual.* 2nd edition. New York, NY: Cold Spring Harbor Laboratory Press; 2001.

doi:10.1186/1475-2859-12-34

Cite this article as: Raimondi et al.: Thermal adaptability of *Kluyveromyces marxianus* in recombinant protein production. *Microbial Cell Factories* 2013 **12**:34.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

