



ON SUMS OF EIGENVALUES OF ELLIPTIC OPERATORS ON MANIFOLDS

Ahmad El Soufi, Evans Harrell, Said Ilias, Joachim Stubbe

► **To cite this version:**

Ahmad El Soufi, Evans Harrell, Said Ilias, Joachim Stubbe. ON SUMS OF EIGENVALUES OF ELLIPTIC OPERATORS ON MANIFOLDS. Journal of Spectral Theory, 2016. <hal-01174814v2>

HAL Id: hal-01174814

<https://hal.archives-ouvertes.fr/hal-01174814v2>

Submitted on 16 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON SUMS OF EIGENVALUES OF ELLIPTIC OPERATORS ON MANIFOLDS

AHMAD EL SOUFI, EVANS M. HARRELL II, SAÏD ILIAS, AND JOACHIM STUBBE

ABSTRACT. We use the averaged variational principle introduced in a recent article on graph spectra [10] to obtain upper bounds for sums of eigenvalues of several partial differential operators of interest in geometric analysis, which are analogues of Kröger's bound for Neumann spectra of Laplacians on Euclidean domains [15]. Among the operators we consider are the Laplace-Beltrami operator on compact subdomains of manifolds. These estimates become more explicit and asymptotically sharp when the manifold is conformal to homogeneous spaces (here extending a result of Strichartz [26] with a simplified proof). In addition we obtain results for the Witten Laplacian on the same sorts of domains and for Schrödinger operators with confining potentials on infinite Euclidean domains. Our bounds have the sharp asymptotic form expected from the Weyl law or classical phase-space analysis. Similarly sharp bounds for the trace of the heat kernel follow as corollaries.

1. INTRODUCTION

In this article we consider the eigenvalues of self-adjoint, second-order elliptic partial differential operators defined on a subdomain of a Riemannian manifold (M, g) of dimension $\nu \geq 2$. The model for the operators we are able to treat is the Laplacian on a domain with Neumann boundary conditions, defined in the weak sense, i.e. via the Laplacian energy

$$\frac{\int_{\Omega} (|\nabla^g \varphi(\mathbf{x})|^2) dv_g}{\int_{\Omega} |\varphi(\mathbf{x})|^2 dv_g}$$

on functions $\varphi \in H^1(\Omega)$, but the class treated includes a large variety of Schrödinger operators, even with weights. Specifically, the eigenvalues we shall discuss are operationally defined by the min-max procedure applied to expressions of the general form

$$\mathcal{R}(\varphi) := \frac{\int_{\Omega} (|\nabla^g \varphi(\mathbf{x})|^2 + V(\mathbf{x})|\varphi(\mathbf{x})|^2) e^{-2\theta(\mathbf{x})} dv_g}{\int_{\Omega} |\varphi(\mathbf{x})|^2 e^{-2\rho(\mathbf{x})} dv_g}.$$

For convenience we set $w = e^{2(\rho-\theta)}$ so that \mathcal{R} takes on the form

$$\mathcal{R}(\varphi) = \frac{\int_{\Omega} (|\nabla^g \varphi(\mathbf{x})|^2 + V(\mathbf{x})|\varphi(\mathbf{x})|^2) w(\mathbf{x}) e^{-2\rho(\mathbf{x})} dv_g}{\int_{\Omega} |\varphi(\mathbf{x})|^2 e^{-2\rho(\mathbf{x})} dv_g}. \quad (1)$$

Here $\rho \in C^1(\Omega)$, $0 < C \leq w(\mathbf{x}) \in C^0(\Omega)$, and $V \in \text{Lip}(\Omega)$ are real-valued functions. We define the Neumann eigenvalues of (1) by the min-max principle [3, 27], i.e.,

$$\mu_{\ell} := \max_{\{\text{subspace } \mathfrak{S} : \dim(\mathfrak{S})=\ell\}} \min_{\{\varphi \in H^1(\Omega) : \varphi \perp \mathfrak{S}, \|\varphi\|_{L^2}=1\}} \mathcal{R}(\varphi). \quad (2)$$

2010 *Mathematics Subject Classification.* 58J50, 35P15, 47A75.

Key words and phrases. Manifold with density; Weighted Laplacian; Schrödinger operator, Witten Laplacian; Eigenvalue; Upper bound, phase space, Weyl law, homogeneous space, conformal.

Of course, μ_ℓ depends on the domain Ω as well as the choice of the metric g , the density $e^{-2\rho}$ and weight w , and the potential V , but dependence on these quantities will not be indicated explicitly unless necessary.

Under suitable regularity assumptions on Ω and V , the sequence $\{\mu_\ell\}$ is nothing but the spectrum of the eigenvalue problem

$$H\varphi = \mu\varphi \quad \text{in } \Omega \quad (3)$$

with Neumann boundary conditions if $\partial\Omega \neq \emptyset$, where

$$H\varphi = -e^{2\rho} \operatorname{div}_g (we^{-2\rho} \nabla^g \varphi) + Vw\varphi \quad (4)$$

$$= w \left\{ \Delta_g \varphi + 2(\nabla^g \theta, \nabla^g \varphi)_g + V\varphi \right\} \quad (5)$$

where $\Delta_g \varphi := -\operatorname{div}_g(\nabla^g \varphi)$ is the Laplace Beltrami operator associated with g .

In the following sections we derive semiclassically sharp phase-space upper bounds for the sums of the first k eigenvalues associated with (1). We also obtain bounds for the corresponding Riesz means and heat trace. The following inequalities, which are valid for any bounded domain $\Omega \subset \mathbb{R}^\nu$, provide a sampling of these bounds :

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{4\pi^2 \nu}{\nu+2} \left(\frac{k}{|\Omega| \omega_\nu} \right)^{\frac{2}{\nu}} \int_{\Omega} w(\mathbf{x}) d^\nu x + \int_{\Omega} \tilde{V}(\mathbf{x}) w(\mathbf{x}) d^\nu x$$

and

$$\sum_{j \geq 0} e^{-t\mu_j} \geq \frac{|\Omega|}{(4\pi t)^{\frac{\nu}{2}}} \left(\int_{\Omega} w(\mathbf{x}) d^\nu x \right)^{-\frac{\nu}{2}} e^{-t \int_{\Omega} \tilde{V}(\mathbf{x}) w(\mathbf{x}) d^\nu x},$$

where $\tilde{V}(\mathbf{x}) := V(\mathbf{x}) + |\nabla \rho|^2(\mathbf{x})$, $|\Omega|$ is the volume of Ω , ω_ν is the volume of the unit ball in \mathbb{R}^ν and, for every $f \in L^1(\Omega)$, $\int_{\Omega} f(\mathbf{x}) d^\nu x = \frac{1}{|\Omega|} \int_{\Omega} f(\mathbf{x}) d^\nu x$ is the mean value of f with respect to Lebesgue measure.

When appropriate we remark on the simpler consequences that apply under assumptions on ρ , w , and V . The case where $V = \rho = 0$ and $w = 1$ identically, and $M = \mathbb{R}^\nu$ reduces to the situation treated by Kröger in his ground-breaking work [15], and this result was already extended to subdomains of general homogeneous spaces by Strichartz [26] (see also [8]). The upper bounds in [15, 26] are notable for being sharp in the sense of agreeing with the ‘‘semiclassical’’ Weyl law, with the optimal constant. For the background and context of Weyl-sharp bounds on sums of Laplacian eigenvalues, we refer to [16, 17].

In this article a new, simplified proof is used, and we considerably enlarge the family of self-adjoint elliptic operators for which semiclassical upper bounds are proved. Even when $V = 0$, new cases of interest that are treated include the Witten Laplacian, for which $w = 1$; the Laplacian of a conformal metric $\tilde{g} = \alpha^{-2}g$, for which $e^{-2\rho} = \alpha^{-n}$ and $w = \alpha^2$; and the vibrating membrane with variable density $\gamma(\mathbf{x})$, for which $\gamma(\mathbf{x}) = e^{-2\rho}$ and $w = e^{2\rho}$.

In the last part of the paper, we focus on domains of compact homogeneous Riemannian spaces. We revisit the inequality of Strichartz ([26, Theorem 2.2]) in the light of this new approach and obtain extensions of Strichartz’s inequality to the case where the Laplace operator is penalized by a potential in the presence of weights. For example, we prove that if Ω is a domain of a compact homogeneous Riemannian manifold (M, g) , then the eigenvalues μ_l associated with (1) on Ω satisfy

$$\sum_{j \geq 0} (z - \mu_j)_+ \geq \frac{|\Omega|_g}{|M|_g} \sum_{j \geq 0} (z - \tilde{\lambda}_j)_+$$

for all $z \in \mathbb{R}$, and

$$\sum_{j \geq 0} e^{-\mu_j t} \geq \frac{|\Omega|_g}{|M|_g} \sum_{j \geq 0} e^{-\tilde{\lambda}_j t}$$

for all $t > 0$, where $\tilde{\lambda}_j = \lambda_j \int_{\Omega} w dv_g + \int_{\Omega} \tilde{V} w dv_g$, and where the λ_j 's are the eigenvalues of the Laplacian on the whole manifold M (see Theorem 5.1 and Corollary 5.2). The extension (stricto sensu) of Strichartz inequality is given in Theorem 5.2 and takes the following form when $\Omega = M$:

$$\sum_{j=0}^{k-1} \mu_j \leq \sum_{j=0}^{k-1} \tilde{\lambda}_j.$$

It is known that without assumptions of regularity, these variationally defined Neumann eigenvalues for the Laplacian may have finite points of accumulation of a quite arbitrary sort, as entertainingly discussed in [12]. In this case the definition (2) would yield $\mu_\ell = \inf(\sigma_{\text{ess}})$ for all ℓ greater than some value, and the bounds we shall provide would become uninteresting. We note that, for example, the spectrum of the Neumann Laplacian is guaranteed to have no finite points of accumulation if the boundary is piecewise smooth [12].

Remark 1.1. *Before closing this section, we make some further technical remarks about how to define the Dirichlet and Neumann problems for these elliptic operators in the weak, or quadratic-form, sense. In this regard we follow Edmunds and Evans [5], where in Chapter VII it is shown that uniformly elliptic quadratic forms, on arbitrary open sets in Euclidean space, determine unique operators via the Friedrichs extension, which, when the domain is sufficiently regular, reduce to the classically defined operators for the Dirichlet and Neumann problems. (See also [28, 24].) In particular, defining the quadratic form (1) initially on the Sobolev space $W_0^{1,2}(\Omega)$ corresponds to Dirichlet boundary conditions, whereas defining it initially on the restrictions to Ω of functions in the space $W_0^{1,2}(\mathbb{R}^\nu)$ corresponds to Neumann conditions. (For domains allowing a Sobolev extension property the latter set coincides with $W^{1,2}(\Omega)$.) It is not in general possible to say that the operators thus defined satisfy boundary conditions in a classical sense, or to guarantee regularity at the boundary. However, in cases where the boundary is sufficiently regular, integration by parts transforms expressions $\langle H\varphi, \varphi \rangle$ where H is a classically defined operator into a quadratic form of the type (1) for φ in a dense subset of the Sobolev spaces corresponding to Dirichlet or respectively Neumann conditions.*

There are certainly significant questions of regularity of the eigenfunctions in the case when Ω is an arbitrary open set, treated for example in [2], but they will play no role in the present article.

The extension of the analysis of [5] from \mathbb{R}^ν to manifolds is straightforward, because only Hilbert-space structures and locally defined properties of functions and their gradients are used.

2. THE AVERAGED VARIATIONAL PRINCIPLE

In this section we recall the averaged variational principle which will be foundational for this article. The following is mainly a restatement of Theorem 3.1 of Harrell-Stubbe [10], along with a characterization of the case of equality.

Theorem 2.1. *Consider a self-adjoint operator H on a Hilbert space \mathcal{H} , the spectrum of which is discrete at least in its lower portion, so that $-\infty < \mu_0 \leq \mu_1 \leq \dots$. The corresponding orthonormalized eigenvectors are denoted $\{\psi^{(\ell)}\}$. The closed quadratic form corresponding to H is denoted $Q(\varphi, \varphi)$ for vectors φ in the quadratic-form domain $\mathcal{Q}(H) \subset \mathcal{H}$. Let $f_\zeta \in \mathcal{Q}(H)$ be a*

family of vectors indexed by a variable ζ ranging over a measure space $(\mathfrak{M}, \Sigma, \sigma)$. Suppose that \mathfrak{M}_0 is a subset of \mathfrak{M} . Then for any $z \in \mathbb{R}$,

$$\sum_j (z - \mu_j)_+ \int_{\mathfrak{M}} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \geq \int_{\mathfrak{M}_0} (z \|f_\zeta\|^2 - Q(f_\zeta, f_\zeta)) d\sigma, \quad (6)$$

provided that the integrals converge. Moreover, equality holds in (6) for $z \in \mathbb{R}$ if and only if up to sets of measure 0,

$$\{f_\zeta; \zeta \in \mathfrak{M}_0\} \subset E(z) \quad \text{and} \quad \{f_\zeta; \zeta \in \mathfrak{M} \setminus \mathfrak{M}_0\} \perp E_0(z),$$

where $E(z) = \bigoplus_{\mu \leq z} \ker(H - \mu I)$ and $E_0(z) = \bigoplus_{\mu < z} \ker(H - \mu I)$.

Taking $z = \mu_k$ in (6) we obtain

$$\begin{aligned} \mu_k \left(\int_{\mathfrak{M}_0} \|f_\zeta\|^2 d\sigma - \sum_{j=0}^{k-1} \int_{\mathfrak{M}} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \right) \\ \leq \int_{\mathfrak{M}_0} Q(f_\zeta, f_\zeta) d\sigma - \sum_{j=0}^{k-1} \mu_j \int_{\mathfrak{M}} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma, \end{aligned} \quad (7)$$

Remarks 2.1. 1. The averaged variational principle is an abstract version and sharpening of ideas appearing in various place in the literature, including not only [15], but also work of Lieb and others on coherent states and trace inequalities [20, 22]. In special cases, similar use of averaging and tight frames for the study of eigenvalue sums and related quantities has also been made by Laugesen and Siudeja [18].

2. We point out that the normalization of the test function f_ζ could be incorporated into the measure, so that, for example, Eq. (6) could alternatively be written in terms of integrals of expectation values such as

$$\int \left(\frac{|\langle \psi^{(j)}, f_\zeta \rangle|^2}{\|f_\zeta\|^2} \right) d\sigma, \quad (8)$$

i.e., over norms of projections of the eigenfunctions. Despite the suggestiveness of these alternatives, an advantageous feature of (7)-(6) that we shall later exploit is that useful identities are available for averages of norms of some choices of f_ζ . Still, if the test functions f_ζ and the measure space \mathfrak{M} constitute a tight frame, in the sense of satisfying a generalized Parseval identity [13], then alternative forms of the inequalities imply appealing variational principles for sums and Riesz means of eigenvalues, as captured in the next corollary.

Corollary 2.1. Under the assumptions of the Theorem, suppose further that f_ζ is a nonvanishing family of test functions with the property that for all $\phi \in \mathcal{H}$,

$$\int_{\mathfrak{M}} \frac{|\langle \phi, f_\zeta \rangle|^2}{\|f_\zeta\|^2} d\sigma = A \|\phi\|^2$$

for a fixed constant $A > 0$. Then for any $\mathfrak{M}_0 \subset \mathfrak{M}$ such that $(|\mathfrak{M}_0| - Ak) \mu_k \geq 0$,

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{1}{|\mathfrak{M}_0|} \int_{\mathfrak{M}_0} \left(\frac{Q(f_\zeta, f_\zeta)}{\|f_\zeta\|^2} \right) d\sigma. \quad (9)$$

For Riesz means,

$$\sum_j (z - \mu_j)_+ \geq \frac{1}{A} \int_{\mathfrak{M}_0} \left(z - \frac{Q(f_\zeta, f_\zeta)}{\|f_\zeta\|^2} \right) d\sigma. \quad (10)$$

The proof of Corollary 2.1 is immediate; see [10] for more in this connection. To make our exposition self-contained, we provide here the proof of the inequality (6) in Theorem 2.1 before discussing the case of equality.

Proof of Theorem 2.1. For every integer $l \geq 0$, we denote by P_l the orthogonal projector onto the subspace spanned by $\{\psi^{(j)}, j \leq l\}$, i.e. $P_l f = \sum_{j=0}^l \langle \psi^{(j)}, f \rangle \psi^{(j)}$. Let $z \in \mathbb{R}$, $z > \mu_0$ (the inequality (6) being obvious for $z \leq \mu_0$), and let k be the smallest integer such that $z \leq \mu_k$ (that is $z \in (\mu_{k-1}, \mu_k]$). Then

$$z\|f - P_{k-1}f\|^2 \leq \mu_k\|f - P_{k-1}f\|^2 \leq Q(f - P_{k-1}f, f - P_{k-1}f), \quad (11)$$

and, after direct computations,

$$z(\|f\|^2 - \|P_{k-1}f\|^2) \leq Q(f, f) - Q(P_{k-1}f, P_{k-1}f).$$

With $\|P_{k-1}f\|^2 = \sum_{j=0}^{k-1} \langle \psi^{(j)}, f \rangle^2$ and $Q(P_{k-1}f, P_{k-1}f) = \sum_{j=0}^{k-1} \mu_j \langle \psi^{(j)}, f \rangle^2$, this yields

$$z\|f\|^2 - Q(f, f) \leq \sum_{j=0}^{k-1} (z - \mu_j) \langle \psi^{(j)}, f \rangle^2.$$

Applying this last inequality to f_ζ , $\zeta \in \mathfrak{M}_0$, and integrating over \mathfrak{M}_0 we get

$$\begin{aligned} z \int_{\mathfrak{M}_0} \|f_\zeta\|^2 d\sigma - \int_{\mathfrak{M}_0} Q(f_\zeta, f_\zeta) d\sigma &\leq \sum_{j=0}^{k-1} (z - \mu_j) \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \\ &= \sum_{j \geq 0} (z - \mu_j)_+ \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma. \end{aligned} \quad (12)$$

The inequality (6) follows from (12) and the obvious inequality

$$\sum_{j \geq 0} (z - \mu_j)_+ \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \leq \sum_{j \geq 0} (z - \mu_j)_+ \int_{\mathfrak{M}} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma. \quad (13)$$

Assume now that equality holds in (6). This implies that equality holds in (13) and in (11) for $f = f_\zeta$ for almost all $\zeta \in \mathfrak{M}_0$. Equality in (11) holds for f either when $z < \mu_k$ and $f = P_{k-1}f$ or if $z = \mu_k$ and $H(f - P_{k-1}f) = \mu_k(f - P_{k-1}f)$, which implies in both cases that $f \in E(z)$. On the other hand, equality in (13) implies that, for almost all $\zeta \in \mathfrak{M} \setminus \mathfrak{M}_0$ and all $j \in \mathbb{N}$ such that $\mu_j < z$, f_ζ is orthogonal to $\text{span}\{\psi^{(0)}, \dots, \psi^{(j)}\}$, which means that f_ζ is orthogonal to $E_0(z)$.

Conversely, under the conditions of the statement,

$$\begin{aligned} \sum_j (z - \mu_j)_+ \int_{\mathfrak{M}} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma &= \sum_j (z - \mu_j)_+ \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \\ &= \sum_{\mu_j \leq z} z \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma - \sum_{\mu_j \leq z} \mu_j \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \\ &= \sum_{j=0}^{+\infty} z \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma - \sum_{j=0}^{+\infty} \mu_j \int_{\mathfrak{M}_0} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d\sigma \\ &= \int_{\mathfrak{M}_0} (z\|f_\zeta\|^2 - Q(f_\zeta, f_\zeta)) d\sigma. \end{aligned}$$

□

As the guiding example for this article, when Ω is a bounded subdomain of \mathbb{R}^ν , we may use test functions of the form

$$f_\zeta(\mathbf{x}) := \frac{1}{(2\pi)^{\nu/2}} e^{i\mathbf{p}\cdot\mathbf{x}},$$

where ζ has been equated to \mathbf{p} , which ranges over $\mathfrak{M} = \mathbb{R}^\nu$ with Lebesgue measure (The reason for distinguishing ζ logically from \mathbf{p} will be made clear in Theorem 4.1.). Indeed, $\|f_\zeta\|^2 = \frac{|\Omega|}{(2\pi)^\nu}$ for all ζ , where $|\Omega|$ is the Euclidean volume of Ω , and Parseval's identity gives

$$\int_{\mathbb{R}^\nu} |\langle \phi, f_\zeta \rangle|^2 d^\nu p = \|\phi\|^2.$$

Hence, applying Corollary 2.1 with $\mathfrak{M}_0 \subset \mathfrak{M}$ taken to be the Euclidean ball of radius $2\pi \left(\frac{k}{|\Omega|\omega_\nu}\right)^{\frac{1}{\nu}}$, we recover Kröger's inequality for Neumann eigenvalues of the Euclidean Laplacian (here ω_ν stands for the volume of the ν -dimensional Euclidean unit ball). Indeed, in this case, the Rayleigh quotient of f_ζ is simply given by $\mathcal{R}(f_\zeta) = |\mathbf{p}|^2$ and (9) yields

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{1}{|\mathfrak{M}_0|} \int_{\mathfrak{M}_0} |\mathbf{p}|^2 d^\nu p = \frac{4\pi^2\nu}{\nu+2} \left(\frac{k}{|\Omega|\omega_\nu}\right)^{\frac{2}{\nu}}.$$

This approach can be applied to easily extend Kröger's inequality to Neumann eigenvalues on a bounded subdomain of \mathbb{R}^ν in the presence of nontrivial potential and weights.

Example 2.1. *Let $\mu_0 \leq \mu_1 \leq \dots$ be the variationally defined Neumann eigenvalues (2) on a bounded open set $\Omega \subset \mathbb{R}^\nu$ endowed with the standard Euclidean metric, where w , ρ , and V satisfy the assumptions stated above. Then*

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{4\pi^2\nu}{\nu+2} \left(\frac{k}{|\Omega|\omega_\nu}\right)^{\frac{2}{\nu}} \int_{\Omega} w(\mathbf{x}) d^\nu x + \int_{\Omega} \tilde{V}(\mathbf{x}) w(\mathbf{x}) d^\nu x, \quad (14)$$

where $\tilde{V}(\mathbf{x}) := V(\mathbf{x}) + |\nabla\rho|^2(\mathbf{x})$ and, for every $f \in L^1(\Omega)$, $\int_{\Omega} f(\mathbf{x}) d^\nu x = \frac{1}{|\Omega|} \int_{\Omega} f(\mathbf{x}) d^\nu x$ is the mean value of f with respect to Lebesgue measure.

Example 2.1 sets the stage for a more general result that we obtain in Section 3 in the context of Riemannian manifolds. We also stress that these estimates will be improved in Section 4, with the aid of a coherent-state analysis relating the upper bounds to phase-space volumes.

Upper bounds for individual Neumann eigenvalues μ_j are also obtainable from the averaged variational principle. In order to somewhat simplify the bound, let us define the shifted Neumann eigenvalues

$$\tilde{\mu}_j := \mu_j - \int_{\Omega} \tilde{V}(\mathbf{x}) w(\mathbf{x}) d^\nu x. \quad (15)$$

In terms of these quantities, we will be able to show that (see Corollary 3.1)

$$\tilde{\mu}_k \leq 4\pi^2 \left(\frac{k}{|\Omega|\omega_\nu}\right)^{\frac{2}{\nu}} \left(1 + 2\sqrt{\frac{1-S_k}{\nu+2}}\right) \int_{\Omega} w(\mathbf{x}) d^\nu x, \quad (16)$$

where

$$S_k := \frac{\frac{1}{k} \sum_{j=0}^{k-1} \tilde{\mu}_j}{\frac{4\pi^2\nu}{\nu+2} \left(\frac{k}{|\Omega|\omega_\nu}\right)^{\frac{2}{\nu}} \int_{\Omega} w(\mathbf{x}) d^\nu x} \leq 1. \quad (17)$$

3. BOUNDS FOR NEUMANN EIGENVALUES ON DOMAINS OF RIEMANNIAN MANIFOLDS

Let (M, g) be a Riemannian manifold of dimension $\nu \geq 2$ and let Ω be a bounded subdomain of M . Of course, when M is a closed manifold, Ω can be equal to the whole of M .

Let $F : (M, g) \rightarrow \mathbb{R}^N$, be an isometric embedding (whose existence for sufficiently large N is guaranteed by Nash's embedding Theorem). To any function $u \in L^2(\Omega)$, we associate the function $\hat{u}_F : \mathbb{R}^N \rightarrow \mathbb{R}$ defined by

$$\hat{u}_F(\mathbf{p}) = \int_{\Omega} u(\mathbf{x}) e^{i\mathbf{p} \cdot F(\mathbf{x})} dv_g, \quad (18)$$

where the dot stands for the Euclidean inner product in \mathbb{R}^N (i.e., \hat{u}_F is the Fourier transform of the signed measure $F_*(udv_g)$ supported by $F(\Omega)$). It is well-known, since the works of Hörmander, Agmon-Hörmander and others (see [1, Theorem 2.1] [14, Theorem 7.1.26], [25, Corollary 5.2]), that there exists a constant $C_{F(\Omega)}$ such that, $\forall u \in L^2(\Omega)$ and $\forall R > 0$,

$$\int_{B_R} |\hat{u}_F(\mathbf{p})|^2 d^N p \leq C_{F(\Omega)} R^{N-\nu} \|u\|^2, \quad (19)$$

where B_R is the Euclidean ball of radius R in \mathbb{R}^N centered at the origin and $\|u\|^2 = \int_{\Omega} u^2 dv_g$. In other words the Fourier functions appearing in (18) constitute a frame that is not generally tight.

We define the Riemannian constant H_{Ω} by

$$H_{\Omega} = \inf_{N \geq \nu} \inf_{F \in I(M, \mathbb{R}^N)} \left(\frac{\nu + 2}{N + 2} \right)^{\frac{\nu}{2}} \frac{1}{\omega_N} C_{F(\Omega)}, \quad (20)$$

where $I(M, \mathbb{R}^N)$ is the set of isometric embeddings from (M, g) to \mathbb{R}^N .

When Ω is a domain of \mathbb{R}^{ν} , we may take for F the identity map so that, $\forall u \in L^2(\Omega)$, \hat{u}_I is nothing but the Fourier transform of u extended by zero outside Ω . Using Parseval's identity we get $\forall R > 0$,

$$\int_{B_R} |\hat{u}_I(\mathbf{p})|^2 d^{\nu} p \leq \int_{\mathbb{R}^{\nu}} |\hat{u}_I(\mathbf{p})|^2 d^{\nu} p = (2\pi)^{\nu} \|u\|^2.$$

Thus $C_{I(\Omega)} = (2\pi)^{\nu}$ and

$$H_{\Omega} \leq \frac{(2\pi)^{\nu}}{\omega_{\nu}}. \quad (21)$$

In all the sequel, the notation $|\Omega|_g$ will designate the Riemannian volume of Ω with respect to g . We will also use the notation $\int_{\Omega} f dv_g$ to represent the mean value of a function $f \in L^1(\Omega)$ with respect to the Riemannian measure dv_g . (I.e., $\int_{\Omega} f dv_g = \frac{1}{|\Omega|_g} \int_{\Omega} f dv_g$.)

Theorem 3.1. *Let (M, g) be a Riemannian manifold of dimension $\nu \geq 2$. Let $\mu_l = \mu_l(\Omega, g, \rho, w, V)$, $l \in \mathbb{N}$, be the eigenvalues defined by (2) on a bounded open set $\Omega \subset M$, where w , ρ , and V satisfy the assumptions stated above. Then*

(1) *For all $z \in \mathbb{R}$,*

$$\sum_{j \geq 0} (z - \mu_j)_+ \geq \frac{2 |\Omega|_g}{(\nu + 2) H_{\Omega}} \left(\int_{\Omega} w dv_g \right)^{-\frac{\nu}{2}} \left(z - \int_{\Omega} \tilde{V} w dv_g \right)_+^{1+\frac{\nu}{2}}, \quad (22)$$

where $\tilde{V} = V + |\nabla^g \rho|^2$.

(2) For all $k \in \mathbb{N}$,

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{\nu}{\nu+2} \left(\frac{H_\Omega}{|\Omega|_g} k \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g + \int_{\Omega} \tilde{V} w dv_g. \quad (23)$$

(3) For all $t > 0$,

$$\sum_{j \geq 0} e^{-t(\mu_j - \int_{\Omega} \tilde{V} w dv_g)} \geq \left(\frac{\pi}{t} \right)^{\frac{\nu}{2}} \frac{|\Omega|_g}{\omega_\nu H_\Omega} \left(\int_{\Omega} w dv_g \right)^{-\frac{\nu}{2}}. \quad (24)$$

Proof. Let $F : (M, g) \rightarrow \mathbb{R}^N$ be an isometric embedding. For simplicity, we identify the domain Ω with its image $F(\Omega) \subset \mathbb{R}^N$ and any function $u : \Omega \rightarrow \mathbb{R}$ with $u \circ F^{-1} : F(\Omega) \rightarrow \mathbb{R}$.

We apply Theorem 2.1 using test functions of the form

$$f_\zeta(\mathbf{x}) := e^{i\mathbf{p} \cdot \mathbf{x} + \rho(\mathbf{x})},$$

where ζ has been equated to \mathbf{p} , which ranges over $\mathfrak{M} = B_R \subset \mathbb{R}^N$ endowed with Lebesgue measure, where B_R is a Euclidean N -dimensional ball whose radius R is to be determined later. Our Hilbert space here is $L^2(\Omega, e^{-2\rho} dv_g)$ (endowed with the norm $\|u\|^2 = \int_{\Omega} u^2 e^{-2\rho} dv_g$). Hence, for all ζ ,

$$\|f_\zeta\|^2 = |\Omega|_g$$

and consequently

$$\int_{\mathfrak{M}} \|f_\zeta\|^2 d^N p = |\Omega|_g \omega_N R^N. \quad (25)$$

On the other hand, in our case the quadratic form is

$$Q(f_\zeta, f_\zeta) = \int_{\Omega} (|\nabla^\tau f_\zeta|^2 + V|f_\zeta|^2) w(\mathbf{x}) e^{-2\rho(\mathbf{x})} dv_g,$$

where $\nabla^\tau f_\zeta$ is the tangential part of the gradient of f_ζ (more generally, for all $v \in \mathbb{R}^N$, v^τ will designate the tangential vector field induced on Ω by orthogonal projection of v). Thus, with $|f_\zeta|^2 = e^{2\rho}$ and $|\nabla^\tau f_\zeta|^2 = |\mathbf{p}^\tau + \nabla^\tau \rho|^2$,

$$Q(f_\zeta, f_\zeta) = \int_{\Omega} (|\mathbf{p}^\tau|^2(\mathbf{x}) + 2\mathbf{p} \cdot \nabla^\tau \rho(\mathbf{x})) w(\mathbf{x}) dv_g + \int_{\Omega} (V + |\nabla^\tau \rho|^2) w(\mathbf{x}) dv_g.$$

Observe that for symmetry reasons, for all $v \in \mathbb{R}^N \setminus \{0\}$,

$$\int_{B_R} \mathbf{p} \cdot v d^N p = 0$$

and, after elementary calculations,

$$\int_{B_R} (\mathbf{p} \cdot v)^2 d^N p = \frac{|v|}{N} \int_{B_R} |\mathbf{p}|^2 d^N p = \frac{|v|}{N+2} \omega_N R^{N+2}.$$

Thus, if $\{v_1, \dots, v_\nu\}$ is an orthonormal basis of the tangent space of Ω at a point \mathbf{x} , then

$$\int_{\Omega} |\mathbf{p}^\tau|^2(\mathbf{x}) d^N p = \sum_{j \leq \nu} \int_{B_R} (\mathbf{p} \cdot v_j)^2 d^N p = \frac{\nu}{N+2} \omega_N R^{N+2}.$$

This leads to

$$\int_{\mathfrak{M}} Q(f_\zeta, f_\zeta) d^N p = \frac{\nu}{N+2} \omega_N R^{N+2} \int_{\Omega} w dv_g + \omega_N R^N \int_{\Omega} \tilde{V} w dv_g. \quad (26)$$

It remains to deal with the integrals $\int_{\mathfrak{M}} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d^N p$, where $\{\psi^{(j)}\}$ is an $L^2(\Omega, e^{-2\rho} dv_g)$ -orthonormal basis of eigenfunctions associated to $\{\mu_j\}$. Setting $\phi^{(j)} = e^{-\rho} \psi^{(j)}$,

$$\langle \psi^{(j)}, f_\zeta \rangle = \int_{\Omega} f_\zeta \psi^{(j)} e^{-2\rho} dv_g = \int_{\Omega} e^{i\mathbf{p}\cdot\mathbf{x}} \psi^{(j)} e^{-\rho} dv_g = \hat{\phi}_F^{(j)}(\mathbf{p}).$$

Using (19) we obtain

$$\begin{aligned} \int_{B_R} |\langle \psi^{(j)}, f_\zeta \rangle|^2 d^N p &= \int_{B_R} |\hat{\phi}_F^{(j)}(\mathbf{p})|^2 d^N p \leq C_{F(\Omega)} R^{N-\nu} \int_{\Omega} |\phi^{(j)}|^2 dv_g \\ &= C_{F(\Omega)} R^{N-\nu} \int_{\Omega} |\psi^{(j)}|^2 e^{-2\rho} dv_g = C_{F(\Omega)} R^{N-\nu}. \end{aligned} \quad (27)$$

We put (25), (26), and (27) into (6) after choosing $\mathfrak{M}_0 = \mathfrak{M} = B_R$, and obtain for all $R > 0$ and $z \in \mathbb{R}$

$$\sum_{j \geq 0} (z - \mu_j)_+ C_{F(\Omega)} \geq |\Omega|_g \omega_N R^\nu \left(z - \frac{\nu R^2}{N+2} \int_{\Omega} w dv_g - \int_{\Omega} \tilde{V} w dv_g \right). \quad (28)$$

The right side of this inequality is optimized when $R = 0$ if $z \leq \int_{\Omega} \tilde{V} w dv_g$ and when $R^2 = \frac{N+2}{\nu+2} \left(z - \int_{\Omega} \tilde{V} w dv_g \right) / \int_{\Omega} w dv_g$ otherwise. Thus

$$\begin{aligned} \sum_{j \geq 0} (z - \mu_j)_+ C_{F(\Omega)} &\geq \\ &|\Omega|_g \omega_N \left(\frac{N+2}{\nu+2} \right)^{\frac{\nu}{2}} \frac{2}{\nu+2} \left(\int_{\Omega} w dv_g \right)^{-\frac{\nu}{2}} \left(z - \int_{\Omega} \tilde{V} w dv_g \right)_+^{1+\frac{\nu}{2}}. \end{aligned} \quad (29)$$

Taking the infimum with respect to F and N we get (22).

To prove (23) we first observe that taking $z = \mu_k$ in (28) gives

$$k\mu_k - \sum_{j=0}^{k-1} \mu_j \geq \frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} \left(\mu_k - \frac{\nu R^2}{N+2} \int_{\Omega} w dv_g - \int_{\Omega} \tilde{V} w dv_g \right) \quad (30)$$

for all $R > 0$, or

$$\sum_{j=0}^{k-1} \mu_j \leq \left(k - \frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} \right) \mu_k + \frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} \left(\frac{\nu R^2}{N+2} \int_{\Omega} w dv_g + \int_{\Omega} \tilde{V} w dv_g \right).$$

Choosing R such that $\frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} = k$ we get

$$\sum_{j=0}^{k-1} \mu_j \leq k \left(\frac{\nu}{N+2} \left(\frac{C_{F(\Omega)} k}{|\Omega|_g \omega_N} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g + \int_{\Omega} \tilde{V} w dv_g \right), \quad (31)$$

which leads to (23) after taking the infimum with respect to F and N .

The inequality (24) is a consequence of (22) and the following identity relating the heat trace to the Laplace transform of the Riesz mean :

$$\sum_{j \geq 0} e^{-\mu_j t} = t^2 \int_0^\infty e^{-zt} \sum_{j \geq 0} (z - \mu_j)_+ dz. \quad (32)$$

□

Remarks 3.1. 1. In [19, Theorem 1.2], Li and Tang obtained for the Laplacian (i.e. in the case $V = \rho = 0$, $w = 1$) an inequality which is similar to but weaker than (31). Indeed, instead of the term $\frac{\nu}{N+2}$ in the right side their inequality appears with $\frac{N}{N+2}$.

2. It is possible to derive (23) from (22) using Legendre transform. Indeed, the Legendre transform of a function f of the form $f(z) = A(z - B)_+^{1+\frac{\nu}{2}}$ with $A > 0$ is given by

$$f^\wedge(p) = \sup_{z \geq 0} (pz - f(z)) = \left(\frac{2}{A}\right)^{\frac{2}{\nu}} \frac{\nu}{(\nu+2)^{1+\frac{2}{\nu}}} p^{1+\frac{2}{\nu}} + Bp,$$

while the Legendre transform of $g(z) = \sum_{j \geq 0} (z - \mu_j)_+$ is

$$g^\wedge(p) = \sum_{j=0}^{\lfloor p \rfloor - 1} \mu_j + (p - \lfloor p \rfloor) \mu_{\lfloor p \rfloor}.$$

(Indeed, for $z \in [\mu_{k-1}, \mu_k]$, $pz - g(z) = (p - k)z + \sum_{j=0}^{k-1} \mu_j$ which is nondecreasing as soon as $k \leq \lfloor p \rfloor$.) Hence, it suffices to apply Legendre transform to both sides of (22) taking into account that such a transform is inequality-reversing.

Corollary 3.1. Under the assumptions of Theorem 3.1, for any positive integer k ,

$$\widetilde{\mu}_k \leq \left(1 + 2\sqrt{\frac{1 - S_k}{\nu + 2}}\right) \left(\frac{H_\Omega}{|\Omega|_g} k\right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g, \quad (33)$$

where

$$\widetilde{\mu}_k = \mu_k - \int_{\Omega} \widetilde{V} w dv_g \quad \text{and} \quad S_k := \frac{\frac{1}{k} \sum_{j=0}^{k-1} \widetilde{\mu}_j}{\left(\frac{H_\Omega k}{|\Omega|_g}\right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g}.$$

Notice that according to Theorem 3.1 (2), $S_k \leq 1$.

Proof of Corollary 3.1. We take back the proof of Theorem 3.1 and rewrite (30) as follows: For every positive R ,

$$k\widetilde{\mu}_k - \sum_{j=0}^{k-1} \widetilde{\mu}_j \geq \frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} \left(\widetilde{\mu}_k - \frac{\nu R^2}{N+2} \int_{\Omega} w dv_g \right), \quad (34)$$

which yields

$$\left(\frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} - k \right) \widetilde{\mu}_k \leq \frac{\nu}{N+2} \frac{|\Omega|_g \omega_N R^{\nu+2}}{C_{F(\Omega)}} \int_{\Omega} w dv_g - \sum_{j=0}^{k-1} \widetilde{\mu}_j.$$

With the change of variable $\sigma := \frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)} k}$ the last inequality reads

$$(\sigma - 1) \widetilde{\mu}_k \leq \frac{\nu}{N+2} \left(\frac{C_{F(\Omega)} k}{|\Omega|_g \omega_N} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g \sigma^{1+\frac{2}{\nu}} - \frac{1}{k} \sum_{j=0}^{k-1} \widetilde{\mu}_j$$

for all $\sigma > 1$. Taking the infimum with respect to F and N and using (20), we get

$$\begin{aligned} (\sigma - 1) \widetilde{\mu}_k &\leq \left(\frac{H_\Omega k}{|\Omega|_g} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g \sigma^{1+\frac{2}{\nu}} - \frac{1}{k} \sum_{j=0}^{k-1} \widetilde{\mu}_j \\ &= \left(\frac{H_\Omega k}{|\Omega|_g} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g \left(\sigma^{1+\frac{2}{\nu}} - S_k \right). \end{aligned}$$

That is

$$\widetilde{\mu}_k \leq \left(\frac{H_\Omega k}{|\Omega|_g} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g \frac{\sigma^{1+\frac{2}{\nu}} - S_k}{\sigma - 1}. \quad (35)$$

This inequality can be explicitly optimized with respect to $\sigma \in [1, +\infty)$ only when $\nu = 2$, and we then obtain $\sigma_+ = 1 + \sqrt{1 - S_k}$, yielding the desired bound. For general $\nu \geq 2$ we introduce a new change of variable as follows : $\sigma = 1 + \alpha z_k$, where $z_k = (1 - S_k)^{\frac{1}{p}}$, $p = \frac{\nu + 2}{\nu}$, and α is a free positive parameter. Then the bound (35) reads:

$$\widetilde{\mu}_k \leq \left(\frac{H_\Omega k}{|\Omega|_g} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g \frac{(1 + \alpha z_k)^p - 1 + z_k^p}{\alpha z_k}.$$

Since $1 < p \leq 2$ for all $\nu \geq 2$, it follows that

$$\begin{aligned} \frac{1}{p} \frac{(1 + \alpha z_k)^p - 1 + z_k^p}{\alpha z_k} &= \frac{1}{\alpha z_k} \int_0^{\alpha z_k} (1 + s)^{p-1} ds + \frac{z_k^{p-1}}{p\alpha} \\ &\leq \frac{1}{\alpha z_k} \int_0^{\alpha z_k} (1 + (p-1)s) ds + \frac{z_k^{p-1}}{p\alpha} \\ &= 1 + \frac{(p-1)\alpha z_k}{2} + \frac{z_k^{p-1}}{p\alpha}. \end{aligned}$$

Optimizing with respect to α leads to choose $\alpha^2 = \frac{2z_k^{p-2}}{p(p-1)}$. Thus,

$$\frac{1}{p} \frac{(1 + \alpha z_k)^p - 1 + z_k^p}{\alpha z_k} \leq 1 + \sqrt{2} \sqrt{\frac{p-1}{p}} z_k^{\frac{p}{2}} = 1 + 2 \frac{\sqrt{1 - S_k}}{\sqrt{\nu + 2}},$$

which implies the the desired inequality. \square

Corollary 3.2. *Under the assumptions of Theorem 3.1, for any integer $k \in \mathbb{N}$ such that $\sum_{j=0}^{k-1} \mu_j \geq 0$,*

$$\mu_k \left(1 - \frac{\int_{\Omega} \widetilde{V} w dv_g}{\mu_k} \right)_+^{1+\frac{2}{\nu}} \leq \left(\frac{\nu + 2}{2} \right)^{\frac{2}{\nu}} \left(\frac{H_\Omega k}{|\Omega|_g} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g. \quad (36)$$

In particular,

$$\mu_k \leq \max \left\{ 2 \int_{\Omega} \widetilde{V} w dv_g ; 2(\nu + 2)^{\frac{2}{\nu}} \left(\frac{H_\Omega k}{|\Omega|_g} \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g \right\}. \quad (37)$$

Proof. From the inequality (30) in the proof of Theorem 3.1, we deduce with $\sum_{j=0}^{k-1} \mu_j \geq 0$ that for all $R \geq 0$,

$$k\mu_k - \frac{|\Omega|_g \omega_N R^\nu}{C_{F(\Omega)}} \left(\mu_k - \frac{\nu R^2}{N+2} \int_{\Omega} w dv_g - \int_{\Omega} \widetilde{V} w dv_g \right) \geq 0. \quad (38)$$

The left side achieves its minimum when $R = 0$ if $\mu_k \leq \int_{\Omega} \widetilde{V} w dv_g$ and otherwise when $R^2 = \frac{N+2}{\nu+2} \left(\mu_k - \int_{\Omega} \widetilde{V} w dv_g \right) / \int_{\Omega} w dv_g$. Since (36) is obviously satisfied when $\mu_k \leq \int_{\Omega} \widetilde{V} w dv_g$, we shall assume $\mu_k > \int_{\Omega} \widetilde{V} w dv_g$ and get

$$k\mu_k - \frac{|\Omega|_g \omega_N}{C_{F(\Omega)}} \left(\frac{N+2}{\nu+2} \right)^{\frac{\nu}{2}} \frac{2}{\nu+2} \left(\mu_k - \int_{\Omega} \widetilde{V} w dv_g \right)^{1+\frac{\nu}{2}} \left(\int_{\Omega} w dv_g \right)^{-\frac{\nu}{2}} \geq 0$$

which gives

$$\left(\mu_k - \int_{\Omega} \tilde{V} w dv_g \right)^{1+\frac{\nu}{2}} \leq \frac{C_{F(\Omega)}}{|\Omega|_g \omega_N} \left(\frac{\nu+2}{N+2} \right)^{\frac{\nu}{2}} \frac{\nu+2}{2} \left(\int_{\Omega} w dv_g \right)^{\frac{\nu}{2}} k \mu_k.$$

Therefore,

$$\mu_k^{\frac{\nu}{2}} \left(1 - \frac{\int_{\Omega} \tilde{V} w dv_g}{\mu_k} \right)^{1+\frac{\nu}{2}} \leq \frac{C_{F(\Omega)}}{|\Omega|_g \omega_N} \left(\frac{\nu+2}{N+2} \right)^{\frac{\nu}{2}} \frac{\nu+2}{2} \left(\int_{\Omega} w dv_g \right)^{\frac{\nu}{2}} k.$$

Raising to the power $\frac{2}{\nu}$ and taking the infimum with respect to F and N we obtain (36).

To prove (37) we observe that if $\mu_k > 2 \int_{\Omega} \tilde{V} w dv_g$, then $1 - \frac{\int_{\Omega} \tilde{V} w dv_g}{\mu_k} > \frac{1}{2}$, so we can deduce from (36)

$$\left(\frac{1}{2} \right)^{1+\frac{2}{\nu}} \mu_k \leq \left(\frac{\nu+2}{2} \right)^{\frac{2}{\nu}} \left(\frac{H_{\Omega}}{|\Omega|_g} k \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g.$$

□

Note that when $\rho = V = 0$, the inequality (36) of Corollary 3.2 produces

$$\mu_k \leq \left(\frac{\nu+2}{2} \right)^{\frac{2}{\nu}} \left(\frac{H_{\Omega}}{|\Omega|_g} k \right)^{\frac{2}{\nu}} \int_{\Omega} w dv_g,$$

which coincides with Kröger's estimate [15, Corollary 2] when Ω is a Euclidean domain and $w = 1$ (just replace H_{Ω} by $\frac{(2\pi)^{\nu}}{\omega_{\nu}}$).

Let us highlight some consequences of Theorem 3.1 on Schrödinger operators, Witten Laplacians, and Laplacians associated with conformally Euclidean metrics.

Example 3.1 (Schrödinger operators). *From (23) in Theorem 3.1 and (37) in Corollary 3.2 we deduce that for any Schrödinger operator $\Delta_g + V$ on Ω and any integer $k \geq 0$ we find (with $\rho = 0$ and $w = 1$)*

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j(\Delta_g + V) \leq \frac{\nu}{\nu+2} \left(\frac{H_{\Omega}}{|\Omega|_g} k \right)^{\frac{2}{\nu}} + \int_{\Omega} V dv_g. \quad (39)$$

Furthermore, if $\sum_{j=0}^{k-1} \mu_j(\Delta_g + V) \geq 0$,

$$\mu_k(\Delta_g + V) \leq \max \left\{ 2 \int_{\Omega} V dv_g ; 2(\nu+2)^{\frac{2}{\nu}} \left(\frac{H_{\Omega}}{|\Omega|_g} k \right)^{\frac{2}{\nu}} \right\}. \quad (40)$$

These estimates are to be compared with [6, Theorem 2.2 and Corollary 2.8], [7, Theorem 2.1], and the results by Grigor'yan, Netrusov and Yau [9, Theorem 5.15 and (1.14)] by which, under the assumption $\mu_0(\Delta_g + V) \geq 0$,

$$\mu_k(\Delta_g + V) \leq C(\Omega)k + \frac{1}{\varepsilon(\Omega)} \int_{\Omega} V dv_g \quad (41)$$

where $C(\Omega) > 0$ and $\varepsilon(\Omega) \in (0, 1)$ are two Riemannian constants that do not depend on V or k . They ask whether such an estimate holds true with $\varepsilon(\Omega) = 1$. The inequality (39) answers this question for the eigenvalue sums $\sum_{j=0}^{k-1} \mu_j$ in the affirmative, without any positivity condition.

On the other hand, unlike the upper bound in (41), our estimates (39) and (40) are consistent with the Weyl law regarding the power of k . Notice that (41) has been recently improved by A. Hassannezhad [11] who obtained

$$\mu_k(\Delta_g + V) \leq C(\Omega) + A_\nu \int_{\Omega} V dv_g + B_\nu \left(\frac{k}{|\Omega|_g} \right)^{\frac{2}{\nu}}$$

under the same assumption of positivity of $\mu_0(\Delta_g + V)$, where $A_\nu > 1$ and B_ν are two constants that only depend on the dimension ν , and $C(\Omega)$ is a Riemannian constant that does not depend on V or k . Our estimates are valid, however, under weaker assumptions and, moreover, the coefficient in front of $\int_{\Omega} V dv_g$ in (40) is equal to 1 while the other coefficient is explicitly computable at least in the elementary case where Ω is conformally Euclidean.

Example 3.2 (Witten Laplacians). Let Ω be a bounded domain of a Riemannian manifold (M, g) and let Δ_ρ be the Witten Laplacian associated with the density $e^{-2\rho}$, that is,

$$\Delta_\rho \varphi = \Delta_g \varphi + 2(\nabla^g \rho, \nabla^g \varphi).$$

The Neumann eigenvalues $\{\mu_l\}$ of Δ_ρ in Ω satisfy the following estimates :

(1) For all $z \in \mathbb{R}$,

$$\sum_{j \geq 0} (z - \mu_j)_+ \geq \frac{2 |\Omega|_g}{(\nu + 2) H_\Omega} \left(z - \int_{\Omega} |\nabla^g \rho|^2 dv_g \right)_+^{1 + \frac{\nu}{2}}. \quad (42)$$

(2) For all $k \in \mathbb{N}^*$,

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{\nu}{\nu + 2} \left(\frac{H_\Omega}{|\Omega|_g} k \right)^{\frac{2}{\nu}} + \int_{\Omega} |\nabla^g \rho|^2 dv_g. \quad (43)$$

(3) For all $k \in \mathbb{N}^*$,

$$\mu_k \left(1 - \frac{\int_{\Omega} |\nabla^g \rho|^2 dv_g}{\mu_k} \right)_+^{1 + \frac{2}{\nu}} \leq \left(\frac{\nu + 2}{2} \right)^{\frac{2}{\nu}} \left(\frac{H_\Omega}{|\Omega|_g} k \right)^{\frac{2}{\nu}}. \quad (44)$$

In particular,

$$\mu_k \leq \max \left\{ 2 \int_{\Omega} |\nabla^g \rho|^2 dv_g ; 2(\nu + 2)^{\frac{2}{\nu}} \left(\frac{H_\Omega}{|\Omega|_g} k \right)^{\frac{2}{\nu}} \right\}. \quad (45)$$

This last inequality is to be compared with the estimates obtained in [4].

For example, when Ω is a bounded domain of \mathbb{R}^ν endowed with the Gaussian density $e^{-|x|^2/2}$, we have for the corresponding Witten Laplacian

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{\nu}{\nu + 2} \left(4\pi^2 \left(\frac{k}{|\Omega| \omega_\nu} \right)^{\frac{2}{\nu}} + \frac{\omega_\nu}{|\Omega|} R^{\nu+2} \right),$$

where R is chosen so that Ω is contained in the Euclidean ball B_R .

Example 3.3 (Laplacian associated with a conformally Euclidean metric). Let Ω be a bounded domain of \mathbb{R}^ν and let $g = \alpha^{-2} g_E$ be a Riemannian metric that is conformal to the Euclidean metric g_E . The Neumann eigenvalues $\{\mu_l\}$ of the Laplacian Δ_g in Ω satisfy the following estimates in which $|\Omega|$ denotes the Euclidean volume of Ω :

(1) For all $z \in \mathbb{R}$,

$$\sum_{j \geq 0} (z - \mu_j)_+ \geq \frac{2\omega_\nu |\Omega|}{(\nu + 2)(2\pi)^\nu} \left(\int_{\Omega} \alpha^2 d^\nu x \right)^{\frac{\nu}{2}} \left(z - \frac{\nu^2}{4} \int_{\Omega} |\nabla \alpha|^2 d^\nu x \right)_+^{1 + \frac{\nu}{2}}. \quad (46)$$

(2) For all $k \in \mathbb{N}$,

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{4\pi^2\nu}{\nu+2} \left(\frac{k}{\omega_\nu|\Omega|} \right)^{\frac{2}{\nu}} \int_{\Omega} \alpha^2 d^\nu x + \frac{\nu^2}{4} \int_{\Omega} |\nabla\alpha|^2 d^\nu x. \quad (47)$$

(3) For all $k \in \mathbb{N}$,

$$\mu_k \left(1 - \frac{\nu^2}{4} \frac{\int_{\Omega} |\nabla\alpha|^2 d^\nu x}{\mu_k} \right)_+^{1+\frac{2}{\nu}} \leq 4\pi^2 \left(\frac{\nu+2}{2} \right)^{\frac{2}{\nu}} \left(\frac{k}{\omega_\nu|\Omega|} \right)^{\frac{2}{\nu}} \int_{\Omega} \alpha^2 d^\nu x. \quad (48)$$

In particular,

$$\mu_k \leq \max \left\{ \frac{\nu^2}{2} \int_{\Omega} |\nabla\alpha|^2 d^\nu x ; 8\pi^2 (\nu+2)^{\frac{2}{\nu}} \left(\frac{k}{\omega_\nu|\Omega|} \right)^{\frac{2}{\nu}} \int_{\Omega} \alpha^2 d^\nu x \right\}. \quad (49)$$

Note that a domain of the hyperbolic space \mathbf{H}^ν can be identified with a domain of the Euclidean unit ball endowed with the metric $g = \left(\frac{2}{1-|x|^2} \right)^2 g_E$. For such a domain we get, with $\alpha = \frac{1-|x|^2}{2}$, $\int_{\Omega} \alpha^2 d^\nu x \leq \frac{1}{4}$, and $\int_{\Omega} |\nabla\alpha|^2 d^\nu x = \int_{\Omega} |x|^2 d^\nu x$,

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j \leq \frac{\pi^2\nu}{\nu+2} \left(\frac{k}{\omega_\nu|\Omega|} \right)^{\frac{2}{\nu}} + \frac{\nu^2}{4} \int_{\Omega} |x|^2 d^\nu x. \quad (50)$$

$$\mu_k \left(1 - \frac{\nu^2}{4} \frac{\int_{\Omega} |x|^2 d^\nu x}{\mu_k} \right)_+^{1+\frac{2}{\nu}} \leq \pi^2 \left(\frac{\nu+2}{2} \right)^{\frac{2}{\nu}} \left(\frac{k}{\omega_\nu|\Omega|} \right)^{\frac{2}{\nu}} \quad (51)$$

and

$$\mu_k \leq \max \left\{ \frac{\nu^2}{2} \int_{\Omega} |x|^2 d^\nu x ; 2\pi^2 (\nu+2)^{\frac{2}{\nu}} \left(\frac{k}{\omega_\nu|\Omega|} \right)^{\frac{2}{\nu}} \right\}. \quad (52)$$

4. SUMS OF NEUMANN EIGENVALUES ON DOMAINS CONFORMAL TO EUCLIDEAN SETS, AND PHASE-SPACE VOLUMES

A phase-space analysis can considerably sharpen the upper bounds on sums of eigenvalues from the previous sections so that they become sharp in the semiclassical regime. Following physical tradition, it is shown in [21] how this may be achieved in some circumstances with the aid of coherent states. We carry out such an analysis in this section for (1) when $(M, g) = (\mathbb{R}^\nu, d^\nu x)$. We must first introduce a few quantities that will be helpful to relate spectral estimates to phase-space volumes. To avoid complications we assume that the potential energy V is Lipschitz continuous and bounded from below. We do not assume that Ω is necessarily bounded, but if it is not, we require V to be confining in the sense that there is a radial function $V_{\text{rad}}(r)$ tending to $+\infty$ as $r \rightarrow \infty$ with $V(\mathbf{x}) \geq V_{\text{rad}}(|\mathbf{x}|)$ for all $\mathbf{x} \notin \Omega$. This condition is sufficient to ensure that the eigenvalues form a discrete sequence tending to $+\infty$.

Definition 4.1. The effective potential incorporating a correction for the conformal transformation will be denoted $\tilde{V}(\mathbf{x}) := V(\mathbf{x}) + |\nabla\rho|^2(\mathbf{x})$, and the maximal Lipschitz constant of $\tilde{V}(\mathbf{x})$ on the region $\Omega \cap \{\mathbf{x} : \tilde{V}(\mathbf{x}) \leq \Lambda\}$ will be denoted $\text{Lip}(\Lambda)$.

The L^2 -normalized ground-state Dirichlet eigenfunction for the ball of geodesic radius r in M will be denoted h_r and $\mathcal{K}(h_r) := \int_{B_r} |\nabla h_r(\mathbf{x})|^2 d^\nu x$. I.e., in this section where $M = \mathbb{R}^\nu$, h is a scaled Bessel function and

$$\mathcal{K}(h_r) = \frac{j_{\frac{\nu}{2}-1,1}^2}{r^2}.$$

Remark 4.1. *The function h_r will ensure that some coherent-state functions to be defined below are localized in configuration space. Its specific form is but one of many plausible choices.*

We next recall some quantities that arise in phase-space analysis.

Definition 4.2. *The Euclidean phase-space volume for energy Λ is defined as*

$$\Phi_1(\Lambda) := \frac{1}{(2\pi)^\nu} \left| \{(\mathbf{x}, \mathbf{p}) : |\mathbf{p}|^2 + \tilde{V}(\mathbf{x}) \leq \Lambda\} \right| = \frac{\omega_\nu}{(2\pi)^\nu} \int_{\Omega} \left(\Lambda - \tilde{V}(\mathbf{x}) \right)_+^{\frac{\nu}{2}} d^\nu x,$$

according to a standard calculation to be found, for example, in [21]. If the weight in (1) is not constant, we make use of a weighted phase-space volume,

$$\Phi_w(\Lambda) = \frac{\omega_\nu}{(2\pi)^\nu} \int_{\Omega} \left(\Lambda - \tilde{V}(\mathbf{x}) \right)_+^{\frac{\nu}{2}} w(\mathbf{x}) d^\nu x.$$

The total energy associated with this quantity is correspondingly

$$\begin{aligned} E_w(\Lambda) &:= \frac{1}{(2\pi)^\nu} \int_{\{(\mathbf{x}, \mathbf{p}) : \mathbf{x} \in \Omega, |\mathbf{p}|^2 + \tilde{V}(\mathbf{x}) \leq \Lambda\}} \left(|\mathbf{p}|^2 + \tilde{V}(\mathbf{x}) \right) w(\mathbf{x}) d^\nu x d^\nu p \\ &= \frac{\nu}{\nu + 2} \frac{\omega_\nu}{(2\pi)^\nu} \int_{\Omega} \left(\Lambda - \tilde{V}(\mathbf{x}) \right)_+^{1 + \frac{\nu}{2}} w(\mathbf{x}) d^\nu x. \end{aligned} \quad (53)$$

We note that according to (53),

$$\frac{dE_w}{d\Lambda}(\Lambda) = \Phi_w(\Lambda), \quad (54)$$

and that Φ_w increases strictly monotonically in Λ , implying that E_w is strictly convex.

Theorem 4.1. *Let $\mu_0 \leq \mu_1 \leq \dots$ be the variationally defined Neumann eigenvalues (2) on an open set $\Omega \in \mathbb{R}^\nu$, where w, ρ , and V satisfy the assumptions stated above, and define $\Lambda(k)$ as the minimal value of Λ for which $\Phi_1(\Lambda) \geq (2\pi)^\nu k$. Then*

$$\sum_{j=0}^{k-1} \mu_j \leq E_w(\Lambda(k)) + 3 \left(2j_{\nu-1,1}^2 \text{Lip}(\Lambda(k)) \right)^{\frac{1}{3}} \Phi_w \left(\Lambda(k) + \left(2j_{\nu-1,1}^2 \text{Lip}(\Lambda(k)) \right)^{\frac{1}{3}} \right). \quad (55)$$

The Riesz-mean form of the inequality reads

$$\sum_{j=0}^{k-1} (z - \mu_j)_+ \geq \frac{2}{\nu + 2} \frac{\omega_\nu}{(2\pi)^\nu} \int (z - V(y))_+^{1 + \frac{\nu}{2}} dy \quad (56)$$

$$- \left(\frac{\omega_\nu}{(2\pi)^\nu} \int (z - V(y))_+^{\frac{\nu}{2}} dy \right) \left(\|\nabla h_r\|^2 + \int |\mathbf{x}| h_r^2 \right). \quad (57)$$

Remarks 4.1. 1. *We call attention to the fact that the condition in this theorem defining Λ uses the Euclidean phase space, whereas weighted phase-space quantities appear in (55).*

2. *The dominant term in the semiclassical regime can be identified by introducing a small parameter α as a coefficient of $|\nabla\varphi|^2$ in (1), i.e.,*

$$\mathcal{R}_\alpha(\varphi) := \frac{\int_{\Omega} (\alpha |\nabla\varphi(\mathbf{x})|^2 + V(\mathbf{x}) |\varphi(\mathbf{x})|^2) w(\mathbf{x}) e^{-2\rho(\mathbf{x})} dv_g}{\int_{\Omega} |\varphi(\mathbf{x})|^2 e^{-2\rho(\mathbf{x})} dv_g}. \quad (58)$$

The result, in the Riesz-mean form after choosing a convenient relationship between r and α , is

$$\begin{aligned} \sum_{j=0}^{k-1} (z - \mu_j)_+ &\geq \alpha^{-\frac{\nu}{2}} \frac{2}{\nu + 2} \frac{\omega_\nu}{(2\pi)^\nu} \int (z - V(y))_+^{1+\frac{\nu}{2}} dy \\ &\quad - \left(\alpha^{\frac{1}{3}-\frac{\nu}{2}} \frac{\omega_\nu}{(2\pi)^\nu} \int (z - V(y))_+^{\frac{\nu}{2}} dy \right) \left(\|\nabla h_1\|^2 + \int |\mathbf{x}| h_1^2 \right), \end{aligned} \quad (59)$$

in which the leading term is precisely the expected semiclassical expression, in contrast to results of the previous section such as Example 3.3.

3. Inequalities of the type (55) imply estimates of quantities including trace of the heat kernel (= the partition function in quantum physics) and the spectral zeta functions by simple transforms. For instance (55) implies for the Riesz mean $R_1(z) := \sum_j (z - \mu_j)_+$ that

$$R_1(z) \geq \frac{1}{(2\pi)^\nu} \left(z \Phi_w(\Lambda(z)) - E_w(\Lambda(k)) + 3 (2j_{\nu-1,1}^2 \text{Lip}(\Lambda(k)))^{\frac{1}{3}} \Phi_w \left(\Lambda(k) + (2j_{\nu-1,1}^2 \text{Lip}(\Lambda(k)))^{\frac{1}{3}} \right) \right). \quad (60)$$

The Riesz mean is in turn related to the heat trace by the Laplace transform (32).

Proof of Theorem 4.1. We apply Theorem 2.1 to the Neumann eigenvalues of (1) as defined by (2), using for test functions “coherent states” [21, 27] of the form:

$$f_\zeta(\mathbf{x}) := \frac{1}{(2\pi)^{\nu/2}} e^{i\mathbf{p}\cdot(\mathbf{x})+\rho(\mathbf{x})} h_r(\mathbf{x} - \mathbf{y}).$$

In this formula, $\zeta = (\mathbf{p}, \mathbf{y})$ ranges over the phase space $\mathfrak{M} = \mathbb{R}^{2\nu}$ with Lebesgue measure. The radius r will be chosen below.

We note that the inner product that appears is a Fourier transform with respect to the variable \mathbf{x} , viz.,

$$\langle \phi, f_\zeta \rangle = \mathfrak{F}[h_r(\mathbf{x} - \mathbf{y}) e^{-\rho(\mathbf{x})} \phi(\mathbf{x})],$$

where if Ω is a strict subset of \mathbb{R}^ν , then ϕ is extended by 0 outside Ω . Thus, with the Parseval identity,

$$\int_{\mathbb{R}^{2\nu}} |\langle \phi, f_\zeta \rangle|^2 d^\nu p d^\nu y = \int_{\mathbb{R}^\nu} \int_{\mathbb{R}^\nu} h_r(\mathbf{x} - \mathbf{y})^2 |\phi|^2 e^{-2\rho} d^\nu y d^\nu x = \int_{\mathbb{R}^\nu} |\phi|^2 e^{-2\rho} d^\nu x = \|\phi\|^2. \quad (61)$$

The set \mathfrak{M}_0 in Theorem 2.1 must be taken large enough so that

$$k \leq \int_{\mathfrak{M}_0} \|f_\zeta\|^2 d\sigma = \frac{1}{(2\pi)^\nu} \int_{\mathfrak{M}_0} \int_{\Omega} h_r^2(\mathbf{x} - \mathbf{y}) e^{2\rho(\mathbf{x})-2\rho(\mathbf{y})} d^\nu x d^\nu p d^\nu y \leq \frac{1}{(2\pi)^\nu} |\mathfrak{M}_0|, \quad (62)$$

in which case

$$\begin{aligned} \sum_{j=0}^{k-1} \mu_j &\leq \int_{\mathfrak{M}_0} \mathcal{R}(f_\zeta) d^\nu p d^\nu y = \\ &= \frac{1}{(2\pi)^\nu} \int_{\mathfrak{M}_0 \times \Omega} \left((|\mathbf{p}|^2 + V(\mathbf{x})) h_r^2(\mathbf{x} - \mathbf{y}) + |\nabla h_r(\mathbf{x} - \mathbf{y}) + h_r(\mathbf{x} - \mathbf{y}) \nabla \rho(\mathbf{x})|^2 \right) w(\mathbf{x}) d^\nu x d^\nu p d^\nu y. \end{aligned} \quad (63)$$

We now make the ansatz that $\mathfrak{M}_0 = \mathfrak{M}_0(\Lambda) := \{(\mathbf{x}, \mathbf{p}) : \mathbf{x} \in \Omega, |\mathbf{p}|^2 + \tilde{V}(\mathbf{x}) \leq \Lambda\}$, where $\Lambda \geq \Lambda(k)$, defined as the minimum value of Λ for (62) to be valid. Thus the upper bound in (62) becomes $\frac{1}{(2\pi)^\nu} \Phi_1(\Lambda)$, whence the condition in the theorem.

Since the support of h is restricted to a ball of radius r , the x -integral may be restricted to the set $\{\mathbf{x} \in \Omega : \exists \mathbf{y}, |\mathbf{y} - \mathbf{x}| \leq r, |\mathbf{p}|^2 + \tilde{V}(\mathbf{x}) \leq \Lambda\} \subset \mathfrak{M}_0(\Lambda + \text{Lip}(\Lambda)r)$. Thus, integrating first in \mathbf{y} , the right side of (63) is bounded above by

$$\begin{aligned} & \frac{1}{(2\pi)^\nu} \int_{\{\mathbf{p}:|\mathbf{p}|^2 \leq \Lambda\}} \int_{\{\mathbf{x}:\tilde{V}(\mathbf{x}) \leq \Lambda + \text{Lip}(\Lambda)r - |\mathbf{p}|^2\}} \int_{\mathbb{R}^\nu} \left((|\mathbf{p}|^2 + V(\mathbf{x}))h_r^2(\mathbf{x} - \mathbf{y}) \right. \\ & \quad \left. + |\nabla h_r(\mathbf{x} - \mathbf{y}) + h_r(\mathbf{x} - \mathbf{y})\nabla \rho(\mathbf{x})|^2 \right) w(\mathbf{x})d^\nu y d^\nu x d^\nu p \\ & = \frac{1}{(2\pi)^\nu} \int_{\{\mathbf{p}:|\mathbf{p}|^2 \leq \Lambda\}} \int_{\{\mathbf{x}:\tilde{V}(\mathbf{x}) \leq \Lambda + \text{Lip}(\Lambda)r - |\mathbf{p}|^2\}} \int_{\mathbb{R}^\nu} \left((|\mathbf{p}|^2 + \tilde{V}(\mathbf{x}))h_r^2(\mathbf{x} - \mathbf{y}) \right. \\ & \quad \left. + |\nabla h_r(\mathbf{x} - \mathbf{y})|^2 + \nabla \rho(\mathbf{x}) \cdot \nabla h_r^2(\mathbf{x} - \mathbf{y}) \right) w(\mathbf{x})d^\nu y d^\nu x d^\nu p. \end{aligned}$$

The last contribution vanishes because

$$\int_{\mathbb{R}^\nu} \nabla \rho(\mathbf{x}) \cdot \nabla h_r^2(\mathbf{x} - \mathbf{y})d^\nu y = \int_{\mathbb{R}^\nu} \nabla \rho(\mathbf{x}) \cdot \nabla 1d^\nu y = 0, \quad (64)$$

leaving

$$\sum_{j=0}^{k-1} \mu_j \leq \frac{1}{(2\pi)^\nu} \int_{\mathfrak{M}_0(\Lambda + \text{Lip}(\Lambda)r)} (|\mathbf{p}|^2 + \tilde{V}(\mathbf{x}) + \mathcal{K}(h_r))w(\mathbf{x})d^\nu x d^\nu p \quad (65)$$

for all values of $r > 0$. The upper bound (65) is of the form

$$\begin{aligned} & \frac{1}{(2\pi)^\nu} (E_w(\Lambda + \text{Lip}(\Lambda)r) + \Phi_w(\Lambda + \text{Lip}(\Lambda)r)\mathcal{K}(h_r)) \\ & \leq \frac{1}{(2\pi)^\nu} \left(E_w(\Lambda) + \left(\frac{j_{\nu-1,1}^2}{r^2} + \text{Lip}(\Lambda)r \right) \Phi_w(\Lambda + \text{Lip}(\Lambda)r) \right), \end{aligned}$$

where we have made use of (54) and the monotonicity of Φ_w in a first-order expansion of E_w .

Choosing the optimal value $r = \left(\frac{2j_{\nu-1,1}^2}{L} \right)^{\frac{1}{3}}$, we get the claim (55).

The derivation of (56) proceeds similarly. \square

Remark 4.2. We note the following special cases of particular interest.

1. Laplace operators with Neumann conditions on a compact Euclidean domain ($V = \rho = 0$, $w = 1$). In this case $\text{Lip}(\Lambda) = 0$, $\Lambda^{\frac{\nu}{2}} := \frac{(2\pi)^\nu k}{B_\nu |\Omega|}$, and we recover the inequality of Kröger, that

$$\sum_{j=0}^{k-1} \mu_j \leq \frac{\nu}{\nu + 2} \frac{\omega_\nu}{(2\pi)^\nu} |\Omega| \Lambda^{1+\frac{\nu}{2}} = \frac{\nu}{\nu + 2} (2\pi)^2 \omega_\nu^{-\frac{2}{\nu}} \frac{k^{\frac{\nu+2}{\nu}}}{|\Omega|^{\frac{2}{\nu}}}.$$

Indeed, without the potential V , the introduction of the function h_r is not needed for the proof.

2. Nonhomogeneous problems with $\rho = V = 0$, but w is variable, under Neumann conditions:

$$\sum_{j=0}^{k-1} \mu_j \leq \frac{\nu}{\nu + 2} \frac{\omega_\nu}{(2\pi)^\nu} \left(\int_{\Omega} w(\mathbf{x})d^\nu x \right) \Lambda^{1+\frac{\nu}{2}}.$$

The eigenvalue bounds of Corollary 3.1 are sharp as k tends to infinity. Indeed since $\widetilde{\mu}_k = \mu_k$ we get

$$\mu_k \leq 4\pi^2 \int_{\Omega} w(\mathbf{x})d^\nu x \left(\frac{k}{|\Omega|\omega_\nu} \right)^{\frac{2}{\nu}} \left(1 + 2\sqrt{\frac{1 - S_k}{\nu + 2}} \right)$$

with S_k as given in Corollary 3.1 :

$$S_k = \frac{\frac{\nu+2}{\nu} \frac{1}{k} \sum_{j=0}^{k-1} \mu_j}{4\pi^2 \int_{\Omega} w(\mathbf{x}) d^\nu x \left(\frac{k}{|\Omega| \omega_\nu} \right)^{\frac{2}{\nu}}} \leq 1.$$

5. BOUNDS FOR NEUMANN EIGENVALUES ON SUBDOMAINS OF COMPACT HOMOGENEOUS SPACES

In this section, we deal with the case where the ambient space is a compact homogeneous Riemannian manifold (M, g) with isomorphism group denoted G . In particular, we shall recover Strichartz's result [26] with a more efficient proof and extend it to a wider class of operators. We begin with bounds in the spirit of Theorem 3.1 and then derive a phase-space bound analogous to Theorem 4.1.

Let us denote by

$$\text{spec}(M) = \{0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots\}$$

the spectrum of the Laplace-Beltrami operator Δ_g on M (each eigenvalue is repeated according to its multiplicity). Although 0 is a simple eigenvalue, all the other eigenvalues are degenerate owing to the transitive action of the isometry group G (recall that the eigenspaces are invariant under the action of G .)

Given a regular domain $\Omega \subset M$ endowed with densities $e^{-2\rho}$ and $e^{-2\theta} = we^{-2\rho}$, and a potential V , we consider the eigenvalues $\mu_l(\Omega, g, \rho, w, V)$, $l \in \mathbb{N}$, defined by (1) and (2) and seek for relationships between the μ_l 's and the λ_l 's. As before, we will use the notation $\tilde{V} = V + |\nabla^g \rho|^2$. We also need the following subspaces introduced in Theorem 2.1

$$E_0(R) = \bigoplus_{\mu < R} \ker(H - \mu I) \quad \text{and} \quad E(R) = \bigoplus_{\mu \leq R} \ker(H - \mu I)$$

where $H = H(\Omega, g, \rho, w, V)$ is the operator defined by (4). The corresponding subspaces associated with the Laplacian Δ_g on M will be denoted

$$F_0(R) = \bigoplus_{\lambda < R} \ker(\Delta_g - \lambda I) \quad \text{and} \quad F(R) = \bigoplus_{\lambda \leq R} \ker(\Delta_g - \lambda I).$$

Theorem 5.1. *Let (M, g) be a compact homogeneous Riemannian manifold. Let $\mu_l = \mu_l(\Omega, g, \rho, w, V)$, $l \in \mathbb{N}$, be the eigenvalues defined by (2) on a bounded open set $\Omega \subset M$. Then, for all $z \in \mathbb{R}$,*

$$\sum_{j \geq 0} (z - \mu_j)_+ \geq \frac{|\Omega|_g}{|M|_g} \sum_{j \geq 0} (z - \tilde{\lambda}_j)_+, \quad (66)$$

where $\tilde{\lambda}_j = \lambda_j \int_{\Omega} w dv_g + \int_{\Omega} \tilde{V} w dv_g$. Equality holds in (66) for some $z \in \mathbb{R}$ if and only if

$$E_0(z) \subset e^\rho F(\tilde{z}) \subset E(z),$$

with $\tilde{z} = \frac{1}{\int_{\Omega} w dv_g} \left(z - \int_{\Omega} \tilde{V} w dv_g \right)$.

Proof. Let $\{y_\lambda : \lambda \in \text{spec}(M)\}$ be an orthonormal basis of $L^2(M, g)$ with $\Delta_g y_\lambda = \lambda y_\lambda$. The proof relies on Theorem 2.1 in which we take $\mathfrak{M} = \text{spec}(M)$ endowed with the uniform discrete measure, and use test functions of the form

$$f_\lambda = y_\lambda e^\rho.$$

For any function $\psi \in L^2(\Omega, e^{-2\rho} dv_g)$ (endowed with the norm $\|\psi\|^2 = \int_{\Omega} \psi^2 e^{-2\rho} dv_g$),

$$\begin{aligned} \int_{\mathfrak{M}} \langle f_{\lambda}, \psi \rangle^2 d\lambda &= \sum_{\lambda \in \text{spec}(M)} \left(\int_{\Omega} f_{\lambda} \psi e^{-2\rho} dv_g \right)^2 = \sum_{\lambda \in \text{spec}(M)} \left(\int_{\Omega} y_{\lambda} \psi e^{-\rho} dv_g \right)^2 \\ &= \sum_{\lambda \in \text{spec}(M)} \left(\int_M y_{\lambda} \psi e^{-\rho} dv_g \right)^2 = \int_M \psi^2 e^{-2\rho} dv_g = \|\psi\|^2, \end{aligned}$$

where we used the same notation ψ to designate the extension of ψ by zero outside Ω .

Let $R > 0$ and let $\mathfrak{M}_0 = \{\lambda \in \mathfrak{M} : \lambda \leq R\}$. Due to the transitive action of the isometry group G on M , for every eigenvalue Λ of Δ_g , with multiplicity m_{Λ} , the basis $\{y_{\lambda} : \lambda = \Lambda\}$ of the corresponding eigenspace is such that $\sum_{\lambda=\Lambda} y_{\lambda}^2$ is constant on M . Integrating over M , we get

$$\sum_{\lambda=\Lambda} y_{\lambda}^2 = \frac{m_{\Lambda}}{|M|_g}. \quad (67)$$

Moreover, $0 = \frac{1}{2} \Delta_g (\sum_{\lambda=\Lambda} y_{\lambda}^2) = \sum_{\lambda=\Lambda} (\Lambda y_{\lambda}^2 - |\nabla^g y_{\lambda}|^2)$, that is

$$\sum_{\lambda=\Lambda} |\nabla^g y_{\lambda}|^2 = \frac{m_{\Lambda}}{|M|_g} \Lambda. \quad (68)$$

Therefore,

$$\int_{\mathfrak{M}_0} \|f_{\lambda}\|^2 d\lambda = \sum_{\lambda \leq R} \int_{\Omega} y_{\lambda}^2 dv_g = \sum_{\Lambda \leq R} \frac{|\Omega|_g}{|M|_g} m_{\Lambda} = \frac{|\Omega|_g}{|M|_g} N(R).$$

where $N(R)$ is the number of eigenvalues of Δ_g on M that are less or equal to R (counted with multiplicity). On the other hand, using (67) and (68), we get for every Λ ,

$$\begin{aligned} \sum_{\lambda=\Lambda} |\nabla^g f_{\lambda}|^2 &= e^{2\rho} \sum_{\lambda=\Lambda} (|\nabla^g y_{\lambda}|^2 + y_{\lambda}^2 |\nabla^g \rho|^2 + g(\nabla^g \rho, \nabla^g y_{\lambda}^2)) \\ &= \frac{m_{\Lambda}}{|M|_g} e^{2\rho} (\Lambda + |\nabla^g \rho|^2). \end{aligned}$$

Thus

$$\begin{aligned} \int_{\mathfrak{M}_0} Q(f_{\lambda}, f_{\lambda}) d\lambda &= \sum_{\lambda \leq R} \int_{\Omega} (|\nabla^g f_{\lambda}|^2 + V f_{\lambda}^2) w e^{-2\rho} dv_g \\ &= \sum_{\Lambda \leq R} \int_{\Omega} \frac{m_{\Lambda}}{|M|_g} (\Lambda + |\nabla^g \rho|^2 + V) w dv_g \frac{\int_{\Omega} w dv_g}{|M|_g} \sum_{\lambda \leq R} \lambda + \frac{\int_{\Omega} \tilde{V} w dv_g}{|M|_g} N(R). \end{aligned}$$

Inserting into (6), we get for every $z \in \mathbb{R}$ and $R > 0$,

$$\begin{aligned} \sum_{j \geq 0} (z - \mu_j)_+ &\geq z \frac{|\Omega|_g}{|M|_g} N(R) - \frac{\int_{\Omega} w dv_g}{|M|_g} \sum_{\lambda \leq R} \lambda - \frac{\int_{\Omega} \tilde{V} w dv_g}{|M|_g} N(R) \\ &= \frac{|\Omega|_g}{|M|_g} \sum_{\lambda \leq R} \left(z - \lambda \int_{\Omega} w dv_g - \int_{\Omega} \tilde{V} w dv_g \right). \end{aligned} \quad (69)$$

Notice that the RHS is negative if $z \leq \int_{\Omega} \tilde{V} w dv_g$. Now, when $z > \int_{\Omega} \tilde{V} w dv_g$, we can choose $R = \tilde{z} = \frac{z - \int_{\Omega} \tilde{V} w dv_g}{\int_{\Omega} w dv_g}$ so that the last sum is taken over all eigenvalues λ for which the involved

terms are nonnegative, thus

$$\sum_{j \geq 0} (z - \mu_j)_+ \geq \frac{|\Omega|_g}{|M|_g} \sum_{j \geq 0} \left(z - \lambda_j \int_{\Omega} w dv_g - \int_{\Omega} \tilde{V} w dv_g \right)_+.$$

Regarding the case of equality, it follows from Theorem 2.1 that equality holds in (66) if and only if $f_{\lambda} \in E(z)$ for $\lambda \leq \tilde{z}$ and f_{λ} is orthogonal to $E_0(z)$ for $\lambda > \tilde{z}$. Equivalently, $e^{\rho} F(\tilde{z}) \subset E(z)$ and, since $\text{Span}\{f_{\lambda} : \lambda > \tilde{z}\}$ is the orthogonal complement of $e^{\rho} F(\tilde{z})$, $E_0(z) \subset e^{\rho} F(\tilde{z})$. \square

As we have seen in the previous sections, our technique allows obtaining bounds on eigenvalue sums. In order to simplify the statement of these bounds, we introduce the following notation : Given any sequence $(a) = (a_k)_{k \geq 0}$ of real numbers, we set for $p \in [1, +\infty)$,

$$\mathfrak{S}_{(a)}(p) = \sum_{j=0}^{\lfloor p \rfloor - 1} a_j + (p - \lfloor p \rfloor) a_{\lfloor p \rfloor},$$

so that when p is an integer, $\mathfrak{S}_{(a)}(p)$ is nothing but the sum of the first p terms a_0, \dots, a_{p-1} of the sequence (a) .

Theorem 5.2. *Let (M, g) be a compact homogeneous Riemannian manifold. Let $(\mu) = (\mu_l)_{l \geq 0}$ be the sequence of eigenvalues defined by (2) on an open set $\Omega \subset M$. Then, for every $p \in [1, +\infty)$,*

$$\mathfrak{S}_{(\mu)}(p) \leq \frac{|\Omega|_g}{|M|_g} \mathfrak{S}_{(\tilde{\lambda})} \left(\frac{|M|_g}{|\Omega|_g} p \right), \quad (70)$$

where $(\tilde{\lambda}) = (\tilde{\lambda}_l)_{l \geq 0}$ is the sequence defined by $\tilde{\lambda}_l = \lambda_l \int_{\Omega} w dv_g + \int_{\Omega} \tilde{V} w dv_g$. Moreover, equality holds in (70) for some $p = k \in \mathbb{N}^*$ if and only if

$$E_0(\mu_k) \subset e^{\rho} F_0(\lambda_{\check{k}}) \quad \text{and} \quad e^{\rho} F(\lambda_{\hat{k}-1}) \subset E(\mu_k). \quad (71)$$

with $\check{k} = \lfloor \frac{|M|_g}{|\Omega|_g} k \rfloor$ and $\hat{k} = \lceil \frac{|M|_g}{|\Omega|_g} k \rceil$, where $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ denote the floor and the ceiling functions, respectively.

Observe that we have $F_0(\lambda_{\check{k}}) \subset F(\lambda_{\hat{k}-1})$ with equality if and only if $\lambda_{\hat{k}-1} < \lambda_{\check{k}}$.

Proof of Theorem 5.2. As mentioned in Remark 3.1, Legendre's transform enables us to obtain (70) from (66). Alternatively, we can prove (70) using the averaged principle, which has the advantage of allowing us to characterize the case of equality. Indeed, taking $z = \mu_k$ in (69) we immediately get, $\forall R > 0$

$$\sum_{j=0}^{k-1} \mu_j \leq \frac{|\Omega|_g}{|M|_g} \sum_{j=0}^{N(R)-1} \tilde{\lambda}_j + \left(k - \frac{|\Omega|_g}{|M|_g} N(R) \right) \mu_k. \quad (72)$$

Denote by $1 = N_0 < N_1 < N_2 < \dots < N_j < \dots$ the values taken by the function $N(R)$, $R \in \mathbb{R}$, that is $N_j = m_0 + m_1 + \dots + m_j$. The sequence of eigenvalues of Δ_g on M is then numbered as follows :

$$\begin{aligned} 0 = \lambda_0 < \lambda_1 = \lambda_2 = \dots = \lambda_{N_1-1} < \lambda_{N_1} = \dots = \lambda_{N_2-1} < \lambda_{N_2} = \dots \\ = \lambda_{N_j-1} < \lambda_{N_j} = \dots = \lambda_{N_{j+1}-1} < \lambda_{N_{j+1}} = \dots \end{aligned}$$

Let $q \in \mathbb{N}$ such that

$$N_q \leq \frac{|M|_g}{|\Omega|_g} k < N_{q+1}.$$

We consider the inequality (72) with first $N(R) = N_q$ and, then, $N(R) = N_{q+1}$. We multiply the first inequality by $\alpha = (N_{q+1} - \frac{|M|_g}{|\Omega|_g}k)/m_{q+1}$ and add the second inequality multiplied by $1 - \alpha = (\frac{|M|_g}{|\Omega|_g}k - N_q)/m_{q+1}$ to get

$$\begin{aligned} \sum_{j=0}^{k-1} \mu_j &\leq \frac{|\Omega|_g}{|M|_g} \sum_{j=0}^{N_q-1} \tilde{\lambda}_j + (1 - \alpha) \frac{|\Omega|_g}{|M|_g} m_{q+1} \tilde{\lambda}_{N_q} + k\mu_k - \frac{|\Omega|_g}{|M|_g} (\alpha N_q + (1 - \alpha) N_{q+1}) \mu_k \\ &= \frac{|\Omega|_g}{|M|_g} \sum_{j=0}^{N_q-1} \tilde{\lambda}_j + \left(k - \frac{|\Omega|_g}{|M|_g} N_q \right) \tilde{\lambda}_{N_q} = \frac{|\Omega|_g}{|M|_g} \left(\sum_{j=0}^{N_q-1} \tilde{\lambda}_j + \left(\frac{|M|_g}{|\Omega|_g} k - N_q \right) \tilde{\lambda}_{N_q} \right) \end{aligned}$$

since α is chosen such that $\alpha N_q + (1 - \alpha) N_{q+1} = \frac{|M|_g}{|\Omega|_g} k$. Now, from the definition of N_q we have $\lambda_{N_q} = \lambda_{N_q+1} = \dots = \lambda_{\lfloor \frac{|M|_g}{|\Omega|_g} k \rfloor}$ and, then,

$$\sum_{j=0}^{N_q-1} \tilde{\lambda}_j + \left(\frac{|M|_g}{|\Omega|_g} k - N_q \right) \tilde{\lambda}_{N_q} = \mathfrak{S}_{(\tilde{\lambda})} \left(\frac{|M|_g}{|\Omega|_g} k \right),$$

which yields

$$\sum_{j=0}^{k-1} \mu_j \leq \frac{|\Omega|_g}{|M|_g} \mathfrak{S}_{(\tilde{\lambda})} \left(\frac{|M|_g}{|\Omega|_g} k \right).$$

This means that (70) holds for all $p \in \mathbb{N}$. Since the functions $\mathfrak{S}_{(\mu)}$ and $\mathfrak{S}_{(\tilde{\lambda})}$ are piecewise-affine in p , the extension of (70) to all positive p is immediate.

Let k be a positive integer. The equality is achieved in (70) for $p = k$ if and only if one of the following holds :

- $\frac{|M|_g}{|\Omega|_g} k = N_q$ and equality holds in (72) for R such that $N(R) = N_q$, i.e. for $R = \lambda_{N_q-1}$
- $\frac{|M|_g}{|\Omega|_g} k > N_q$ and equality holds in (72) for the values of R such that $N(R) = N_q$ and $N(R) = N_{q+1}$, i.e. for both $R = \lambda_{N_q-1}$ and $R = \lambda_{N_{q+1}-1}$.

The first case corresponds to the case of equality in Theorem 2.1 with $z = \mu_k$, $\mathfrak{M} = \text{spec}(M)$, $\mathfrak{M}_0 = \{\lambda \in \text{spec}(M) ; \lambda \leq \lambda_{N_q-1}\}$. As in the proof of Theorem 5.1, this situation occurs if and only if $E_0(\mu_k) \subset e^\rho F(\lambda_{N_q-1}) \subset E(\mu_k)$, with $N_q = \frac{|M|_g}{|\Omega|_g} k$. Since $\lambda_{N_q-1} < \lambda_{N_q}$, $F(\lambda_{N_q-1}) = F_0(\lambda_{N_q})$, and the last conditions can be written as follows :

$$E_0(\mu_k) \subset e^\rho F_0(\lambda_{\frac{|M|_g}{|\Omega|_g} k}) \text{ and } F(\lambda_{\frac{|M|_g}{|\Omega|_g} k-1}) \subset E(\mu_k). \quad (73)$$

which is equivalent to (71).

In the second case, similar considerations show that equality holds if and only if

$$E_0(\mu_k) \subset e^\rho F(\lambda_{N_q-1}) \subset e^\rho F(\lambda_{N_{q+1}-1}) \subset E(\mu_k). \quad (74)$$

Since $N_q < \frac{|M|_g}{|\Omega|_g} k < N_{q+1}$, it is clear that

$$N_q \leq \left\lfloor \frac{|M|_g}{|\Omega|_g} k \right\rfloor \leq N_{q+1} - 1$$

and

$$N_q \leq \left\lceil \frac{|M|_g}{|\Omega|_g} k \right\rceil - 1 \leq N_{q+1} - 1.$$

Thus,

$$\lambda_{N_q} = \lambda_{N_{q+1}-1} = \lambda \left[\frac{|M|_g k}{|\Omega|_g} \right] = \lambda \left[\frac{|M|_g}{|\Omega|_g} k \right]_{-1}.$$

Consequently,

$$F(\lambda_{N_{q+1}-1}) = F \left(\lambda \left[\frac{|M|_g}{|\Omega|_g} k \right]_{-1} \right)$$

and, since $\lambda_{N_q-1} < \lambda_{N_q}$,

$$F(\lambda_{N_q-1}) = F_0(\lambda_{N_q}) = F_0 \left(\lambda \left[\frac{|M|_g}{|\Omega|_g} k \right] \right).$$

Therefore, (74) is equivalent to (71). \square

Remarks 5.1. 1. The particular case of (70) in which $w = 1$ and $\rho = V = 0$ corresponds to the inequality obtained by Strichartz [26, Theorem 2.2].

In the same paper [26], Strichartz also proved, following Gallot [8, Proposition 2.9], that for Dirichlet eigenvalues μ_l^D of the Laplacian on a domain Ω of a compact homogeneous Riemannian manifold (M, g) , the reverse inequality

$$\mathfrak{S}_{(\mu^D)}(p) \geq \frac{|\Omega|_g}{|M|_g} \mathfrak{S}_{(\lambda)} \left(\frac{|M|_g}{|\Omega|_g} p \right)$$

holds. Contrary to what was found for Neumann eigenvalues in Theorem 5.2, a straightforward extension of the latter inequality to Dirichlet eigenvalues of a Laplacian with potential cannot hold in general. Indeed, such an extension would imply for $p = 1$ that $\mu_0^D(\Delta_g + V) \geq \frac{1}{|M|_g} \int_{\Omega} V dv_g$, which is not always true (for example, if Ω is a spherical cap of radius r and if u is a positive first eigenfunction of the Dirichlet Laplacian on Ω , we can take the family of continuous potentials V_{ε} with $V_{\varepsilon} = \frac{1}{u^2}$ on the spherical cap of radius $(1 - \varepsilon)r$, and V_{ε} is constant on the complement, then, using u as a test function, it is easy to see that $\mu_0^D(\Delta_g + V_{\varepsilon}) \leq \mu_0^D(\Delta_g) + |\Omega|$ while $\int_{\Omega} V_{\varepsilon} dv_g$ tends to infinity as $\varepsilon \rightarrow 0$.)

2. Assume that $|\Omega| \geq \frac{2}{3}|M|$, then an immediate consequence of Theorem 5.2 and the fact that the first positive eigenvalue λ_1 of the Laplacian on a homogeneous manifold (M, g) has multiplicity at least 2, is the following inequality

$$\mu_0 + \mu_1 \leq \frac{|\Omega|_g}{|M|_g} \left(2\lambda_1 \int_{\Omega} w dv_g + 3 \int_{\Omega} \tilde{V} w dv_g \right)$$

which yields for the Neumann Laplacian (with $\rho = V = 0$ and $w = 1$)

$$\mu_1 \leq 2 \frac{|\Omega|_g}{|M|_g} \lambda_1.$$

In the case where Ω is equal to the whole of M , Theorem 5.2 leads to the following

Corollary 5.1. Let (M, g) be a compact homogeneous Riemannian manifold. Let μ_l , $l \in \mathbb{N}$, be the eigenvalues defined by (2) on M . Then, for every $k \in \mathbb{N}^*$,

$$\sum_{j=0}^{k-1} \mu_j \leq \sum_{j=0}^{k-1} \tilde{\lambda}_j, \quad (75)$$

where equality holds if and only if

$$E_0(\mu_k) \subset e^{\rho} F_0(\lambda_k) \quad \text{and} \quad e^{\rho} F(\lambda_{k-1}) \subset E(\mu_k).$$

In particular, if m_1 is the multiplicity of λ_1 , then equality holds in (75) for $k \leq m_1$ if and only if $(V + |\nabla^g \rho|^2)w - \operatorname{div}_g(w \nabla \rho)$ is constant on M and $\mu_j = \tilde{\lambda}_j$ for $j = 0, 1, \dots, k-1$.

Proof of Corollary 5.1. Assume that equality holds in (75) for $k \leq m_1$. Then we have $E_0(\mu_k) \subset e^\rho F_0(\lambda_k)$. Since $k \leq m_1$, $F_0(\lambda_k) = F(\lambda_0) = \text{span}\{1\}$. It follows that $E_0(\mu_k)$ has dimension 1, that is $E_0(\mu_k) = E(\mu_0) = \text{span}\{e^\rho\}$. Consequently, $\mu_1 = \mu_2 = \dots = \mu_k$, and e^ρ is an eigenfunction of H associated with μ_0 . Thus

$$\begin{aligned} H e^\rho &= e^{2\rho} \text{div}_g (w e^{-2\rho} \nabla^g e^\rho) + V w e^\rho = \dots, \\ &= ((V + |\nabla^g \rho|^2) w - \text{div}_g(w \nabla \rho)) e^\rho = \mu_0 e^\rho \end{aligned}$$

which implies that $(V + |\nabla^g \rho|^2) w - \text{div}_g(w \nabla \rho) = \mu_0$. Integrating, we get $\mu_0 = \int_\Omega \tilde{V} w \, dv_g = \tilde{\lambda}_0$. Now,

$$\mu_0 + (k-1)\mu_1 = \sum_{j=0}^{k-1} \mu_j = \sum_{j=0}^{k-1} \tilde{\lambda}_j = \tilde{\lambda}_0 + (k-1)\tilde{\lambda}_1$$

and, consequently, $\mu_1 = \tilde{\lambda}_1$. □

Remarks 5.2. 1. An immediate consequence of Corollary 5.1 is that for any potential V on a compact homogeneous (M, g) , one has for every positive k ,

$$\frac{1}{k} \sum_{j=0}^{k-1} \mu_j (\Delta_g + V) \leq \frac{1}{k} \sum_{j=0}^{k-1} \mu_j (\Delta_g) + \int_M V \, dv_g,$$

to be compared with the results of [6].

2. We know that for any $k \geq 2$, either $\lambda_{k-1} = \lambda_k$ or else $\lambda_{k-1} = \lambda_{k-2}$. Notice that if $\lambda_{k-1} = \lambda_k$ and if equality holds in (75) for k , then, necessarily, $\tilde{\lambda}_{k-1} = \mu_{k-1} = \mu_k = \tilde{\lambda}_k$. (This follows directly from the combination of $\sum_{j=0}^{k-1} \mu_j = \sum_{j=0}^{k-1} \tilde{\lambda}_j$ with $\sum_{j=0}^k \mu_j \leq \sum_{j=0}^k \tilde{\lambda}_j$ and $\sum_{j=0}^{k-2} \mu_j \leq \sum_{j=0}^{k-2} \tilde{\lambda}_j$.) Consequently, the equality also holds in (75) for $k-1$ and $k+1$.

Moreover, if $\mu_k > \mu_{k-1}$, then the equality holds in (75) for k if and only if $\lambda_k > \lambda_{k-1}$ and $E(\mu_{k-1}) = e^\rho F(\lambda_{k-1})$. (Indeed, in this case, $\dim E_0(\mu_k) = \dim E(\mu_{k-1}) = k$ and $\dim F_0(\lambda_k) = \dim F_0(\lambda_{k-1}) = k$.)

Applying the Laplace transform to both sides of (66), we obtain the following comparison of the heat traces (see (32)):

Corollary 5.2. Let (M, g) be a compact homogeneous Riemannian manifold. Let $\mu_l, l \in \mathbb{N}$, be the eigenvalues defined by (2) on a bounded open set $\Omega \subset M$. Then, for all $t > 0$,

$$\sum_{j \geq 0} e^{-\mu_j t} \geq \frac{|\Omega|_g}{|M|_g} \sum_{j \geq 0} e^{-\tilde{\lambda}_j t}, \quad (76)$$

where $\tilde{\lambda}_j = \lambda_j \int_\Omega w \, dv_g + \int_\Omega \tilde{V} w \, dv_g$.

Let us define the theta function via:

$$\Theta(t) = \frac{1}{4\pi t} \sum_{(p,q) \in \mathbb{Z}^2} e^{-\frac{p^2 + q^2 + pq}{4t}}.$$

Corollary 5.3. Let $\Gamma = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \subset \mathbb{R}^2$ be a lattice, where $\{e_1, e_2\}$ is a basis of \mathbb{R}^2 . Let $\rho, w > 0, V$ be Γ -periodic functions on \mathbb{R}^2 and denote by $\mu_l = \mu_l(\rho, w, V), l \in \mathbb{N}$, the eigenvalues of the operator $H(\rho, w, V)$ defined by (4), acting on Γ -periodic functions on \mathbb{R}^2 . Then, for all $t > 0$,

$$\sum_{j \geq 0} e^{-\mu_j t} \geq \Theta \left(\frac{\int_\Omega w \, dx}{|\Omega|} t \right) e^{-t \int_\Omega \tilde{V} w \, dx}, \quad (77)$$

where Ω is a fundamental domain for the action of Γ on \mathbb{R}^2 .

Proof. This result is a direct consequence of (76) combined with Poisson's formula and Montgomery's Theorem [23]. \square

We turn now to the phase space analysis taking into account the form of the potential V , and allowing conformal transformations and nontrivial weights. Let us denote by

$$0 = \Lambda_0 < \Lambda_1 < \Lambda_2 < \dots < \Lambda_l < \dots$$

the increasing sequence of eigenvalues of the Laplacian of the compact homogeneous space (M, g) . The multiplicity of Λ_l is denoted m_l , and we designate by $\{y_{l,1}, y_{l,2}, \dots, y_{l,m_l}\}$ an L^2 -orthonormal basis of the eigenspace associated with Λ_l .

In the case of a domain Ω in a manifold $X \simeq (M, e^{-2\rho}g)$ that is conformally equivalent to (M, g) , we shall use coherent-state test functions of the form:

$$f_\zeta(\mathbf{x}) := y_{\ell m}(\mathbf{x})e^{\rho(\mathbf{x})}h_{\mathbf{y}}(\mathbf{x}). \quad (78)$$

In this formula, $h(\mathbf{x})$ is a nonnegative H^1 function supported in the geodesic ball of radius r in the canonical metric on M , with $\int_{B_r} h_r^2(\mathbf{x})d^\nu x = 1$, and \mathbf{y} ranges over the isometry group G . As before we choose it specifically as the ground-state Dirichlet eigenfunction on the geodesic ball of radius r and set $\mathcal{K}(h_r) := \int_{B_r} |\nabla h(\mathbf{x})|^2 d^\nu x$, which is thus the fundamental Dirichlet eigenvalue for the Laplacian on the geodesic disk of radius r . Denoting by $T_{\mathbf{y}}(\mathbf{x})$ the action by the group element \mathbf{y} on the point \mathbf{x} , we let

$$h_{\mathbf{y}}(\mathbf{x}) := h(T_{\mathbf{y}}(\mathbf{x})).$$

Recall that one can designate an arbitrary point of M as 0 and cover M with translates $T_{\mathbf{y}}(0)$. We normalize the uniform measure $d\gamma$ on G so that for any $f \in L^\infty(M)$, $\int_G f(T_{\mathbf{y}}(\mathbf{x}))d\gamma(\mathbf{y}) = \int_M f(\mathbf{x})dv_g$. The index $\zeta = (\ell, m, \mathbf{y})$ ranges over $\mathfrak{M} = \mathcal{J} \times G$, where \mathcal{J} is the set of all pairs of integer indices for the normalized eigenfunctions $y_{\ell m}(\mathbf{x})$, and the associated measure $d\sigma$ is the product of the counting measure on \mathcal{J} with $d\gamma$.

As in Section 2, we find it helpful to define:

Definition 5.1. *As before,*

$$\tilde{V}(\mathbf{x}) := V(\mathbf{x}) + |\nabla\rho|^2,$$

The weighted phase-space volume is

$$\begin{aligned} \Phi_w^h(\Lambda) &:= \left| \{(\ell, m, \mathbf{y}) : m \leq m_\ell, T_{\mathbf{y}}(0) \in \Omega, \Lambda_\ell + \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\} \right| \\ &= \int_{\{\mathbf{y}: T_{\mathbf{y}}(0) \in \Omega, \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}} \left(\sum_{\{\ell: \Lambda_\ell + \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}} m_\ell \right) d\gamma(\mathbf{y}). \end{aligned} \quad (79)$$

The total energy associated with this phase-space volume is correspondingly

$$E_w^h(\Lambda) := \int_{\{\mathbf{y}: T_{\mathbf{y}}(0) \in \Omega, \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}} \left(\sum_{\{\ell: \Lambda_\ell + \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}} m_\ell (\Lambda_\ell + \tilde{V}(T_{\mathbf{y}}(0))) \right) d\gamma(\mathbf{y}). \quad (80)$$

Theorem 5.3. *Let $\mu_0 \leq \mu_1 \leq \dots$ be the variationally defined Neumann eigenvalues (2) on a bounded open set $\Omega \subset M$, where w, ρ , and V satisfy the assumptions stated in Section 1. Then*

for all $r > 0$,

$$\sum_{j=0}^{k-1} \mu_j \leq E_w^h(\Lambda + \text{Lip}(\Lambda)r) + \mathcal{K}(h_r)\Phi_w^h(\Lambda + \text{Lip}(\Lambda)r). \quad (81)$$

Proof. Note that

$$\langle \phi, f_\zeta \rangle_\Omega = \left\langle e^{-\rho(\mathbf{x})} h_{\mathbf{y}}(\mathbf{x}) \phi(\mathbf{x}), y_{\ell m} \right\rangle_M,$$

where if Ω is a strict subset of M , then ϕ is extended by 0 outside Ω . Thus, by the Fourier completeness relation,

$$\begin{aligned} \int_G \left(\sum_{\ell, m} |\langle \phi, f_\zeta \rangle_\Omega|^2 \right) d\gamma(\mathbf{y}) &= \int_G \|e^{-\rho(\mathbf{x})} h_{\mathbf{y}}(\mathbf{x}) \phi(\mathbf{x})\|_{L^2(M, dv_g)}^2 d\gamma(\mathbf{y}) \\ &= \int_\Omega |\phi|^2 e^{-2\rho} \left(\int_G h_{\mathbf{y}}(\mathbf{x})^2 d\gamma(\mathbf{y}) \right) dv_g \\ &= \int_\Omega |\phi|^2 e^{-2\rho} dv_g = \|\phi\|^2. \end{aligned} \quad (82)$$

To apply the theorem, choose \mathfrak{M}_0 of the form $\{(\ell, m, \mathbf{y}) : m \leq m_\ell, T_{\mathbf{y}}(0) \in \Omega, \Lambda_\ell + \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}$ for a finite Λ large enough so that

$$k \leq \int_{\mathfrak{M}_0} \|f_\zeta\|_{L^2(\Omega)}^2 d\sigma = \int_\Omega \int_{\mathfrak{M}_0} h_{\mathbf{y}}^2(\mathbf{x}) e^{2\rho(\mathbf{x}) - 2\rho(\mathbf{x})} d\sigma dv_g = |\Omega| \Phi_w^h(\Lambda). \quad (83)$$

We define $\Lambda(k)$ as the minimal value of Λ for which (83) is valid and henceforth choose $\mathfrak{M}_0 = \{(\ell, m, \mathbf{y}) : m \leq m_\ell, T_{\mathbf{y}}(0) \in \Omega, \Lambda_\ell + \tilde{V}(\mathbf{y}) \leq \Lambda(k)\}$. Then

$$\begin{aligned} \sum_{j=0}^{k-1} \mu_j &\leq \int_{\mathfrak{M}_0} \mathcal{R}(f_\zeta) d\sigma(\zeta) \\ &= \int_\Omega \int_{\mathfrak{M}_0} w(\mathbf{x}) \left(y_{\ell m}^2 \left(h_{\mathbf{y}}^2(\mathbf{x}) \tilde{V}(\mathbf{x}) + |\nabla h_{\mathbf{y}}(\mathbf{x})|^2 + \nabla \rho(\mathbf{x}) \cdot \nabla h_{\mathbf{y}}^2(\mathbf{x}) \right) \right. \\ &\quad \left. + h_{\mathbf{y}}^2(\mathbf{x}) |\nabla y_{\ell m}|^2 + 2h_{\mathbf{y}}(\mathbf{x}) \nabla h_{\mathbf{y}}(\mathbf{x}) \cdot y_{\ell m} \nabla y_{\ell m} \right) d\sigma dv_g \\ &\leq \int_\Omega \int_{\{\mathbf{y} : \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}} w(\mathbf{x}) \left(\sum_{\{\ell : \Lambda_\ell + \tilde{V}(T_{\mathbf{y}}(0)) \leq \Lambda\}} \left(\frac{m_\ell}{|M|} \right) \left(h_{\mathbf{y}}^2(\mathbf{x}) (\Lambda_\ell + \tilde{V}(\mathbf{x})) + |\nabla h_{\mathbf{y}}(\mathbf{x})|^2 \right) \right) d\gamma(\mathbf{y}), \end{aligned}$$

by dint of (67) and (68). (The final cross term dropped out because it was proportional to the gradient of a constant function (67), in analogy with (64).) Because h is supported in a ball of radius r , we restrict the x -integration to $\mathbf{x} : \text{dist}(\mathbf{x}, \mathbf{y}) \leq r$ with $(\ell, m, \mathbf{y}) \in \mathfrak{M}_0$ and estimate the integral in analogy with (65), obtaining

$$\sum_{j=0}^{k-1} \mu_j \leq \left(\frac{1}{|M|} \right) \int_{(\ell, m, \mathbf{x}) \in \mathfrak{M}_0(\Lambda + \text{Lip}(\Lambda)r)} w(\mathbf{x}) \left(\Lambda_\ell + \tilde{V}(\mathbf{x}) + \mathcal{K}(h_r) \right) d\sigma, \quad (84)$$

which yields the statement in the Theorem. \square

Acknowledgments E.H. is grateful to the Université F. Rabelais and to École Polytechnique Fédérale de Lausanne for hospitality that supported this collaboration.

REFERENCES

- [1] S. Agmon and L. Hörmander. Asymptotic properties of solutions of differential equations with simple characteristics. *J. Analyse Math.*, 30:1–38, 1976.
- [2] S. Agmon. Lectures on elliptic boundary value problems. Prepared for publication by B. Frank Jones, Jr. with the assistance of George W. Batten, Jr. Van Nostrand Mathematical Studies, No. 2. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London, 1965.
- [3] E. F. Beckenbach and R. Bellman. Inequalities. Second revised printing. Ergebnisse der Mathematik und ihrer Grenzgebiete. Neue Folge, Band 30. Springer-Verlag, New York, Inc., 1965.
- [4] B. Colbois, A. El Soufi, and A. Savo. Eigenvalues of the Laplacian on a compact manifold with density. *Comm. Anal. Geom.*, 23(3):639–670, 2015.
- [5] D. E. Edmunds and W. D. Evans. Spectral theory and differential operators. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1987. Oxford Science Publications.
- [6] A. El Soufi and S. Ilias. Majoration de la seconde valeur propre d’un opérateur de Schrödinger sur une variété compacte et applications. *J. Funct. Anal.*, 103(2):294–316, 1992.
- [7] A. El Soufi and S. Ilias. Second eigenvalue of Schrödinger operators and mean curvature. *Comm. Math. Phys.*, 208(3):761–770, 2000.
- [8] S. Gallot. Inégalités isopérimétriques et analytiques sur les variétés riemanniennes. *Astérisque*, (163-164):5–6, 31–91, 281 (1989), 1988. On the geometry of differentiable manifolds (Rome, 1986).
- [9] A. Grigor’yan, Yu. Netrusov, and S. -T. Yau. Eigenvalues of elliptic operators and geometric applications. In Surveys in differential geometry. Vol. IX, Surv. Differ. Geom., IX, pages 147–217. Int. Press, Somerville, MA, 2004.
- [10] E. M. Harrell II and J. Stubbe. On sums of graph eigenvalues. *Linear Algebra Appl.*, 455:168–186, 2014.
- [11] A. Hassannezhad. Eigenvalues of perturbed Laplace operators on compact manifolds. *Pacific J. Math.*, 264(2):333–354, 2013.
- [12] R. Hempel, L. A. Seco, and B. Simon. The essential spectrum of Neumann Laplacians on some bounded singular domains. *J. Funct. Anal.*, 102(2):448–483, 1991.
- [13] C. E. Heil and D. F. Walnut. Continuous and discrete wavelet transforms. *Siam Review* 31(4):628–666, 1989.
- [14] L. Hörmander. The analysis of linear partial differential operators. I, volume 256 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1983. Distribution theory and Fourier analysis.
- [15] P. Kröger. Upper bounds for the Neumann eigenvalues on a bounded domain in Euclidean space. *J. Funct. Anal.*, 106(2):353–357, 1992.
- [16] A. Laptev. Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces *J. Funct. Anal.*, 151:531–545, 1992.
- [17] A. Laptev. Recent improvements of Berezin–Li & Yau type inequalities 2013 BIRS lecture available at <http://www.birs.ca/workshops/2013/13w5059/files/Laptev.pdf>
- [18] R. S. Laugesen and B. A. Siudeja. Sums of Laplace eigenvalues–rotationally symmetric maximizers in the plane *J. Funct. Anal.*, 260(6):1795–1823, 2011.
- [19] L. Li and L. Tang. Some upper bounds for sums of eigenvalues of the Neumann Laplacian. *Proc. Amer. Math. Soc.*, 134(11):3301–3307 (electronic), 2006.
- [20] E. H. Lieb. Coherent States as a Tool for Obtaining Rigorous Bounds, In: D. H. Feng and J. Klauder, eds., Coherent states: Past, present, and future. Proceedings, Oak Ridge, Singapore: World Scientific, 1994, pp. 267–278. Reprinted as pp 377–388 in Inequalities, Selecta of Elliott H. Lieb, M. Loss and M. B. Ruskai, eds., Springer, 2002.
- [21] E. H. Lieb and M. Loss. Analysis, volume 14 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, second edition, 2001.
- [22] E. H. Lieb and J. P. Solovej. Quantum coherent operators: A generalization of coherent states, *Lett. Math. Phys.*, 22, 145–154, 1991, Reprinted as pp 367–376 in Inequalities, Selecta of Elliott H. Lieb, M. Loss and M. B. Ruskai, eds., Springer, 2002.
- [23] H. L. Montgomery. Minimal theta functions. *Glasgow Math. J.*, 30(1):75–85, 1988.
- [24] M. Reed and B. Simon. Methods of modern mathematical physics. IV. Analysis of operators. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1978.
- [25] R. S. Strichartz. Fourier asymptotics of fractal measures. *J. Funct. Anal.*, 89(1):154–187, 1990.
- [26] R. S. Strichartz. Estimates for sums of eigenvalues for domains in homogeneous spaces. *J. Funct. Anal.*, 137(1):152–190, 1996.

- [27] W. Thirring. Quantum mathematical physics. Springer-Verlag, Berlin, second edition, 2002. Atoms, molecules and large systems, Translated from the 1979 and 1980 German originals by Evans M. Harrell II.
- [28] J. Weidmann. Linear operators in Hilbert spaces, volume 68 of Graduate Texts in Mathematics. Springer-Verlag, New York-Berlin, 1980. Translated from the German by Joseph Szücs.

UNIVERSITÉ DE TOURS, LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UMR-CNRS 6083,
PARC DE GRANDMONT, 37200 TOURS, FRANCE.

E-mail address: `elsoufi@univ-tours.fr`

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA GA 30332-0160, USA.

E-mail address: `harrell@math.gatech.edu`

UNIVERSITÉ DE TOURS, LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UMR-CNRS 6083,
PARC DE GRANDMONT, 37200 TOURS, FRANCE.

E-mail address: `ilias@univ-tours.fr`

EPFL, MATH-GEOM, STATION 8, CH-1015 LAUSANNE, SWITZERLAND

E-mail address: `Joachim.Stubbe@epfl.ch`