



”The minimality of the map $x \mapsto |x|$ for weighted energy”.

Jean-Christophe Bourgain

► **To cite this version:**

Jean-Christophe Bourgain. ”The minimality of the map $x \mapsto |x|$ for weighted energy”.. Calculus of Variation and P.D.E's, 2006, 25, pp.469-489. <hal-00022138>

HAL Id: hal-00022138

<https://hal.archives-ouvertes.fr/hal-00022138>

Submitted on 3 Apr 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The minimality of the map $\frac{x}{\|x\|}$ for weighted energy.

Jean-Christophe Bourgoin

LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUES THÉORIQUES, UNIVERSITÉ DE TOURS, PARC GRANDMONT 37200 TOURS

MSC(2000) : 58E20; 53C43.

abstract

In this paper, we investigate the minimality of the map $\frac{x}{\|x\|}$ from the euclidean unit ball \mathbf{B}^n to its boundary \mathbb{S}^{n-1} for weighted energy functionals of the type $E_{p,f} = \int_{\mathbf{B}^n} f(r) \|\nabla u\|^p dx$, where f is a non-negative function. We prove that in each of the two following cases :

i) $p = 1$ and f is non-decreasing,

ii) p is an integer, $p \leq n - 1$ and $f = r^\alpha$ with $\alpha \geq 0$,

the map $\frac{x}{\|x\|}$ minimizes $E_{p,f}$ among the maps in $W^{1,p}(\mathbf{B}^n, \mathbb{S}^{n-1})$ which coincide with $\frac{x}{\|x\|}$ on $\partial\mathbf{B}^n$. We also study the case where $f(r) = r^\alpha$ with $-n+2 < \alpha < 0$ and prove that $\frac{x}{\|x\|}$ does not minimize $E_{p,f}$ for α close to $-n + 2$ and when $n \geq 6$, for α close to $4 - n$.

Keys Words : minimizing map, p -harmonic map, p -energy, weighted energy.

0.1 Introduction and statement of results

For $n \geq 3$, the map $u_0(x) = \frac{x}{\|x\|} : \mathbf{B}^n \longrightarrow \mathbb{S}^{n-1}$ from the unit ball \mathbf{B}^n of \mathbb{R}^n to its boundary \mathbb{S}^{n-1} plays a crucial role in the study of certain natural energy functionals. In particular, since the works of Hildebrandt, Kaul and Widman ([13]), this map is considered as a natural candidate to realize, for each real number $p \in [1, n)$ the minimum of the p -energy functional,

$$E_p(u) = \int_{\mathbf{B}^n} \|\nabla u\|^p dx$$

among the maps $u \in W^{1,p}(\mathbf{B}^n, \mathbb{S}^{n-1}) = \{u \in W^{1,p}(\mathbf{B}^n, \mathbb{R}^n; \|u\| = 1 \text{ a.e.}\}$ satisfying $u(x) = x$ on \mathbb{S}^{n-1} .

This question was first treated in the case $p = 2$. Indeed, the minimality of u_0 for E_2 was established by Jäger and Kaul ([16]) in dimension $n \geq 7$ and by Brezis, Coron and Lieb in dimension 3 ([2]). In [5], Coron and Gulliver proved the minimality of u_0 for E_p for any integer $p \in \{1, \dots, n-1\}$ and any dimension $n \geq 3$.

Lin ([17]) has introduced the use of the elegant null Lagrangian method (or calibration method) in this topic. Avellaneda and Lin showed the efficiency of this method in [1] where they give a simpler alternative proof to the Coron-Gulliver result. Note that several results concerning the minimizing properties of p -harmonic diffeomorphisms were also obtained in this way in particular by Coron, Helein and El Soufi, Sandier ([4], [12], [7] and [6]).

The case of non-integer p seemed to be rather difficult. It is only ten years after the Coron-Gulliver article [5], that Hardt, Lin and Wang ([10]) succeeded to prove that, for all $n \geq 3$, the map u_0 minimizes E_p for $p \in [n-1, n)$. Their proof is based on a deep studies of singularities of harmonic and minimizing maps made in the last two decades. In dimension $n \geq 7$, Wang ([20]) and Hong ([14]) have independently proved the minimality of u_0 for any $p \geq 2$ satisfying $p + 2\sqrt{p} \leq n - 2$.

In [15], Hong remarked that the minimality of the p -energy E_p , $p \in (2, n-1]$, is related to the minimization of the following weighted 2-energy :

$$\tilde{E}_p(u) = \int_{\mathbf{B}^n} r^{2-p} \|\nabla u\|^2 dx$$

where $r = \|x\|$. Indeed, using Hölder inequality, it is easy to see that if the map u_0 minimizes \tilde{E}_p , then it also minimizes E_p (see [15], p.465). Unfortunately, as we will see in Corollary 1.1 below, for many values of $p \in (2, n)$, the map u_0 is not a minimizer of \tilde{E}_p . Therefore, Theorem 6 of ([15]), asserting that u_0 minimizes \tilde{E}_p seems to be not correct and the question of whether u_0 is a minimizing map of the p -energy E_p for non-integer $p \in (2, n-1)$ is still open ¹

The aim of this paper is to study the minimizing properties of the map

¹We suspect a problem in Theorem 6 p.464 of [15]. Indeed the author claims that the quantity $G_{\varphi_1^0, \dots, \varphi_{n-1}^0}(v, p)$, which represents a weighted energy of the map v on the 3-dimensional cone \mathcal{C}_0 in \mathbf{B}^n , is uniformly proportional to the weighted energy on the euclidian ball \mathbf{B}^3 . There is no reason for this fact to be true, the orthogonal projection of \mathcal{C}_0 on to \mathbf{B}^n being not homothetic.

u_0 in regard to some weighted energy functionals of the form :

$$E_{p,f}(u) = \int_{\mathbf{B}^n} f(r) \|\nabla u\|^p dx,$$

where $p \in \{1, \dots, n-1\}$ and $f: [0, 1] \rightarrow \mathbb{R}$ is a non-negative non-decreasing continuous function. For $p = 1$, the map u_0 minimizes $E_{1,f}$ for a large class of weights. Indeed, we have the following

Theorem 0.1 *Suppose that f is a non-negative differentiable non-decreasing function. Then the map $u_0 = \frac{x}{\|x\|}$ is a minimizer of the energy $E_{1,f}$, that is, for any u in $W^{1,1}(\mathbf{B}^n, \mathbb{S}^{n-1})$ with $u(x) = x$ on \mathbb{S}^{n-1} , we have*

$$\int_{\mathbf{B}^n} f(r) \|\nabla u_0\| dx \leq \int_{\mathbf{B}^n} f(r) \|\nabla u\| dx,$$

Moreover, if f has no critical points in $(0, 1)$, then the map $u_0 = \frac{x}{\|x\|}$ is the unique minimizer of the energy $E_{1,f}$, that is, the equality in the last inequality holds if and only if $u = u_0$.

For $p \geq 2$, we restrict ourselves to power functions $f(r) = r^\alpha$,

Theorem 0.2 *For any $\alpha \geq 0$ and any integer $p \in \{1, \dots, n-1\}$, the map $u_0 = \frac{x}{\|x\|}$ is a minimizer of the energy E_{p,r^α} that is, for any u in $W^{1,p}(\mathbf{B}^n, \mathbb{S}^{n-1})$ with $u(x) = x$ on \mathbb{S}^{n-1} , we have,*

$$\int_{\mathbf{B}^n} r^\alpha \|\nabla u_0\|^p dx \leq \int_{\mathbf{B}^n} r^\alpha \|\nabla u\|^p dx.$$

Moreover, if $\alpha > 0$, then the map $u_0 = \frac{x}{\|x\|}$ is the unique minimizer of the energy E_{p,r^α} , that is the equality in the last inequality holds if and only if $u = u_0$.

The proof of these two theorems is given in section 2. It is based on a construction of an adapted null-Lagrangian. The case of $p = 1$ can be obtained passing through more direct ways and will be treated independently.

The case of weights of the form $f(r) = r^\alpha$, with $\alpha < 0$, is treated in section 3. The weighted energy $\int_{\mathbf{B}^n} r^\alpha \|\nabla u_0\|^2 dx$ of $u_0 = \frac{x}{\|x\|}$ is finite for $\alpha > -n + 2$. Hence we consider the family of maps,

$$u_a(x) = a + \lambda_a(x)(x - a), \quad a \in \mathbf{B}^n,$$

where $\lambda_a(x) \in \mathbb{R}$ is chosen such that $u_a(x) \in \mathbb{S}^{n-1}$ (that is $u_a(x)$ is the intersection point of \mathbb{S}^{n-1} with the half-line of origin a passing by x).

We study the energy $E_{2,r^\alpha}(u_a)$ of these maps and deduce the following theorem.

Theorem 0.3 Suppose that $n \geq 3$.

(i) For any $a \in \mathbf{B}^n, a \neq 0$, there exists a negative real number $\alpha_0 \in (-n + 2, 0)$, such that, for any $\alpha \in (-n + 2, \alpha_0]$ we have

$$\int_{\mathbf{B}^n} r^\alpha \|\nabla u_0\|^2 dx > \int_{\mathbf{B}^n} r^\alpha \|\nabla u_a\|^2 dx \quad .$$

(ii) For any integer $n \geq 6$, there exists $\alpha_0 \in (4 - n, 5 - n)$ such that, for any $\alpha \in (4 - n, \alpha_0)$, there exists $a \in \mathbf{B}^n$ such that,

$$\int_{\mathbf{B}^n} r^\alpha \|\nabla u_0\|^2 dx > \int_{\mathbf{B}^n} r^\alpha \|\nabla u_a\|^2 dx \quad .$$

Replacing in Theorem 0.3 α by $2 - p, p \in (2, n)$, we obtain the following corollary :

Corollary 0.1 For any $n \geq 6$, there exists $p_0 \in (n - 3, n - 2)$ such that, for any $p \in (p_0, n - 2)$ the map $u_0 = \frac{x}{\|x\|}$ does not minimize the functional $\int_{\mathbf{B}^n} r^{2-p} \|\nabla u\|^2 dx$ among the maps $u \in W^{1,2}(\mathbf{B}^n, \mathbb{S}^{n-1})$ satisfying $u(x) = x$ on \mathbb{S}^{n-1} .

acknowledgements. The author would express his grateful to Professor Ahmad El Soufi and Professor Etienne Sandier for their helpful advice.

0.2 Proof of theorems 0.1 and 0.2

Consider an integer $p \in \{1, \dots, n-1\}$ and f a differentiable, non-negative, increasing, and non-identically zero map. We can suppose without loss of generality, that $f(1) = 1$.

For any subset $I = \{i_1, \dots, i_p\} \subset \{1, \dots, n-1\}$ with $i_1 < i_2 \dots < i_p$ and for any map,

$$u = (u_1, \dots, u_n) : \mathbf{B}^n \longrightarrow \mathbb{S}^{n-1} \quad \text{in } \mathcal{C}^\infty(\mathbf{B}^n, \mathbb{S}^{n-1}) \quad \text{with } u(x) = x \text{ on } \mathbb{S}^{n-1},$$

we consider the n-form :

$$\omega_I(u) = dx_1 \wedge \dots \wedge d(f(r)u_{i_1}) \wedge \dots \wedge d(f(r)u_{i_k}) \wedge \dots \wedge dx_n$$

Lemma 0.1 We have the identity :

$$\int_{\mathbf{B}^n} \omega_I(u) = \int_{\mathbf{B}^n} \omega_I(Id) \quad \forall x \in \mathbf{B}^n \quad \text{where } Id(x) = x.$$

Proof By Stokes theorem, we have :

$$\begin{aligned}
\int_{\mathbf{B}^n} \omega_I(u) &= \int_{\mathbf{B}^n} dx_1 \wedge \cdots \wedge d(f(r)u_{i_1}) \wedge \cdots \wedge d(f(r)u_{i_p}) \wedge \cdots \wedge dx_n \\
&= \int_{\mathbf{B}^n} (-1)^{i_1-1} d\left(f(r)u_{i_1} dx_1 \wedge \cdots \wedge d(\widehat{f(r)u_{i_1}}) \wedge \right. \\
&\quad \left. \cdots \wedge d(f(r)u_{i_p}) \wedge \cdots \wedge dx_n\right) \\
&= \int_{\mathbb{S}^{n-1}} (-1)^{i_1-1} x_{i_1} dx_1 \wedge \cdots \wedge d(\widehat{f(r)u_{i_1}}) \wedge \\
&\quad \cdots \wedge d(f(r)u_{i_p}) \wedge \cdots \wedge dx_n.
\end{aligned}$$

Indeed, on \mathbb{S}^{n-1} , we have $f(r)u_{i_1} = x_{i_1}$ ($r = 1, f(1) = 1$ and $u(x) = x$). Iterating, we get the designed identities. Consider the n-form :

$$S(u) = \sum_{|I|=p} w_I(u)$$

By Lemma 0.1, we have :

$$\int_{\mathbf{B}^n} S(u) = \sum_{|I|=p} \int_{\mathbf{B}^n} w_I(u) = \sum_{|I|=p} \int_{\mathbf{B}^n} dx = C_n^p \frac{|\mathbb{S}^{n-1}|}{n},$$

where $|\mathbb{S}^{n-1}|$ is the Lebesgue measure of the sphere.

Lemma 0.2 *The n-form $S(u)$ is $O(n)$ -equivariant, that is, for any rotation R in $O(n)$, we have :*

$$S({}^tRuR)({}^tRx) = S(u)(x) \quad \forall x \in \mathbf{B}^n.$$

Proof Consider $S(u)(x)(e_1, \dots, e_n)$ where (e_1, \dots, e_n) is the standart basis of \mathbb{R}^n and notice that it is equal to $(-1)^n$ times the $(p+1)^{th}$ coefficient of the polynomial $P(\lambda) = \det(\text{Jac}(fu)(x) - \lambda Id)$ which does not change when we replace fu by tRfuR .

For any $x \in \mathbf{B}^n$, let $R \in O(n)$ be such that ${}^tRu(x) = e_n = (0, \dots, 0, 1)$. Consider $y = {}^tRx$, $v = {}^tRuR$, so that :

$$v(y) = e_n, \quad d({}^tRuR)(y)(\mathbb{R}^n) \subset e_n^\perp \quad \text{that is} \quad \frac{\partial v_n}{\partial x_j}(y) = 0 \quad \forall j \in \{1, \dots, n\}.$$

Lemma 0.3 *Let a_1, \dots, a_n be n non-negative numbers, and $p \in \{1, \dots, n-1\}$. Then :*

$$\sum_{i_1 < \dots < i_p} a_{i_1} \cdots a_{i_p} \leq \frac{1}{(n-1)^p} C_{n-1}^p \left(\sum_{j=1}^{n-1} a_j \right)^p.$$

Proof See for instance Hardy coll.[4], theorem 52.

Let $I = \{i_1, \dots, i_p\} \subset \{1, \dots, n\}$. We have :
if $i_p \neq n$,

$$\begin{aligned} \omega_I(v)(y) &= (dx_1 \wedge \cdots \wedge d(f(r)v_{i_1}) \wedge \cdots \wedge d(f(r)v_{i_k}) \wedge \cdots \wedge dx_n)(y) \\ &= |f(r)|^p (dx_1 \wedge \cdots \wedge dv_{i_1} \wedge \cdots \wedge dv_{i_k} \wedge \cdots \wedge dx_n)(y). \end{aligned}$$

Indeed, $\forall j \leq n-1$, $d(f(r)v_j(y)) = d(f(r))v_j(y) + f(r)dv_j(y) = f(r)dv_j(y)$
since $v(y) = e_n$.

If $i_p = n$,

$$\omega_I(v)(y) = |f(r)|^{p-1} (dx_1 \wedge \cdots \wedge dv_{i_1} \wedge \cdots \wedge df)(y).$$

Indeed, $d(f(r)v_n)(y) = df(y)v_n(y) + f(r)dv_n(y) = df(y)$ (as $dv(y) \subset e_n^\perp$). The Hadamard inequality gives :

$$\begin{aligned} |S(v)(y)| &= \left| \sum_{|I|=p} \omega_I(v)(y) \right| \leq |f(r)|^p \sum_{1 \leq i_1 < i_2 < \dots < i_p \leq n-1} \|dx_1\| \cdots \|dv_{i_1}\| \\ &\quad \cdots \|dv_{i_p}\| \cdots \|dx_n\|(y) \\ &+ |f(r)|^{p-1} \sum_{1 \leq i_1 < i_2 < \dots < i_{p-1} \leq n-1} \|dx_1\| \cdots \|dv_{i_1}\| \\ &\quad \cdots \|dv_{i_p}\| \cdots \|df\|(y) \\ &\leq |f(r)|^p \left(\sum_{1 \leq i_1 < i_2 < \dots < i_p \leq n-1} \|dx_1\|^2 \cdots \|dv_{i_1}\|^2 \cdots \right. \\ &\quad \left. \cdots \|dv_{i_p}\|^2 \cdots \|dx_n\|^2(y) \right)^{\frac{1}{2}} (C_n^p)^{\frac{1}{2}} \\ &+ f'(r)f(r)^{p-1} \sum_{1 \leq i_1 < i_2 < \dots < i_{p-1} \leq n-1} \|dx_1\| \cdots \|dv_{i_1}\| \\ &\quad \cdots \|dv_{i_p}\|(y). \end{aligned}$$

The Hardy inequality gives, after integration and using the fact that $\|\nabla u\| = \|\nabla v\|$,

$$\begin{aligned} \frac{C_n^p}{n} |\mathbb{S}^{n-1}| &\leq \frac{C_{n-1}^p}{(n-1)^{p/2}} \int_{\mathbf{B}^n} f^p(r) \|\nabla u\|^p dx \\ &\quad + \frac{C_{n-1}^{p-1}}{(n-1)^{\frac{p-1}{2}}} \int_{\mathbf{B}^n} f'(r) f^{p-1}(r) \|\nabla u\|^{p-1} dx. \quad (1) \end{aligned}$$

Remark : If f' is positive and if equality holds in (1), then, $\forall i \leq n-1$, $y_i = 0$ and $y_n = \pm \frac{x}{\|x\|}$, which implies that $u(x) = \pm \frac{x}{\|x\|}$.

Proof of the Theorem 1.1 Inequality (1) give

$$|\mathbb{S}^{n-1}| \leq \sqrt{n-1} \int_{\mathbf{B}^n} f(r) \|\nabla u\| dx + \int_{\mathbf{B}^n} f'(r) dx.$$

Hence :

$$\begin{aligned} \int_{\mathbf{B}^n} f \|\nabla u\| dx &\geq \frac{|\mathbb{S}^{n-1}|}{\sqrt{n-1}} \left(1 - \int_0^1 f'(r) r^{n-1} dr\right) \\ \int_{\mathbf{B}^n} f \|\nabla u\| dx &\geq \sqrt{n-1} |\mathbb{S}^{n-1}| \int_0^1 f(r) r^{n-2} dr = \int_{\mathbf{B}^n} f(r) \|\nabla u_0\| dx. \end{aligned}$$

To see the uniqueness it suffices to refer to the remark above. It gives that for any $x \in \mathbf{B}^n$, $u(x) = \frac{x}{\|x\|}$ or $u(x) = -\frac{x}{\|x\|}$. As $u(x) = x$ on the unit sphere, we have, for any $x \in \mathbf{B}^n \setminus \{0\}$, $u(x) = \frac{x}{\|x\|}$. \blacksquare

Proof of the Theorem 1.2. Let α be a positive real number. From inequality (1) we have :

$$\frac{C_n^p}{n} |\mathbb{S}^{n-1}| \leq \frac{C_{n-1}^p}{(n-1)^{p/2}} \int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u\|^p dx + \alpha \frac{C_{n-1}^{p-1}}{(n-1)^{\frac{p-1}{2}}} \int_{\mathbf{B}^n} r^{\alpha p-1} \|\nabla u\|^{p-1} dx.$$

By Hölder inequality, we have, setting $q = \frac{p}{p-1}$:

$$\begin{aligned} \frac{C_n^p}{n} |\mathbb{S}^{n-1}| &\leq \frac{C_{n-1}^p}{(n-1)^{p/2}} \int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u\|^p dx \\ &\quad + \alpha \frac{C_{n-1}^{p-1}}{(n-1)^{\frac{p-1}{2}}} \left(\int_{\mathbf{B}^n} r^{p(\alpha-1)} dx \right)^{1/p} \left(\int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u\|^p dx \right)^{1/q} \\ &\leq \frac{C_{n-1}^p}{(n-1)^{p/2}} \int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u\|^p dx \\ &\quad + \alpha \frac{C_{n-1}^{p-1}}{(n-1)^{\frac{p-1}{2}}} \frac{|\mathbb{S}^{n-1}|^{1/p}}{(n+p(\alpha-1))^{1/p}} \left(\int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u\|^p dx \right)^{1/q} \end{aligned}$$

Consider the polynomial function :

$$P(t) = \frac{C_{n-1}^p}{(n-1)^{p/2}} t^q + \alpha \frac{C_{n-1}^{p-1}}{(n-1)^{\frac{p-1}{2}}} \frac{|\mathbb{S}^{n-1}|^{1/p}}{(n+p(\alpha-1))^{1/p}} t - \frac{C_n^p}{n} |\mathbb{S}^{n-1}|.$$

Setting $A = (\int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u\|^p)^{1/q}$ and $B = (\int_{\mathbf{B}^n} r^{\alpha p} \|\nabla u_0\|^p)^{1/q}$, we get $P(A) \geq 0$ while

$$\begin{aligned} P(B) &= \frac{C_{n-1}^{p-1}}{n+p(\alpha-1)} |\mathbb{S}^{n-1}| + \alpha \frac{C_{n-1}^{p-1}}{n+p(\alpha-1)} |\mathbb{S}^{n-1}| - \frac{C_n^p}{n} |\mathbb{S}^{n-1}| \\ &= \frac{C_{n-1}^{p-1}}{n+p(\alpha-1)} |\mathbb{S}^{n-1}| \left(\frac{n-p}{n} + \alpha - \frac{C_n^p}{nC_{n-1}^{p-1}} (n+p(\alpha-1)) \right) \\ &= 0. \end{aligned}$$

On the other hand, $\forall t \geq 0$, $P'(t) > 0$. Hence, P is increasing in $[0, +\infty)$ and is equal to zero only for B . Necessarily, we have $A \geq B$.

Moreover, if $\alpha > 0$, $A = B$ implies that equality in the inequality (1) holds. Referring to the remark above, and as $u_0(x) = x$ on the sphere, we have $u = u_0 = \frac{x}{\|x\|}$. Replacing α by α/p we finish the prove of the theorem. \blacksquare

0.3 The energy of a natural family of maps.

Let $a = (\theta, \dots, 0)$ be a point of \mathbf{B}^n with $0 < \theta < 1$ and consider the map,

$$u_a(x) = a + \lambda_a(x)(x - a),$$

where $\lambda_a(x) > 0$ is chosen so that $u_a(x) \in \mathbb{S}^{n-1}$ for any $x \in \mathbf{B}^n \setminus \{0\}$,

$$\lambda_a(x) = \frac{\sqrt{\Delta_a(x)} - (a|x-a|)}{\|x-a\|^2}$$

and

$$\Delta_a(x) = (1 - \|a\|^2)\|x-a\|^2 + (a|x-a|)^2.$$

Notice that $u_a(x) = x$ as soon as x is on the sphere. If we denote by $\{e_i\}_{i \in \{1, \dots, n\}}$ the standard basis of \mathbb{R}^n , then, $\forall i \leq n$, we have,

$$\begin{aligned}
\|du_a(x).e_i\|^2 &= \left(\frac{\sqrt{\Delta_a} - (a|x-a)}{\|x-a\|^2} \right)^2 \\
&+ \left[-2 \frac{(x-a|e_i)}{\|x-a\|^4} \left(\sqrt{\Delta_a} - (a|x-a) \right) \right. \\
&+ \frac{(1-\|a\|^2)(x-a|e_i) + (x-a|a)(a|e_i)}{\sqrt{\Delta_a}\|x-a\|^2} \\
&\left. - \frac{(a|e_i)}{\|x-a\|^2} \right]^2 \|x-a\|^2 \\
&+ 2 \left(\frac{\sqrt{\Delta_a} - (a|x-a)}{\|x-a\|^2} \right) \left(-2 \frac{(x-a|e_i)}{\|x-a\|^4} \left(\sqrt{\Delta_a} - (a|x-a) \right) \right) \\
&+ \frac{(1-\|a\|^2)(x-a|e_i) + (x-a|a)(a|e_i)}{\sqrt{\Delta_a}\|x-a\|^2} \\
&\left. - \frac{(a|e_i)}{\|x-a\|^2} \right) (x-a|e_i).
\end{aligned}$$

Let us prove that, for each $\alpha \in (-n, 0)$, $\int_{\mathbf{B}^n} r^\alpha \|\nabla u_a\| dx$ is finite. Consider the map :

$$\begin{aligned}
F : \mathbb{R}^+ \times \mathbb{S}^{n-1} &\longrightarrow \mathbb{R}^n \\
(r, s) &\longmapsto a + rs = x.
\end{aligned}$$

Then, we have,

$$F^*(\|\nabla u_a\|^2 dx) = \frac{1}{r^2} \sum_{i=1}^n H_{i,a}(s) r^{n-1} dr \wedge ds,$$

where $H_{i,a}(s)$ is given on the sphere by,

$$\begin{aligned}
H_{i,a}(s) &= ((1 - \|a\|^2 + (a|s)^2)^{1/2} - (a|s))^2 \\
&+ \left[-2(s|e_i) ((1 - \|a\|^2 + (a|s)^2)^{1/2} - (s|a)) \right. \\
&+ \left. \frac{(1 - \|a\|^2)(s|e_i) + (a|e_i)(s|a)}{(1 - \|a\|^2 + (a|s)^2)^{1/2}} - (a|e_i) \right]^2 \\
&+ 2 \left((1 - \|a\|^2 + (a|s)^2)^{1/2} - (a|s) \right) \\
&\left(-2(s|e_i) ((1 - \|a\|^2 + (a|s)^2)^{1/2} - (s|a)) \right. \\
&+ \left. \frac{(1 - \|a\|^2)(s|e_i) + (a|e_i)(s|a)}{(1 - \|a\|^2 + (a|s)^2)^{1/2}} - (a|e_i) \right) (s|e_i).
\end{aligned}$$

It is clear that $H_{i,a}(s)$ is continuous on \mathbb{S}^{n-1} . Therefore, near the point a , as $n \geq 3$, the map $\|x\|^\alpha \|\nabla u_a\|$ is integrable. Furthermore, near the point 0 , as $\alpha > -n$, this map is also integrable. In conclusion, for any $\alpha \in (-n, 0)$, the energy $E_{r^\alpha, 2}(u_a)$ is finite.

Proof of Theorem 1.3(i). Since we have

$$E_{2,r^\alpha}(u_0) = \int_{\mathbf{B}^n} \|x\|^\alpha \|\nabla u_0\|^2 dx = \frac{|\mathbb{S}^{n-1}|(n-1)}{n+\alpha-2},$$

the energy $E_{2,r^\alpha}(u_0)$ goes to infinity as $\alpha \rightarrow -n+2$. On the other hand, as the energy $E_{2,r^\alpha}(u_a)$ is continuous in α , there exists a real number $\alpha_0 \in (-n+2, 0)$ such that, $\forall \alpha, 2-n < \alpha \leq \alpha_0$,

$$\int_{\mathbf{B}^n} \|x\|^\alpha \|\nabla u_0\|^2 dx > \int_{\mathbf{B}^n} \|x\|^\alpha \|\nabla u_a\|^2 dx.$$

Proof of Theorem 1.3(ii). Since $a = (\theta, 0, \dots, 0)$, we will study the function,

$$G(\theta) = E_{2,r^\alpha}(u_a) = \int_{\mathbf{B}^n} r^\alpha \|\nabla u_a\|^2 dx.$$

Precisely, we will show that for any $\alpha \in (5-n, 4-n)$, G is two times differentiable at $\theta = 0$ with $\frac{dG}{d\theta}(0) = 0$ and, when α is sufficiently close to $4-n$, $\frac{d^2G}{d\theta^2}(0) < 0$. Assertion (ii) of Theorem 1.3 then follows immediately.

We have,

$$\begin{aligned}
H_{i,a}(s) = H_{i,\theta}(s) &= \left(\sqrt{1-\theta^2+\theta^2s_1^2} - \theta s_1 \right)^2 \\
&+ \left(-2s_i(\sqrt{1-\theta^2+\theta^2s_1^2} - \theta s_1) + \frac{(1-\theta^2)s_i + \delta_{i1}\theta^2s_1}{\sqrt{1-\theta^2+\theta^2s_1^2}} - \delta_{i1}\theta \right)^2 \\
&+ 2 \left(\sqrt{1-\theta^2+\theta^2s_1^2} - \theta s_1 \right) \left(-2s_i(\sqrt{1-\theta^2+\theta^2s_1^2} - \theta s_1) \right. \\
&\quad \left. + \frac{(1-\theta^2)s_i + \delta_{i1}\theta^2s_1}{\sqrt{1-\theta^2+\theta^2s_1^2}} - \delta_{i1}\theta \right) s_i,
\end{aligned}$$

where $\delta_{ij} = 0$ if $i \neq j$ and 1 else.

We notice that $H_{i,\theta}(s)$ is bounded on $[0, 1] \times \mathbb{S}^{n-1}$. Indeed, for all $x, y, z \in [0, 1]$, excepting $(x, y) = (0, 1)$, we have,

$$\left| \frac{x}{\sqrt{1-y^2+y^2x^2}} \right| \leq 1 \quad \text{and} \quad \left| \frac{(1-y^2)z}{\sqrt{1-y^2+y^2x^2}} \right| \leq 1.$$

Then, for almost all $(s, \theta) \in \mathbb{S}^{n-1} \times [0, 1]$, we have,

$$\left| \frac{(1-\theta^2)s_i + \delta_{i1}\theta^2s_1}{\sqrt{1-\theta^2+\theta^2s_1^2}} \right| \leq 1,$$

and the others terms are continuous in $[0, 1] \times \mathbb{S}^{n-1}$.

We have,

$$\begin{aligned}
E_{2,r^\alpha}(u_a) &= \int_{\mathbf{B}^n} \|x\|^\alpha \|\nabla u_a\|^2 dx = \int_{\mathbf{B}^n} \|a + rs\|^\alpha r^{n-3} H(\theta, s) dr ds \\
&= \int_{\mathbb{S}^{n-1}} H(\theta, s) \left(\int_0^{\gamma_\theta(s)} ((r + \theta s_1)^2 + \theta^2(1 - s_1^2))^{\alpha/2} r^{n-3} dr \right) ds,
\end{aligned}$$

where $\gamma_\theta(s) = \sqrt{1-\theta^2+\theta^2s_1^2} - \theta s_1$ and $H(\theta, s) = \sum_{i=1}^n H_{i,\theta}(s)$. We notice that

$H(\theta, s)$ is indefinitely differentiable in $(-1/2, 1/2) \times \mathbb{S}^{n-1}$. Let C_n be a positive real number so that, $\forall(\theta, s) \in (-1/2, 1/2) \times \mathbb{S}^{n-1}$

$$|H(\theta, s)| \leq C_n, \quad \left| \frac{\partial H(\theta, s)}{\partial \theta} \right| \leq C_n, \quad \left| \frac{\partial^2 H(\theta, s)}{\partial \theta^2} \right| \leq C_n.$$

Furthermore, we have,

$$H(\theta, s) = (n-1) - 2(n-1)s_1\theta + ((2n-3)s_1^2 - n + 2)\theta^2 + o(\theta^2). \quad (A)$$

Let us set $\rho = r + \theta s_1$, $\beta(\theta, s) = \sqrt{1 - \theta^2 + \theta^2 s_1^2}$ and

$$F(\theta, s) = \int_{\theta s_1}^{\beta(\theta, s)} (\rho - \theta s_1)^{n-3} (\rho^2 + \theta^2(1 - s_1^2))^{\alpha/2} d\rho.$$

Notice that $\rho \in [-1, 3]$. Then, $G(\theta) = \int_{\mathbb{S}^{n-1}} H(\theta, s) F(\theta, s) ds$. Let us set $g(\rho, \theta, s) = (\rho - \theta s_1)^{n-3} (\rho^2 + \theta^2(1 - s_1^2))^{\alpha/2}$.

Lemma 0.4 *The map $\theta \mapsto G(\theta)$ is continuous on $(-1/2, 1/2)$ and continuously differentiable on $(-1/2, 1/2) \setminus \{0\}$ for any $\alpha > 3 - n$.*

Proof We have, $\forall s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$\frac{(\rho - \theta s_1)^2}{(\rho^2 + \theta^2(1 - s_1^2))} \leq \frac{2}{1 - s_1^2} \quad (1.1)$$

Indeed, $(1 - s_1^2)(\rho - \theta s_1)^2 \leq 2(1 - s_1^2)(\rho^2 + \theta^2) \leq 2(\rho^2 + \theta^2(1 - s_1^2))$. And then,

$$g(\rho, \theta, s) \leq \frac{2^{\frac{n-3}{2}}}{(1 - s_1^2)^{\frac{n-3}{2}}} (\rho^2 + \theta^2(1 - s_1^2))^{\frac{\alpha+n-3}{2}}. \quad (1.2)$$

Since $\alpha > 3 - n$ we deduce that the map $(\rho, \theta) \rightarrow g(\rho, \theta, s)$ is continuous on $(-1/2, 1/2) \times [-1, 3]$. Hence, the map $z \mapsto \int_0^z g(\rho, \theta, s) d\rho$ is differentiable on $[-1, 3]$ and,

$$\frac{\partial}{\partial z} \int_0^z g(\rho, \theta, s) d\rho = g(z, \theta, s).$$

Furthermore, for any $\rho \in [-1, 3]$, the map $\theta \mapsto g(\rho, \theta, s)$ is differentiable and

$$\begin{aligned} \frac{\partial g}{\partial \theta}(\rho, \theta, s) &= -(n-3)s_1(\rho - \theta s_1)^{n-4} (\rho^2 + \theta^2(1 - s_1^2))^{\frac{\alpha}{2}} \\ &+ \frac{\alpha}{2} (\rho - \theta s_1)^{n-3} 2\theta(1 - s_1^2) (\rho^2 + \theta^2(1 - s_1^2))^{\frac{\alpha}{2}-1}. \end{aligned}$$

Let a, b be two real in $(0, 1/2)$ with $a < b$. We have for any $|\theta| \in (a, b)$, for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$\begin{aligned} \left| \frac{\partial g}{\partial \theta}(\rho, \theta, s) \right| &\leq (n-3)4^{n-4} (a^2(1 - s_1^2))^{\frac{\alpha}{2}} \\ &+ |\alpha|4^{n-3} (1 - s_1^2) (a^2(1 - s_1^2))^{\frac{\alpha}{2}-1}. \quad (1.3) \end{aligned}$$

This shows that $\theta \mapsto \int_0^z g(\rho, \theta, s) d\rho$ is differentiable on $(-1/2, 1/2) \setminus \{0\}$ and

$$\frac{\partial}{\partial \theta} \int_0^z g(\rho, \theta, s) d\rho = \int_0^z \frac{\partial g}{\partial \theta}(\rho, \theta, s) d\rho.$$

Moreover the map $(z, \theta) \mapsto \int_0^z \frac{\partial g}{\partial \theta}(\rho, \theta, s) d\rho$ is continuous in $[-1, 3] \times (-1/2, 1/2) \setminus \{0\}$. Indeed, $\theta \mapsto \frac{\partial g}{\partial \theta}(\rho, \theta, s)$ is clearly continuous on $(-1/2, 1/2) \setminus \{0\}$ and from (1.3) and by Lebesgue Theorem, $\theta \mapsto \int_0^z \frac{\partial g}{\partial \theta}(\rho, \theta, s) d\rho$ is continuous on $(-1/2, 1/2) \setminus \{0\}$. Then, for any $\epsilon > 0$, we will have for any sufficiently small h, k ,

$$\begin{aligned} \left| \int_0^{z+h} \frac{\partial g}{\partial \theta}(\rho, \theta + k, s) d\rho - \int_0^z \frac{\partial g}{\partial \theta}(\rho, \theta, s) d\rho \right| &\leq \left| \int_0^z \frac{\partial g}{\partial \theta}(\rho, \theta + k, s) d\rho \right. \\ &\quad \left. - \int_0^z \frac{\partial g}{\partial \theta}(\rho, \theta, s) d\rho \right| \\ &\quad + \left| \int_z^{z+h} \frac{\partial g}{\partial \theta}(\rho, \theta + k, s) d\rho \right| \\ &\leq \epsilon. \end{aligned}$$

The map $(z, \theta) \mapsto \int_0^z g(\rho, \theta, s) d\rho$ is differentiable on $[-1, 3] \times (-1/2, 1/2) \setminus \{0\}$ and the map $\theta \mapsto F(\theta, s)$ is differentiable in $(-1/2, 1/2) \setminus \{0\}$ and for any $\theta \in (-1/2, 1/2) \setminus \{0\}$,

$$\begin{aligned} \frac{\partial F}{\partial \theta}(\theta, s) &= \frac{\partial \beta}{\partial \theta}(\theta, s) g(\beta(\theta, s), \theta, s) - s_1 g(\theta s_1, \theta, s) + \int_{\theta s_1}^{\beta(\theta, s)} \frac{\partial g}{\partial \theta}(\rho, \theta, s) d\rho \\ &= \frac{\theta(s_1^2 - 1)}{(1 - \theta^2 + \theta^2 s_1^2)^{1/2}} ((1 - \theta^2 + \theta^2 s_1^2)^{1/2} - \theta s_1)^{n-3} \\ &\quad + \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho + \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho, \end{aligned}$$

where,

$$g_1(\rho, \theta, s) = -(n-3)s_1(\rho - \theta s_1)^{n-4}(\rho^2 + \theta^2(1 - s_1^2))^{\frac{\alpha}{2}}$$

and

$$g_2(\rho, \theta, s) = \frac{\alpha}{2}(\rho - \theta s_1)^{n-3} 2\theta(1 - s_1^2)(\rho^2 + \theta^2(1 - s_1^2))^{\frac{\alpha}{2}-1}.$$

Now, the map $\theta \mapsto F(\theta, s)$ is continuous on $(-1/2, 1/2)$. Indeed, since the map $\theta \mapsto g(\rho, \theta, s) d\rho$ is continuous on $(-1/2, 1/2)$ and from (1.2) $\theta \mapsto \int_0^z g(\rho, \theta, s) d\rho$ is continuous on $(-1/2, 1/2)$. Then, for any $\epsilon > 0$, we have $\forall h, k$ sufficiently small,

$$\begin{aligned} \left| \int_0^{z+h} g(\rho, \theta + k, s) d\rho - \int_0^z g(\rho, \theta, s) d\rho \right| &\leq \left| \int_0^z g(\rho, \theta + k, s) d\rho \right. \\ &\quad \left. - \int_0^z g(\rho, \theta, s) d\rho \right| \\ &\quad + \left| \int_z^{z+h} g(\rho, \theta + k, s) d\rho \right| \\ &\leq \epsilon. \end{aligned}$$

Then, the map $(z, \theta) \mapsto \int_0^z g(\rho, \theta, s) d\rho$ is continuous on $[-1, 3] \times (-1/2, 1/2)$ and consequently $\theta \mapsto F(\theta, s)$ is continuous on $(-1/2, 1/2)$.

Now, we know that $\theta \mapsto H(\theta, s)F(\theta, s)$ is continuous on $(-1/2, 1/2)$ and differentiable on $(-1/2, 1/2) \setminus \{0\}$. Furthermore from (1.2), we have, for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$|H(\theta, s)F(\theta, s)| \leq 3.2^{\frac{n-3}{2}} 10^{\frac{\alpha+n-3}{2}} C_n \cdot \frac{1}{(1-s_1^2)^{\frac{n-3}{2}}}. \quad (1.4)$$

$$\left| \frac{\partial H}{\partial \theta}(\theta, s)F(\theta, s) \right| \leq 3.2^{\frac{n-3}{2}} 10^{\frac{\alpha+n-3}{2}} C_n \cdot \frac{1}{(1-s_1^2)^{\frac{n-3}{2}}}. \quad (1.5)$$

Consider the map $\eta : (\theta, s) \mapsto \eta(\theta, s) = \frac{\theta(s_1^2-1)}{\sqrt{1-\theta^2+\theta^2 s_1^2}} ((1-\theta^2+\theta^2 s_1^2)^{1/2} - \theta s_1)^{n-3}$. This map is indefinitely differentiable on $(-1/2, 1/2) \times \mathbb{S}^{n-1}$. Let B_n be a positive real number so that, $\forall (\theta, s) \in (-1/2, 1/2) \times \mathbb{S}^{n-1}$,

$$|\eta(\theta, s)| \leq B_n \quad \left| \frac{\partial \eta}{\partial \theta}(\theta, s) \right| \leq B_n.$$

Considering $a, b \in (0, 1/2)$ with $a < b$ we have, for any $\theta \in (a, b)$, for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$\begin{aligned} \left| H(\theta, s) \frac{\partial F}{\partial \theta}(\theta, s) \right| &\leq \left(B_n + 3(n-3) \cdot 4^{n-4} \cdot a^\alpha (1-s_1^2)^{\frac{\alpha}{2}} \right. \\ &\quad \left. + |3\alpha| \cdot 4^{n-3} a^{\alpha-1} (1-s_1^2)^{\frac{\alpha}{2}} \right) C_n. \end{aligned} \quad (1.6)$$

Since the maps $s \mapsto \frac{1}{(1-s_1^2)^{\frac{n-3}{2}}}$ and $s \mapsto (1-s_1^2)^{\frac{\alpha}{2}}$ are integrable on \mathbb{S}^{n-1} , we deduce that $\theta \mapsto G(\theta)$ is continuous on $(-1/2, 1/2)$ and continuously differentiable on $(-1/2, 1/2) \setminus \{0\}$.

Lemma 0.5 *The map $\theta \mapsto G(\theta)$ is differentiable at 0 and $\frac{dG}{d\theta}(0) = 0$.*

Proof Since for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$, $\theta \mapsto F(\theta, s)$ is continuous on $(-1/2, 1/2)$ from (A) we have,

$$\frac{\partial H}{\partial \theta}(\theta, s)F(\theta, s) \xrightarrow{\theta \rightarrow 0} \frac{\partial H}{\partial \theta}(0, s)F(0, s) = -2(n-1)s_1 \int_0^1 \rho^{n-3+\alpha} d\rho = \frac{-2(n-1)s_1}{n-2+\alpha}.$$

From (1.5) and Lebesgue Theorem we have,

$$\int_{\mathbb{S}^{n-1}} \frac{\partial H}{\partial \theta}(\theta, s)F(\theta, s) ds \xrightarrow{\theta \rightarrow 0} \int_{\mathbb{S}^{n-1}} \frac{-2(n-1)s_1}{n-2+\alpha} ds = 0.$$

Moreover, it is clear that,

$$\int_{\mathbb{S}^{n-1}} H(\theta, s)\eta(\theta, s)ds \xrightarrow{\theta \rightarrow 0} 0.$$

Let $J(m, n)$ be the integral,

$$J(m, n) = \int_{\frac{s_1}{\sqrt{1-s_1^2}}}^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} (\sqrt{1-s_1^2}t - s_1)^m (t^2 + 1)^n dt.$$

Notice that $J(m, n)$ converges as θ goes to 0 if and only if $m + 2n < -1$. Consider the change of variables $\rho = t\theta\sqrt{1-s_1^2}$ if $\theta > 0$. If $\theta < 0$, then we set $\rho = -t\theta\sqrt{1-s_1^2}$ and conclusion will be the same. Hence, we assume that $\theta > 0$. Then,

$$\int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho = -(n-3)s_1(1-s_1^2)^{\frac{1+\alpha}{2}}\theta^{n-3+\alpha}J(n-4, \frac{\alpha}{2}).$$

$$\int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho = \alpha\theta^{n-3+\alpha}(1-s_1^2)^{\frac{1+\alpha}{2}}J(n-3, \frac{\alpha}{2}-1).$$

First case : $\alpha \geq 4 - n$.

$J(n-4, \frac{\alpha}{2})$ and $J(n-3, \frac{\alpha}{2}-1)$ go to $+\infty$ as $\theta \rightarrow 0$. Furthermore, we have,

$$J(n-4, \frac{\alpha}{2}) \underset{0}{\sim} (1-s_1^2)^{\frac{n-4}{2}} \int_{\frac{s_1}{\sqrt{1-s_1^2}}}^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} t^{n-4+\alpha} dt$$

$$J(n-4, \frac{\alpha}{2}) \underset{0}{\sim} \frac{1}{n-3+\alpha} \frac{1}{\theta^{n-3+\alpha}} (1-s_1^2)^{\frac{-1-\alpha}{2}}.$$

Since $t^{n+\alpha-5}$ may be equal to zero at zero, we write,

$$J(n-3, \frac{\alpha}{2}-1) = \int_{\frac{s_1}{\sqrt{1-s_1^2}}}^1 (\sqrt{1-s_1^2}t - s_1)^{n-3} (t^2 + 1)^{\frac{\alpha}{2}-1} dt$$

$$+ \int_1^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} (\sqrt{1-s_1^2}t - s_1)^{n-3} (t^2 + 1)^{\frac{\alpha}{2}-1} dt.$$

We have,

$$J(n-3, \frac{\alpha}{2}-1) \underset{0}{\sim} (1-s_1^2)^{\frac{n-3}{2}} \int_1^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} t^{n-5+\alpha} dt.$$

Then, if $\alpha \neq 4 - n$,

$$J(n-3, \frac{\alpha}{2} - 1) \underset{0}{\sim} \frac{1}{n-4+\alpha} \frac{1}{\theta^{n-4+\alpha}} (1-s_1^2)^{\frac{1-\alpha}{2}},$$

and note that if $\alpha = 4 - n$, $J(n-3, \frac{\alpha}{2} - 1) \underset{0}{\sim} -(1-s_1^2)^{\frac{n-3}{2}} \ln(\theta^2(1-s_1^2))$. Hence, by (A) we have,

$$\begin{aligned} H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho &= -H(\theta, s)(n-3)s_1(1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} I_1 \\ &\xrightarrow{\theta \rightarrow 0} -\frac{(n-3)(n-1)}{n-3+\alpha} s_1, \end{aligned}$$

and

$$H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho = H(\theta, s)\alpha(1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} I_2 \xrightarrow{\theta \rightarrow 0} 0.$$

Observe that $\frac{|s_1|}{\sqrt{1-s_1^2}} \leq \sqrt{\frac{1}{\theta^2(1-s_1^2)} - 1}$. Indeed, $s_1^2 \theta^2 \leq 1 - \theta^2 + \theta^2 s_1^2$. It follows from (1.1) that

$$(\rho - \theta s_1)^{n-4} \leq \frac{2^{\frac{n-4}{2}}}{(1-s_1^2)^{\frac{n-4}{2}}} (\rho^2 + \theta^2(1-s_1^2))^{\frac{n-4}{2}}.$$

Recall that $\rho = t\theta\sqrt{1-s_1^2}$. Since $\alpha \geq 4 - n$, we have, for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$\begin{aligned} \left| H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho \right| &\leq 2C_n(n-3)2^{\frac{n-4}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\ &\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)} - 1}} (t^2 + 1)^{\frac{n-4+\alpha}{2}} dt \\ &\leq C_n(n-3)2^{\frac{n-2}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\ &\quad \times \sqrt{\frac{1}{\theta^2(1-s_1^2)} - 1} \left(\frac{1}{\theta^2(1-s_1^2)} \right)^{\frac{n-4+\alpha}{2}} \\ &\leq C_n(n-3)2^{\frac{n-2}{2}} (1-s_1^2)^{\frac{-n+4}{2}} \sqrt{1 - \theta^2(1-s_1^2)} \\ &\leq C_n(n-3)2^{\frac{n-1}{2}} (1-s_1^2)^{\frac{-n+4}{2}}. \end{aligned}$$

Since $s \mapsto (1-s_1^2)^{\frac{-n+4}{2}}$ is integrable on \mathbb{S}^{n-1} , by Lebesgue Theorem we have,

$$\int_{\mathbb{S}^{n-1}} H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho ds \xrightarrow{\theta \rightarrow 0} - \int_{\mathbb{S}^{n-1}} \frac{(n-3)(n-1)}{n-3+\alpha} s_1 ds = 0.$$

Moreover, we have, for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$, since $\alpha + n - 5 \geq 0$,

$$\begin{aligned}
\left| H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho \right| &\leq 2C_n |\alpha| 2^{\frac{n-3}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\
&\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} (t^2+1)^{\frac{n-5+\alpha}{2}} dt \\
&\leq C_n |\alpha| 2^{\frac{n-2}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\
&\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} \frac{1}{(t^2+1)} dt \left(\frac{1}{\theta^2(1-s_1^2)} \right)^{\frac{n-3+\alpha}{2}} \\
&\leq C_n |\alpha| 2^{\frac{n-2}{2}} \frac{\pi}{2} (1-s_1^2)^{\frac{-n+4}{2}}.
\end{aligned}$$

Then, by Lebesgue Theorem,

$$\int_{\mathbb{S}^{n-1}} H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho ds \xrightarrow{\theta \rightarrow 0} 0.$$

Second case : $3 - n < \alpha < 4 - n$.

For the same reasons that when $\alpha \geq 4 - n$, we have,

$$H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho \xrightarrow{\theta \rightarrow 0} -\frac{(n-3)(n-1)}{n-3+\alpha} s_1.$$

Furthermore, as $4 - n > \alpha > 3 - n$, $\forall s \in \mathbb{S}^{n-1} \setminus \{(-1, 0, \dots, 0), (1, 0, \dots, 0)\}$,

$$\begin{aligned}
\left| H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho \right| &\leq 2C_n (n-3) 2^{\frac{n-4}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\
&\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} (t^2+1)^{\frac{n-4+\alpha}{2}} dt \\
&\leq C_n (n-3) 2^{\frac{n-2}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\
&\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} (t^2)^{\frac{n-4+\alpha}{2}} dt \\
&\leq \frac{C_n (n-3) 2^{\frac{n-2}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha}}{n-3+\alpha} \left(\frac{1}{\theta^2(1-s_1^2)} - 1 \right)^{\frac{n-3+\alpha}{2}}. \\
&\leq \frac{C_n (n-3) 2^{\frac{2n-7+\alpha}{2}} (1-s_1^2)^{\frac{4-n}{2}}}{n-3+\alpha}.
\end{aligned}$$

Then, by Lebesgue Theorem,

$$\int_{\mathbb{S}^{n-1}} H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho ds \xrightarrow{\theta \rightarrow 0} -\int_{\mathbb{S}^{n-1}} \frac{(n-3)(n-1)}{n-3+\alpha} s_1 ds = 0.$$

Moreover, $J(n-3, \frac{\alpha}{2}-1)$ is finite when $\theta \rightarrow 0$ then, as $\alpha > 3-n$, Furthermore,

$$H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho ds \xrightarrow{\theta \rightarrow 0} 0.$$

$$\begin{aligned} \left| H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho \right| &\leq 2C_n |\alpha| 2^{\frac{n-3}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\ &\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} (t^2+1)^{\frac{n+\alpha-5}{2}} dt \\ &\leq C_n |\alpha| 2^{\frac{n-1}{2}} (1-s_1^2)^{\frac{\alpha+1}{2}} \theta^{n-3+\alpha} \\ &\quad \times \int_0^{\sqrt{\frac{1}{\theta^2(1-s_1^2)}-1}} \frac{1}{(t^2+1)} dt \left(\frac{1}{\theta^2(1-s_1^2)} \right)^{\frac{n-3+\alpha}{2}} \\ &\leq C_n |\alpha| 2^{\frac{n-1}{2}} (1-s_1^2)^{\frac{-n+4}{2}} \int_0^{+\infty} \frac{1}{(t^2+1)} dt. \end{aligned}$$

Then, by Lebesgue Theorem,

$$\int_{\mathbb{S}^{n-1}} H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho ds \xrightarrow{\theta \rightarrow 0} 0.$$

Finally, we have

$$\frac{dG}{d\theta}(\theta) \xrightarrow{\theta \rightarrow 0} 0.$$

By Lemma 1.4 we deduce that G is differentiable at 0 and $\frac{dG}{d\theta}(0) = 0$.

Lemma 0.6 *The map $\theta \rightarrow G(\theta)$ is two times differentiable on $(-1/2, 1/2) \setminus \{0\}$.*

Proof We know that the map $\theta \rightarrow \frac{\partial H}{\partial \theta}(\theta, s)F(\theta, s)$ is differentiable on $(-1/2, 1/2) \setminus \{0\}$. The maps $\theta \rightarrow \eta(\theta, s)$, $\theta \rightarrow g_1(\rho, \theta, s)$, $\theta \rightarrow g_2(\rho, \theta, s)$ are differentiable on $(-1/2, 1/2) \setminus \{0\}$. We have,

$$\begin{aligned} \frac{\partial \eta}{\partial \theta}(\theta, s) &= \frac{(s_1^2-1)\sqrt{1-\theta^2+\theta^2 s_1^2} - \theta(s_1^2-1)\frac{\theta(s_1^2-1)}{\sqrt{1-\theta^2+\theta^2 s_1^2}}}{1-\theta^2+\theta^2 s_1^2} \\ &\quad \times (\sqrt{1-\theta^2+\theta^2 s_1^2} - \theta s_1)^{n-3} \\ &\quad + \frac{(n-3)\theta(s_1^2-1)}{\sqrt{1-\theta^2+\theta^2 s_1^2}} \left(\frac{\theta(s_1^2-1)}{\sqrt{1-\theta^2+\theta^2 s_1^2}} - s_1 \right) \\ &\quad \times (\sqrt{1-\theta^2+\theta^2 s_1^2} - \theta s_1)^{n-4}. \end{aligned}$$

$$\begin{aligned}\frac{\partial g_1}{\partial \theta}(\rho, \theta, s) &= (n-3)(n-4)s_1^2(\rho - \theta s_1)^{n-5}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}} \\ &\quad - \alpha(n-3)s_1(1-s_1^2)\theta(\rho - \theta s_1)^{n-4}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-1}.\end{aligned}$$

$$\begin{aligned}\frac{\partial g_2}{\partial \theta}(\rho, \theta, s) &= -\alpha(n-3)s_1(1-s_1^2)\theta(\rho - \theta s_1)^{n-4}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-1} \\ &\quad + \alpha(\alpha-2)(1-s_1^2)^2\theta^2(\rho - \theta s_1)^{n-3}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-2} \\ &\quad + \alpha(1-s_1^2)(\rho - \theta s_1)^{n-3}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-1}.\end{aligned}$$

We set,

$$\begin{aligned}g_{11}(\rho, \theta, s) &= (n-3)(n-4)s_1^2(\rho - \theta s_1)^{n-5}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}}, \\ g_{12}(\rho, \theta, s) &= -2\alpha(n-3)s_1(1-s_1^2)\theta(\rho - \theta s_1)^{n-4}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-1}, \\ g_{21}(\rho, \theta, s) &= \alpha(\alpha-2)(1-s_1^2)^2\theta^2(\rho - \theta s_1)^{n-3}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-2}, \\ g_{22}(\rho, \theta, s) &= \alpha(1-s_1^2)(\rho - \theta s_1)^{n-3}(\rho^2 + \theta^2(1-s_1^2))^{\frac{\alpha}{2}-1}.\end{aligned}$$

Let $a, b \in (0, 1/2)$ with $a < b$. We have, $\forall s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$\left| \frac{\partial g_1}{\partial \theta}(\rho, \theta, s) \right| \leq (n-3)(n-4)4^{n-5}a^\alpha(1-s_1^2)^{\frac{\alpha}{2}} + |\alpha|(n-3)4^{n-4}a^{\alpha-1}(1-s_1^2)^{\frac{\alpha}{2}}. \quad (1.7)$$

$$\begin{aligned}\left| \frac{\partial g_2}{\partial \theta}(\rho, \theta, s) \right| &\leq |\alpha(\alpha-2)|4^{n-3}a^{\alpha-2}(1-s_1^2)^{\frac{\alpha}{2}} \\ &\quad + |\alpha|4^{n-3}a^{\alpha-1}(1-s_1^2)^{\frac{\alpha}{2}} \\ &\quad + |\alpha|(n-3)4^{n-4}a^{\alpha-1}(1-s_1^2)^{\frac{\alpha}{2}}. \quad (1.8)\end{aligned}$$

Then, for any $i \in \{1, 2\}$, the maps $\theta \mapsto \int_0^z g_i(\rho, \theta, s)d\rho$ is differentiable on $(0, 1/2)$, and

$$\frac{\partial}{\partial \theta} \int_0^z g_i(\rho, \theta, s)d\rho = \int_0^z \frac{\partial g_i}{\partial \theta}(\rho, \theta, s)d\rho.$$

Furthermore, for any $i \in \{1, 2\}$, $\theta \mapsto \frac{\partial g_i}{\partial \theta}(\rho, \theta, s)$ is continuous on $(-1/2, 1/2) \setminus \{0\}$, then, $\theta \mapsto \int_0^z \frac{\partial g_i}{\partial \theta}(\rho, \theta, s)d\rho$, is continuous on $(-1/2, 1/2) \setminus \{0\}$. Hence, for any $i \in \{1, 2\}$ and for any $\epsilon > 0$, we have $\forall h, k$ two sufficiently small,

$$\begin{aligned}\left| \int_0^{z+h} \frac{\partial g_i}{\partial \theta}(\rho, \theta+k, s)d\rho - \int_0^z \frac{\partial g_i}{\partial \theta}(\rho, \theta, s)d\rho \right| &\leq \left| \int_0^z \frac{\partial g_i}{\partial \theta}(\rho, \theta+k, s)d\rho \right. \\ &\quad \left. - \int_0^z \frac{\partial g_i}{\partial \theta}(\rho, \theta, s)d\rho \right| \\ &\quad + \left| \int_z^{z+h} \frac{\partial g_i}{\partial \theta}(\rho, \theta+k, s)d\rho \right| \\ &\leq \epsilon.\end{aligned}$$

This proves that for any $i \in \{1, 2\}$, $(z, \theta) \mapsto \int_0^z \frac{\partial g_i}{\partial \theta}(\rho, \theta, s) d\rho$ is continuous on $[-1, 3] \times (-1/2, 1/2) \setminus \{0\}$. Moreover, for any $i \in \{1, 2\}$ the map $\rho \mapsto g_i(\rho, \theta, s)$ is continuous on $[-1, 3]$ for any $\theta \in (-1/2, 1/2) \setminus \{0\}$. Then, $z \mapsto \int_0^z g_i(\rho, \theta, s) d\rho$ is differentiable on $[-1, 3]$ for any $\theta \in (-1/2, 1/2) \setminus \{0\}$ and $\frac{\partial}{\partial z} \int_0^z g_i(\rho, \theta, s) d\rho = g_i(z, \theta, s)$.

Since $(z, \theta) \mapsto g_i(z, \theta)$ is continuous on $[-1, 3] \times (-1/2, 1/2) \setminus \{0\}$ we finally deduce that for any $i \in \{1, 2\}$, $\theta \mapsto \int_{\theta s_1}^{\beta(\theta, s)} g_i(\rho, \theta, s) d\rho$ is differentiable on $(-1/2, 1/2) \setminus \{0\}$ and,

$$\begin{aligned} \sum_{i=1}^2 \frac{\partial}{\partial \theta} \int_{\theta s_1}^{\beta(\theta, s)} g_i(\rho, \theta, s) d\rho &= \frac{\theta(s_1^2 - 1)}{\sqrt{1 - \theta^2 + \theta^2 s_1^2}} \\ &\quad \times \left(-(n-3)s_1(\sqrt{1 - \theta^2 + \theta^2 s_1^2} - \theta s_1)^{n-4} \right. \\ &\quad \left. + \alpha\theta(1 - s_1^2)(\sqrt{1 - \theta^2 + \theta^2 s_1^2} - \theta s_1)^{n-3} \right) \\ &\quad + \sum_{i=1}^2 \int_{\theta s_1}^{\beta(\theta, s)} \frac{\partial^2 g_i}{\partial^2 \theta}(\rho, \theta, s) d\rho. \end{aligned}$$

We deduce that $\theta \mapsto \frac{\partial F}{\partial \theta}$ is differentiable in $(-1/2, 1/2) \setminus \{0\}$. Moreover, we see that the map,

$$\begin{aligned} \theta \mapsto \lambda(\theta, s) &= \frac{\theta(s_1^2 - 1)}{\sqrt{1 - \theta^2 + \theta^2 s_1^2}} \left(-(n-3)s_1(\sqrt{1 - \theta^2 + \theta^2 s_1^2} - \theta s_1)^{n-4} \right. \\ &\quad \left. + \alpha\theta(1 - s_1^2)(\sqrt{1 - \theta^2 + \theta^2 s_1^2} - \theta s_1)^{n-3} \right) \end{aligned}$$

is indefinitely differentiable on $(-1/2, 1/2) \times \mathbb{S}^{n-1}$. Then, by (1.1), (1.2), (1.8), (1.7), (1.3) and (A), for any $a, b \in (0, 1/2)$, $a < b$ there exists constants $K_{1,n,ab,\alpha}$, $K_{2,n,ab,\alpha}$, $K_{3,n,ab,\alpha}$ so that, for any $|\theta| \in (a, b)$, for any $s \in \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$\left| \frac{\partial^2 HF}{\partial \theta^2}(\theta, s) \right| \leq K_{1,n,ab,\alpha}(1 - s_1^2)^{\frac{\alpha}{2}} + K_{2,n,ab,\alpha}(1 - s_1^2)^{\frac{3-n}{2}} + K_{3,n,ab,\alpha}.$$

We deduce by Lebesgue Theorem that the map $\theta \mapsto E(\theta)$ is two times differentiable on $(-1/2, 1/2) \setminus \{0\}$ and,

$$\frac{d^2 G}{d\theta^2}(\theta) = \int_{\mathbb{S}^{n-1}} \frac{\partial^2 HF}{\partial \theta^2}(\theta, s) ds.$$

Lemma 0.7 *If $5 - n > \alpha > 4 - n$, the map $\theta \mapsto G(\theta)$ is two times differentiable at 0.*

Proof Suppose that $\alpha \in (4 - n, 5 - n)$. As in Lemma 1.5, we can see that,

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} \frac{\partial^2 H}{\partial \theta^2}(\theta, s) F(\theta, s) ds &\xrightarrow{\theta \rightarrow 0} \int_{\mathbb{S}^{n-1}} \frac{1}{2} \frac{(2n-3)s_1^2 - (n-2)}{n-2+\alpha} ds = \frac{-n^2 + 4n - 3}{2n(n-2+\alpha)} |\mathbb{S}^{n-1}|, \\ \int_{\mathbb{S}^{n-1}} \frac{\partial H}{\partial \theta}(\theta, s) \eta(\theta, s) ds &\xrightarrow{\theta \rightarrow 0} 0, \\ \int_{\mathbb{S}^{n-1}} \frac{\partial H}{\partial \theta}(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_1(\rho, \theta, s) d\rho &\xrightarrow{\theta \rightarrow 0} \int_{\mathbb{S}^{n-1}} \frac{2(n-3)(n-1)}{n-3+\alpha} s_1^2 = \frac{2(n-3)(n-1)}{n(n-3+\alpha)} |\mathbb{S}^{n-1}|, \\ \int_{\mathbb{S}^{n-1}} \frac{\partial H}{\partial \theta}(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_2(\rho, \theta, s) d\rho &\xrightarrow{\theta \rightarrow 0} 0, \\ \int_{\mathbb{S}^{n-1}} H(\theta, s) \frac{\partial \eta}{\partial \theta}(\theta, s) ds &\xrightarrow{\theta \rightarrow 0} \int_{\mathbb{S}^{n-1}} (n-1)(s_1^2 - 1) ds = \frac{-(n-1)^2}{n} |\mathbb{S}^{n-1}|, \end{aligned}$$

and

$$\int_{\mathbb{S}^{n-1}} H(\theta, s) \lambda(\theta, s) ds \xrightarrow{\theta \rightarrow 0} 0.$$

As in Lemma 1.5, we set $\rho = \sqrt{1 - s_1^2} \theta t$ if $\theta > 0$. Hence,

$$\int_{\theta s_1}^{\beta(\theta, s)} g_{11}(\rho, \theta, s) d\rho = (n-3)(n-4) s_1^2 (1 - s_1^2)^{\frac{1+\alpha}{2}} \theta^{n-4+\alpha} J\left(n-5, \frac{\alpha}{2}\right).$$

$$\int_{\theta s_1}^{\beta(\theta, s)} g_{12}(\rho, \theta, s) d\rho = -2\alpha(n-3) s_1 (1 - s_1^2)^{\frac{1+\alpha}{2}} \theta^{n-4+\alpha} J\left(n-4, \frac{\alpha}{2} - 1\right)$$

$$\int_{\theta s_1}^{\beta(\theta, s)} g_{21}(\rho, \theta, s) d\rho = \alpha(\alpha-2) (1 - s_1^2)^{\frac{1+\alpha}{2}} \theta^{n-4+\alpha} J\left(n-3, \frac{\alpha}{2} - 2\right)$$

$$\int_{\theta s_1}^{\beta(\theta, s)} g_{22}(\rho, \theta, s) d\rho = \alpha(1 - s_1^2)^{\frac{1+\alpha}{2}} \theta^{n-4+\alpha} J\left(n-3, \frac{\alpha}{2} - 1\right)$$

Since $\alpha \in (4 - n, 5 - n)$, the integrals $J\left(n-5, \frac{\alpha}{2}\right)$ and $J\left(n-3, \frac{\alpha}{2} - 1\right)$ are infinite and we have,

$$J\left(n-5, \frac{\alpha}{2}\right) \underset{0}{\sim} \frac{(1 - s_1^2)^{-\frac{1-\alpha}{2}} \theta^{-n-\alpha+4}}{n-4+\alpha}, \quad J\left(n-3, \frac{\alpha}{2} - 1\right) \underset{0}{\sim} \frac{(1 - s_1^2)^{\frac{1-\alpha}{2}} \theta^{-n-\alpha+4}}{n-4+\alpha}$$

And the integrals

$J(n-4, \frac{\alpha}{2}-1)$ and $J(n-3, \frac{\alpha}{2}-2)$ are finite. Then,

$$\int_{\theta s_1}^{\beta(\theta, s)} g_{11}(\rho, \theta, s) d\rho \xrightarrow{\theta \rightarrow 0} \frac{(n-3)(n-4)s_1^2}{n-4+\alpha}, \quad \int_{\theta s_1}^{\beta(\theta, s)} g_{22}(\rho, \theta, s) d\rho \xrightarrow{\theta \rightarrow 0} \frac{\alpha(1-s_1^2)}{n-4+\alpha}.$$

$$\int_{\theta s_1}^{\beta(\theta, s)} g_{12}(\rho, \theta, s) d\rho \xrightarrow{\theta \rightarrow 0} 0, \quad \int_{\theta s_1}^{\beta(\theta, s)} g_{21}(\rho, \theta, s) d\rho \xrightarrow{\theta \rightarrow 0} 0$$

Moreover, we can see that, for any $i, j \in \{1, 2\}$, for any $(\theta, s) \in (-1/2, 1/2) \times \mathbb{S}^{n-1} \setminus \{(\pm 1, 0, \dots, 0)\}$,

$$H(\theta, s) \int_{\theta s_1}^{\beta(\theta, s)} g_{ij}(\rho, \theta, s) d\rho \leq C_{n, \alpha} (1-s_1^2)^{\frac{5-n}{2}} + D_{n, \alpha} (1-s_1^2)^{\frac{\alpha+1}{2}}.$$

where $C_{n, \alpha}$ and $D_{n, \alpha}$ are two constants independent of θ . By Lebesgue Theorem we deduce that,

$$\int_{\mathbb{S}^{n-1}} H(\theta, s) \frac{\partial^2 F}{\partial \theta^2}(\theta, s) ds \xrightarrow{\theta \rightarrow 0} \frac{-(n-1)^2}{n} |\mathbb{S}^{n-1}|$$

$$+ (n-1) \frac{(n-3)(n-4) + \alpha(n-1)}{n(n-4+\alpha)} |\mathbb{S}^{n-1}|.$$

By Lemmas 1.1, 1.2, 1.3, $\theta \mapsto G(\theta) \in \mathcal{C}^1((-1/2, 1/2), \mathbb{R})$ and is two times differentiable on $(-1/2, 1/2) \setminus \{0\}$. Furthermore, when $\alpha \in (4-n, 5-n)$, as the limit of $\frac{d^2 G}{d\theta^2}(\theta)$ exists as $\theta \rightarrow 0$, we have $\theta \mapsto G(\theta)$ is two times differentiable on $(-1/2, 1/2)$.

Proof of ii). Assume that $\alpha \in (4-n, 5-n)$, by Lemma 1.1, 1.2, 1.3, 1.4, we have,

$$G(\theta) = G(0) + \frac{1}{2} \frac{d^2 G}{d\theta^2}(0) + o(\theta^2).$$

Furthermore we have,

$$\frac{d^2 G}{d\theta^2}(0) = \frac{-n^2 + 4n - 3}{2n(n-2+\alpha)} |\mathbb{S}^{n-1}| + \frac{2(n-3)(n-1)}{n(n-3+\alpha)} |\mathbb{S}^{n-1}|$$

$$+ \frac{-(n-1)^2}{n} |\mathbb{S}^{n-1}| + (n-1) \frac{(n-3)(n-4) + \alpha(n-1)}{n(n-4+\alpha)} |\mathbb{S}^{n-1}|.$$

We have, for any $n \geq 6$.

$$(n-3)(n-4) + \alpha(n-1) \xrightarrow{\alpha \rightarrow 4-n} -2(n-4) < 0.$$

Then,

$$\frac{(n-3)(n-4) + \alpha(n-1)}{n(n-4+\alpha)} \xrightarrow{\alpha \rightarrow > 4-n} -\infty, \quad \text{and} \quad \frac{d^2 G}{d^2 \theta}(0) \xrightarrow{\alpha \rightarrow > 4-n} -\infty.$$

Hence, there is α_0 such that, for any $\alpha \in (4 - n, \alpha_0)$, $G(\theta) < G(0)$ for θ sufficiently small, that is,

$$G(\theta) = E_{2,r^\alpha}(u_a) = \int_{\mathbf{B}^n} r^\alpha \|\nabla u_a\|^2 dx < G(0) = \int_{\mathbf{B}^n} r^\alpha \|\nabla u_0\|^2 dx. \quad \blacksquare$$

Références

- [1] M. Avellaneda, F.-H. Lin, Null-Lagrangians and minimizing $\int |\nabla u|^p$, *C. R. Acad. Sci. Paris*, **306**(1988), 355-358.
- [2] H. Brezis, J.-M. Coron, E.-H. Lieb, Harmonic Maps with Defects, *Commun. Math. Phys.*, **107** (1986), 649-705.
- [3] B. Chen, Singularities of p -harmonic mappings, *Thesis, University of Minnesota*, (1989).
- [4] J.-M. Coron, F. Helein, Harmonic diffeomorphisms, minimizing harmonic maps and rotational symmetry, *Compositio Math.*, **69** (1989), 175-228.
- [5] J.-M. Coron, R. Gulliver, Minimizing p -harmonic maps into spheres, *J. reine angew. Math.*, **401** (1989), 82-100.
- [6] A. El Soufi, A. Jeune, Indice de Morse des applications p -harmoniques, *Ann. Inst. Henri Poincaré*, **13**(1996), 229-250.
- [7] A. El Soufi, E. Sandier, p -harmonic diffeomorphisms, *Calc. of Var.*, **6**(1998), 161-169.
- [8] B. Chen, R. Hardt, Prescribing singularities for p -harmonic mappings, *Indiana University Math. J.*, **44**(1995), 575-601. bibitemHL1R. Hardt, F.-H. Lin, Mapping minimizing the L^p norm of the gradient, *Comm. P.A.M.*, **15**(1987), 555-588.
- [9] R. Hardt, F.-H. Lin, Singularities for p -energy minimizing unit vectorfields on planar domains, *Calculus of Variations and Partial Differential Equations*, **3**(1995), 311-341.
- [10] R. Hardt, F.-H. Lin, C.-Y. Wang, The p -energy minimality of $\frac{x}{\|x\|}$, *Communications in analysis and geometry*, **6**(1998), 141-152.
- [11] R. Hardt, F.-H. Lin, C.-Y. Wang, Singularities of p -Energy Minimizing Maps, *Comm. P.A.M.*, **50**(1997), 399-447.
- [12] F. Helein, Harmonic diffeomorphisms between an open subset of \mathbb{R}^3 and a Riemannian manifold, *C. R. Acad. Sci. Paris*, **308**(1989), 237-240.
- [13] S. Hildebrandt, H. Kaul, K.-O. Wildman, An existence theorem for harmonic mappings of Riemannian manifold, *Acta Math.*, **138**(1977), 1-16.
- [14] M.-C. Hong, On the Jäger-Kaul theorem concerning harmonic maps, *Ann. Inst. Poincaré, Analyse non-linéaire*, **17** (2000), 35-46.
- [15] M.-C. Hong, On the minimality of the p -harmonic map $\frac{x}{\|x\|} : \mathbf{B}^n \rightarrow \mathbf{S}^{n-1}$, *Calc. Var.*, **13** (2001), 459-468.

- [16] W. Jager, H. Kaul, Rotationally symmetric harmonic maps from a ball into a sphere and the regularity problem for weak solutions of elliptic systems, *J.Reine Angew. Math.*, **343**(1983), 146-161.
- [17] F.-H. Lin, Une remarque sur l'application $x/\|x\|$, *C.R. Acad. Sci. Paris* **305**(1987), 529-531.
- [18] R. Shoen, K. Uhlenbeck, A regularity theory for harmonic maps *J. Differential Geom.* **12**(1982), 307-335.
- [19] R. Shoen, K. Uhlenbeck, Boundary theory and the Dirichlet problem for harmonic maps, *J. Differential Geom.*, **18**(1983), 253-268.
- [20] C.Wang, Minimality and perturbation of singularities for certain p -harmonic maps., *Indiana Univ. Math.J.*, **47**(1998), 725-740.