

Probing the Hofmeister Effect with Ultrafast Core-Hole Spectroscopy

Zhong Yin^{a,b,}, Ivan Rajkovic^{a,c}, Katharina Kubicek^{a,b}, Wilson Quevedo^d, Annette Pietzsch^d,
Philippe Wernet^d, Alexander Föhlisch^{d,e} and Simone Techert^{a,b,f,*}*

^aMax Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Göttingen, Germany

^bDeutsches Elektron-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany

^cPaul Scherrer Institut, 5232 Villigen PSI, Switzerland

^dInstitute for Methods and Instrumentation for Synchrotron Radiation Research, Helmholtz-Zentrum Berlin GmbH, Albert-Einstein-Str. 15, 12489 Berlin, Germany

^eUniversity of Potsdam, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany

^fUniversity of Göttingen, Institute for X-ray Physics, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany



ABSTRACT: In the current work, X-ray emission spectra of aqueous solutions of different inorganic salts within the Hofmeister series are presented. The results reflect the direct interaction of the ions with the water molecules and therefore reveal general properties of the salt – water interactions. Within the experimental precision a significant effect of the ions on the water structure has been observed but no ordering according to the structure maker / structure

breaker concept could be mirrored in the results indicating that the Hofmeister effect - if existent - may be caused by more complex interactions.

KEYWORDS: Hofmeister Series, X-ray Spectroscopy, Liquid Jet, Core-Hole Spectroscopy, Cation, Anion, Aqueous Solution, Resonant Inelastic Scattering

INTRODUCTION

At the end of the 19th century Franz Hofmeister discovered that common anions and cations could be classified according to their ability to salt in / salt out proteins, in detail to enhance aggregation and precipitation of proteins dissolved in aqueous solution, see Scheme 1.¹



Scheme 1: A partial list of salts, which are ordered according to their kosmotrop/chaotropic characteristics.

By now, the so-called Hofmeister effects have been observed in a number of systems ranging from atmospheric aerosols up to whole biological entities such as cells, proteins etc.^{2,3} The salting in/out effect even plays an essential role in the metabolism of eukaryotic cells and in the case of mammals it can lead to critical aggregation effects which were discussed as various pre-stages of diseases.^{4,5} However, the question concerning the effect's origin remains unanswered. Two common models to explain it are: (i) water interacting with the hydrated ions and causing precipitation or (ii) a consequence of the direct binding of the salt ions to the

biomolecule by electrostatic interactions.⁶

The first model leads to the concept of structure makers / structure breakers (SMB), depending on the anions and cations used. According to this concept, cations and anions stabilize or destabilize the long range water structure in aqueous solutions,⁷ which is conventionally interpreted as strengthening and weakening of the hydrogen bond structure.⁸ Theories based on molecular dynamic calculations have been introduced to explain the nature of the effect, some supporting the concept of structure breaking and making,⁹⁻¹¹ some contradicting.¹²⁻¹⁵ However, there is a lack of understanding on the molecular level, since SMB is based on experimental observations of macroscopic properties.^{16,17} Several experiments have been performed recently regarding the validation of SMB on the molecular level, which reveal objecting results.^{8,18-30} Yet, the influence of salt ions on the electronic redistribution of water is unclear. In order to probe the electronic influence of salt ions on the water structure and therefore if the SMB concept can be pictured on the molecular orbitals of the aqueous systems, core-hole X-ray emission spectroscopic (XES) measurements have been performed on the oxygen K-edge of pure water and aqueous salt solutions.

In XES, a core electron is excited with a monochromatic X-ray source to an unoccupied molecular orbital or above the ionization threshold to create a vacancy. The excited system relaxes via valence electrons filling the core-hole under emission of a photon containing information about the electronic structure of the molecules under investigation. XES is further divided into resonant XES (RIXS) and non-resonant XES. In RIXS, the core electron excitation to an unoccupied state and the following fluorescence are coupled events.^{31,32} The non-resonant XES is a two-step process, where the absorption and the emission are decoupled processes.^{31,32} An illustration of the molecular orbital and the difference between resonant and non-resonant

XES are shown in Figure 1.




Figure 1: Schematic description of RIXS and non-resonant XES. The molecular orbitals were calculated using the RHF/ 6-31G** basis set from Gaussian. The transition from $2a_1$ to the core level is unlikely due to selection rules.

XES is sensitive to the element and the chemical environment.³²⁻⁴¹ Moreover, photons offer bulk information as the penetration length is much larger compared to electrons. The aim of the current study is to further elucidate if and how the presence of salt ions can influence the water clusters and the origin of the Hofmeister effect.

The experiments presented here are the first XES measurements on aqueous salt solutions from the Hofmeister series using the liquid jet technique.^{42,43}

EXPERIMENTAL SECTION

The experiments were performed at the beamline U49/2-PGM 1 of BESSY II synchrotron at the Helmholtz Zentrum Berlin using the FlexRIXS endstation. For excitation, a monochromatic X-ray beam with $\sim 10^{12}$ photons per second on the sample has been utilized. A detailed description of the liquid jet system and the FlexRIXS end station can be found elsewhere.^{42,44} For sample supply a 20 μm diameter liquid jet has been used. The jet system consists of a mixer, a degasser and an HPLC pump (JASCO). The implementation of the liquid jet provides continuously fresh samples and avoids radiation damage during the experiment. In contrast to a liquid flow cell⁴⁵, the liquid jet also prevents interaction of X-ray and samples with the cell membrane and also prevents further loss of photon flux.

The 2 M NaCl and 1 M MgCl₂ concentrated aqueous salt solutions were prepared from commercially available salts of highest purity from Sigma - Aldrich at least 12 hours prior to the experiment, since Na⁺ is single and Mg²⁺ double charged. The resonant and non-resonant excitation energies have been selected from the X-ray absorption spectra (XAS) of water and aqueous solutions recorded in the total fluorescence yield mode. XAS and XES spectra of pure water were recorded periodically to confirm the stability of the setup. The monochromator slit was set to 150 μm corresponding to an energy bandwidth of approx. 0.4 eV. The resolution of the grating spectrometer was 0.4 eV at 530 eV (oxygen K-edge). The elastic peaks were used for energy calibration and the absolute energy scale in this experiment has been adjusted to Tokushima et al.³⁸

RESULTS AND DISCUSSION

Figure 2 shows the X-ray emission spectra for non-resonant as well as resonant excitation at the

oxygen K-edge for pure liquid water and aqueous NaCl solution with 2 M concentration.




Figure 2: X-ray emission spectra of pure water and 2 M NaCl_{aq.}.

(a) Spectra for resonant excitation energy. (b) Spectra for non-resonant excitation energy.

Resonant excitation at 537.45 eV (Fig. 2a) corresponds to a promotion of a core electron into the second LUMO (Lowest Unoccupied Molecular Orbital) of oxygen in water, the 2b₂ orbital. Non-resonant excitation at 540 eV (Fig. 2b) leads to the ionization of the water molecule, see Figure 1. The emission spectra have been area-normalized to provide an accurate comparison of relative intensities. All emission spectra show a multi-peak structure. Peak A can be correlated to the lone pair electron 1b₁, peak B and C to the covalent bonding electron pairs in the 3a₁ and 1b₂ orbitals, respectively (see Figure 1). For both excitation energies the intensities in peak A' of the 2 M salt solution are equal to water but a significant drop in intensities are observed at peak A'' compared to pure water.

With a 2 M salt concentration and a maximum water concentration of 55.5 M, every salt molecule is surrounded by 27-28 water molecules. As X-ray diffraction experiments have shown Na^+ and Cl^- are coordinated by six water molecules each.¹⁸ This means that twelve out of 28 water molecules are in the first solvation shell and in direct interaction with either the cation or the anion. The 16 water molecules left are in the second solvation shell of either the cation or anion, where they are potentially able to attract up to 12 water molecules each.

In the case of 1 M MgCl_2 solution, there are 55-56 water molecules for one salt moiety. Considering the coordination number for Mg^{2+} ,¹⁸ 18 water molecules are in the first solvation shell of the ions, resulting in 37-38 water molecules forming the second solvation shell.

The anions are bonded to water molecules via the hydrogen; hence the oxygen lone pair electrons of water are not directly interacting with the anions. The positively charged cations however are attracting the negative polar charged oxygen in water.⁴⁶ Thus, the oxygen lone pair electrons are responsible for the water coordination around the cations and changes in the oxygen emission spectra of the aqueous salt solutions, especially in the lone pair region are anticipated. The presence of 2 M NaCl and 1 M MgCl_2 seem not to affect the covalent bonding electrons (peak B and C), as can be seen in Figure 2 and 3.

Measurements of $\text{NaCl}_{(\text{aq})}$ and $\text{MgCl}_{2(\text{aq})}$ at lower concentrations of 500 mM showed no effect in the spectra, see Figure 4. For 500 mM NaCl and MgCl_2 solutions there are around 110 water molecules for one salt moiety. Since only twelve H_2O are in the first solvation shell for $\text{NaCl}_{(\text{aq})}$ and 18 for $\text{MgCl}_{2(\text{aq})}$, around 100 or 90 water molecules, respectively, are left to form higher solvation shells and separate the ions from each other. Therefore, we assume changes on the electronic structure caused by cooperativity to be negligible in the case of NaCl and MgCl_2 solutions.

An interesting feature is the pronounced splitting of the lone pair $1b_1$ line, both for pure water and aqueous solutions. According to Guo et al.³⁵ the splitting of the oxygen lone pair orbitals was reported previously for $\text{NaCl}_{(\text{aq.})}$ and $\text{AlCl}_3_{(\text{aq.})}$, but not for pure water. Consequently, it has been concluded that the origin of this effect is caused by the presence of salt ions. However, in contrast to Guo et al.³⁵ later XES experiments showed a clear double peak even for pure water³⁶⁻⁴⁰ so this cannot be an effect caused by the salt. The exact nature of the splitting is still an ongoing debate.




Figure 3: Resonantly excited emission spectrum of pure water compared to 1 M MgCl_2 solution.

Within the statistical uncertainty the peak position and intensity remain constant.

The fine structure is either ascribed to a dissociation into OH^- , which is a dynamical effect during the core-hole lifetime of around 4 fs^{37,47-49} or to two phases of water molecules with tetrahedral

and distorted structural motifs.^{38,40,50}

Furthermore, molecular dynamic (MD) calculations can qualitatively reproduce the fine structure without including a two component model of liquid water, rather based on ultrafast dynamics during the core-hole lifetime.^{48,49}

Conclusively, this would predict that the splitting in A is due to different molecular orbitals. Forsberg et al.³⁹ reported an angular dependence of A' and A'' which supports the theoretical concept described by Odelius,^{48,49} whereas Tokushima et al.⁴⁰ reported the negligible effect of different polarization in the XES spectra. Regardless of the double peak structure's origin, the presented data of aqueous salt solutions are interpreted within both frameworks. Within our experimental resolution, significant changes in relative intensities of the double peak features have been observed.




Figure 4: Non resonant X-ray emission spectra on the oxygen K-edge of pure water compared to

(a) 500 mM NaCl and (b) 500 mM MgCl₂ solution. No significant changes are observed, therefore a negligible cooperative effect on the electronic structure is deduced.

If the splitting is rooted in the dynamical process during the core-hole lifetime, where A' is from intact water molecules and A'' from dissociated OH⁻, we would expect that Na⁺ reduces the acceptor characteristic of the water molecules due to the steric hindrance and thus lead to less dissociations, which would be mirrored in less intensity of A''. The anion Cl⁻ is connected to the hydrogen site of water. Therefore, the water molecules around the anion still can act as donor and acceptor. The spectra in Figure 2 show a drop of peak A'' of the salt solution compared to pure water, which would indicate a reduced rate of dissociations of H₂O. In the concept of structure maker and structure breaker, Cl⁻ is a structure breaker, Na⁺ is a borderline, but also conventionally referred as a breaker⁶ and Mg²⁺ is a known structure maker ion.⁵¹ According to the Hofmeister categorization, Cl⁻ and Na⁺ break the water structure and thus decrease the number of dissociations, which is in agreement with Figure 2. With a structure maker like Mg²⁺ an enhancement of hydrogen bonding and therefore an increasing rate of dissociations is expected. Figure 3 shows the O K-edge emission spectra of 1 M concentrated MgCl_{2(aq.)} solution in comparison to pure water. The splitting in A is clearly observable. The intensities of A' and A'' are roughly the same in comparison to pure water, which contradicts the characteristics of MgCl_{2(aq.)} as a structure maker, where a significant change in the emission spectra of aqueous salt solution compared to pure water is anticipated. If the SMB concept is valid, the peak A'' would increase further, since a strengthening of hydrogen bonds would lead to a higher rate of dissociations. One reason for a lack of changes in the X-ray emission spectrum might be the lower concentration of MgCl_{2(aq.)}. But this is also in contradiction to the SMB concept, since the latter is conventionally assumed to change the long-range structure of water.⁶

In case of the fine structure A' and A'' being rooted in the two structural motifs, which explains the origin of peak A'' from tetrahedral coordinated water molecules and peak A' from distorted order, the conclusion regarding the validation of SMB doesn't change. Within this framework, the lower intensities of A'' in NaCl_(aq.) can be understood by having less water molecules in tetrahedral formation due to the coordination of the salt, which is in agreement to Na⁺ being a structure breaker. The presence of Mg²⁺ as a structure maker is supposed to increase intact hydrogen bonding and thus reduce the number of distorted water molecules resulting in a significant lower intensity of A' with respect to A''. But this is not reflected in the data, see Figure 3. Noticeably, no significant influence of the MgCl₂ is observable.

By choosing the same kind of anion Cl⁻, the effect presented in Figure 2 can be fully deduced to the influence of changing the cationic species of Na⁺ and Mg²⁺ in solution. Since the number of Cl⁻ is the same for NaCl_(aq.) and MgCl_{2(aq.)}, cooperative effects can be neglected. This conclusion is compliant to former experiments which report no significant effects of Cl⁻ on the hydrogen bond network.^{25,52}

We also conclude that the Hofmeister series based on the structure maker and breaker concept is not mirrored in the electronic structure of the water molecules.

SUMMARY

We have presented the first X-ray emission spectroscopy data of two aqueous salt solutions from the Hofmeister series on the oxygen K-edge using the liquid jet method. The experimental results reflect the direct influence of salt ions on the water molecules. X-ray emission spectra of oxygen K-edge from pure water, 2 M NaCl and 1 M MgCl₂ aqueous salt solutions show significant

differences within the experimental conditions. We conclude that the presence of salt ions in aqueous solution do not cause the peak splitting at peak A and that $\text{NaCl}_{(\text{aq})}$ (i) reduces dissociations in the water structure or (ii) cause less tetrahedral coordinated water molecules, which in both cases is an effect caused by Na^+ and not by Cl^- . Additionally, this indicates a breaking of hydrogen bonds in agreement to the conventional picture of Na^+ as a structure breaker. However, no significant changes are observed in the emission spectra of the 1 M MgCl_2 solution in disagreement with the characterisation of MgCl_2 as a structure maker. The salting in / salting out of proteins in aqueous salt solutions seems not to originate in the structure maker and structure breaker concept. The microscopic understanding (the “local” view) of precipitation of macromolecules caused by salts needs further investigation.

Corresponding Author

*simone.techert@desy.de (Simone Techert)

*zhong.yin@desy.de (Zhong Yin)

Author Contributions

Z.Y. analysed the data and prepared the manuscript. Z.Y., K.K., I.R., A.P. and W.Q. performed the experiment. P.W., A.F. and S.T. designed the FlexRIXS endstation. All authors discussed the results and commented on the manuscript.

Funding Sources

This works was supported by SFB755 “Nanoscale Photonic Imaging” and SFB 1073 “Atomic Scale Control of Energy Conversion” of the German Science Foundation (DFG), the Advanced

Study Group of the Max Planck Society, Deutsches Elektronen-Synchrotron and Helmholtz Zentrum Berlin. S.T. is grateful to the Funds of the Chemical Industry. K.K. thanks the Volkswagen foundation for its support under the Peter Paul Ewald fellowship program I/85832. I.R. has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement n.°290605 (PSI- FELLOW/COFUND).

Notes

ACKNOWLEDGMENT

We thank the scientific staff at U49 for their support. Z.Y. thanks Dr. R. Kia, Dr. S. Thekku Veedu and Dr. L. Inhester for scientific discussions.

ABBREVIATIONS

XES, SMB, XAS, RIXS, LUMO

REFERENCES

- (1) Hofmeister, F. Zur Lehre von der Wirkung der Salze, *Arch. Exp. Pathol. Pharmacol.* **1888**, *24*, 247-260.
- (2) Jungwirth, P.; Tobias, D. J. Specific Ion Effects at the Air/water Interface. *Chem. Rev.* **2006**, *106*, 1259–1281.
- (3) Baldwin, R. L. How Hofmeister Ion Interactions Affect Protein Stability. *Biophys. J.* **1996**, *71*, 2056–2063.
- (4) Wilson, E.K. Hofmeister Still Mystifies, *Chemical & Engineering News* **2012**, July 16.
- (5) Lo Nostro, P.; Ninham, B. W. Hofmeister Phenomena: An Update on Ion Specificity in Biology. *Chem. Rev.* **2012**, *112*, 2286–2322.

- (6) Zhang, Y.; Cremer, P. S. Interactions between Macromolecules and Ions: The Hofmeister Series. *Curr. Opin. Chem. Biol.* **2006**, *10*, 658–663.
- (7) Tobias, D. J.; Hemminger, J. C. Getting Specific About Specific Ion Effects. *Science* **2008**, *319*, 1197–1198.
- (8) Omta, A. W.; Kropman, M. F.; Woutersen, S.; Bakker, H. J. Negligible Effect of Ions on the Hydrogen-Bond Structure in Liquid Water. *Science* **2003**, *301*, 347–349.
- (9) Obst, S.; Bradaczek, H. Molecular Dynamics Study of the Structure and Dynamics of the Hydration Shell of Alkaline and Alkaline-Earth Metal Cations. *J. Phys. Chem.* **1996**, *100*, 15677–15687.
- (10) Tongraar, A.; Liedl, K. R.; Rode, B. M. Born–Oppenheimer Ab Initio QM/MM Dynamics Simulations of Na⁺ and K⁺ in Water: From Structure Making to Structure Breaking Effects. *J. Phys. Chem. A* **1998**, *102*, 10340–10347.
- (11) Tongraar, A.; Hannongbua, S.; Rode, B. M. QM/MM MD Simulations of Iodide Ion (I⁻) in Aqueous Solution: A Delicate Balance between Ion–Water and Water–Water H-Bond Interactions. *J. Phys. Chem. A* **2010**, *114*, 4334–4339.
- (12) Xie, W. J.; Gao, Y. Q. A Simple Theory for the Hofmeister Series. *J. Phys. Chem. Lett.* **2013**, *4*, 4247–4252.
- (13) Xie, W. J.; Gao, Y. Q. Ion Cooperativity and the Effect of Salts on Polypeptide Structure – a Molecular Dynamics Study of BBA5 in Salt Solutions. *Faraday Discuss.* **2013**, *160*, 191–206.
- (14) Thomas, A. S.; Elcock, A. H. Molecular Dynamics Simulations of Hydrophobic Associations in Aqueous Salt Solutions Indicate a Connection between Water Hydrogen Bonding and the Hofmeister Effect. *J. Am. Chem. Soc.* **2007**, *129*, 14887–14898.

- (15) Lightstone, F. C.; Schwegler, E.; Hood, R. Q. A First Principles Molecular Dynamics Simulation of the Hydrated Magnesium Ion. **2001**, *343*, 549–555.
- (16) Gurney, R.W. Ionic Processes in Solution (McGraw- Hill, New York, 1953).
- (17) Marcus, Y., Ion Solvation (Wiley, Chichester, UK, 1985).
- (18) Ohtaki, H.; Radnai, T. Structure and Dynamics of Hydrated Ions. *Chem. Rev.* **1993**, *93*, 1157–1204.
- (19) Schmidt, D. A.; Funkner, S.; Born, B. P.; Gnanasekaran, R.; Schwaab, G. W.; Leitner, D. M.; Havenith, M. Rattling in the Cage : Ions as Probes of Sub-Picosecond Water. **2009**, 18512–18517.
- (20) Rønne, C.; Åstrand, P.-O.; Keiding, S. THz Spectroscopy of Liquid H₂O and D₂O. *Phys. Rev. Lett.* **1999**, *82*, 2888–2891.
- (21) Wachter, W.; Kunz, W.; Buchner, R.; Hefter, G. Is There an Anionic Hofmeister Effect on Water Dynamics ? Dielectric Spectroscopy of. *J. Phys. Chem. A* **2005**, *109*, 8675–8683.
- (22) Cappa, C. D.; Smith, J. D.; Messer, B. M.; Cohen, R. C.; Saykally, R. J. Effects of Cations on the Hydrogen Bond Network of Liquid Water: New Results from X-Ray Absorption Spectroscopy of Liquid Microjets. *J. Phys. Chem. B* **2006**, *110*, 5301–5309.
- (23) Kulik, H. J.; Marzari, N.; Correa, A. a; Prendergast, D.; Schwegler, E.; Galli, G. Local Effects in the X-Ray Absorption Spectrum of Salt Water. *J. Phys. Chem. B* **2010**, *114*, 9594–9601.
- (24) Ramos, A. S. F.; Techert, S. Influence of the Water Structure on the Acetylcholinesterase Efficiency. *Biophys. J.* **2005**, *89*, 1990–2003.

- (25) Näslund, L.-A.; Edwards, D. C.; Wernet, P.; Bergmann, U.; Ogasawara, H.; Pettersson, L. G. M.; Myneni, S.; Nilsson, A. X-Ray Absorption Spectroscopy Study of the Hydrogen Bond Network in the Bulk Water of Aqueous Solutions. *J. Phys. Chem. A* **2005**, *109*, 5995–6002.
- (26) Paterova, J.; Rembert, K. B.; Heyda, J.; Kurra, Y.; Okur, H. I.; Liu, W. R.; Hilty, C.; Cremer, P. S.; Jungwirth, P. Reversal of the Hofmeister Series : Specific Ion Effects on Peptides. *J. Phys. Chem. B* **2013**, *117*, 8150–8158.
- (27) Otten, D. E.; Shaffer, P. R.; Geissler, P. L.; Saykally, R. J. Elucidating the Mechanism of Selective Ion Adsorption to the Liquid Water Surface. *Proc. Natl. Acad. Sci.* **2012**, *109*, 3190–3190.
- (28) Marcus, Y. Effect of Ions on the Structure of Water. *Pure Appl. Chem.* **2010**, *82*, 1889–1899.
- (29) Tielrooij, K. J.; van der Post, S. T.; Hunger, J.; Bonn, M.; Bakker, H. J. Anisotropic Water Reorientation around Ions. *J. Phys. Chem. B* **2011**, *115*, 12638–12647.
- (30) Okur, H. I.; Kherb, J.; Cremer, P. S. Cations Bind Only Weakly to Amides in Aqueous Solutions. *J. Am. Chem. Soc.* **2013**, *135*, 5062–5067.
- (31) Gel'mukhanov, F.; Agren, H. Raman, Non-Raman, and Anti-Raman Dispersion in Resonant X-Ray Scattering Spectra of Molecules. *Phys. Rev. A* **1996**, *54*, 3960–3970.
- (32) de Groot, F. High-Resolution X-Ray Emission and X-Ray Absorption Spectroscopy. *Chem. Rev.* **2001**, *101*, 1779–1808.
- (33) Guo, J.-H.; Luo, Y.; Augustsson, a.; Rubensson, J.-E.; Sånthe, C.; Ågren, H.; Siegbahn, H.; Nordgren, J. X-Ray Emission Spectroscopy of Hydrogen Bonding and Electronic Structure of Liquid Water. *Phys. Rev. Lett.* **2002**, *89*, 1–4.

- (34) Nilsson, a.; Pettersson, L. G. M. Perspective on the Structure of Liquid Water. *Chem. Phys.* **2011**, *389*, 1–34.
- (35) Guo, J.-H.; Augustsson, a.; Kashtanov, S.; Spångberg, D.; Nordgren, J.; Hermansson, K.; Luo, Y. The Interaction of Cations and Liquid Water Studied by Resonant Soft-X-Ray Absorption and Emission Spectroscopy. *J. Electron Spectros. Relat. Phenomena* **2005**, *144-147*, 287–290.
- (36) Lange, K. M.; Könecke, R.; Soldatov, M.; Golnak, R.; Rubensson, J.-E.; Soldatov, A.; Aziz, E. F. On the Origin of the Hydrogen-Bond-Network Nature of Water: X-Ray Absorption and Emission Spectra of Water-Acetonitrile Mixtures. *Angew. Chemie* **2011**, *123*, 10809–10813.
- (37) Fuchs, O.; Zharnikov, M.; Weinhardt, L.; Blum, M.; Weigand, M.; Zubavichus, Y.; Bär, M.; Maier, F.; Denlinger, J.; Heske, C.; et al. Isotope and Temperature Effects in Liquid Water Probed by X-Ray Absorption and Resonant X-Ray Emission Spectroscopy. *Phys. Rev. Lett.* **2008**, *100*, 1–4.
- (38) Tokushima, T.; Harada, Y.; Takahashi, O.; Senba, Y.; Ohashi, H.; Pettersson, L. G. M.; Nilsson, A.; Shin, S. High Resolution X-Ray Emission Spectroscopy of Liquid Water: The Observation of Two Structural Motifs. *Chem. Phys. Lett.* **2008**, *460*, 387–400.
- (39) Forsberg, J.; Gråsjö, J.; Brena, B.; Nordgren, J.; Duda, L.-C.; Rubensson, J.-E. Angular Anisotropy of Resonant Inelastic Soft X-Ray Scattering from Liquid Water. *Phys. Rev. B* **2009**, *79*, 1–3.
- (40) Tokushima, T.; Horikawa, Y.; Arai, H.; Harada, Y.; Takahashi, O.; Pettersson, L. G. M.; Nilsson, A.; Shin, S. Polarization Dependent Resonant X-Ray Emission

Spectroscopy of D2O and H2O Water: Assignment of the Local Molecular Orbital Symmetry. *J. Chem. Phys.* **2012**, *136*, 044517, 1-7.

- (41) Guillemin, R.; Stolte, W. C.; Journal, L.; Carniato, S.; Piancastelli, M. N.; Lindle, D. W.; Simon, M. Angular and Dynamical Properties in Resonant Inelastic X-Ray Scattering: Case Study of Chlorine-Containing Molecules. *Phys. Rev. A* **2012**, *86*, 0134071–0134079.
- (42) Rajkovic, I.; Hallmann, J.; Grübel, S.; More, R.; Quevedo, W.; Petri, M.; Techert, S. Development of a Multipurpose Vacuum Chamber for Serial Optical and Diffraction Experiments with Free Electron Laser Radiation. *Rev. Sci. Instrum.* **2010**, *81*, 045105, 1-6.
- (43) Faubel, M.; Schlemmer, S.; Tönnies, J. P. A Molecular Beam Study of the Evaporation of Water from a Liquid Jet. *Zeitschrift für Phys. D, Atoms, Mol. Clust.* **1988**, *10*, 269–277.
- (44) Kunnus, K.; Rajkovic, I.; Schreck, S.; Quevedo, W.; Eckert, S.; Beye, M.; Suljoti, E.; Weniger, C.; Kalus, C.; Grübel, S.; et al. A Setup for Resonant Inelastic Soft X-Ray Scattering on Liquids at Free Electron Laser Light Sources. *Rev. Sci. Instrum.* **2012**, *83*, 123109, 1-8.
- (45) Forsberg, J.; Duda, L.-C.; Olsson, a; Schmitt, T.; Andersson, J.; Nordgren, J.; Hedberg, J.; Leygraf, C.; Aastrup, T.; Wallinder, D.; et al. System for in Situ Studies of Atmospheric Corrosion of Metal Films Using Soft X-Ray Spectroscopy and Quartz Crystal Microbalance. *Rev. Sci. Instrum.* **2007**, *78*, 083110, 1-7
- (46) Kortüm, G. *Lehrbuch der Elektrochemie* (Weinheim Verlag Chemie GmbH, 1952).

- (47) Nishizawa, K.; Kurahashi, N.; Sekiguchi, K.; Mizuno, T.; Ogi, Y.; Horio, T.; Oura, M.; Kosugi, N.; Suzuki, T. High-Resolution Soft X-Ray Photoelectron Spectroscopy of Liquid Water. *Phys. Chem. Chem. Phys.* **2011**, *13*, 413–417.
- (48) Odelius, M. Information Content in O[1s] K-Edge X-Ray Emission Spectroscopy of Liquid Water. *J. Phys. Chem. A* **2009**, *113*, 8176–8181.
- (49) Odelius, M. Molecular Dynamics Simulations of Fine Structure in Oxygen K-Edge X-Ray Emission Spectra of Liquid Water and Ice. *Phys. Rev. B* **2009**, *79*, 1–6.
- (50) Harada, Y.; Tokushima, T.; Horikawa, Y.; Takahashi, O.; Niwa, H.; Kobayashi, M.; Oshima, M.; Senba, Y.; Ohashi, H.; Wikfeldt, K. T.; et al. Selective Probing of the OH or OD Stretch Vibration in Liquid Water Using Resonant Inelastic Soft-X-Ray Scattering. *Phys. Rev. Lett.* **2013**, *111*, 193001, 1-5.
- (51) Marcus, Y. Effect of Ions on the Structure of Water: Structure Making and Breaking. *Chem. Rev.* **2009**, *109*, 1346–1370.
- (52) Dutkiewicz, E.; Jakubowska, A. Water Activity in Aqueous Solutions of Homogeneous Electrolytes: The Effect of Ions on the Structure of Water. *Chemphyschem* **2002**, *3*, 221–224.