

6-11-2022

## **Inari Large-Bore Mechanical Thrombectomy in Intermediate-High Risk Submassive PE Patients: Case Series and Literature Review**

Mohammad F. Mathbout

Hussam Al Hennawi

Anwar Khedr

Medical University of South Carolina

Thomas M. Todoran

Follow this and additional works at: <https://jdc.jefferson.edu/abingtonfp>



Part of the [Cardiology Commons](#)

**[Let us know how access to this document benefits you](#)**

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Abington Jefferson Health Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).



OPEN ACCESS

Images in cardiology

# Inari large-bore mechanical thrombectomy in intermediate-high risk submassive PE patients: Case series and literature review

Mohammad F. Mathbout<sup>1</sup>, Hussam Al Hennawi<sup>2\*</sup>, Anwar Khedr<sup>3</sup>, Katrina Bidwell<sup>1</sup>, Thomas M. Todoran<sup>1</sup>

## ABSTRACT

Pulmonary embolism (PE) is one of the most common causes of cardiovascular (CV) mortality worldwide. Owing to the associated morbidity and mortality with other treatment modalities, including systemic thrombolysis, a discernible change in the era of acute pulmonary embolism management has been reported. Catheter-directed thrombectomy using the FlowTrier system (Inari Medical; Irvine, CA, USA) was shown to reduce endpoints of interest in patients with acute intermediate-high risk PE and was associated with rapid hemodynamic improvement. In this report, we describe our experience with three cases of patients presenting with submassive PE, whereby immediate pulmonary artery pressure improvement was evident in all cases after successful mechanical thrombectomy. Our experience supports the use of FlowTrier mechanical thrombectomy for the treatment of submassive PE in clinical practice, with a call for further research to establish associated benefits.

<sup>1</sup> Medical University of South Carolina, Department of Cardiology, Charleston, South Carolina, USA

<sup>2</sup> Department of Internal Medicine, Jefferson Abington Hospital, Abington, PA, USA

<sup>3</sup> Department of Critical Care Medicine, Mayo Clinic Health System, Mankato, MN, USA

\*Email: hussamhennawi.md@gmail.com

<https://doi.org/10.21542/gcsp.2022.8>

Received: 22 May 2022

Accepted: 11 June 2022

© 2022 The Author(s), licensee Magdi Yacoub Institute. This is an open access article distributed under the terms of the Creative Commons Attribution license CC BY-4.0, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Mathbout MF, Al Hennawi H, Khedr A, Bidwell K, Todoran TM. Inari large-bore mechanical thrombectomy in intermediate-high risk submassive PE patients: Case series and literature review, Global Cardiology Science and Practice 2022;8 <https://doi.org/10.21542/gcsp.2022.8>

## INTRODUCTION


Followed by acute myocardial infarction and stroke, pulmonary embolism (PE) constitutes the third leading cause of cardiovascular mortality and is a significant cause of in-hospital mortality with 60,000–100,000 deaths per year in the United States<sup>1</sup>. Intermediate-risk (submassive) PE presents in up to 40% of patients characterized by hemodynamic stability with concomitant right ventricular (RV) strain or cardiac injury, or both<sup>2</sup>. Risk stratification of patients with PE is of significant importance, as patients presenting with intermediate-risk PE have up to 21% mortality at three months. Novel interventions have been introduced owing to renewed interest in optimizing PE management, especially among patients presenting with hemodynamic instability. Anticoagulation remains the cornerstone of treatment for PE<sup>3</sup>.

Treatment options can be grouped into three distinct categories: systemic thrombolysis, catheter-directed interventions, and surgical thrombectomy. Randomized clinical trials have demonstrated significant benefits for intermediate- and high-risk PE managed with systemic thrombolysis<sup>4</sup>. Nevertheless, this benefit has been countervailed by an elevated risk of major bleeding and intracranial hemorrhage, restricting its use in clinical practice<sup>4,5</sup>. Among other modalities, catheter-directed thrombolysis involves catheter-mediated direct injection of thrombolytic agents into the affected pulmonary artery. The role of FlowTrieve embolectomy and Indigo thrombectomy system has been investigated in this cohort of patients and was recently approved by the US Food and Drug Administration for clinical use<sup>6</sup>. Nevertheless, there is a lack of solid evidence to evaluate possible mortality benefits in patients undergoing catheter-based thrombectomy using the FlowTrieve system. Therefore, the application of such modality remains subjective to patients according to their risk of bleeding, location of thrombus, and operator skills until well-defined guidelines direct a unified clinical application. In this case series, we present three cases in which FlowTrieve thrombectomy was used to successfully manage submassive PE.

### Case 1

A 37-year-old male patient presented to the emergency department of an outside hospital complaining of squeezing chest pain for two days. The patient had a medical history of hypertension and an episode of PE that occurred five years ago. However, he did not undergo any hematologic workup at that time. The patient was taken off anticoagulants two years after the first PE. He worked as a forklift driver and reported sitting for long periods during his work. During the interview, he experienced an episode of syncope. He underwent a CT pulmonary angiography that revealed bilateral PE with filling defects in the right and left main pulmonary arteries and saddle embolus at the bifurcation (Figure 1A), with evidence of right heart strain confirmed by a transthoracic echocardiogram (Figure 1B). His troponin and pro-BNP levels were elevated.

The patient was admitted to the hospital for further management. In the cardiac catheterization lab, he underwent FlowTrieve thrombectomy. On the procedure table, the patient's systolic blood pressure was 174/95 mmHg, heart rate was 105 bpm, and O<sub>2</sub> saturation was 92% on 4 L/min O<sub>2</sub>. Pulmonary artery catheterization was achieved through the right femoral vein using an 8 French introducer sheath. A pigtail catheter was placed in the main PA, and angiography was performed. A near-occlusive thrombus was observed in the distal right PA. Hemodynamic measurements showed a pulmonary artery (PA) pressure of 80/40 mmHg (mPAP:50 mmHg), a right atrium (RA) pressure of 12 mmHg, and a PA O<sub>2</sub> saturation of 62.8%. A 26 French Gore Dryseal sheath (Gore, Flagstaff, AZ, USA), was introduced to advance the FlowTrieve system (Inari Medical Inc.,



**Figure 1.** Pulmonary CT angiogram showing a large thrombus burden in the distal right main PA before thrombectomy (A). Right ventricular clot burden shown by transthoracic echocardiogram (B). The transthoracic echocardiogram shows large mobile thrombi in the right ventricle. Thrombus removed from the right PA with the Inari device. C: Thrombus removed from the right PA with the Inari device. D: Final pulmonary angiogram after thrombectomy (C).

Irvine, CA, USA), which consists of a trackable 95 cm long, 24 French aspiration catheter used to retrieve the clot mechanically.


Thrombectomy of the right PA was successfully performed, with minimal blood loss and no complications. The 24 French FlowTriever catheter was advanced to the distal right PA and multiple aspirations were performed with significant visible thrombus extracted (Figure 1C). After the procedure, systolic blood pressure normalized to 124/74 mmHg, heart rate decreased to 94, and O<sub>2</sub> saturation was 98% on room air. Repeat PA pressure immediately decreased to 55/18 mmHg (mPAP: 30 mmHg). Repeat CT angiography showed no clot in the distal right PA, with near-complete restoration of pulmonary blood flow (Figure 1D).

The patient tolerated the procedure well and was switched from heparin drip to Eliquis 5 mg twice daily. He was counseled to restart his home medications, and was discharged the following day with a hematology referral to investigate a possible underlying hypercoagulability disorder.

## Case 2

A 79-year-old male patient was transferred to our hospital for advanced management of bilateral submassive PE associated with borderline hemodynamics. The patient had a medical history of coronary artery disease post-3-vessel coronary artery bypass graft, hypertension, hyperlipidemia, and obstructive sleep apnea. He also complained of two years of progressive dyspnea on exertion. However, the patient had not been on home oxygen therapy. On admission, the patient underwent chest CT with contrast using PE





**Figure 2.** Computed tomography angiogram of the chest showing extensive bilateral pulmonary embolism with near occlusive thrombus in the distal right pulmonary artery and non-occlusive thrombus in the left lower lobe and upper lobe pulmonary arteries (A). Axial cardiac view showing an enlarged right ventricle with a right ventricle/left ventricle ratio of 1.68:1, indicating right heart strain (B). Angiogram showing a near-occlusive thrombus in the distal right pulmonary artery (C). Post aspiration angiogram showing restored flow (D). A large amount of clot was extracted, with no haemodynamic instability (E).

protocol (CTPE), which showed a near-occlusive thrombus in the distal right pulmonary artery (Figure 2A) associated with flattening and leftward bowing of the interventricular septum with a right ventricle/left ventricle (RV/LV) ratio nearly 1.68:1 consistent with right heart strain. There was also a non-occlusive thrombus in the left lower lobe and upper lobe pulmonary arteries (Figure 2B).

The patient was transferred to the cardiac catheterization lab. Pulmonary artery catheterization was performed via the right femoral vein using an 8 French introducer sheath, which demonstrated a PA pressure of 60/20 mmHg (mPAP: 32 mmHg), PA O<sub>2</sub> saturation of 67.5%, RA pressure of 10 mmHg, right ventricular (RV) pressure of

60/10 mmHg, and CO/CI 3.56/1.75. A pigtail catheter was advanced to the right PA, and angiography was performed, which showed a near-occlusive thrombus in the distal right PA (Figure 2C). A 26 French Gore Dryseal sheath was introduced to accommodate the FlowTrievers system.

The distal right PA was selectively engaged with the 24 French FlowTrievers catheter, and aspiration thrombectomy was successfully performed after multiple aspirations (Figure 2D). Left-sided thrombectomy was not performed because of decreased burden of the thrombus on angiography. On the table, repeat PA pressure improved to 50/10 mmHg (mPAP:30 mmHg) A pulmonary angiogram showed near-complete restoration of pulmonary blood flow to the right and left sides (Figure 2E). Following the procedure, the patient was initially treated with 6 L of O<sub>2</sub> and was discharged on room air with an O<sub>2</sub> saturation >90%.

### Case 3


A 48-year-old female with a medical history of hypertension presented to an outside hospital complaining of progressive chest pain, shortness of breath, and weakness for two days. She had a recent prolonged hospitalization course because of COVID-19 pneumonia complicated by acute respiratory distress syndrome. She underwent intubation and venovenous extracorporeal membrane oxygenation (ECMO) followed by pulmonary rehabilitation. Initial blood pressure was 130/82 mmHg, heart rate was 116 bpm, and O<sub>2</sub> saturation was 97% on a 2L nasal cannula. Admission CTPE revealed a large bilateral saddle PE. Admission troponin was 0.1 ng/ml, and BNP was 697 pg/ml. She was started on intravenous heparin and was sent to the MUSC for further intervention. On arrival to ICU, the patient remained mildly tachycardic with a heart rate of 110 bpm. Her blood pressure was 108/83 mmHg, and O<sub>2</sub> saturation was 100% on a 3L nasal cannula. The Pulmonary Embolism Response Team was called on arrival, and the patient was transferred to the cardiac catheterization lab for further management.

Pulmonary artery catheterization was performed using an 8 French introducer sheath through the right femoral vein, demonstrating a PA pressure of 50/24 mmHg (mPAP: 30 mmHg), RA pressure of 6 mmHg, RV pressure of 50/6 mmHg, PA O<sub>2</sub> saturation of 51.7, and CO/CI of 3.54/1.62. A pigtail catheter was advanced into the right pulmonary artery, and angiography was performed. A large thrombus was observed in the distal right pulmonary artery (Figure 3A). A 24 French Inari FlowTrievers catheter was advanced to the level of the right interlobar artery and then to the level of the right lobe artery, and aspiration thrombectomy was successfully performed through both arteries (Figure 3B).

The catheter was repositioned into the left pulmonary artery and angiography was performed. A large thrombus was also observed in the distal left pulmonary artery (Figure 3C). The curved 22 French catheter was advanced to the level of the left lower lobe anterobasal and posterobasal branches, and aspiration thrombectomy was successfully performed (Figure 3D). Approximately 300 cc of blood loss was noted, with no other complications. Post-thrombectomy PA pressure was 30/12 mmHg (mPAP: 18 mmHg) and PA O<sub>2</sub> saturation was 54%. The patient was discharged on Eliquis with an O<sub>2</sub> saturation of 97% on room air.

### DISCUSSION

Although better outcomes have been associated with the use of systemic thrombolytics, the risk of emerging intracranial hemorrhage has significantly increased<sup>5</sup>. The clinical implications of local catheter-directed thrombolysis infusion, together with catheter-directed mechanical thrombectomy have expanded considerably over the past years,



**Figure 3. Right pulmonary artery (RPA) and left pulmonary artery (LPA) angiogram.** Angiogram showing large thrombus at the distal right pulmonary artery (A). Post-thrombectomy angiogram showing patency of the right interlobar artery and right lobe artery (B). An angiogram showing a large thrombus in the distal left pulmonary artery (C). Post-thrombectomy of the left lower lobe anterobasal and posterobasal branches was successfully performed (D).

owing to promising outcomes with a substantially decreased risk of intracranial bleeding compared to systemic thrombolysis<sup>7,8</sup>. This was evidenced by the FlowTrieve Pulmonary Embolectomy Clinical Study (FLARE) achieving thrombectomy via the FlowTrieve system in patients presenting with submassive PE patients; the major bleeding risk was 0.9% coupled with no reported events of intracranial hemorrhage. Moreover, FLARE depicted a significant 48 h reduction in RV/LV ratio (1.53–1.15,  $P < 0.0001$ ): an independent predictor of mortality from PE. In addition, the average post-procedural mPAP decreased significantly to 27.8 mmHg from preprocedural pressure of 29.8 mmHg ( $P < 0.0001$ )<sup>9</sup>. In our case series, patients had a considerable decrease in mPAP after the procedure, especially when performed early, which reflects the association of early thrombectomy with optimal pulmonary hypertension treatment in cases of submassive PE.

The FlowTrieve catheter system consists of a large aspiration guide catheter inserted into the PA, through which the FlowTrieve device is advanced. Three self-expanding nitinol disks are unsheathed on the FlowTrieve device to engage, disrupt, and extract the thrombus while aspirating and removing it through a 20 French guiding catheter<sup>10</sup>. In addition to PE, an Inari Flowtriver Catheter (Inari Medical Inc., Irvine, CA, USA) was used for other indications, as shown in (Table 1). This case series reflects our institution's favorable early experience with the use of Inari's novel FlowTrieve large-bore mechanical thrombectomy device for the management of submassive PE.

To date, there are no definitive guidelines regarding the use of mechanical-based thrombectomy in the management of submassive PE. The European Society of Cardiology/European Respiratory Society (ESC/ERS) recommends catheter-based thrombectomy in patients with high-risk PE, if thrombolysis fails or is contraindicated. Catheter-directed treatment in combination with ECMO may be considered if refractory

**Table 1 Overview of cases reporting the use of INARI FlowTrievers mechanical thrombectomy in different clinical implications.**

Case number	Age/Gender	Indications	Outcomes	Complications	References
1	66/F	DVT extending from the left common iliac vein to the left popliteal vein	Patent left lower extremity venous system	N/A	[11]
2	63/F	Phlegmasia Cerulea Dolens in the left lower extremity	immediate clinical resolution, including restoration of pedal pulses 45 min after thrombectomy	Minimal bleeding that resolved with manual compression	[12]
3	57/M	Right upper extremity deep venous thrombosis extending from the brachial vein to the subclavian vein	More than 90% reduction of the thrombus burden after a single pass	N/A	[13]
4	49/F	IVC thrombosis	Complete removal of the thrombus with no recurrent thrombus	N/A	[14]
5	53/M	Acute left ileo-caval DVT	Minimal remaining thrombus, complete resolution of the symptoms except mild left leg swelling with long periods of standing	N/A	[15]
6	29/F	phlegmasia cerulea dolens in the right lower extremity	Extensive thrombus burden removal after 4 passes	N/A	[16]
7	49/F	Left upper extremity DVT and central venous catheter malfunction	Minimal Clot burden with >90% patency of the brachial, axillary, and subclavian veins	Mild contrast extravasation in the proximal axillary vein treated with prolonged low-pressure balloon inflation	[17]
8	91/F	phlegmasia cerulea dolens in the left lower extremity	Nearly complete thrombus removal	N/A	[18]
9	55/F	PE in a patient on veno-arterial extracorporeal membrane oxygenation (VA-ECMO) associated with acute right heart failure	Extraction of several clots bilaterally and immediate improvement in the right ventricular function and dimension	Minimal blood loss	[19]
10	56/M	Aortic mural thrombus	Complete thrombus removal	N/A	[20]
11	68/M	Aortic and left femoropopliteal thrombi	Complete thrombi resolution	N/A	[20]
12	62/F	Left femoral and iliac DVT	Complete resolution of the occlusion after 3 passes	N/A	[21]
13	51/F	Massive PE with acute right heart failure	Removal of most of the thrombus and reestablishment of the blood flow	Persistent right ventricular failure that was managed with right ventricular mechanical support	[22]
14	75/M	sub massive PE	Removal of most of the thrombus burden after 4 passes	N/A	[23]
15	50/M	Left proximal LE DVT	Complete removal of the thrombus with home discharge after 1 day	N/A	[24]
16	88/F	Massive PE and right atrial thrombus	no residual pulmonary clot and substantially reduced right heart thrombus burden	N/A	[25]
17	37/M	Bilateral submassive PE	Near Complete restoration of the blood flow, symptomatic recovery, and post-procedural mPAP decrease to 30 mmHg from 50 mmHg pre-procedure	Minimal blood loss	Case 1 (current)
18	79/M	Bilateral submassive PE	Near Complete restoration of the blood flow, symptomatic recovery, and post-procedural mPAP decrease to 30 mmHg from 32 mmHg pre-procedure	N/A	Case 2 (current)
19	48/F	Bilateral submassive PE	Complete restoration of the blood flow, symptomatic recovery, and post-procedural mPAP decrease to 18 mmHg from 30 mmHg pre-procedure	Minimal blood loss	Case 3 (current)

circulatory collapse or cardiac arrest happens<sup>26</sup>. The American Heart Association (AHA)'s recommendation is dependent on the severity of the PE. In high-risk PE, the AHA recommends the use of interventional devices in cases of rapidly deteriorating hemodynamics to prevent mortality. However, in cases with intermediate-risk PE, the use of such devices is rationalized to hasten symptomatic recovery and prevent possible hemodynamic collapse resulting from progressive right-sided heart dysfunction. Nevertheless, no study has compared the long-term mortality benefits of large-bore mechanical thrombectomy devices in any subtype of PE<sup>2</sup>.

### What we have learned?

- Percutaneous mechanical thrombectomy utilizing the FlowTrievers system is an effective and safe option for acute intermediate-high-risk pulmonary embolism associated with a significant improvement in the RV/LV ratio and minimal major bleeding.



- The FlowTrieveer system is an effective option for mechanical thrombectomy in cases of pulmonary embolism with concomitant clots in transit.
- Early utilization of FlowTrieveer in pulmonary embolism can achieve rapid normalization of pulmonary artery pressure and SpO<sub>2</sub>.
- Mechanical thrombectomy using FlowTrieveer is a safer option for clot removal in patients with contraindications to thrombolysis.
- Further research is needed to identify the long-term clinical benefits of mechanical thrombectomy.

### Conclusion

According to our experience in this case series, the use of the FlowTrieveer device for mechanical thrombectomy in patients with intermediate-risk PE improved patient outcomes. Early intervention leads to a significant decrease in mPAP and the resolution of patient symptoms. The device was also shown to be safe with minimal bleeding complications. Our case series complement the results of the FLARE study, which demonstrated the safety and efficacy of the INARI FlowTrieveer device. Because of its promising results, it is necessary to conduct more research studies on the FlowTrieveer device to compare its effectiveness and long-term mortality benefits to anticoagulants and systemic thrombolytic therapy in this set of patients and adjust management guidelines accordingly. Moreover, the role of large-bore catheter-based mechanical thrombectomy devices in managing acute pulmonary embolism and long-term potential in preventing chronic thromboembolic pulmonary hypertension (CTEPH) needs to be further investigated.

### REFERENCES

- [1] Centers for Disease Control and Prevention (CDC). Venous thromboembolism in adult hospitalizations United States, 2007-2009. *MMWR Morb Mortal Wkly Rep*. 2012;61:401–404.
- [2] Konstantinides SV. 2014 ESC Guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J*. 2014;35:3145–3146.
- [3] Aujesky D, Roy PM, Verschuren F, Righini M, Osterwalder J, Egloff M, Renaud B, Verhamme P, Stone RA, Legall C, Sanchez O, Pugh NA, N'gako A, Cornuz J, Hugli O, Beer HJ, Perrier A, Fine MJ, Yealy DM. Outpatient versus inpatient treatment for patients with acute pulmonary embolism: an international, open-label, randomised, non-inferiority trial. *Lancet*. 2011;378:41–48.
- [4] Giri J, Sista AK, Weinberg I, Kearon C, Kumbhani DJ, Desai ND, Piazza G, Gladwin MT, Chatterjee S, Kobayashi T, Kabrhel C, Barnes GD. Interventional therapies for acute pulmonary embolism: current status and principles for the development of novel evidence: a scientific statement from the American Heart Association. *Circulation*. 2019;140:e774–801.
- [5] Meyer G, Vicaut E, Danays T, Agnelli G, Becattini C, Beyer-Westendorf J, Bluhmki E, Bouvaist H, Brenner B, Couturaud F, Dellas C, Empen K, Franca A, Galiè N, Geibel A, Goldhaber SZ, Jimenez D, Kozak M, Kupatt C, Kucher N, Lang IM, Lankeit M, Meneveau N, Pacouret G, Palazzini M, Petris A, Pruszczyk P, Rugolotto M, Salvi A, Schellong S, Sebbane M, Sobkowicz B, Stefanovic BS, Thiele H, Torbicki A, Verschuren F, Konstantinides SV, PEITHO Investigators. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med*. 2014;370:1402–1411.
- [6] Food and Drug Administration of the U.S. [https://www.accessdata.fda.gov/cdrh\\_docs/pdf19/K192833.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf19/K192833.pdf). December 20, 2019.
- [7] Kucher N, Boekstegers P, Müller OJ, Kupatt C, Beyer-Westendorf J, Heitzer T, Tebbe U, Horstkotte J, Müller R, Blessing E, Greif M, Lange P, Hoffmann RT, Werth S, Barmeyer A, Härtel D, Grünwald H, Empen K, Baumgartner I. Randomized, controlled trial of ultrasound-assisted catheter-directed thrombolysis for acute intermediate-risk pulmonary embolism. *Circulation*. 2014;129:479–486.
- [8] Piazza G, Hohlfelder B, Jaff MR, Ouriel K, Engelhardt TC, Sterling KM, Jones NJ, Gurley JC, Bhatheja R, Kennedy RJ, Goswami N, Natarajan K, Rundback J, Sadiq IR, Liu SK, Bhalla N, Raja ML, Weinstock BS, Cynamon J, Elmasri FF, Garcia MJ, Kumar M, Ayerdi J, Soukas P, Kuo W, Liu PY, Goldhaber SZ, SEATTLE II Investigators. A prospective, single-arm, multicenter trial of ultrasound-facilitated, catheter-directed, low-dose fibrinolysis for acute massive and submassive pulmonary embolism: the SEATTLE II study. *JACC Cardiovasc Interv*. 2015;8:1382–1392.
- [9] Tu T, Toma C, Tapson VF, Adams C, Jaber WA, Silver M, Khandhar S, Amin R, Weinberg M, Engelhardt T, Hunter M, Holmes D, Hoots G, Hamdalla H, Maholic RL, Lilly SM, Ouriel K, Rosenfield K,

- FLARE Investigators. A prospective, single-arm, multicenter trial of catheter-directed mechanical thrombectomy for intermediate-risk acute pulmonary embolism: the FLARE study. *JACC Cardiovasc Interv*. 2019;12:859–869.
- [10] Tukaye DN, McDaniel M, Liberman H, Burkin Y, Jaber W. Percutaneous pulmonary embolus mechanical thrombectomy. *JACC Cardiovasc Interv*. 2017;10(1):94–95. doi: 10.1016/j.jcin.2016.09.029.
- [11] Abdel-Hafez O, Salih M, Aloka F, Patel K. Intravascular ultrasound-guided mechanical thrombectomy using the INARI ClotTrier device of an iatrogenic deep venous thrombosis. *BMJ Case Rep*. 2020;13(7):e235464. Published 2020 Jul 16. doi: 10.1136/bcr-2020-235464.
- [12] Al-Hakim RA, Boscanin A, Prosser DD, Kaufman JA. Management of phlegmasia cerulea dolens with percutaneous mechanical thrombectomy. *Cardiovasc Intervent Radiol*. 2020;43(9):1398–1401. doi: 10.1007/s00270-020-02570-x.
- [13] Harmon D, Dabaja W, Qaqi O. A novel interventional approach to upper extremity swelling. *J Vasc Surg Cases Innov Tech*. 2020;6(2):209–211. Published 2020 Apr 15. doi: 10.1016/j.jvscit.2020.02.010.
- [14] Murali N, Nezami N, Latich I, Brown J, Mojibian H. Simultaneous proximal embolic protection and inferior vena cava mechanical thrombectomy using the FlowTrier system. *Diagn Interv Radiol*. 2020;26(4):345–348. doi: 10.5152/dir.2019.19458.
- [15] Crowner JR, Marston W. Percutaneous thrombectomy using a novel single-session device for acute ilio-caval deep vein thrombosis. *J Vasc Surg Cases Innov Tech*. 2019;5(3):302–304. Published 2019 Jun 29. doi: 10.1016/j.jvscit.2019.03.010.
- [16] Mouawad NJ. Effective single-session percutaneous nonpharmacologic mechanical thrombectomy for phlegmasia cerulea dolens. *J Vasc Surg Cases Innov Tech*. 2020;6(2):212–215. Published 2020 Apr 15. doi: 10.1016/j.jvscit.2020.02.008.
- [17] Agarwal S, Sosnofsky C, Saum J, Aggarwal M, Patel S. Single-session treatment of upper extremity deep venous thrombosis and central venous catheter malfunction using the ClotTrier system. *Cureus*. 2020;12(12):e12071. Published 2020 Dec 14. doi: 10.7759/cureus.12071.
- [18] Nezami N, Jarmakani H, Latich I, Groenwald M, Perez Lozada JC. Vertebroplasty-associated cement leak leading to iatrogenic venous compression and thrombosis. *J Vasc Surg Cases Innov Tech*. 2019;5(4):561–565. Published 2019 Nov 22. doi: 10.1016/j.jvscit.2019.08.009.
- [19] Stadler S, Debl K, Ritzka M, Maier LS, Sossalla S. Bail-out treatment of pulmonary embolism using a large-bore aspiration mechanical thrombectomy device [published online ahead of print, 2021 Aug 24]. *ESC Heart Fail*. 2021; doi: 10.1002/ehf2.13571.
- [20] Silverglate Q, Maldonado TS, Narula N, Garg K. A novel approach to percutaneous aortic thrombectomy. *J Vasc Surg Cases Innov Tech*. 2020;7(1):123–127. Published 2020 Dec 16. doi: 10.1016/j.jvscit.2020.12.001.
- [21] Srivastava A. Acute thrombolysis-resistant occlusive left femoral and iliac venous thrombosis treated with mechanical thrombectomy via the ClotTrier device. *Ann Vasc Surg*. 2020;65:284.e7-284.e12. doi: 10.1016/j.avsg.2019.10.097.
- [22] Agarwal S, Saum J, Chanamalou S, Cole W, Patel SM. When Harry met Sally: single-session INARI FlowTrier and Impella RP. *J Cardiol Cases*. 2020;23(2):57–60. Published 2020 Oct 22. doi: 10.1016/j.jccase.2020.10.002.
- [23] Capanegro J, Quinn E, Arndt M, Sherard D. Successful removal of a Life-threatening PE using the INARI flow triever device. *Radiol Case Rep*. 2021;16(7):1878–1881. Published 2021 May 25. doi: 10.1016/j.radcr.2021.04.045.
- [24] Quinn E, Arndt M, Capanegro J, Sherard D. Successful removal of an acute deep vein thrombosis by the INARI ClotTrier system. *Radiol Case Rep*. 2021;16(6):1433–1437. Published 2021 Apr 10. doi: 10.1016/j.radcr.2021.03.043.
- [25] Nezami N, Latich I, Murali N, Ali R, Lin BA, ShervinRad M, Pollak J, Mojibian H. Right atrial and massive pulmonary artery mechanical thrombectomy under echocardiography guidance using the FlowTrier system. *EJVES Short Rep*. 2019;45:22–25. Published 2019 Nov 7. doi: 10.1016/j.ejvssr.2019.10.001.
- [26] Konstantinides SV, Meyer G, Becattini C, Bueno H, Geersing GJ, Harjola VP, Huisman MV, Humbert M, Jennings CS, Jiménez D, Kucher N, Lang IM, Lankeit M, Lorusso R, Mazzolai L, Meneveau N, Ní Áinle F, Prandoni P, Pruszczyk P, Righini M, Torbicki A, Van Belle E, Zamorano JL, ESC Scientific Document Group. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J*. 2020;41(4):543–603. doi: 10.1093/eurheartj/ehz405.