

REVIEW ARTICLE

Adolescent ambivalence about diabetes technology—The Janus faces of automated care

Fergus J. Cameron^{1,2}  | Michael Arnold³ | John W. Gregory⁴

¹Department of Endocrinology and Diabetes, Murdoch Children's Research Institute, The Royal Children's Hospital, Melbourne, Australia

²Department of Paediatrics, The University of Melbourne, Melbourne, Australia

³School of Historical and Philosophical Studies, The University of Melbourne, Melbourne, Australia

⁴Division of Population Medicine, School of Medicine, Cardiff University, Cardiff, UK

Correspondence

Fergus J. Cameron, Department of Endocrinology and Diabetes, The Royal Children's Hospital, Murdoch Children's Research Institute; Department of Paediatrics, The University of Melbourne, Melbourne, Australia.

Email: fergus.cameron@rch.org.au

Abstract

The Janus face metaphor approach highlights that a technology may simultaneously have two opposite faces or properties with unforeseen paradoxes within human-technology interaction. Suboptimal acceptance and clinical outcomes are sometimes seen in adolescents who use diabetes-related technologies. A traditional linear techno-determinist model of technology use would ascribe these unintended outcomes to suboptimal technology, suboptimal patient behavior, or suboptimal outcome measures. This paradigm has demonstratively not been successful at universally improving clinical outcomes over the last two decades. Alternatively, the Janus face metaphor moves away from a linear techno-determinist model and focuses on the dynamic interaction of the human condition and technology. Specifically, it can be used to understand variance in adoption or successful use of diabetes-related technology and to retrospectively understand suboptimal outcomes. The Janus face metaphor also allows for a prospective exploration of potential impacts of diabetes-related technology by patients, families, and their doctors so as to anticipate and minimize potential subsequent tensions.

KEYWORDS

adolescents, diabetes, technology, uptake

1 | INTRODUCTION

In this digital age we increasingly turn to scientific and technological solutions for the adverse challenges of life. Technological panaceas though are not neutral or without countervailing impacts of their own. Technologies have emergent characteristics and properties (“affordances”) that play their part in our interactions with life-challenges.¹ These affordances are specifically designed to meet and to mitigate the challenges we face, and they may do this to varying degrees. However, there may also be perverse, ironic or paradoxical impacts of technologies. First, the benefits of technologies for one individual may be inconsistent or have unwanted and unforeseen consequences.

Second, the technological solution may change the context of the challenge and the agency or capabilities of the person challenged, thus changing the nature of the problem itself. In other words, the affordances of a particular technology may ameliorate and at the same time exacerbate the challenge, and change the nature of the issue at hand.¹⁻³

Currently, nearly three decades after the findings of the Diabetes Control and Complication Trial (DCCT) were published,⁴ many adolescent patients still struggle to achieve either a current consensus target HbA1C or even the HbA1C levels that were achieved in the DCCT.⁵⁻⁷ At the time of publication of the DCCT results, an accompanying editorial noted the highly selective nature of the DCCT cohort and asked

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Pediatric Diabetes* published by John Wiley & Sons Ltd.

the following question: “Who among the many patients with diabetes is likely to benefit from intensive therapy?”⁸ Intensive therapy now encompasses an increasing array of technologies and interventions: insulin delivery and glucose detection devices, insulin types, diabetes education strategies, and psychological support strategies⁹ and the notion of matching the patient to the therapy remains pertinent. How do adolescent patients and their health care providers best navigate through the complex array of therapeutic options to achieve optimal health outcomes?

The frequently heard cry for more and greater access to technology¹⁰ to achieve better health outcomes assumes a somewhat uncritical view of the published outcome data. Over the last two decades there has been an exponential increase in the use of diabetes-related technology (defined here as automated or semi-automated detection and treatment solutions) in pediatric and adolescent patients.^{11,12} While some groups have shown improved outcomes in metabolic markers,¹³ this is not true of all populations.^{6,14,15} Indeed, despite increased technology use, metabolic control has deteriorated in some instances.^{5,7} These variable responses in outcomes speak to variations in how the technology is used and how acceptable the technology is to various groups.¹⁶ Why would one population of adolescents embrace and excel in the use of technology yet another struggle? The obvious suspects of literacy, numeracy, and socioeconomic factors would not seem to be applicable when one is comparing population registries from countries with similarly developed health care systems in the examples cited above. Variable cultural and individual acceptance of the affordances of diabetes-related technology might instead be determinative of outcome.

The Janus face metaphor has been previously used to describe how people adapt to the various affordances offered by technological solutions.¹ In particular, the Janus face perspective highlights how some affordances may simultaneously have opposite properties and thus create inherent tensions or paradoxes for adopters of technological solutions. In the context of type 1 diabetes in adolescence, these tensions occur against a variable background of sense of self and readiness or ability to change or embrace diabetes-related technology. The purpose of this commentary is to explore tensions and paradoxes created by the Janus faced affordances of diabetes-related technology and consider how these might influence the use of such technology in adolescents with type 1 diabetes.

2 | THE JANUS FACE METAPHOR AND NOTIONS OF PARADOXICAL PROPERTIES AND AFFORDANCE TENSION

Metaphors are useful rhetorical instruments, not just for communicating a phenomenon, but also for generating analysis and understanding of the phenomenon in question.¹⁷ The metaphor of the Janus face relates to a well-known Roman god held to be responsible for transitions, for beginnings and endings, births and deaths, war and peace, leaving and arriving, the sacred and the profane. Janus is usually depicted as having two faces, and with these two faces, he is both blessed and cursed. Janus is blessed to be arriving, yet in the



FIGURE 1 Janus, the two-headed Roman god of beginnings and endings. Credit: Science History Images/Alamy

same movement, is always leaving. Janus simultaneously faces forwards and backwards (Figure 1). This is not a choice, it is his very nature.¹⁸

The Janus face metaphor is used here in an analysis which focuses on the paradoxical or perverse impacts we often observe when trying to understand why a given technology is not achieving what it was designed to achieve. Examples of the Janus face metaphor include traffic congestion often becoming worse when we build more roads; air-conditioning systems creating a hotter global environment as well as a cooler local environment; antibiotics killing bacteria but potentially creating more resistant pathogens; mobile communications devices making us equally close, but also equally distant.¹⁹ When the use of diabetes-related technologies fails to achieve desired clinical outcomes, the Janus face metaphor is again useful to deploy. A traditional linear “techno-determinist” approach (defined here as a unidimensional model of outcome being determined solely by the properties of the technology) assumes that treatment failure results from diabetes-related technologies failing to perform as designed, patients failing to perform as instructed or incorrectly defined clinical outcomes (Figure 2). The corollary of this is that clinical solutions are framed in terms of “better technologies,” “better patients,” or “redefining desired outcomes.”^{20–22} As has been previously noted, this linear approach has failed to transform clinical outcomes in pediatric diabetes over the last 10–20 years. The Janus face metaphor on the other hand, moves away from a linear techno-determinist model and focuses on the dynamic interaction of the human condition and technology. Specifically, the Janus face metaphor approach highlights unforeseen paradoxes, contrariness and ironies within this interaction. These unanticipated properties are as much an “affordance” of the technologies, and determinative of outcome, as the programmed and anticipated properties.




FIGURE 2 Linear techno-determinist model of understanding the performance of diabetes-related technology (adapted from Arnold et al.¹), with consequent reasons for suboptimal performance

3 | APPLYING THE JANUS FACE METAPHOR

To apply the Janus face metaphor to the case of technologies for diabetes one must therefore look not just at the technology and assess the extent to which its pre-determined functions are executed, but instead look at the technology's relationship to its user, and assessing the emergence of new outcomes afforded by that relationship. The effects of technologies for diabetes are not reducible to linear functions, but are emergent in a context where:

- The performance of the diabetes technologies give rise to multiple possibilities for action, at least some of which pull in opposite directions toward contrasting outcomes (Figure 3).
- These contrasting outcomes may occur on the same axis of analysis.
- Importantly, these contrasting outcomes or implications are not a result of a failure in the technology, or a failure on the part of the user of the technology, but are co-dependent and co-productive, and are intrinsic to the operation of the system.

These points imply that the emergence of possibilities when certain diabetes-related technologies are interacting with certain users in certain contexts is not determined by the technology, user or context. Put simply, the addition of diabetes technologies to a person with diabetes, does not result in the same person in the same context, plus technology. The person with diabetes has properties such as personal characteristics, capacities and desires, and the technologies impact on these. The relations are complicated by many influences in the context of the technology and user. In other words:

... technology does not answer this or that question, satisfy this or that demand, or extend this or that capacity. Rather, technology works at a more


fundamental level; it enframes the world such that the question is changed along with the answer, the need is changed along with its gratification, and direction is changed along with the mechanism. The calculator, the word processor, are not more effective, efficient or convivial methods of doing mathematics or writing – they change what it is to do mathematics and to write... (Arnold et al.,¹ p. 236)

In this way all technology, including diabetes-related technology, operates at a metaphysical level not just an instrumental level. The paradoxes, ironies, and contrariness of technology affordances can generate tension. This affordance tension may in turn lead to ambivalence or a lack of acceptance which may ultimately undermine adoption and effective use by the technology user.

4 | TECHNOLOGY AFFORDANCE TENSION IN ADOLESCENTS WITH DIABETES

When diabetes-related technology first arrived in the form of insulin pumps, their novel performances were rapidly observed by clinicians. One eminent colleague quipped in a presentation on insulin pumps at the European Society for Pediatric Endocrinology in 2004 that it was as though patients using pumps were “made of glass.” They had become transparent. There were no more secrets as most of the adolescent's diabetes related behaviors had been automatically recorded and could readily be seen in the pump download. The presenter commented on how useful this was for the clinician but also how confronting it was for many of his adolescent patients (the pump faces both ways). The use of diabetes-related technology profoundly changed the dynamic of the follow-up clinical appointment. Since that time, technological “solutions” have provided more information on patient behavior in increasingly granular and, some might argue,

Affordance tensions associated with diabetes-related technology

**FIGURE 3** Affordance tensions associated with diabetes-related technology (adapted from Arnold et al.¹)

intrusive detail. The ability to forensically dissect the circumstances of any diabetes misadventure (should the clinician wish to do so) may result in family tensions that arise from the clinical review process, when parent-trusted, adolescent self-reporting of behaviors is at odds with irrefutable contemporaneous data.²³ As technology advanced, other affordances have become increasingly underscored, relating to visibility, dependence, immediacy, privacy, and continuity of data flow. As can be seen in Figure 3, many of these affordances have paradoxical properties leading to an overall performance that is in tension with itself.

Transition to independence is one of the major developmental tasks of adolescence. In chronic disease this transition can be more problematic, whereby the burdens imposed by that disease create tensions and difficulties in forward progress.²⁴ When technology is overlaid, the use of that technology may make the adolescent less dependent on their adult care givers, but paradoxically more dependent on the technology itself.²⁵ Thus while the technology may assist the adolescent to “leave the parental nest,” it does not necessarily foster diabetes-related independence, rather it may instead simply foster a transfer of dependence.²⁶ From a parental perspective, the process of their child's independence implicitly involves trust. In a non-diabetes context this trust revolves around the adolescent being mature enough to make good choices in life. In the diabetes context which is often perceived as ever-threatening, the trust dilemma for parents is considerably heightened.²⁷ If their adolescent child adopts a diabetes-related technology option, parents now have to trust both their adolescent's choices and the “black box” of an insulin-delivery algorithm or glycaemia alarm system. Parents who have been caring for their child's diabetes for many years in an experientially informed and nuanced manner, now face the quandary of now having to transfer trust to new care-givers, their adolescent child and opaque technology systems.²⁸ This fragile transfer of trust can be either supported by good outcomes or alternatively undermined by a suboptimal responses to technology usage by their adolescent offspring or poor technology performance (e.g., inaccurate CGM readings, repeated false alarms, performance issues with consumables such as glucose sensors and pump cannulae, etc). Repeated technical failure

experiences are not uncommon^{29–32} and can lead to frustration from the adolescent trying to achieve independence or fear and frustration by the parent trying to encourage independence. Paradoxically, the very technology that might assist in independence also has affordances that may impede independence.²⁵ Arguably, the ultimate manifestation of this trust dilemma is seen in the consumer driven Open Artificial Pancreas System Project (<https://openaps.org/what-is-openaps/>), the foundations of which are built on transparent, patient controlled technology systems.

Another paradox relates to information flow. Having more diabetes data upon which to base decisions may be a useful thing for some adolescents. Yet in others a continual data stream may become a form of white noise and mitigate against engagement with and the use of those data. The diabetes data stream may be linked to alarms that paradoxically either “encourage” or “coerce” adolescent patients to engage with the technology to facilitate a treatment outcome.¹⁶ In the case of the latter, the adolescent experience of a cajoling parent may be transposed by an alarming/cajoling machine. The alignment of technology in addressing an adolescent's perceived needs is determinative in their use of it and response to it.^{33,34} Diabetes-related technologies paradoxically provide both immediate and delayed or summary data. In the context of day-to-day life, the emphasis is on the immediate data with actions having to be taken in response to glycemic perturbations to prevent clinical deterioration and calculations made around prandial insulin dosages. However, best outcomes are achieved if there is also a strategic retrospective assessment of trends and repeated outcomes over time.³⁵ In some respects this seemingly contrasting approach of immediate and delayed responses is true of all diabetes management regimens, but the dominance of real-time decision making afforded by diabetes-related technology, extends this paradox considerably.

Finally, most adolescents wish to blend in with their peer group and resist badges or restrictions that would visibly identify them as being different.³⁶ Insulin pumps theoretically do allow for less planned, more spontaneous activity allowing diabetic adolescents to blend in behaviorally. Yet insulin pumps and other devices are simultaneously present and visible, providing a “badge” that is a visual reminder that diabetic adolescents are “different.”^{26,37–39}

The uptake and acceptance of some technologies such as smart phones and sharing of personal information on digital media appear to be universal among adolescents. For the most part, these technologies facilitate knowledge access, increase ease of transactions and enhance peer group interaction. Their unifying feature is that they are seemingly controlled by the adolescent in what some have termed a “participatory surveillance” paradigm of mutuality, empowerment and sharing.⁴⁰ The performance paradoxes of such technologies are accepted because the technology performance changes the nature of the affordance in a way that is accepted and valued by the user (e.g., with a mobile phone, one is not either distant or connected, one can simultaneously be both⁴¹). On the other hand, technologies that manifestly control adolescent behavior (even if it is in the context of health promotion) appear to be less universally accepted.^{41–47} In cultures that variably value individuality over conformity the paradoxical affordances associated with diabetes-related technology may have varying impacts. For example adolescents who are used to “conforming” may be less troubled by trust in opaque automation, less concerned about the need for independence, less troubled by adherence coercion and less concerned about badges of illness. On the other hand, adolescents from cultures that emphasize individuality and freedom of choice may be more likely to respond negatively to technology automated control, coercion, and dependence. The compact of accepting technology altered affordances might be perceived to be more or less conflicted depending upon contextual cultural norms.

5 | RESOLUTION OR CONTINUANCE OF AFFORDANCE TENSION?

Technology, adolescents and contexts are continuously changing. In order of frequency, the physical and emotional changes of adolescence (monthly) are usually more rapid than iterative technology change (2–3 yearly) which is again more rapid than fluxes in social norms (generational). Psychological and social theory would indicate that in adolescence, individual and social norm changes will have the greatest impact. However, the impact of technology upon the environment (e.g., social media usage) should not be underestimated. Given this state of dynamic flux one would also expect to see associated changes in the affordance tension ascribed to the interplay of device, patient and environment. Affordance tension may resolve, continue, or increase according to individual circumstance. For example, in a diabetes context tensions around data privacy versus data visibility may change according to the adolescent's changing desire for privacy (less or more), the type of data that is shared (summarized or granular), the people with whom it is shared (parents or partners), the reasons why it is shared (requirement for licensing, insurance or occupational fitness) and the acceptance of the adolescent/young adult's peer group (social peers or work colleagues). The consequence of this ever changing dynamic is that technological solutions are far from a “set and forget” proposition. Simply put, the perceived pro's and con's and acceptance of technology use by adolescents and emerging young adults can vary over time⁴⁷ and clinicians should be prepared to

discuss changing levels of readiness for change or acceptance of the status quo.

6 | INCORPORATING THE JANUS FACE METAPHOR INTO CLINICAL PRACTICE

Affordance tensions that might impact upon individual patient engagement need to be acknowledged as part of critical agency of any adolescent engaged in decisions around technology usage.⁴⁸ All the more so when adolescents have experienced living with diabetes for some time and may have developed significant preferences for how they should manage their diabetes. Whereas there may be strongly held views by either clinicians or parents, about the likely benefits that will ensue from introducing new technology, any such plan needs full and honest discussion with the young person themselves.

The principles that underpin communication approaches for facilitating behavior change, such as motivational interviewing, provide a useful framework for embarking on such discussions. These discussions should start by trying to understand the likely benefits and challenges (pros and cons) from the young person's viewpoint of using the technology; in other words, a review of the Janus faces or affordance tensions of the technology concerned. It is important to recognize and acknowledge that a young person may have a different perspective to their parents or the healthcare professionals on these tensions. For example, as mentioned above, the greater transparency afforded by continuous glucose monitoring may be deemed unacceptably intrusive from an adolescent's perspective and concerns such as these should be discussed in full, using an empathic and non-judgmental approach which recognizes the young person's autonomy and agency. It is equally important for treating teams to be aware of their own heuristics and biases⁴⁹ and how these may impact on discussion of the potential advantages and disadvantages of various technologies.

The next step in communication, should be to identify the importance that the young person places on addressing this aspect of care and their associated confidence that they can engage with the technology in question, assessing their so-called “readiness to change.” From the perspective of the young person, if there is little importance attached to improving the aspect of care in question, it is unlikely that there will be much motivation to incorporate the technology in question and make a success of its use. In such circumstances, careers may be better advised to delay using the new technology and to initially focus upon why the issue in question is of little concern to the young person.

If the young person is still engaged with the idea of changing their self-management then the discussions need to review their confidence that they can incorporate the new technology into their care package. It is critical to finish by drawing up an agreed plan of action, including identifying training requirements and any need for ongoing support. Ideally, the young person should be in charge of decisions about both when and how the impact of the technology be judged, recognizing that from their perspective, benefit may not be entirely focused on blood glucose-related outcomes. It is also important, that

the possibility of reversing the decision to introduce the new technology at some point in the future is allowed for, should the experience of using the technology prove counter-productive from the young person's perspective. Critically, the young person themselves should be in charge of agreeing this plan of action, as this is more likely to ensure their engagement with the process. Interestingly in a trial of motivational interviewing communication, those young people with the greatest worry and lowest satisfaction with their diabetes care, experienced the greatest glycemic improvement.⁵⁰ This affirms the notion, that selection of those with the greatest pre-existing concerns about their self-care and greatest "readiness to change" in their patterns of thinking, will likely benefit most from behavior change techniques.

However, when technological interventions are considered for the newly diagnosed who have no previous experience of diabetes self-care, the scenario is slightly different. In these circumstances, young people and their families are likely to look to health-care professionals for advice on how best to manage their diabetes, given their likely previous lack of knowledge and experience. This may partly explain why the impact of interventions such as insulin pumps at diagnosis has proved less beneficial⁵¹ than suggested by some large registry studies.^{52,53} Even at diagnosis, there was a high rate of refusal of the newly diagnosed to take part in a technology trial and a high drop-out rate in those who were not randomized to their preferred treatment.⁵¹ This suggests that even in those with no prior experience of managing their diabetes, careful patient selection is still important to identify those most likely to benefit from technological support.

7 | CONCLUSION

Outcomes in type 1 diabetes remain suboptimal for many if not most adolescents. The addition of diabetes-related technology to self-care over the last two decades has been associated with variable improvements or deterioration in different cohorts. Understandably clinicians have looked for reasons and a deeper understanding of why this is occurring. The traditional linear techno-determinist model implies that the simple application of technology to a patient should lead to an improved outcome and that if this outcome is not achieved there must be a problem somewhere along the clinical line of cause and effect. This may lead to a blame paradigm where bad technology, bad patients or bad measures are variably blamed for bad outcomes. The blame paradigm eschews that in suboptimal outcome situations the technology, the patient or the outcome measure are the things that "need to change." This paradigm though, has demonstratively not been successful at universally improving clinical outcomes after many years of experience.

Clinicians need to understand that human use of technological solutions in particular contexts may generate simultaneous affordances which may be paradoxical. Exploration of these effects requires health care providers to think beyond traditional clinical paradigms. Accordingly, further research enquiry in this area of human interaction with diabetes health care technology requires broader expertise that should encompass consumers, engineers,

clinicians, psychologists, and philosophers. The Janus face metaphor describes the phenomenon of paradoxical affordances and the inherent accompanying tension. It can be used to understand variance in adoption or successful use of diabetes-related technology. In addition to retrospectively understanding suboptimal outcomes, use of the Janus face metaphor allows for a prospective exploration of potential impacts of diabetes-related technology by patients, families and their doctors so as to anticipate and minimize potential subsequent tensions. This in turn will inform a deeper understanding of readiness for change by the adolescent and whether it is timely to introduce diabetes-related technology into their overall self-care package.

AUTHOR CONTRIBUTIONS

All authors contributed to the intellectual input and writing of this manuscript.

ACKNOWLEDGEMENT

Open access publishing facilitated by The University of Melbourne, as part of the Wiley - The University of Melbourne agreement via the Council of Australian University Librarians.

CONFLICT OF INTEREST

There are no conflicts of interest to declare by any of the authors.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/pedi.13423>.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

ORCID

Fergus J. Cameron  <https://orcid.org/0000-0002-1566-7436>

REFERENCES

1. Arnold M. On the phenomenology of technology: the "Janus-faces" of mobile phones. *Inf Organ*. 2003;13:231-256.
2. Cooper S. *Technoculture and Critical Theory: In the Service of the Machine?*. Routledge; 2002.
3. Verbeek P. *What Things Do: Philosophical Reflections on Technology, Agency and Design*. Pennsylvania State University Press; 1992.
4. The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med*. 1993;329(14):977-986.
5. Foster NC, Beck RW, Miller KM, et al. State of type 1 diabetes management and outcomes from the T1D exchange in 2016-2018. *Diabetes Technol Ther*. 2019;21:66-72.
6. Bratke H, Margeirsdottir HD, Assmus J, Njølstad PR, Skriverhaug T. Does current diabetes technology improve metabolic control? A cross-sectional study on the use of insulin pumps and continuous glucose monitoring devices in a nationwide pediatric population. *Diabetes Ther*. 2021 Sep;12(9):2571-2583. doi:10.1007/s13300-021-01127-6

7. Malik FS, Sauder KA, Isom S, et al. Trends in glycemic control among youth and young adults with diabetes: the SEARCH for diabetes in youth study. *Diabetes Care*. 2022;45:285-294.
8. Lasker RD. The diabetes control and complications trial. Implications for policy and practice. *N Engl J Med*. 1993;329:1035-1036.
9. Danne T, Phillip M, Buckingham BA, et al. ISPAD Clinical Practice Consensus Guidelines 2018: insulin treatment in children and adolescents with diabetes. *Pediatr Diabetes*. 2018;19(Suppl 27):115-135.
10. Hirsch IB. Ranting in 2021: could this get any worse? *Diabetes Technol Ther*. 2021;23:157-159.
11. Cameron FJ, Wherrett DK. Care of diabetes in children and adolescents: controversies, changes, and consensus. *Lancet*. 2015;385:2096-2106.
12. Gerhardsson P, Schwandt A, Witsch M, et al. The SWEET project 10-year benchmarking in 19 countries worldwide is associated with improved HbA_{1c} and increased use of diabetes technology in youth with type 1 diabetes. *Diabetes Technol Ther*. 2021;23:491-499.
13. Cardona-Hernandez R, Schwandt A, Alkandari H, et al. Glycemic outcome associated with insulin pump and glucose sensor use in children and adolescents with type 1 diabetes. Data from the international pediatric registry SWEET. *Diabetes Care*. 2021;44:1176-1184.
14. Di Bartolo P, Nicolucci A, Cherubini V, et al. Young patients with type 1 diabetes poorly controlled and poorly compliant with self-monitoring of blood glucose: can technology help? Results of the i-NewTrend randomized clinical trial. *Acta Diabetol*. 2017;54:393-402.
15. Addala A, Auzanneau M, Miller K, et al. A decade of disparities in diabetes technology use and HbA_{1c} in pediatric type 1 diabetes: a transatlantic comparison. *Diabetes Care*. 2021;44:133-140.
16. Kubiak T, Priesterroth L, Barnard-Kelly KD. Psychosocial aspects of diabetes technology. *Diabet Med*. 2020;37:448-454.
17. Schon DA. Generative metaphor: a perspective on problem-setting in social policy. In: Ortony A, ed. *Metaphor and Thought*. Cambridge University Press; 1979.
18. Lopez-Ruiz C, ed. *Gods, Heroes and Monsters: A Sourcebook of Greek, Roman, and Near Eastern Myths in Translation*. Oxford University Press; 2014.
19. Tenner E. *Why Things Bite Back: Technology and the Revenge Effect*. Alfred A. Knof; 1996.
20. Phillip M, Battelino T, Rodriguez H, Danne T, Kaufman F, European Society for Paediatric Endocrinology; Lawson Wilkins Pediatric Endocrine Society; International Society for Pediatric and Adolescent Diabetes; American Diabetes Association; European Association for the Study of Diabetes. Use of insulin pump therapy in the pediatric age-group: consensus statement from the European Society for Paediatric Endocrinology, the Lawson Wilkins Pediatric Endocrine Society, and the International Society for Pediatric and Adolescent Diabetes, endorsed by the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care*. 2007;30:1653-1662.
21. Neylon OM, Skinner TC, O'Connell MA, Cameron FJ. A novel tool to predict youth who will show recommended usage of diabetes technologies. *Pediatr Diabetes*. 2016;17:174-183.
22. Danne T, Nimri R, Battelino T, et al. International consensus on use of continuous glucose monitoring. *Diabetes Care*. 2017;40:1631-1640.
23. Messer LH, Johnson R, Driscoll KA, Jones J. Best friend or spy: a qualitative meta-synthesis on the impact of continuous glucose monitoring on life with type 1 diabetes. *Diabet Med*. 2018;35:409-418.
24. Cameron FJ, Garvey K, Hood KK, Acerini CL, Codner E. ISPAD clinical practice consensus guidelines 2018: diabetes in adolescence. *Pediatr Diabetes*. 2018;19(Suppl 27):250-261.
25. Lupton D. How does health feel? Towards research on the affective atmospheres of digital health. *Digit Health*. 2017;3:1-11.
26. Pals RAS, Hviid P, Cleal B, Grabowski D. Demanding devices—living with diabetes devices as a pre-teen. *Soc Sci Med*. 2021;286:114279.
27. Spencer JE, Cooper HC, Milton B. The lived experiences of young people (13–16 years) with type 1 diabetes mellitus and their parents—a qualitative phenomenological study. *Diabet Med*. 2013;30:e17-e24.
28. Naranjo D, Suttiratana SC, Iturralde E, et al. What end users and stakeholders want from automated insulin delivery systems. *Diabetes Care*. 2017;40:1453-1461.
29. Wheeler BJ, Heels K, Donaghue KC, Reith DM, Ambler GR. Insulin pump-associated adverse events in children and adolescents—a prospective study. *Diabetes Technol Ther*. 2014;16:558-562.
30. Ross PL, Milburn J, Reith DM, Wiltshire E, Wheeler BJ. Clinical review: insulin pump-associated adverse events in adults and children. *Acta Diabetol*. 2015;52:1017-1024.
31. Rabbone I, Minuto N, Bonfanti R, et al. Italian Paediatric Pump Failure Study Group. Insulin pump failures in Italian children with type 1 diabetes: retrospective 1-year cohort study. *Diabet Med*. 2017;34:621-624.
32. MAUDE data 2020, <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfmaude/search.cfm>
33. Konijn EA, Veldhuis J, Plaisier XS, Spekman M, den Hamer A. Adolescent development and psychological mechanisms in interactive media use. In: Sundar SS, ed. *The Handbook of the Psychology of Communication Technology*. 1st ed. John Wiley & Sons, Inc; 2015:332-364.
34. Shapka JD. Adolescent technology engagement: it is more complicated than lack of self control. *Hum Behav Emerg Tech*. 2019;1:103-110.
35. Pihoker C, Forsander G, Fantahun B, et al. ISPAD Clinical Practice Consensus Guidelines 2018: the delivery of ambulatory diabetes care to children and adolescents with diabetes. *Pediatr Diabetes*. 2018;19(Suppl 27):84-104.
36. Patterson JM, Garwick AW. Coping with chronic illness. A family systems perspective on living with diabetes. *Diabetes and the Adolescent*. Miranova Publishers; 1998.
37. Ritholz MD, Atakov-Castillo A, Beste M, et al. Psychosocial factors associated with use of continuous glucose monitoring. *Diabet Med*. 2010;27:1060-1065.
38. Tanenbaum ML, Adams RN, Hanes SJ, et al. Optimal use of diabetes devices: clinician perspectives on barriers and adherence to device use. *J Diabetes Sci Technol*. 2017;11:484-492.
39. Boucher S, Blackwell M, Galland B, et al. Initial experiences of adolescents and young adults with type 1 diabetes and high-risk glycemic control after starting flash glucose monitoring—a qualitative study. *J Diabetes Metab Disord*. 2019;19:37-46.
40. Albrechtslund A. Online social networking as participatory surveillance. *First Monday*. 2008;13(3). <https://firstmonday.org/ojs/index.php/fm/article/view/2142>
41. Diabetes Research in Children Network Study Group, Weinzimer S, Xing D, et al. Prolonged use of continuous glucose monitors in children with type 1 diabetes on continuous subcutaneous insulin infusion or intensive multiple-daily injection therapy. *Pediatr Diabetes*. 2009;10:91-96.
42. Ramchandani N, Arya S, Ten S, Bhandari S. Real-life utilization of real-time continuous glucose monitoring: the complete picture. *J Diabetes Sci Technol*. 2011;5:860-870.
43. Wong JC, Foster NC, Maahs DM, et al. Real-time continuous glucose monitoring among participants in the T1D exchange clinic registry. *Diabetes Care*. 2014;37:2702-2709.
44. Battelino T, Liabat S, Veeze HJ, Castañeda J, Arrieta A, Cohen O. Routine use of continuous glucose monitoring in 10 501 people with diabetes mellitus. *Diabet Med*. 2015;32:1568-1574.
45. Farfel A, Liberman A, Yackobovitch-Gavan M, Phillip M, Nimri R. Executive functions and adherence to continuous glucose monitoring in children and adolescents with type 1 diabetes. *Diabetes Technol Ther*. 2020;22:265-270.

46. Šumník Z, Pavlíková M, Pomahačová R, et al. Use of continuous glucose monitoring and its association with type 1 diabetes control in children over the first 3 years of reimbursement approval: population data from the ČENDA registry. *Pediatr Diabetes*. 2021;22:439-447.
47. Weinstein JM, Kahkoska AR. Association of continuous glucose monitoring use and hemoglobin A1c levels across the lifespan among individuals with type 1 diabetes in the US. *JAMA Netw Open*. 2022;5:e2223942.
48. Rashotte J, Tousignant K, Richardson C, et al. Living with sensor-augmented pump therapy in type 1 diabetes: adolescents' and parents' search for harmony. *Can J Diabetes*. 2014;38:256-262.
49. Cameron FJ, Moore B, Gillam L. Two's company, is three a crowd? Ethical cognition in decision making and the role of industry third parties in pediatric diabetes care. *Pediatr Diabetes*. 2019;20(1):15-22.
50. Channon SJ, Huws-Thomas MV, Rollnick S, et al. A multicenter randomized controlled trial of motivational interviewing in teenagers with diabetes. *Diabetes Care*. 2007;30:1390-1395.
51. Blair JC, McKay A, Ridyard C, et al. Continuous subcutaneous insulin infusion versus multiple daily injection regimens in children and young people at diagnosis of type 1 diabetes: pragmatic randomised controlled trial and economic evaluation. *BMJ*. 2019;365:l1226.
52. Szypowska A, Schwandt A, Svensson J, et al. Insulin pump therapy in children with type 1 diabetes: analysis of data from the SWEET registry. *Pediatr Diabetes*. 2016;17(Suppl 23):38-45.
53. Sherr JL, Hermann JM, Campbell F, et al. Use of insulin pump therapy in children and adolescents with type 1 diabetes and its impact on metabolic control: comparison of results from three large, transatlantic paediatric registries. *Diabetologia*. 2016;59:87-91.

How to cite this article: Cameron FJ, Arnold M, Gregory JW. Adolescent ambivalence about diabetes technology—The Janus faces of automated care. *Pediatr Diabetes*. 2022;1-8. doi:[10.1111/pedi.13423](https://doi.org/10.1111/pedi.13423)