

Heterocycles

How to cite: *Angew. Chem. Int. Ed.* **2021**, *60*, 18882–18887

International Edition: doi.org/10.1002/anie.202107495

German Edition: doi.org/10.1002/ange.202107495


Sydnone Methides—A Forgotten Class of Mesoionic Compounds for the Generation of Anionic N-Heterocyclic Carbenes

Sebastian Mummel, Felix Lederle, Eike G. Hübner, Jan C. Namyslo, Martin Nieger, and Andreas Schmidt*

Abstract: Sydnone methides are described from which only one single example has been mentioned in the literature so far. Their deprotonation gave anions which can be formulated as π -electron rich anionic N-heterocyclic carbenes. Sulfur and selenium adducts were stabilized as their methyl ethers, and mercury, gold as well as rhodium complexes of the sydnone methide carbenes were prepared. Sydnone methide anions also undergo C–C coupling reactions with 1-fluoro-4-iodobenzene under Pd(PPh₃)₄ and CuBr catalysis. ⁷⁷Se NMR resonance frequencies and ¹J_{C–Se} as well as ¹J_{C–H} coupling constants have been determined to gain knowledge about the electronic properties of the anionic N-heterocyclic carbenes. The carbene carbon atom of the sydnone methide anion **3j** resonates at $\delta = 155.2$ ppm in ¹³C NMR spectroscopy at –40°C which is extremely shifted upfield in comparison to classical N-heterocyclic carbenes.

Undoubtedly, sydnones are the best known mesoionic compounds. They are widely applied as masked nitrile imines in 1,3-dipolar cycloadditions (“Huisgen reactions”^[1]) which proceed in a copper-catalyzed,^[2] metal-free^[3] or metal-free strain-promoted click fashion^[4,5] with sub-millisecond intermediates,^[6] as summarized in numerous review articles and compilations about [2+3]-cycloadditions.^[7] Recent developments include the synergistic combination of organocatalysis and visible-light photocatalysis of cycloadditions of sydnones to form pyrazoles.^[8] Apart from cycloadditions, sydnones are versatile precursors for nucleophilic radiofluorination for the

synthesis of [¹⁸F]fluoroarenes,^[9] and co-catalysts or ligands of Pd catalysts for Suzuki–Miyaura reactions under acidic conditions.^[10] They are of ongoing interest as biologically active compounds^[11] from which the sydnone imine^[12] Molsidomine^[13] and Sydnocarb^[14] are in clinical use.^[15] Recently they have been applied for in vitro bioconjugation of purified proteins,^[5] as sialic acid-substituted bioorthogonal reporters,^[4] as imaging reagents,^[16] labeling and imaging tools in live cells,^[17] and fluorogenic compounds.^[18] In view of this remarkable career it is surprising that some members of the sydnone family of compounds (Scheme 1) have remained almost unknown to date. Thus, only one sydnone methide (sydnone methanide) (**2a**) has been prepared,^[19] examined by ¹⁷O,^[20] ¹⁵N, and ¹⁴N NMR spectroscopy,^[21] and repeatedly mentioned in monographs.^[15]




Scheme 1. Some members of the sydnone family of compounds.


Our interest in mesomeric betaines^[22] and their relationship to neutral^[12a,b,23,24] as well as anionic N-heterocyclic carbenes^[25] in combination with the introduction of an energy-based quantitative index of the ease of NHC formation from precursors by deprotonation (carbene relative energy of formation, CREF)^[26] stimulated initial calculations which suggested that sydnone methides are interesting precursors for the generation of unique π -electron-rich anionic N-heterocyclic carbenes. This index supplements other measures of σ -donor and/or π -properties of NHCs such as molecular electrostatic potentials (MESP),^[27] computationally derived ligand electronic parameters (CEP),^[28] Tolman parameters (TEP),^[29] HOMO energies,^[30,31] calculated proton affinities,^[30,32] ¹J_{CH} and ¹J_{CSe} coupling constants^[33] of precursor salts and Se adducts, respectively, and Huynh electronic parameters (HEP)^[34] of Pd complexes (review^[35]). We calculated the CREF value of sydnone methide **2a** to be 0.534 (B3LYP/6-311++G**) which seemed to be a promising value in comparison to other mesomeric betaines as carbene precursors like 1,3-dimethylimidazolium-4-olate and its 4-aminide, and 1,3-dimethyl-6-oxo-pyrimidinium-4-olate (CREFs = 0.576, 0.557, 0.547, resp.).^[24,25] We therefore report here on the syntheses of a series of sydnone methides, their

[*] M. Sc. S. Mummel, Dr. F. Lederle, Prof. Dr. E. G. Hübner, Dr. J. C. Namyslo, Prof. Dr. A. Schmidt
 Clausthal University of Technology, Institute of Organic Chemistry
 Leibnizstrasse 6, D-38678 Clausthal-Zellerfeld (Germany)
 E-mail: schmidt@ioc.tu-clausthal.de

Dr. F. Lederle, Prof. Dr. E. G. Hübner
 Fraunhofer Heinrich Hertz Institute HHI, Fiber Optical Sensor
 Systems
 Am Stollen 19H, D-38640 Goslar (Germany)


Dr. M. Nieger
 University of Helsinki, Department of Chemistry
 P.O. Box 55, FIN-00014 Helsinki (Finland)

 Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202107495>.

 © 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

conversion into anionic NHCs by deprotonation, and trapping reactions with sulfur, selenium, C-electrophiles, mercury, gold, and rhodium as well as C–C coupling reactions under Pd⁰/Cu^I-catalysis.


We first developed a reliable synthetic method which lead to a series of sydnone methides **2a–l** (Scheme 2, Table 1). Thus, we started from the sydnones **1a–l** which we treated with Tf₂O to obtain a mixture of inseparable bis-sydnone ethers and sydnone 5-triflates which proved to be very sensitive towards minute traces of water. Trapping with in situ generated malodinitrile anions for the synthesis of **2b–d**, methyl 2-cyanoacetate anions for **2e–h**, and 2-(methylsulfonyl)acetonitrile anions for the preparation of **2i–l** gave the desired sydnone methides, respectively. These compounds are stable, brilliant yellow to orange in color and slightly fluorescent (see Supporting Information for fluorescence spectra). Sydnone methides can be represented by several



Scheme 2. Syntheses of sydnone methides **2a–l** and generation of anionic N-heterocyclic carbenes **3a–l** derived thereof (for substitution patterns, see Table 1). Selected mesomeric structures **A–C**. Several Li species detectable by ⁷Li NMR due to rapid decomposition of the anions.

Table 1: Substitution pattern and numbering of the sydnone methides **2a–l**

Compd.	Ar	Z	Yield
2a	Ph	CN	53 %
2b	4-C ₆ H ₄ Me	CN	38 %
2c	4-C ₆ H ₄ OMe	CN	28 %
2d	4-C ₆ H ₄ Cl	CN	33 %
2e	Ph	COOMe	61 %
2f	4-C ₆ H ₄ Me	COOMe	65 %
2g	4-C ₆ H ₄ OMe	COOMe	71 %
2h	4-C ₆ H ₄ Cl	COOMe	17 %
2i	Ph	SO ₂ Me	63 %
2j	C ₆ H ₄ Me	SO ₂ Me	37 %
2k	4-C ₆ H ₄ OMe	SO ₂ Me	12 %
2l	4-C ₆ H ₄ Cl	SO ₂ Me	9 %



resonance structures three of which are shown in Scheme 2. It is interesting to note that carbon atom C4 is a site of negative charge according to the rules of resonance as indicated by mesomeric structure **2a-IB**, although it is a site of deprotonation for the formation of anionic N-heterocyclic carbenes. Correspondingly, the C4 carbon atoms resonate at high field between $\delta = 108.2$ ppm (**2j**) and 110.9 ppm (**2f**) in ¹³C NMR spectroscopy. Further DFT calculations predict that the *E* isomer of **2e** is by $\Delta G_{\text{vac}} = 4.6$ kJ mol⁻¹ more stable in vacuo than the corresponding *Z* isomer and that the rotation barrier is $\Delta E_{\text{vac}} = 82.7$ kJ mol⁻¹ (PBE0-d3/ 6-31G**). We also calculated the solvent dependence. In THF, dichloromethane (DCM) and DMSO the rotation barriers decrease to $\Delta E_{\text{THF}} = 65.8$ kJ mol⁻¹, $\Delta E_{\text{DCM}} = 65.0$ kJ mol⁻¹, and $\Delta E_{\text{DMSO}} = 61.6$ kJ mol⁻¹, respectively (see Supporting Information). These results are in agreement with the fact that the bonds C5–C6 of **2e** and **2i** [crystallographic numbering, Figure 1] display a considerable double bond character, as determined by single crystal X-ray analyses. As N2 resonates considerably more upfield in ¹⁵N NMR spectroscopy than N3, the mesomeric structures **2a-IC** can be identified as the most suitable representation for sydnone methides. In the elemental cells, the phenyl ring is twisted by $-52.65(17)^\circ$ (**2e**) and $-12.74(17)^\circ$ (**2i**) out of the plane of the sydnone rings. As determined by HMBC measurements, the ¹J_{C4-H} coupling constants of **2a,e,i** are 214 Hz, 220 Hz, and 219 Hz, respectively, and these values are between those of 1,3-dimesitylimidazolium and 1,3-dimesitylimidazolidinium (¹J_{CH} = 225 Hz and 206 Hz) as precursors of normal N-heterocyclic carbenes, respectively, thus indicating strong σ -donor capacities of the




Figure 1. Molecular drawing of sydnone methide **2e** (above) and **2i** (below) (displacement parameters are drawn at 30% (**2e**) and 50% (**2i**) probability level). Selected bond lengths of **2e** [pm] (crystallographic numbering): N3–C4: 134.12(15), C4–C5: 138.91(15), C5–C6: 139.87(16), C6–C7: 143.83(17), C7–O7: 121.05(15), C9–N9: 114.51(18) pm. Selected torsion angles of **2e** [°]: N2–N3–C10–C15: 128.96(13), C4–C5–C6–C7: $-1.8(2)^\circ$. Selected bond lengths of **2i** [pm]: N3–C4: 135.00(17), C4–C5: 138.10(17), C5–C6: 139.80(18), C6–S7: 172.16(13), S7–O9: 144.04(10), C11–N12: 115.02(18) pm. Selected torsion angles of **2i** [°]: N2–N3–C13–C18: $-12.74(17)^\circ$, C4–C5–C6–S7: $11.6(2)^\circ$.^[46]

corresponding N-heterocyclic carbenes.^[33] We next performed a base screening which revealed that deprotonation of the sydnone methides **2a–l** can best be accomplished by LiHMDS in THF at -10°C . According to the rules of resonance, the resulting anions **3a–l** can be represented by a number of canonical forms, among those the mesomeric structures of abnormal N-heterocyclic carbenes **3a-1A**, as structures possessing two formal negative charges at C4 **3a-1B**, and as normal anionic N-heterocyclic carbenes **3a-1C** (Scheme 2).

Although the anions **3a–l** are unstable and decompose rapidly even at low temperatures, we successfully generated **3j** quantitatively at -50°C and immediately measured NMR spectra at -40°C . The ^1H NMR spectra show the absence of the proton at C4. In the ^{13}C NMR spectra the signal of C4 of the precursor shifted considerably from 108.2 ppm (**2j**) to 155.2 ppm (**3j**). The carbene's resonance frequency is thus extremely shifted upfield in comparison to other NHCs. All chemical shift differences are summarized in Table S2 (Supporting Information). Moreover, the mass of the sydnone methide anion **3j** was confirmed by high resolution electro-spray ionization mass spectrometry in the anion detection mode on spraying a cooled in situ prepared sample of **3j**. To gain insight into the electronic properties of the sydnone methide anions we performed DFT calculations (B3LYP/6-311++G**). In contrast to 1,3-dimesitylimidazol-2-ylidene and 1,3-dimesitylimidazolidin-2-ylidene which we chose as examples, the highest occupied molecular orbitals (HOMOs) of the sydnone methide carbenes **3a**, **3e** and **3i** are π -orbitals with significant atomic orbital coefficients on C4 as legacy of their origin from mesoionic compounds (Figure 2 and Supporting Information). Vice versa, the HOMO-1 display the characteristic geometries of N-heterocyclic carbenes. Their energies are considerably higher than those of the aforementioned N-heterocyclic carbenes of imidazole, but slightly lower than those of N-phenylsydnone (Figure 3). Concerning the substituent effects, the different electron-withdrawing capacities of the COOMe group (Hammett constants^[36] $\sigma_m = 0.37$; $\sigma_p = 0.45$), CN group ($\sigma_m = 0.56$; $\sigma_p = 0.66$) and SO_2Me group ($\sigma_m = 0.60$; $\sigma_p = 0.72$) correlates with the HOMO energies within the series of sydnone methide anions.

Reaction of the in situ generated anionic sydnone methide carbenes **3a,e,i** with sulfur and selenium gave the thioethers **4a–c** and selenium ethers **5a–c** after methylation as stable compounds, respectively (Scheme 3). The ^{77}Se NMR reso-




Figure 2. Calculated frontier orbital profile of the anion of sydnone methide **3a**.





Figure 3. Comparison of selected molecular orbital energies in vacuo of imidazole-2-ylidene, imidazolidin-2-ylidene (left) and members of the sydnone family of carbenes (right) which are π -electron-rich N-heterocyclic carbenes calculated on B3LYP/6-311++G** level. Pictures of selected orbitals and HOMO/LUMO energies in THF are presented in the Supporting Information (see Table S1).




Scheme 3. Chemistry of sydnone methide anionic carbenes with sulfur, selenium, and acyl chlorides, respectively.

nance frequencies of the selenium ethers **5a–c** were detected at $\delta = 86$ ppm, 101 ppm and 109 ppm, respectively, and these values correspond to those measured for Molsidomine [that is, N-(ethoxycarbonyl)-3-(4-morpholino)sydnone imine; $\delta = 97$ ppm].^[37] As expected they are considerably more upfield than those of the selenium ethers of 1,3-dimesitylimidazol-2-ylidene and its imidazolidine derivative which are cations and which resonate at $\delta = 199$ ppm and 271 ppm, respectively. Treatment of **3a,e,i** with acetyl chloride (R = Me) and benzoyl chloride (R = Ph) gave **6a–e** which slowly reconstitute the corresponding sydnone methides on exposure to water. The acetyl derivative of **6f** (Z = SO_2Me , R = Me) could not be isolated as it is unstable under ambient conditions. These reactions correspond to rare trapping reactions of normal^[38,39] as well as abnormal imidazolylenes with acyl chlorides.^[39] Mesoionic compounds such as 1,3-dimesitylimidazolium-4-olates, however, undergo electrophilic heteroaromatic substitutions to give these structures.^[40]

The reaction of **3e,f** with 0.5 equiv of mercury(II)-chloride lead to the predominant formation of the dimeric mercury complexes **7₂(a,b)**, whereas 3–4 equiv of HgCl_2 yielded the monomeric complexes **7₁(a,b)** from **3b,f**.

Chloro(triphenylphosphine)gold(I) converted the sydnone methide carbenes **3c,f,i** into the gold(I) complexes **8a–c** as yellow solids (Scheme 4). We also reacted the sydnone methide carbenes **3a,f,i** with RhCO(PPh₃)Cl which resulted in the formation of the rhodium complexes **9a–c** as pale yellow solids.



Scheme 4. Trapping of the sydnone methide anionic carbenes with mercury, gold, and rhodium complexes, respectively.

Single crystals of the gold complex **8a** were subjected to an X-ray analysis (Figure 4). The results show that complex formation does not influence the starting mesoion's geometry (**2a**) significantly.

Concerning C–C coupling reactions, treatment of **3a,e,i** with CuBr gave in situ generated, non-isolable copper compounds, which coupled with 4-fluoro-1-iodobenzene in the presence of Pd(PPh₃)₄ in acceptable yields (Scheme 5). These reactions proceed in analogy to those of sydnone.^[41]

In summary, we present the syntheses, spectroscopic characterizations, results of calculations^[42,43] and single crystal





Figure 4. Molecular drawing of gold complex **8a** (displacement parameters are drawn at 30% probability level). Selected bond lengths [pm] (crystallographic numbering): N3–C4: 136.2(3), C4–C5: 138.7(3), C5–C6: 140.1(3), C6–C7: 141.3(3), C4–Au1: 204.07(19), C8–N8: 113.7(3) pm. Selected bonding angles [°]: N2–N3–C4: 116.39(17), N3–C4–C5: 102.07(17), C4–C5–C6: 134.7(2)°. Selected torsion angles [°]: N2–N3–C9–C10: 46.6(3), C4–C5–C6–C7: 4.7(4)°.^[46]



Scheme 5. Copper-catalysed C–C coupling reactions.

X-ray analyses^[44,45] of new stable members of the substance class of sydnone methides from which only one single example had been described in 1984. Deprotonation yielded sydnone methide anions which can be formulated by a number of resonance forms, among those mesomeric structures of anionic N-heterocyclic carbenes. The frontier orbital profile sets these N-heterocyclic carbenes apart from other examples of this class of compounds, as their highest occupied molecular orbitals are π -orbitals with considerable atomic orbital coefficients on the carbene carbon atom, and extremely upfield shifted ¹³C NMR resonance frequencies of the carbene carbon atom (**3j**: $\delta = 155.2$ ppm). The sydnone methide anions can be reacted with sulfur and selenium, respectively, and stabilized by S- and Se-methylation. Reaction with acyl chlorides gave sydnone methide ketones, and trapping reactions with mercury, gold, and rhodium gave the corresponding complexes. Finally, we presented a Pd⁰/Cu^I-catalyzed C–C coupling reaction at C4 of the sydnone methide. Our results supplement the knowledge about the sydnone family of compounds as well as about anionic N-heterocyclic carbenes.

Acknowledgements

The Deutsche Forschungsgemeinschaft (DFG) is acknowledged for financial support. Open access funding enabled and organized by Projekt DEAL.

Conflict of Interest

The authors declare no conflict of interest.

Keywords: betaines · carbenes · heterocycles · rhodium · selenium

[1] M. Breugst, H.-U. Reissig, *Angew. Chem. Int. Ed.* **2020**, *59*, 12293–12307; *Angew. Chem.* **2020**, *132*, 12389–12404.

[2] a) H. Liu, D. Audisio, L. Plougastel, E. Decuypere, D.-A. Buisson, O. Koniev, S. Kolodych, A. Wagner, M. Elhabiri, A. Krzyczmonik, S. Forsback, O. Solin, V. Gouverneur, F. Taran, *Angew. Chem. Int. Ed.* **2016**, *55*, 12073–12077; *Angew. Chem.* **2016**, *128*, 12252–12256; b) S. Kolodych, E. Rasolofonjatovo, M. Chaumontet, M.-C. Nevers, C. Créminon, F. Taran, *Angew. Chem. Int. Ed.* **2013**, *52*, 12056–12060; *Angew. Chem.* **2013**, *125*, 12278–12282.

- [3] a) D. L. Browne, M. D. Helm, A. Plant, J. P. A. Harrity, *Angew. Chem. Int. Ed.* **2007**, *46*, 8656–8658; *Angew. Chem.* **2007**, *119*, 8810–8812; b) M. M. Popa, S. Shova, M. Hrubaru, L. Barbu, C. Draghici, F. Dumitrescu, D. E. Dumitrescu, *RSC Adv.* **2020**, *10*, 15656–15664; c) M. N. Kumbar, S. K. J. Shaikh, R. R. Kamble, P. K. Bayannavar, A. A. Kamble, R. K. Hunnur, *ACS Omega* **2019**, *4*, 4955–4962; d) V. Hladíková, J. Vana, J. Hanusek, *Beilstein J. Org. Chem.* **2018**, *14*, 1317–1348; e) H. M. Tao, F. Liu, R. X. Zeng, Z. Z. Shao, L. F. Zou, Y. Cao, J. M. Murphy, K. N. Houk, Y. Liang, *Chem. Commun.* **2018**, *54*, 5082–5085.
- [4] Z. S. Chinoy, C. Bodineau, C. Favre, K. W. Moremen, R. V. Durán, F. Friscourt, *Angew. Chem. Int. Ed.* **2019**, *58*, 4281–4285; *Angew. Chem.* **2019**, *131*, 4325–4329.
- [5] a) C. Favre, L. deCremoux, J. Badaut, F. Friscourt, *J. Org. Chem.* **2018**, *83*, 2058–2066; b) S. Bernard, D. Audisio, M. Riomet, S. Bregant, A. Sallustrau, L. Plougastel, E. Decuypere, S. Gabillet, R. A. Kumar, J. Elyian, M. N. Trinh, O. Koniev, A. Wagner, S. Kolodych, F. Taran, *Angew. Chem. Int. Ed.* **2017**, *56*, 15612–15616; *Angew. Chem.* **2017**, *129*, 15818–15822; c) M. K. Narayanan, Y. Liang, K. N. Houk, J. M. Murphy, *Chem. Sci.* **2016**, *7*, 1257–1261; d) S. Wallace, J. W. Chin, *Chem. Sci.* **2014**, *5*, 1742–1744.
- [6] Y. Qing, G. S. Pulcu, N. A. W. Bell, H. Bayley, *Angew. Chem. Int. Ed.* **2018**, *57*, 1218–1221; *Angew. Chem.* **2018**, *130*, 1232–1235.
- [7] a) F. de Moliner, N. Kielland, R. Lavilla, M. Vendrell, *Angew. Chem. Int. Ed.* **2017**, *56*, 3758–3769; *Angew. Chem.* **2017**, *129*, 3812–3823; b) F. Albota, M. D. Stanescu, *Rev. Roum. Chim.* **2017**, *62*, 711–734; c) D. L. Browne, J. P. Harrity, *Tetrahedron* **2010**, *66*, 553–568; d) *1,3-Dipolar Cycloaddition Chemistry, Vol. 1 and 2*, (Ed.: A. Padwa), Wiley, New York, **1984**; e) *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products* (Eds.: A. Padwa, W. H. Pearson), Wiley, New York, **2002**; f) R. Huisgen, in *1,3-Dipolar Cycloaddition Chemistry, Vol. 1*, (Ed.: A. Padwa), Wiley, New York, **1984**, pp. 1–176.
- [8] C. P. Lakeland, D. W. Watson, J. P. A. Harrity, *Chem. Eur. J.* **2020**, *26*, 155–159.
- [9] M. K. Narayanan, G. Ma, P. A. Champagne, K. N. Houk, J. M. Murphy, *Angew. Chem. Int. Ed.* **2017**, *56*, 13006–13010; *Angew. Chem.* **2017**, *129*, 13186–13190.
- [10] a) L. Pruschinski, A.-L. Lücke, T. Freese, S.-R. Kahnert, S. Mummel, A. Schmidt, *Synthesis* **2020**, *6*, 882–892; b) S. Wiechmann, T. Freese, M. H. H. Drafz, E. G. Hübner, J. C. Namyslo, M. Nieger, A. Schmidt, *Chem. Commun.* **2014**, *50*, 11822–11824.
- [11] a) J. Shum, P. Z. Zhang, L. C.-C. Lee, K. K.-W. Lo, *ChemPlusChem* **2020**, *85*, 1374–1378; b) A. Dorababu, R. R. Kamble, R. Ravindra, S. K. J. Shaikh, S. M. Somagond, P. K. Bayannavar, S. D. Joshi, *J. Heterocycl. Chem.* **2019**, *56*, 2430–2441; c) N. O. Corneli, F. Albota, C. C. Dragomirescu, M. D. Stanescu, *Farmacia* **2018**, *66*, 658–662.
- [12] a) T. Freese, J. C. Namyslo, M. Nieger, A. Schmidt, *RSC Adv.* **2019**, *9*, 4781–4788; b) T. Freese, M. Nieger, J. C. Namyslo, A. Schmidt, *Tetrahedron Lett.* **2019**, *60*, 1272–1276; c) T. Freese, A.-L. Lücke, J. C. Namyslo, M. Nieger, A. Schmidt, *Eur. J. Org. Chem.* **2018**, 1646–1654; d) A. S. Samarskaya, I. A. Cherepanov, I. A. Godovikov, A. O. Dmitrienko, S. K. Moiseev, V. N. Kalinin, E. Hey-Hawkins, *Tetrahedron* **2018**, *74*, 2693–2702; e) I. A. Cherepanov, A. S. Samarskaya, R. G. Nosov, I. A. Godovikov, Y. V. Nelyubina, V. N. Kalinin, *Mendeleev Commun.* **2014**, *24*, 386–387.
- [13] a) R. Ogawa, J. Stachnik, H. Echizen, *Clin. Pharmacokinet.* **2014**, *53*, 1083–1114; b) B. Rosenkranz, B. R. Winkelmann, M. J. Parnham, *Clin. Pharmacokinet.* **1996**, *30*, 372–384; c) J. Reden, *Blood Vessels* **1990**, *27*, 282–294.
- [14] J. A. Gruner, J. R. Mathiasen, D. G. Flood, M. Gasior, *J. Pharm. Exp. Ther.* **2011**, *337*, 380–390.
- [15] T. L. Gilchrist, in *Science of Synthesis, Vol. 13*, Thieme, Stuttgart, **2004**, pp. 109–125.
- [16] E. Decuypère, M. Riomet, A. Sallustrau, S. Bregant, R. Thai, G. Pieters, G. Clavier, D. Audisio, F. Taran, *Chem. Commun.* **2018**, *54*, 10758–10761.
- [17] L. C.-C. Lee, H. H.-M. Cheung, H.-W. Liu, K. K.-W. Lo, *Chem. Eur. J.* **2018**, *24*, 14064–14068.
- [18] C. Favre, F. Friscourt, *Org. Lett.* **2018**, *20*, 4213–4217.
- [19] a) S. Araki, J. Mizuya, Y. Butsugan, *Chem. Lett.* **1984**, *13*, 1045–1046; b) S. Araki, J. Mizuya, Y. Butsugan, *J. Chem. Soc. Perkin Trans. 1* **1985**, 2439–2441.
- [20] H. Dahn, U. T. M. Ngoc, *Helv. Chim. Acta* **1988**, *71*, 241–248.
- [21] J. Jaźwiński, O. Staszewska, J. W. Wiench, L. Stefaniak, S. Araki, G. A. Webb, *Magn. Reson. Chem.* **2000**, *38*, 617–626.
- [22] a) N. L. Ahlburg, T. Freese, S. Kolb, S. Mummel, A. Schmidt, D. B. Werz, *Eur. J. Org. Chem.* **2021**, 1603–1610; b) C. F. Otto, M. Liu, C. Herzberger, J. C. Namyslo, M. Nieger, E. G. Hübner, F. Lederle, T. Freese, A. Schmidt, *Tetrahedron* **2020**, *76*, 131627; c) A. Schmidt, S. Batsyts, A. Smeyanov, T. Freese, E. G. Hübner, M. Nieger, *J. Org. Chem.* **2016**, *81*, 4202–4209; d) A. Schmidt, B. Snovydyovych, *Synthesis* **2008**, 2798–2804; e) A. Schmidt, M. K. Kindermann, *J. Org. Chem.* **1998**, *63*, 4636–4644.
- [23] a) S. Deev, S. Batsyts, E. Sheina, T. S. Shestakova, I. Khalimbadzha, M. A. Kiskin, V. Charushin, O. Chupakhin, A. S. Paramonov, Z. O. Shenkarev, J. C. Namyslo, A. Schmidt, *Eur. J. Org. Chem.* **2020**, 450–465; b) S. Batsyts, R. Vedmid, J. C. Namyslo, M. Nieger, A. Schmidt, *Eur. J. Org. Chem.* **2019**, 1301–1310; c) M. Liu, M. Nieger, E. Hübner, A. Schmidt, *Chem. Eur. J.* **2016**, *22*, 5416–5424; d) A. Schmidt, N. Münster, A. Dreger, *Angew. Chem. Int. Ed.* **2010**, *49*, 2790–2793; *Angew. Chem.* **2010**, *122*, 2851–2854.
- [24] Reviews: a) A. Schmidt, S. Wiechmann, C. F. Otto, *Adv. Heterocycl. Chem.* **2016**, *119*, 143–172; b) A. Schmidt, S. Wiechmann, T. Freese, *ARKIVOC* **2013**, 424–469.
- [25] a) Review: A. Nasr, A. Winkler, M. Tamm, *Coord. Chem. Rev.* **2016**, *316*, 68–124; b) recent publications: J. Frosch, M. Koneczny, T. Bannenberg, M. Tamm, *Chem. Eur. J.* **2021**, *27*, 4349–4363; c) A. A. Grineva, D. A. Valyaev, V. César, O. A. Filippov, V. N. Khrustalev, S. E. Nefedov, N. Lugan, *Angew. Chem. Int. Ed.* **2018**, *57*, 7986–7991; *Angew. Chem.* **2018**, *130*, 8118–8123; d) F. Medici, G. Gontard, E. Derat, G. Lemiere, L. Fensterbank, *Organometallics* **2018**, *37*, 517–520.
- [26] a) C. A. Ramsden, W. P. Oziminski, *J. Org. Chem.* **2016**, *81*, 10295–10301; b) C. A. Ramsden, W. P. Oziminski, *J. Org. Chem.* **2017**, *82*, 12485–12491; c) W. P. Oziminski, C. A. Ramsden, *RSC Adv.* **2018**, *8*, 14833–14837.
- [27] J. Mathew, C. H. Suresh, *Inorg. Chem.* **2010**, *49*, 4665–4669.
- [28] L. Perrin, E. Clot, O. Eisenstein, J. Loch, R. H. Crabtree, *Inorg. Chem.* **2001**, *40*, 5806–5811.
- [29] C. A. Tolman, *J. Am. Chem. Soc.* **1970**, *92*, 2953–2956.
- [30] A. A. Tukov, A. T. Normand, M. S. Nechaev, *Dalton Trans.* **2009**, 7015–7028.
- [31] D. Bourissou, O. Guerret, F. P. Gabbaï, G. Bertrand, *Chem. Rev.* **2000**, *100*, 39–91.
- [32] J. C. Bernhammer, G. Frison, H. V. Huynh, *Chem. Eur. J.* **2013**, *19*, 12892–12905.
- [33] K. Verlinden, H. Buhl, W. Frank, C. Ganter, *Eur. J. Inorg. Chem.* **2015**, 2416–2425.
- [34] a) H. V. Huynh, Y. Han, R. Jothibas, J. A. Yang, *Organometallics* **2009**, *28*, 5395–5404; b) Q. Teng, H. V. Huynh, *Dalton Trans.* **2017**, *46*, 614–627.
- [35] H. V. Huynh, *Chem. Rev.* **2018**, *118*, 9457–9492.
- [36] C. Hansch, A. Leo, R. W. Taft, *Chem. Rev.* **1991**, *91*, 165–195.
- [37] T. Freese, A.-L. Lücke, C. A. S. Schmidt, M. Polamo, M. Nieger, J. C. Namyslo, A. Schmidt, *Tetrahedron* **2017**, *73*, 5350–5357.

- [38] a) M. Begtrup, *J. Chem. Soc. Chem. Commun.* **1975**, 334–335;
b) C. L. Deardorff, R. E. Sikma, C. P. Rhodes, T. W. Hudnall,
Chem. Commun. **2016**, 52, 9024–9027.
- [39] D. Mendoza-Espinosa, B. Donnadiou, G. Bertrand, *J. Am. Chem. Soc.* **2010**, 132, 7264–7265.
- [40] V. César, V. Mallardo, A. Nano, G. Dahm, N. Luga, G. Lavigne, S. Bellemin-Laponnaz, *Chem. Commun.* **2015**, 51, 5271–5274.
- [41] X. Bantreil, N. Pétry, F. Lamaty, *Dalton Trans.* **2019**, 48, 15753–15761.
- [42] Alex A. Granovsky, Firefly version 8, <http://classic.chem.msu.su/gran/firefly/index.html>.
- [43] M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, T. L. Windus, M. Dupuis, J. A. Montgomery, *J. Comput. Chem.* **1993**, 14, 1347–1363.
- [44] G. M. Sheldrick, *Acta Crystallogr. Sect. A* **2015**, 71, 3–8.
- [45] G. M. Sheldrick, *Acta Crystallogr. Sect. C* **2015**, 71, 3–8.
- [46] Deposition Number(s) 1055127, 1055128, and 1055129 contain(s) the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service www.ccdc.cam.ac.uk/structures.

Manuscript received: June 5, 2021

Accepted manuscript online: June 21, 2021

Version of record online: July 14, 2021