



This is a repository copy of *Chronic kidney disease as cardiovascular risk factor in routine clinical practice: a position statement by the Council of the European Renal Association*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/192427/>

Version: Published Version

Article:

Ortiz, A., Wanner, C., Gansevoort, R. et al. (12 more authors) (2022) Chronic kidney disease as cardiovascular risk factor in routine clinical practice: a position statement by the Council of the European Renal Association. *European Journal of Preventive Cardiology*. ISSN 2047-4873

<https://doi.org/10.1093/eurjpc/zwac186>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Chronic kidney disease as cardiovascular risk factor in routine clinical practice: a position statement by the Council of the European Renal Association

Alberto Ortiz ^{1,2}, Christoph Wanner ³, and Ron Gansevoort ^{4*},
ERA Council

¹IIS-Fundacion Jimenez Diaz- UAM, Madrid, Spain; ²Department of Medicine, Universidad Autonoma de Madrid, Madrid, Spain; ³Department of Internal Medicine I, University Hospital Würzburg, Würzburg, Germany; and ⁴Department of Nephrology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

Received 23 April 2022; revised 31 July 2022; accepted 20 August 2022

Abstract

The European Society of Cardiology 2021 guideline on cardiovascular (CV) disease (CVD) prevention in clinical practice has major implications for both CV risk screening and kidney health of interest to primary care physicians, cardiologists, nephrologists, and other professionals involved in CVD prevention. The proposed CVD prevention strategies require as first step the categorization of individuals into those with established atherosclerotic CVD, diabetes, familiar hypercholesterolaemia, or chronic kidney disease (CKD), i.e. conditions that are already associated with a moderate to very-high CVD risk. This places CKD, defined as decreased kidney function or increased albuminuria as a starting step for CVD risk assessment. Thus, for adequate CVD risk assessment, patients with diabetes, familiar hypercholesterolaemia, or CKD should be identified by an initial laboratory assessment that requires not only serum to assess glucose, cholesterol, and creatinine to estimate the glomerular filtration rate, but also urine to assess albuminuria. The addition of albuminuria as an entry-level step in CVD risk assessment should change clinical practice as it differs from the current healthcare situation in which albuminuria is only assessed in persons already considered to be at high risk of CVD. A diagnosis of moderate or severe CKD requires a specific set of interventions to prevent CVD. Further research should address the optimal method for CV risk assessment that includes CKD assessment in the general population, i.e. whether this should remain opportunistic screening or whether systematic screening.

Keywords

GFR • albuminuria • cardiovascular risk • risk prediction • prevention

Introduction

The recent publication of the European Society of Cardiology (ESC) 2021 Guideline on cardiovascular (CV) disease (CVD) prevention in clinical practice introduces chronic kidney disease (CKD) risk categories that are based on the integration of estimated glomerular filtration rate (eGFR) and urinary albumin-to-creatinine ratio (UACR) data. Furthermore, this guideline indicates that assessment of eGFR and UACR (i.e. a urine analysis) are needed for correct patient

categorization according to CVD risk that will guide preventive measures. This has major implications for CVD risk screening as well as kidney health, which is of interest to primary care physicians, cardiologists, nephrologists, and other professionals involved in CVD prevention. In this Position Paper, we first provide an overview of the concept and disease burden of CKD and of the key messages on CKD embedded in the 2021 ESC guideline and then provide a call for action to implement this guideline in order to improve the outcomes of patients with CKD.

* Corresponding author. Department of Nephrology, UMC Groningen, Hanzeplein 1,9713 GZ Groningen, the Netherlands. Tel: +31-50-3616161, Fax: +31-50-3619310, Email: r.t.gansevoort@umcg.nl

This article has been co-published with permission in *Clinical Kidney Journal*, *Nephrology Dialysis Transplantation*, and *European Journal of Preventive Cardiology*. © The Author(s) 2022. Published by Oxford University Press on behalf of the ERA and European Society of Cardiology. All rights reserved. The articles are identical except for minor stylistic and spelling differences in keeping with each journal's style. Either citation can be used when citing this article.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com.

			Albuminuria Categories (mg/g)		
			A1	A2	A3
			<30	30-299	≥300
GFR Categories (mL/min/1.73 m ²)	G1	>90	55.6%	1.9%	0.4%
	G2	60-89	32.9%	2.2%	0.3%
	G3a	45-59	3.6%	0.8%	0.2%
	G3b	30-44	1.0%	0.4%	0.2%
	G4	15-29	0.2%	0.1%	0.1%
	G5	<15	<0.1%	<0.1%	<0.1%

What is known?

CKD severity

- Mild CKD
- Moderate CKD
- Severe CKD

What is new?

ESC 2021 CVD risk class

- High CVD risk
- Very high CVD risk

Figure 1 Translation of chronic kidney disease risk classes (as defined by Kidney Disease: Improving Global Outcomes in 2012) into cardiovascular disease risk classes as defined by the European society of cardiology in the 2021 guideline on cardiovascular disease prevention in clinical practice. Numbers within cells represent prevalence in the general population.

The concept of chronic kidney disease

In the 2012 KDIGO (abbreviation for Kidney Disease: Improving Global Outcomes), the global organization developing clinical practice guidelines for nephrology, defined CKD as abnormalities of kidney structure or function present for ≥ 3 months, with implications for health.^{1,2} In clinical practice, the main diagnostic criteria for CKD are the presence of an eGFR below 60 mL/min/1.73 m², and/or elevated albuminuria, i.e. UACR over 30 mg/g. Chronic kidney disease is staged according to six eGFR and three UACR categories (Figure 1), with higher categories being associated with increased risk of adverse outcomes including CKD progression, kidney failure, and fatal and non-fatal CVD. Combinations of specific eGFR and UACR cells are used to define three global risk classes: CKD with mildly increased risk, CKD with moderately increased risk, and CKD with severely increased risk (Figure 1).

A key message from the 2012 KDIGO guideline was that a low eGFR is not a prerequisite for CKD diagnosis; for example, a person with normal eGFR can be diagnosed with CKD if albuminuria is elevated. This enables CKD identification at an early stage, and thus allows timely start of effective prevention of CKD progression towards kidney failure.

Current and future burden of chronic kidney disease

Globally, around 10–12% of the general population has CKD, implying that this involves 850 million persons.³ Despite being largely preventable and treatable, the prevalence of CKD as well as mortality

due to CKD are increasing.⁴ The Global Burden of Disease study projects that worldwide by 2040, CKD will become the fifth most common cause of death.⁵ The years of life lost due to CKD will double. This represents the fastest predicted increase among the major causes of death, after Alzheimer's disease.⁵ Notably, the increasing burden of CKD may be magnified in countries with changing demography due to an ageing population, as is the case for several European countries. For example, it is projected that CKD will become the second leading cause of death in Spain before the end of the century.⁶

KDIGO 2012 guideline and the concept of chronic kidney disease

Awareness of the concept of CKD, and of the possible consequences, is key to promoting both CV and kidney health. However, in routine clinical practice, the implementation of diagnosing CKD is far from desirable. For instance, in several countries screening for eGFR and albuminuria in patients at high risk for CKD lags seriously behind what is advised in guidelines.⁷ Currently, the invisibility of CKD as a clinical diagnosis is a barrier to the implementation of CV risk mitigation strategies as CKD patients are not identified as such. In recent cohort studies, only 23–39% of patients with CKD actually had a CKD diagnosis registered in the electronic health records by their treating physician.^{8,9} This has consequences, as for instance patients lacking CKD diagnoses were more frequently prescribed nephrotoxic medications.⁸

Notably, there is still confusion regarding the concept of CKD among some physicians. For instance, as recently as 2018, a seminal medical journal published a manuscript that referred to all patients with an eGFR above 60 mL/min/1.73 m² as having CKD G1/G2,

although in most albuminuria or structural kidney abnormalities were not assessed.²

ESC 2021 guideline on cardiovascular disease prevention in clinical practice

In 2021, the ESC issued an update of its seminal guideline on CVD prevention in clinical practice.¹⁰ This update was made in consensus with 12 medical societies, including the European Renal Association (ERA).¹⁰ As the prior 2016 edition, the 2021 guideline recognizes CKD, diagnosed as a low eGFR (<60 mL/min/1.73 m²), as a CV risk factor.¹¹ However, from the perspective of the ERA, this guideline incorporates four main novelties:

- (1) The concept of moderate and severe CKD is expanded by incorporating albuminuria into the definition. Thus, moderate CKD is no longer limited to persons with an eGFR 30–59 mL/min/1.73 m² and severe CKD is no longer limited to an eGFR <30 mL/min/1.73 m², as they were in 2016. Rather, moderate, and severe CKD now follow the KDIGO global risk classes to encompass a combination of eGFR and UACR values that is associated with high or very-high CVD risk (Figure 1). Thus, now CKD diagnosed by not only low eGFR, but also high UACR is acknowledged as a strong and independent risk factor for CVD. Although the 2016 guideline mentioned albuminuria, it did not integrate it into an integrated CV risk score.
- (2) Assessment of CKD, based on either elevated albuminuria (>30 mg/g) or low eGFR (<60 mL/min/1.73 m²), should be part of routine CV risk assessment (see below).
- (3) When CKD is present, not only an ACE inhibitor or an ARB, but also an SGLT2 inhibitor should be considered if non-diabetic and is recommended if the patient has Type 2 diabetes to prevent progression of CKD and CV events (Table 1). This advice was based on the results of the DAPA-CKD trial, in which patients with CKD and with or without Type 2 diabetes were included.¹² Recently, the EMPA-KIDNEY Trial, that included a similar patient population, was stopped early because of positive efficacy that met the study's pre-specified threshold for early termination.¹³ This has strengthened the evidence supporting this advice.
- (4) Nomenclature for CKD classes is proposed, which is easier to understand than the original one by KDIGO. It is based on the associated overall risk of adverse health outcomes: mild CKD, moderate CKD (associated with high CV risk), and severe CKD (associated with very-high CV risk) (Figure 1). This simplified nomenclature may facilitate uptake of the CKD concept, and its incorporation into clinical diagnostic codes and risk assessment tools.

Implications for screening

The ESC 2021 guideline on CVD Prevention in Clinical Practice has major implications for both CV risk screening and kidney health, stating that not only blood pressure, cholesterol, and glucose should be

Table 1 Interventions suggested or recommended for primary prevention in persons with chronic kidney disease according to the ESC 2021 guideline on cardiovascular disease prevention in clinical practice

In persons with moderate or severe CKD, there is no need to estimate 10-year fatal and non-fatal CVD risk with either SCORE2 or SCORE2-OP. The CVD risk is already high (moderate CKD) or very high (severe CKD), respectively, warranting preventive interventions according to the level of risk, such as:

- For all patients with CKD:
 - Lifestyle advice including diet and appropriate weight.
 - Blood pressure control, preferably by RAS inhibitors.
- For CKD patients with specific conditions:
 - LDL-cholesterol reduction to target [LDL-C <1.8 mmol/L (70 mg/dL) in high-risk patients and <1.4 mmol/L (55 mg/dL) in very-high-risk patients].
 - RAS inhibition in case of severely elevated albuminuria, independent of blood pressure.
 - SGLT2 inhibition for those with Type 2 diabetes, or with diabetic as well as non-diabetic kidney disease and an eGFR higher than 25 mL/min/1.73 m², or with heart failure.
 - In addition, for consideration to prevent CVD in specific patients (not part of the ESC guideline): (nonsteroidal) mineralocorticoid receptor antagonists and/or PCSK9 inhibitors.

assessed to estimate CV risk in an individual, but also albuminuria and eGFR. This goes beyond the traditional view that eGFR and UACR should only be assessed in populations at high risk for CKD, i.e. those with proven hypertension, proven diabetes, or a CVD history. In this regard, every healthcare system should promote that individuals should know their ABCDE profile regarding CV risk—i.e. albuminuria, blood pressure, cholesterol, diabetes status, and eGFR. This ABCDE profile should be considered along with other risk factors that are self-evident for the individual, such as obesity and smoking. Screening strategies should begin at an age at which absolute risk for outcomes is high enough to warrant preventive strategies, and that allows sufficient lead time for preventive approaches to be successful. The ESC 2021 guideline on CVD prevention in clinical practice supports considering opportunistic and even systematic CV risk assessment among the general population, in men >40 years of age and in women who are postmenopausal or >50 years of age, even when these subjects have no known atherosclerotic CVD risk factors (Class IIb, level C). As discussed above such screening should include eGFR and UACR, and could be repeated after 5 years, or sooner if risk was close to treatment thresholds (Class IIb, level C).

The design of a health check-up directed at prevention of CVD includes CKD awareness and should be adapted according to local resources and characteristics of the population. It may range from opportunistic screening by general practitioners, as done nowadays in most countries, to an extensive health check-up performed at healthcare centres, including a medical history, physical exam, and

basic serum and urine biochemistry. From a CKD perspective, the future may even bring home-based urine screening for albuminuria. Such screening could be linked to currently existent colon cancer screening campaigns that in some countries are implemented around the assessment of occult faecal blood in a sample collected at home.¹⁴ Addition of a urine sample will take advantage of an ongoing screening effort and would only add the cost of albuminuria assessment. This would identify target organ injury (i.e. kidney injury) that could result from various known or unknown pre-existing conditions, and thereby identify patients who require a more in-depth assessment and optimized therapy. Such a strategy is likely cost-effective to delay or prevent dialysis and CVD.¹⁵

Emphasizing that screening for CVD risk factors should include CKD markers is timely for several reasons. First, because we can now estimate risk better. Adding albuminuria and eGFR to the SCORE2 and SCORE2-OP algorithms improves CVD risk prediction in subjects with CKD^{16,17} (see [Supplementary material online, Figure S1](#)). Precise information about prognosis may stimulate patients and physicians to start preventive treatments ([Table 1](#)). Second, because we can now treat patients better. In the last decade, various agents have demonstrated cardio- and renoprotection on top of standard-of-care including RAS inhibitors, GLP1 analogs, SGLT2 inhibitors, and the nonsteroidal MRA finerenone.^{12,18–21} Notwithstanding this progress in knowledge, it should also be noted that CKD patients have been excluded from several CVD trials.²² For some interventions it is therefore still unresolved what their efficacy-safety ratio is in specifically subjects with CKD. Including subgroups of CKD patients in CVD trials yet to be started will therefore be important.

A call to action by the Council of the European Renal Association

The ESC emphasizes in their guideline that subjects with CKD are at high risk for CVD, and has given CKD a prominent place in CVD prevention. The guideline now requires several stages of dissemination and implementation into clinical practice. If correctly planned this may also generate novel information regarding the optimal implementation strategies:

- (1) Wide dissemination of the concept that assessment of serum creatinine (eGFR) and albuminuria (UACR) should be part of any evaluation of CV risk. The message should reach all key stakeholders: healthcare system payers, healthcare personnel, patients, and the general population. To achieve this goal, nephrologists should liaise with primary care physicians, CVD physicians, and cardiologists as key healthcare stakeholders. For instance, they could work out this concept in national guidelines for management of CKD and for CVD prevention, develop educational tools, and organize post-graduate medical education for the various specialties involved. To reach other healthcare personnel, healthcare system payers and the general population the organization of CKD awareness campaigns are indicated. The ERA has templates for such campaigns, for instance to be organized around World Kidney Day.²³
- (2) Europe-wide evaluation of optimal screening strategies to identify CKD. Given the differences in baseline risk for CVD

and CKD, and the different healthcare systems, it is unlikely that a single strategy fits all and that the type of screening will depend on local financial resources and logistic possibilities. On a national level initiatives should be started how to best screen for CKD, as for instance the THOMAS study in the Netherlands that investigates the cost-effectiveness of two different ways of home-based albuminuria screening of the general population.²⁴ A prospective pan-European evaluation of different screening strategies may allow definition of the optimal strategy for local conditions. This could be achieved via the organization of an invitational conference on this topic to be organized by the ERA, KDIGO, and/or ESC.

- (3) Chronic kidney disease patients should be included in CVD trials yet to be started. Integration of cohorts into registries may generate pragmatic trials that test novel CV protective strategies identified through clinical trials from which CKD patients were excluded. Conversely, clinical trials focused on CKD populations should also have CV end points as CVD constitutes a major proportion of morbidity and mortality in this patient population.

In conclusion, the ESC 2021 guideline on CVD Prevention in Clinical Practice identifies albuminuria and eGFR as CV risk factors that need assessment. Ample dissemination of this guideline among primary care physicians and specialists offers the opportunity for an earlier diagnosis and treatment of CKD and is expected to improve both CV and kidney outcomes in affected subjects.

Supplementary material

[Supplementary material](#) is available at *European Journal of Preventive Cardiology* online.

Conflict of interest: none declared.

ERA council

Alberto Ortiz^{1,2}, Christoph Wanner³, Ron T. Gansevoort⁴, Mario Cozzolino⁵, Danilo Fliser⁶, Giovanni Gambaro⁷, Albert Ong⁸, Alexander R. Rosenkranz⁹, Ivan Rychlik¹⁰, Pantelis Sarafidis¹¹, Roser Torra¹², Serhan Tuglular¹³

¹IIS-Fundacion Jimenez Diaz- UAM, Madrid, Spain.

²Department of Medicine, Universidad Autonoma de Madrid, Madrid, Spain.

³Department of Internal Medicine I, University Hospital Würzburg, Würzburg, Germany.

⁴Department of Nephrology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

⁵Department of Health Sciences, University of Milan, Renal Division, ASST Santi Paolo e Carlo, Milan, Italy.

⁶Internal Medicine IV, Renal and Hypertensive Disease, University Medical Center, Homburg, Saar, Germany.

⁷Division of Nephrology and Dialysis, Department of Medicine, Ospedale Maggiore, Verona, Italy.

⁸Kidney Genetics Group, Academic Nephrology Unit, Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield Medical School, Sheffield, UK.

⁹Department of Internal Medicine, Division of Nephrology, Medical University of Graz, Graz, Austria.

¹⁰1st Department of Internal Medicine, Third Faculty of Medicine, Charles University, Faculty Hospital Krvskinohrady, Prague, Czech Republic.

¹¹Department of Nephrology, Hippokraton Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece.

¹²Inherited Kidney Diseases Nephrology Department, Fundació Puigvert Instituto de Investigaciones Biomédicas Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain. REDINREN, Barcelona, Spain.

¹³Division of Nephrology, Department of Internal Medicine, School of Medicine, Marmara University, Istanbul, Turkey.

References

- Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 2013;**3**:1–150.
- Perez-Gomez MV, Bartsch LA, Castillo-Rodriguez E, Fernandez-Prado R, Fernandez-Fernandez B, Martin-Cleary C, Gracia-Iguacel C, Ortiz A. Clarifying the concept of chronic kidney disease for non-nephrologists. *Clin Kidney J* 2019;**12**:258–226.
- Jager KJ, Kovesdy C, Langham R, Rosenberg M, Jha V, Zoccali C. A single number for advocacy and communication-worldwide more than 850 million individuals have kidney diseases. *Nephrol Dial Transplant* 2019;**34**:1803–1805.
- GBD Chronic Kidney Disease Collaboration. Global, regional, and national burden of chronic kidney disease, 1990–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2020;**395**:709–733.
- Foreman KJ, Marquez N, Dolgert A, Fukutaki K, Fullman N, McGaughey M, Pletcher MA, Smith AE, Tang K, Yuan CW, Brown JC, Friedman J, He J, Heuton KR, Holmberg M, Patel DJ, Reidy P, Carter A, Cercy K, Chapin A, Douwes-Schultz D, Frank T, Goettsch F, Liu PY, Nandakumar V, Reitsma MB, Reuter V, Sadat N, Sorensen RJD, Srinivasan V, Updike RL, York H, Lopez AD, Lozano R, Lim SS, Mokdad AH, Vollset SE, Murray CJL. Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality for 250 causes of death: reference and alternative scenarios for 2016–40 for 195 countries and territories. *Lancet* 2018;**392**:2052–2090.
- Shin JI, Chang AR, Grams ME, Coresh J, Ballew SH, Surapaneni A, Matsushita K, Bilo HJG, Carrero JJ, Chodick G, Daratha KB, Jassal SK, Nadkarni GN, Nelson RG, Nowak C, Stempniewicz N, Sumida K, Traynor JP, Woodward M, Sang Y, Gansevoort RT for the CKD Prognosis Consortium. Albuminuria testing in hypertension and diabetes: an individual-participant data meta-analysis in a global consortium. *Hypertension* 2021;**78**:1042–1052.
- Ortiz A, Sanchez-Niño MD, Crespo-Barrio M, De-Sequera-Ortiz P, Fernández-Giráldez E, García-Maset R, Macía-Heras M, Pérez-Fontán M, Rodríguez-Portillo M, Salgueira-Lazo M, Sánchez-Álvarez E, Santamaria-Olmo R, Simal-Blanco F, Pino-Pino MD. The Spanish society of nephrology (SENEFRO) commentary to the Spain GBD 2016 report: keeping chronic kidney disease out of sight of health authorities will only magnify the problem. *Nefrología* 2019;**39**:29–34.
- Bosi A, Xu Y, Gasparini A, Wettermark B, Barany P, Bellocco R, Inker LA, Chang AR, McAdams-DeMarco M, Grams ME, Shin JI, Carrero JJ. Use of nephrotoxic medications in adults with chronic kidney disease: parallel cohort studies in Swedish and U.S. Routine care. *Clin Kidney J* 2021;**15**:442–445.
- Wang HY, Ding GH, Lin H, Sun X, Yang C, Peng S, Wang J, Du J, Zhao Y, Chen Z, Bao B, Kong G, Zhang L. Influence of doctors' perception on the diagnostic status of chronic kidney disease: results from 976 409 individuals with electronic health records in China. *Clin Kidney J* 2021;**14**:2428–2436.
- Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Böck M, Benetos A, Biffi A, Boavida JM, Capodanno D, Cosyns B, Crawford C, Davos CH, Desormais I, Di Angelantonio E, Franco OH, Halvorsen S, Hobbs FDR, Hollander M, Jankowska EA, Michal M, Sacco S, Sattar N, Tokgozoglu L, Tonstad S, Tsioufias KP, van Dis I, van Gelder IC, Wanner C, Williams B; ESC Scientific Document Group. 2021 ESC guidelines on cardiovascular disease prevention in clinical practice. *Eur J Prev Cardiol* 2022;**29**:5–115.
- Authors/Task Force Members, Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, Cooney MT, Corrà U, Cosyns B, Deaton C, Graham I, Hall MS, Hobbs FD, Løchen ML, Löllgen H, Marques-Vidal P, Perk J, Prescott E, Redon J, Richter DJ, Sattar N, Smulders Y, Tiberi M, van der Worp HB, van Dis I, Verschuren WM; Additional Contributor: Simone Binno (Italy); Document Reviewers; De Backer G, Roffi M, Aboyans V, Bachl N, Bueno H, Carerj S, Cho L, Cox J, De Sutter J, Egidi G, Fisher M, Fitzsimons D, Franco OH, Guenoun M, Jennings C, Jug B, Kirchhof P, Koteva K, Lip GY, Mach F, Mancía G, Bermudo FM, Mezzani A, Niessner A, Ponikowski P, Rauch B, Rydén L, Stauder A, Turc G, Wiklund O, Windecker S, Zamorano JL. 2016 European guidelines on cardiovascular disease prevention in clinical practice: the sixth joint task force of the European society of cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of 10 societies and by invited experts): developed with the special contribution of the European association for cardiovascular prevention & rehabilitation (EACPR). *Eur J Prev Cardiol* 2016;**23**:1–96.
- Heerspink HJL, Stefansson BV, Correa-Rotter R, Chertow GM, Greene T, Hou FF, Mann JFE, McMurray JJV, Lindberg M, Rossing P, Sjöström CD, Toto RD, Langkilde AM, Wheeler DC, DAPA-CKD Trial Committees and Investigators. Dapagliflozin in patients with chronic kidney disease. *N Engl J Med* 2020;**383**:1436–1446.
- <https://www.medscape.com/viewarticle/970434>
- Senore C, Basu P, Anttila A, Ponti A, Tomatis M, Vale DB, Ronco G, Soerjomataram I, Primic-Zakelj M, Riggi E, Dillner J, Elfström MK, Lönnberg S, Sankaranarayanan R, Segnan N. Performance of colorectal cancer screening in the European union member states: data from the second European screening report. *Gut* 2019;**68**:1232–1244.
- Crews DC, Boulware LE, Gansevoort RT, Jaar BG. Albuminuria: is it time to screen the general population? *Adv Chronic Kidney Dis* 2011;**18**:249–257.
- Matsushita K, Jassal SK, Sang Y, Ballew SH, Grams ME, Surapaneni A, Arnlöv J, Bansal N, Bozic M, Brenner H, Brunskill NJ, Chang AR, Chinnadurai R, Cirillo M, Correa A, Ebert N, Eckardt KU, Gansevoort RT, Gutierrez O, Hadaegh F, He J, Hwang SJ, Jafar TH, Kayama T, Kovesdy CP, Landman GW, Levey AS, Lloyd-Jones DM, Major RW, Miura K, Muntner P, Nadkarni GN, Naimark DM, Nowak C, Ohkubo T, Pena MJ, Polkinghorne KR, Sabanayagam C, Sairenchi T, Schneider MP, Shalev V, Shlipak M, Solbu MD, Stempniewicz N, Tollitt J, Valdivielso JM, van der Leeuw J, Wang AY, Wen CP, Woodward M, Yamagishi K, Yatsuya H, Zhang L, Schaeffner E, Coresh J. Incorporating kidney disease measures into cardiovascular risk prediction: development and validation in 9 million adults from 72 datasets. *EClinicalMedicine* 2020;**27**:100552.
- Matsushita K, Kaptoge S, Hageman SHJ, Sang Y, Ballew SH, Grams ME, Surapaneni A, Sun L, Arnlöv J, Bozic M, Brenner H, Brunskill NJ, Chang AR, Chinnadurai R, Cirillo M, Correa A, Ebert N, Eckardt KU, Gansevoort RT, Gutierrez O, Hadaegh F, He J, Hwang SJ, Jafar TH, Jassal SK, Kayama T, Kovesdy CP, Landman GW, Levey AS, Lloyd-Jones DM, Major RW, Miura K, Muntner P, Nadkarni GN, Nowak C, Ohkubo T, Pena MJ, Polkinghorne KR, Sairenchi T, Schaeffner E, Schneider MP, Shalev V, Shlipak MG, Solbu MD, Stempniewicz N, Tollitt J, Valdivielso JM, van der Leeuw J, Wang AYM, Wen CP, Woodward M, Yamagishi K, Yatsuya H, Zhang L, Dorresteijn JAN, Di Angelantonio E, Visseren FLJ, Pennells L, Coresh J. Adding albuminuria and eGFR to the SCORE2 algorithms to improve CVD risk prediction in subjects with chronic kidney disease. *Eur J Prev Cardiol* 2022. doi: 10.1093/eurjpc/zwac176
- Pitt B, Filippatos G, Agarwal R, Anker SD, Bakris GL, Rossing P, Joseph A, Kolkhof P, Nowack C, Schloemer P, Ruilope LM, FIGARO-DKD Investigators. Cardiovascular events with finerenone in kidney disease and type 2 diabetes. *N Engl J Med* 2021;**385**:2252–2263.
- Bakris GL, Agarwal R, Anker SD, Pitt B, Ruilope LM, Rossing P, Kolkhof P, Nowack C, Schloemer P, Joseph A, Filippatos G, FIDELIO-DKD Investigators. Effect of finerenone on chronic kidney disease outcomes in type 2 diabetes. *N Engl J Med* 2020;**383**:2219–2229.
- Perkovic V, Jardine MJ, Neal B, Bompoint S, Heerspink HJL, Charytan DM, Edwards R, Agarwal R, Bakris G, Bull S, Cannon CP, Capuano G, Chu PL, de Zeeuw D, Greene T, Levin A, Pollock C, Wheeler DC, Yavin Y, Zhang H, Zinman B, Meininger G, Brenner BM, Mahaffey KW, CREDESCENCE Trial Investigators. Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019;**380**:2295–2306.
- Sarafidis P, Ferro CJ, Morales E, Ortiz A, Malyszko J, Hojs R, Khazim K, Ekart R, Valdivielso J, Fouque D, London GM, Massy Z, Ruggenenti P, Porrini E, Wiecek A, Zoccali C, Mallamaci F, Hornum M. SGLT-2 inhibitors and GLP-1 receptor agonists for nephroprotection and cardioprotection in patients with diabetes mellitus and chronic kidney disease. A consensus statement by the EURECA-m and the DIABESITY working groups of the ERA-EDTA. *Nephrol Dial Transplant* 2019;**34**:208–230.
- Zoccali C, Blankestijn PJ, Bruchfeld A, Capasso G, Fliser D, Fouque D, Goumenos D, Ketteler M, Massy Z, Rychlik I, Jose Soler M, Stevens K, Spasovski G, Wanner C. Children of a lesser god: exclusion of chronic kidney disease patients from clinical trials. *Nephrol Dial Transplant* 2019;**34**:1112–1114.
- <https://www.era-online.org/en/strongkidneys/european-awareness-campaign/ideasforactivities/>.
- <https://ClinicalTrials.gov/ct2/show/NCT04295889>.