



TITLE:

Evaluation of Oxidative Potential of Pyrenequinone Isomers by the Dithiothreitol (DTT) Assay

AUTHOR(S):

Okubo, Rikito; Kameda, Takayuki; Tohno, Susumu

CITATION:

Okubo, Rikito ...[et al]. Evaluation of Oxidative Potential of Pyrenequinone Isomers by the Dithiothreitol (DTT) Assay. *Polycyclic Aromatic Compounds* 2022, 42(8): 5152-5159

ISSUE DATE:


2022

URL:

<http://hdl.handle.net/2433/276241>

RIGHT:

This is an Accepted Manuscript of an article published by Taylor & Francis in *Polycyclic Aromatic Compounds* on 12 May 2021, available online: <http://www.tandfonline.com/10.1080/10406638.2021.1925711>; The full-text file will be made open to the public on 12 May 2022 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ください。



Evaluation of the oxidative potential of pyrenequinone isomers by the dithiothreitol (DTT) assay

Journal:	<i>Polycyclic Aromatic Compounds</i>
Manuscript ID	PAC-0034-2020.R1
Manuscript Type:	Special Issue
Date Submitted by the Author:	n/a
Complete List of Authors:	Okubo, Rikito; Kyoto University, Graduate School of Energy Science Kameda, Takayuki; Kyoto University, Graduate School of Energy Science Tohno, Susumu; Kyoto University, Graduate School of Energy Science
Keywords:	DTT assay, particulate matter, oxidation potential, pyrenequinone isomers

SCHOLARONE™
Manuscripts

1
2
3 **Evaluation of the oxidative potential of pyrenequinone isomers by the**
4 **dithiothreitol (DTT) assay**
5
6

7
8 Rikito Okubo, Takayuki Kameda*, and Susumu Tohno
9

10
11 *Graduate School of Energy Science, Kyoto University, Yoshida-Honmachi, Sakyo-ku,*
12 *Kyoto 606-8501, Japan*
13

14 * Corresponding author: tkameda@energy.kyoto-u.ac.jp
15
16

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Evaluation of oxidative potential of pyrenequinone isomers by the dithiothreitol (DTT) assay

Abstract

Atmospheric quinoid polycyclic aromatic hydrocarbons (PAHQs) have adverse health effects as redox-active species in particulate matter (PM). Several PAHQs are known to be very reactive in the dithiothreitol (DTT) assay; however, it is unclear if pyrenequinones, their parent pyrene is one of the most abundant polycyclic aromatic hydrocarbons (PAHs) in the atmosphere, contribute to the loss of DTT in PM extracts. Herein, by employing the DTT assay, we evaluated oxidative potentials of three pyrenequinone isomers (4,5-pyrenequinone [4,5-PyQ] and a mixture of 1,6-/1,8-pyrenequinones [1,6-/1,8-PyQ]), along with 9,10-phenanthrenequinone (PQN), 1,2-naphthoquinone (1,2-NQ), and 1,4-naphthoquinone (1,4-NQ), of which DTT loss rates were examined previously. Our results demonstrate that the DTT consumption by ortho-type PAHQs is fast, particularly by 4,5-PyQ. The order of DTT loss rate by the PAHQs tested in this study was as follows: 4,5-PyQ ~ PQN > 1,2-NQ > 1,4-NQ ~ 1,6-/1,8-PyQ. 4,5-PyQ was confirmed to be active in the DTT assay for the first time in this study. The DTT consumption rate by 4,5-PyQ is 14.6 ± 0.8 mol/min/PAHQ-mol, which is comparable to that of PQN (14.4 ± 0.1 mol/min/PAHQ-mol) known as the most active PAHQs by the DTT assay so far.

Keywords: DTT assay; particulate matter; oxidation potential, pyrenequinone isomers

1 Introduction

Exposure to airborne particulate matter (PM), one of the air pollutants, can have adverse health effects on humans, regardless of the duration of exposure.¹⁻⁴ The harmful health effects of PM on humans include those caused by a single component of PM⁵ and those caused by the synergistic effects of multiple components.⁶ They are regarded as a multifaceted phenomenon, involving many chemical components, such as aliphatic hydrocarbons, polycyclic aromatic hydrocarbons (PAHs), and metals.⁷ In recent years, there has been much interest and research on the health effects of oxidative stress due to the generation of reactive oxygen species (ROS) caused by PM.^{8,9} ROS are reactive chemical species based on oxygen atoms, such as superoxide radical (O_2^-), hydroxy radical (OH^\cdot), hydrogen peroxide (H_2O_2), and singlet oxygen (1O_2). It is known that

1
2
3 excessive production of ROS, such as superoxide radicals, or the lack of an
4 antioxidant's ability to eliminate them *in vivo*, leads to the oxidization and damage of
5 biomolecules like fats, proteins/enzymes, and DNA molecules responsible for genetic
6 information. This results in various conditions like inflammation, cancer, cardiovascular
7 diseases, and neurodegenerative diseases.
8

9
10 The dithiothreitol (DTT; HSCH₂(CH(OH))₂CH₂SH) assay is a method used to
11 quantitatively evaluate the oxidative ability of PM without the use of cells. Many
12 studies on the DTT assay method have been reported.¹⁰⁻¹³ In this assay, compounds
13 contained in PM that have the ability to oxidize DTT to a disulfide structure when
14 mixed with DTT are analyzed, and the rate of oxidation is considered an indicator of the
15 oxidative ability of the compound. Studies using the DTT assay have been reported for
16 the analysis of metals, such as copper and manganese, and substances, such as some
17 quinoid PAHs (PAHQs), which contribute to the oxidative ability of PM.¹² In particular,
18 PAHQs, which have a ketone group at the ortho position, have been reported to catalyze
19 a redox reaction by a redox cycle and generate excessive ROS (Fig. 1).¹³
20
21

22 **Insert Fig. 1 here**

23
24 However, there are many PAHQs that are known to exist in PM, whose oxidative ability
25 has not yet been investigated. For example, pyrenequinones are compounds derived
26 from pyrene which is one of the major PAHs present in atmospheric PM, yet their
27 oxidative ability has not been investigated so far. Therefore, we evaluated the oxidative
28 ability of two pyrenequinone isomers, 4,5-pyrenequinone (4,5-PyQ) and 1,6-/1,8-
29 pyrenequinone (1,6-/1,8-PyQ), by DTT assay along with those of 9,10-
30 phenanthrenequinone (PQN), 1,2-naphthoquinone (1,2-NQ), and 1,4-naphthoquinone
31 (1,4-NQ), which have already been evaluated by the DTT assay.
32
33

34 **2 Materials and methods**

35 **2.1 Chemicals and reagents**

36
37
38 PQN, 1,2-NQ, and 4,5-PyQ were purchased from Sigma-Aldrich. 1,4-NQ was
39 purchased from Tokyo Chemical Industry Co., Ltd. 1,6-/1,8-PyQ (a mixture of 1,6-PyQ
40 and 1,8-PyQ) was provided by Kanto Chemical Co., Inc. DTT, Tris-HCl buffer (pH
41 9.0), and dimethyl sulfoxide (DMSO) were supplied by FUJIFILM Wako Pure
42 Chemical Co. Dithiobisnitrobenzoic acid (DTNB) was obtained from DOJINDO
43 MOLECULAR TECHNOLOGIES, Inc. Acetonitrile, potassium dihydrogen phosphate
44 (KH₂PO₄), and dipotassium hydrogen phosphate (K₂HPO₄) were procured from
45 NACALAI TESQUE, Inc.
46
47

48 **2.2 Preparation of reagents for the DTT assay**

49
50 Solutions of PQN, 1,2-NQ, and 4,5-PyQ at 5 μM concentrations, and those of 1,4-NQ
51 and 1,6-/1,8-PyQ at 25 μM concentrations, were prepared in DMSO. The
52 concentrations of the reagents used in this study were determined within the range
53 where a linear relationship between the concentration and the DTT consumption rate
54 was observed.
55
56
57
58
59
60

2.3 DTT assay

Among different methods of measurement, we followed the method reported by Li et al.¹⁴ In a conical tube immersed in a bath maintained at 37 °C (DIGITAL WATER BATH SB-1000: EYELA), 1425 μL of a 79 μM DTT solution in a 0.05 M potassium phosphate buffer (15 mM K₂HPO₄ and 35 mM KH₂PO₄; pH 7.2) and 75 μL of each test sample were mixed well. PQN, 1,2-NQN, and 4,5-PyQ were reacted for 10 min, while 1,4-NQN and 1,6-/1,8-PyQ were reacted for 20 min. In the blank test, DMSO which was used as a solvent for the preparation of test samples was used. 100 μL of the reacted mixture was collected after a predetermined reaction time and mixed with 20 μL of a 3 mM DTNB solution in a 0.05 M potassium phosphate buffer (pH 7.2) in a disposable spectrophotometer cell. 5-Mercapto-2-nitrobenzoic acid (TNB) was produced by the reaction of DTT and DTNB. The generated TNB was quantified using a UV-Vis spectrophotometer (UVmini-1240: SHIMADZU) at a wavelength of 412 nm. A mixture of the 0.05 M potassium phosphate buffer (pH 7.2) and the DTNB solution was used as a reference for absorbance measurement. The corresponding TNB reduction was calculated as the DTT consumption by each PAHQ after each reaction time. Each sample was analyzed thrice and the results of the three measurements were averaged and divided by reaction time and sample concentration to calculate the DTT consumption rate per unit sample amount (mol/min/PAHQ-mol).

2.4 HPLC analysis

The reaction mixtures of PyQs with DTT were analyzed by HPLC. A 6 mM DTT solution was prepared by diluting the 1 M DTT solution with water. Then, 4,5-PyQ and 1,6-/1,8-PyQ were dissolved in DMSO at prescribed concentrations, followed by the dissolution of these solutions in 250 mM Tris-HCl to a final concentration of 15 μM PyQ (the DMSO concentration in 4,5-PyQ and 1,6-/1,8-PyQ was 5.9 and 7.5%, respectively). Then, 200 μL of the 6 mM DTT solution and 200 μL of each test sample (15 μM 4,5-PyQ or 1,6-/1,8-PyQ solution) were mixed and reacted. After 15 min of reaction, the solution was analyzed using an HPLC system with a HITACHI L-4200 UV-Vis detector. The PyQs were separately eluted from an ODS-P column (4.6 × 250 mm i.d., 5 μm particle size, GL Science) using H₂O/CH₃CN (1/1, v/v) as the mobile phase at a flow rate of 1.0 mL/min, and the compounds were detected at the wavelength of 254 nm. For reference, each sample solution was also analyzed by HPLC before its reaction with DTT.

3 Results and Discussion

3.1 DTT consumption rate by individual quinone compounds

Insert Fig. 2 here

As shown in Fig. 2, the order of the DTT consumption rate of the five quinones analyzed in this study is 4,5-PyQ ~ PQN > 1,2-NQ > 1,4-NQ ~ 1,6-/1,8-PyQ. The activities of three of these quinones, viz., PQN, 1,2-NQ, and 1,4-NQ, on DTT have already been reported by Xiong et al.¹⁰, and this order is consistent with their results.

Charrier et al. also studied the activities of PQN, 1,2-NQ, and 1,4-NQ¹², and reported a DTT consumption rate ratio of 111: 32: 4. Meanwhile, 4,5-PyQ was confirmed to be active in the DTT assay for the first time in this study. The DTT consumption rate by 4,5-PyQ is 14.6 ± 0.8 mol/min/PAHQ-mol, which is comparable to that of PQN (14.4 ± 0.1 mol/min/PAHQ-mol) known as the most active PAHQs^{10, 12} by the DTT assay so far.

3.2 HPLC analysis of 4,5-PyQ and 1,6-/1,8-PyQ

The DTT assay indicated that the oxidation abilities of 1,6-/1,8-PyQ and 4,5-PyQ are remarkably different, although they are positional isomers having similar structures. In order to verify the key factor that affects the DTT consumption rate, the residual amount of the PyQs in the DTT assay samples were analyzed by HPLC.

Insert Figs. 3 and 4 here

After the reaction with DTT, 58% of the initial amount of 4,5-PyQ remained (Fig. 3), whereas only 1% of the initial amount of 1,6-/1,8-PyQ remained at the retention time of ~8 min (Fig. 4), which is much lower than the residual amount of 4,5-PyQ. In addition, after the reaction of 1,6-/1,8-PyQ with DTT, two peaks different from those of 1,6-/1,8-PyQ were observed at the retention time of ~5 min.

Motoyama et al. reported that ortho-PAHQs have higher oxidative ability than para-PAHQs because the oxidation–reduction reaction of PAHQs occurs cyclically, leading to the generation of an excessive amount of ROS, such as H₂O₂, while they oxidize and consume DTT.¹³ It is assumed that the two-electron reduced form of 4,5-PyQ is extremely unstable, and is rapidly oxidized after its formation. It is then converted to 4,5-PyQ via the one-electron reduced form, a semiquinone radical, which continues to catalytically consume DTT. After the reaction with DTT, 58% of 4,5-PyQ remained, although the amount of DTT consumed in the reaction was the largest among the five PAHQs examined in this study. This result indicates that the redox reaction is actively cycled and ROS is generated excessively in the reaction of DTT with 4,5-PyQ. By contrast, 1,6-/1,8-PyQ is considered to have been consumed immediately by the reaction with DTT, because the two-electron reductant is relatively stable, and no redox cycle is formed. The peaks observed in Fig. 4 at retention times of 4.9 and 5.3 min are presumed to be derived from dihydroxypyrenes, the two-electron reduced forms of 1,6-/1,8-PyQ. These results support the hypothesis that the high ROS-generation ability of PAHQs is related to the stability of the two-electron reduced form of the PAHQ compounds.¹⁵

3.3 Atmospheric implications

Table 1 shows the concentrations of various PAHQs present in four reference PMs: NIST SRM1648a (urban PM), SRM1649b (urban dust), SRM2786 (fine PM), and NIES CRM No. 28 (urban aerosol), as reported by Toriba et al.¹⁶

We estimated the oxidative ability of each PAHQ in the reference materials by multiplying the molar concentration of each PAHQ per particle weight by the DTT consumption rate obtained in this study (Table 2). 1,6-PyQ and 1,8-PyQ were assumed

1
2
3 to be present at equal proportions in the standard sample solution analyzed by the DTT
4 assay.
5

6 The oxidative ability of 4,5-PyQ in the reference materials is equivalent to that of PQN.
7 The contribution of the oxidative ability of PQN in PM samples was reported by
8 Charrier et al. to be 17%,¹² but 4,5-PyQ was not considered as a contributing factor. Our
9 result suggests that 4,5-PyQ is also an important substance in terms of the oxidative
10 ability exhibited by PM as well as PQN.
11
12

13 14 **4 Conclusion**

15
16 We evaluated the oxidative abilities of five PAHQs by the DTT assay. As a result, in
17 addition to the well-known oxidants, PQN, 1,2-NQN, and 1,4-NQN, another, 4,5-PyQ,
18 was found to have oxidative ability. The DTT consumption rate by 4,5-PyQ was found
19 to be comparable to that of PQN. On the other hand, the DTT consumption rate of 1,6-
20 /1,8-PyQ was the lowest of the five PAHQs evaluated in this study. In order to examine
21 the cause of the difference in DTT activity between 4,5-PyQ and 1,6-/1,8-PyQ, we used
22 HPLC to analyze the solutions of 4,5-PyQ and 1,6-/1,8-PyQ reacted with DTT. The
23 results revealed that 58% of the initial amount of 4,5-PyQ remained after the reaction
24 with DTT, whereas only 1% of the initial amount of 1,6-/1,8-PyQ remained after the
25 reaction. The result suggests that a redox cycle was formed during the reaction of DTT
26 with 4,5-PyQ, and the ROS were excessively produced catalytically. It is speculated that
27 1,6-/1,8-PyQ was transformed to a more stable two-electron reductant by the reaction.
28 We speculate that the activity of PAHQs in the DTT assay is related to the stability of
29 the two-electron reduced form of the PAHQs. Based on the results of the DTT assay in
30 this study, and the concentration of PAHQs in the reference materials of airborne
31 particulates, we found that 4,5-PyQ makes a large contribution to the ROS production
32 ability of PM, similar to PQN. However, the information obtained from the DTT assay
33 is not sufficient to evaluate the biological effect of 4,5-PyQ. To clarify the oxidative
34 potential of the substances tested in this study, not only other acellular assays such as
35 ascorbic acid (AA) assay¹⁷ and Glutathione (GSH) assay¹⁸, although they usually
36 measure the capability of metals present in PM to deplete antioxidants¹⁹⁻²², but also
37 cell-based assays such as dichloro-dihydro-fluorescein diacetate (DCFH-DA)²³ should
38 be used.
39
40
41
42
43

44 **Acknowledgments**

45 This work was supported in part by the Japan Society for the Promotion of Science
46 Grants-in-Aid for Scientific Research (JSPS KAKENHI), Grant Number JP17H01906.
47
48

49 **Declaration of Interest**

50 There are no conflicts of interest to declare.
51
52

53
54
55 Word Count: 2193 words
56
57
58
59
60

References

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- ¹Hem H. Dholakia, Dhiman Bhadra, and Amit Garg, “Short term association between ambient air pollution and mortality and modification by temperature in five Indian cities,” *Atmospheric Environment* 99, (2014): 168–174.
- ²Regina Hampel, Johanna Lepeule, Alexandra Schneider, Sébastien Bottagisi, Marie-Aline Charles, Pierre Ducimetière, Annette Peters, and Rémy Slama, “Short-term impact of ambient air pollution and air temperature on blood pressure among pregnant women,” *Epidemiology* 22, no. 5 (2011): 671–679.
- ³Kristin A. Miller, David S. Siscovick, Lianne Sheppard, Kristen Shepherd, Jeffrey H. Sullivan, Garnet L. Anderson, and Joel D. Kaufman, “Long-term exposure to air pollution and incidence of cardiovascular events in women,” *The New England Journal of Medicine* 356, no. 5 (2007): 447–458.
- ⁴C. Arden Pope, Richard T. Burnett, George D. Thurston, Michael J. Thun, Eugenia E. Calle, Daniel Krewski, and John J. Godleski, “Cardiovascular mortality and long-term exposure to particle air pollution: Epidemiological evidence of general pathophysiological pathways of disease,” *Circulation* 109, no. 1 (2004): 71–77.
- ⁵Jesus A. Araujo, and Andre E. Nel, “Particulate matter and atherosclerosis: Role of particle size, composition and oxidative stress,” *Particle and Fibre Toxicology* 24 (2009): 6–24.
- ⁶Oyewale M. Morakinyo, Matlou I. Mokgobu, Murembiwa S. Mukhola, and Raymond P. Hunter, “Health outcomes of exposure to biological and chemical components of inhalable and respirable particulate matter,” *International Journal of Environmental Research and Public Health* 13, no. 6 (2016): 592.
- ⁷Muhammad Abu-Elmagd, Mansour A. Alghamdi, Madgy Shamy, Mamdouh I. Khoder, Max Costa, Mourad Assidi, Roaa Kadam, Haneen Alsehli, Mamdooh Gari, Peter N. Pushparaj, et al. “Evaluation of the effects of airborne particulate matter on bone marrow-mesenchymal stem cells (BM-MSCs): cellular, molecular and systems biological approaches,” *International Journal of Environmental Research and Public Health* 14 (2017): 440.
- ⁸Hirotohi Shima, Eiko Koide, Ritsuko Shinohara, and Takahiro Kobayashi, “Oxidative ability and toxicity of n-Hexane insoluble fraction of diesel exhaust particles,” *Toxicological Sciences* 91, no. 1 (2006): 218–226.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- ⁹Yoshito Kumagai, Sachie Koide, Keiko Taguchi, Akiko Endo, Yumi Nakai, Toshikazu Yoshikawa, and Nobuhiro Shimojo, “Oxidation of proximal protein sulfhydryls by phenanthraquinone, a component of diesel exhaust particles,” *Chemical Research in Toxicology* 15 (2002): 483–489.
- ¹⁰Qianshan Xiong, Haoran Yu, Runran Wang, Jinlai Wei, and Vishal Verma, “Rethinking dithiothreitol-based particulate matter oxidative potential: measuring dithiothreitol consumption versus reactive oxygen species generation,” *Environmental Science and Technology* 51 (2007): 6507–6514.
- ¹¹Wing Y. Tuet, Yunle Chen, Shierly Fok, Dong Gao, Rodney J. Weber, Julia A. Champion, and Nga L. Ng, “Chemical and cellular oxidant production induced by naphthalene secondary organic aerosol (SOA): Effect of redox-active metals and photochemical aging,” *Scientific Reports* 7 (2017): 15157.
- ¹²Jessica G. Charrier, and Cort Anastasio, “On dithiothreitol (DTT) as a measure of oxidative potential for ambient particles: Evidence for the importance of soluble transition metals,” *Atmospheric Chemistry and Physics* 12 (2012): 9321–9333.
- ¹³Yumi Motoyama, Kanae Bekki, Sang W. Chung, Ning Tang, Takayuki Kameda, Akira Toriba, Keiko Taguchi, and Kazuichi Hayakawa, “Oxidative stress more strongly induced by ortho- than para-quinoid polycyclic aromatic hydrocarbons in A549 cells,” *Journal of Health Science* 55, no. 5 (2009): 845–850.
- ¹⁴Qiangeng Li, Anna Wyatt, and Richard M. Kamens, “Oxidant generation and toxicity enhancement of aged-diesel exhaust,” *Atmospheric Environment* 43, no. 5 (2009): 1037–1042.
- ¹⁵Keiko Taguchi, Megumi Shimada, Sayako Fujii, Daigo Sumi, Xiaoqing Pan, Shigeru Yamano, Takahito Nishiyama, Akira Hiratsuka, Masayuki Yamamoto, Arthur K. Cho, et al. “Redox cycling of 9,10-phenanthraquinone to cause oxidative stress is terminated through its monoglucuronide conjugation in human pulmonary epithelial A549 cells,” *Free Radical Biology and Medicine* 44, no. 8 (2008): 1645–1655.
- ¹⁶Akira Torida, Chiharu Homma, Masahiro Kita, Waka Uozaki, Yaowatat Boongla, Walaiorn Orakij, Ning Tang, Takayuki Kameda, and Kazuichi Hayakawa, “Simultaneous determination of polycyclic aromatic hydrocarbon quinones by gas chromatography-tandem mass spectrometry, following a one-pot reductive trimethylsilyl derivatization,” *Journal of Chromatography A* 1459 (2016): 89–100.
- ¹⁷Ting Fang, Vishal Verma, Josephine T. Bates, Joseph Abrams, Mitchel Klein, Matthew J. Strickland, Stefanie E. Sarnat, Howard H. Chang, James A. Mulholland,

1
2
3 Paige E. Tolbert, Armistead G. Russell, and Rodney J. Weber, “Oxidative potential of
4 ambient water-soluble PM_{2.5} in the southeastern United States: Contrasts in sources
5 and health associations between ascorbic acid (AA) and dithiothreitol (DTT) assays”
6 *Atmospheric Chemistry and Physics* 16 (2016): 3865–3879.
7
8

9
10 ¹⁸Pourya Shahpoury, Tom Harner, Gerhard Lammel, Steven Lelieveld, Haijie Tong, and
11 Jake Wilson, “Development of an antioxidant assay to study oxidative potential of
12 airborne particulate matter” *Atmospheric Measurement Techniques* 12 (2019): 6529–
13 6539.
14
15

16
17 ¹⁹Kaspar R. Daellenbach, Gaëlle Uzu, Jianhui Jiang, Laure-Estelle Cassagnes, Zaira
18 Leni, Athanasia Vlachou, Giulia Stefenelli, Francesco Canonaco, Samuël Weber, Arjo
19 Segers et al., “Sources of particulate -matter air pollution and its oxidative potential in
20 Europe” *Nature* 587 (2020): 414–419.
21
22

23
24 ²⁰Maria C. Pietrogrande, Ilaria Bertoli, Francesco Manarini, and Mara Russo,
25 “Ascorbate assay as a measure of oxidative potential for ambient particles: Evidence
26 for the importance of cell -free surrogate lung fluid composition” *Atmospheric*
27 *Environment* 211 (2019): 103–112.
28
29

30
31 ²¹Lanfang Rao, Luying Zhang, Xingzi Wang, Tingting Xie, Shumin Zhou, Senlin Lu,
32 Xinchun Liu, Hui Lu, Kai Xiao, Weiqian Wang, and Qingyue Wang, “Oxidative
33 potential induced by ambient particulate matters with acellular assays: A review”
34 *Processes* 8 (2020): 1410.
35
36

37
38 ²²Josephine T. Bates, Ting Fang, Vishal Verma, Linghan Zeng, Rodney J. Weber, Paige
39 E. Tolbert, Joseph Y. Abrams, Stefanie E. Sarnat, Mitchel Klein, James A.
40 Mulholland, and Armistead G. Russell, “Review of acellular assays of ambient
41 particulate matter oxidative potential: Methods and relationships with composition,
42 sources, and health effects” *Environmental Science and Technology* 53 (2019): 4003–
43 4019.
44
45
46

47
48 ²³Prasanna Venkatachari, Philip K. Hopke, Brett D. Grover, and Delbert J. Eatough,
49 “Measurement of particle-bound reactive oxygen species in rubidoux aerosols”
50 *Journal of Atmospheric Chemistry* 50 (2005): 49–58.
51
52
53
54
55
56
57
58
59
60

Table 1. PAHQ concentrations in the reference materials of airborne particulates (unit: $\mu\text{mol}/\text{kg}$)¹⁶

	SRM1649b	SRM1648a	SRM2786	CRM No.28
4,5-PyQ	4.6	3.7	2.7	3.5
1,6-PyQ	0.47	0.44	0.76	0.67
1,8-PyQ	0.57	0.31	0.92	0.65
PQN	3.7	3.8	2.7	3.4
1,2-NQ	0.49	0.79	1.4	1.0
1,4-NQ	0.53	0.35	0.80	0.43

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Estimated oxidative ability of PAHQs in the reference materials of airborne particulates (unit: DTT- μ mol/min/kg)

	SRM1649b	SRM1648a	SRM2786	CRM No.28
4,5-PyQ	67	54	40	51
1,6-PyQ	0.70	0.65	1.1	0.99
1,8-PyQ	0.67	0.36	1.1	0.76
PQN	53	54	39	49
1,2-NQ	1.2	2.0	3.6	2.6
1,4-NQ	0.77	0.52	1.2	0.63

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures




Fig. 1




Fig. 2

Peer Review Only




Fig. 3

Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 4

1
2
3 **Figure captions**
4

5 Figure 1. Redox cycle causing the overproduction of H₂O₂ by ortho-PAHQ.
6

7
8 Figure 2. Blank-corrected rate of DTT consumption by five PAHQs: 4,5-PyQ, PQN,
9 1,2-NQ, 1,4-NQ and 1,6-/1,8-PyQ.
10

11
12 Figure 3. HPLC analysis of 4,5-PyQ in the test solution, before DTT assay (black line)
13 and after DTT assay (red line).
14

15
16
17 Figure 4. HPLC analysis of 1,6-/1,8-PyQ in the test solution, before DTT assay (black
18 line) and after DTT assay (red line).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60