



TITLE:

A complete dietary review of Japanese birds with special focus on molluscs

AUTHOR(S):

Morii, Yuta; Kitazawa, Munehiro; Squires, Theodore E.;
Watanabe, Megumi; Watanabe, Yoshiaki; Saito, Takumi;
Yamazaki, Daishi; Uchida, Akitomo; Machida, Yoshiyasu

CITATION:

Morii, Yuta ...[et al]. A complete dietary review of Japanese birds with special focus on molluscs. *Scientific Data* 2021, 8: 19.

ISSUE DATE:

2021

URL:

<http://hdl.handle.net/2433/276502>

RIGHT:

© The Author(s) 2021; This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

SCIENTIFIC DATA



OPEN

DATA DESCRIPTOR

A complete dietary review of Japanese birds with special focus on molluscs

Yuta Morii^{1,2}✉, Munehiro Kitazawa³, Theodore E. Squires⁴, Megumi Watanabe⁵, Yoshiaki Watanabe⁵, Takumi Saito⁶, Daishi Yamazaki⁷, Akitomo Uchida⁸ & Yoshiyasu Machida⁹

Birds often hold important positions in the food webs of ecosystems. As a result, interactions between birds and their prey have attracted attention not only in ecology, but also in fields like agriculture and conservation. Avian food resources are well researched in Japan, however there is no database critically reviewing molluscs as a food resource for birds. Here, we present a new database reviewing dietary information for all Japanese bird species. In addition to addressing general diet categories and specific food habits for each bird, we include detailed data on the molluscan prey observed for all species that consume them. The information within this database was collected through intense literary review to provide a complete look at bird species historically present around the country. We also include new information on snail species found in the upper digestive tract of harvested wild birds. This database is publicly available in the Zenodo repository. The information should aid research around the Japanese archipelago, especially projects involving birds or molluscs.

Background & Summary

Ornithology has long been one of the main fields of naturalist study and significant knowledge has been accumulated during its long history^{1,2}. Birds have some advantages for scientific research: (1) compared to other taxa, birds are easier to detect and identify via vocalization and are often less cryptic with fewer overall species than other macroorganisms³; (2) the occurrence, abundance, and reproductive success of birds has been shown to respond to environmental changes over many spatial scales³; (3) birds are often very popular flagship species for conservation and important for government policies^{4,5}; and (4) the large volume of accumulated ornithological knowledge allows for strong review and cross-referencing of new research^{6,7}.

Birds assume many fundamental ecological functions in the maintenance of local to continental-scale ecosystems. Their roles include dispersion of seeds, pollination, and long distance transfer and/or deposition of nutrients⁸⁻¹². At the same time, birds are often in important positions as keystone and/or umbrella species in various community food webs¹³⁻¹⁷. In particular, the predator-prey interactions between birds and insects are well known; more than 50% of avian species are predominantly insectivorous, and nearly 75% prey on invertebrates at least occasionally^{2,8,16}. Birds can reduce the densities of herbivorous insects in agriculture and may exert top-down effects on primary producers to the extent that their removal instigates trophic cascades^{18,19}. Various investigations have shown that birds contribute significantly to agricultural success especially where they prey upon insects that are detrimental to human activities^{17,20}. Thus, it is important to understand basic information about the food preferences of each bird species not only for ecological science but also for practical applications.

¹The Hakubi Project, Kyoto University, Oiwakecho, Kitashirakawa, Sakyo-ward, Kyoto, 6068501, Japan.

²Laboratory of Animal Ecology, Department of Zoology, Graduate School of Science, Kyoto University, Oiwakecho, Kitashirakawa, Sakyo-ward, Kyoto, 6068502, Japan. ³Graduate School of Agriculture, Hokkaido University, Nishi 9, Kita 9, Kita-ward, Sapporo, Hokkaido, 0608589, Japan. ⁴Kenkyu Services, 12511 84th Ave NE, Kirkland, WA, 98034, United States of America. ⁵Komabakita 4-5-5, Abashiri, Hokkaido, 0930033, Japan. ⁶Department of Biology, Faculty of Science, Toho University, Miyama 2-2-1, Funabashi, Chiba, 2748510, Japan. ⁷Center for Northeast Asian Studies, Tohoku University, 41 Kawauchi, Aoba-ward, Sendai, Miyagi, 9808576, Japan. ⁸Tokiwacho 3-16-71, Kitami, Hokkaido, 0900817, Japan. ⁹Bihoro Museum, Midori 253-4, Bihoro, Hokkaido, 0920002, Japan. ✉e-mail: yutamorii@gmail.com

Ten variables treated in the database	Explanations of factor labels for each variable
1. the distribution and breeding status	The bird distribution and residency in each area of Japan. RB : resident breeder, MB : migrant breeder, WV : winter visitor, PV : passage visitor, FB : former bred, or —: not distributed, rare or unknown (see also Fig. 1).
2. the endemism in Japan	Endemic , or —: not endemic to Japan.
3. the species status in the Japanese Red List	The seven categories of species status based on the 2020, 4th Version of the Japanese Red Lists. EX : extinct, CR : critically endangered, EN : endangered, VU : vulnerable, NT : near threatened, DD : data deficient, or —: common species or not listed.
4. main habitat	Main habitat. Terrestrial , Freshwater , and/or Marine , or Unknown .
5. dietary categories	I : carnivore, II : herbivore, IV : omnivore, or Unknown (see also Fig. 2).
6. main diet(s)	Main diet(s). I : some of animals, II : some of plants, I-i : fishes, I-ii : vertebrates, I-iii : arthropods, I-iv : molluscs, I-v : unknown or other animals, II-fr : plants (fruits and/or seeds), III : scavenger, or Unknown (see also Fig. 2).
7. all recorded food habits	All recorded food habits of each bird species. I-i : fishes, I-ii : vertebrates, I-iii : arthropods, I-iv : molluscs, I-v : unknown or other animals, II-fr : plants (fruits and/or seeds), II-le : plants (leaves and/or others), or III : scavenger (see also Fig. 2).
8. molluscs as avian food resources	Six categories of molluscs as avian food resources (I-iv). iv-t : terrestrial molluscs, iv-f : freshwater molluscs, iv-mg : marine gastropods, iv-mb : marine bivalves, iv-mc : marine cephalopods, or iv-o : others or unknown molluscs (see also Fig. 2).
9. descriptions of molluscan prey in literature	The descriptions of molluscan prey and their taxonomies in literature.
10. referenced bibliographies	Referenced papers and books (shown in the reference list on Zenodo).

Table 1. Description of each variable, and factor levels.

It is well established that birds prey on molluscs in various environments. Many species of sea ducks (Anatidae, tribe Mergini) prey on bivalves and have economic impacts on mussel farms worldwide²¹. The Snail Kite (*Rostrhamus sociabilis*; Accipitridae, Accipitriformes) and Limpkin's (*Aramus guarauna*; Aramidae, Gruiformes) diet consists almost entirely of freshwater snails in the Americas^{22–26}. Even for land snails, omnivorous birds are considered as regular predators (e.g. *Turdus* spp.; Muscicapidae, Passeriformes)^{27–30}. Another example, the Great Tit (*Parus major*; Paridae, Passeriformes) and some other forest passerines in Eurasia take land snails as the main calcium source for eggshell production²⁸. Despite these well documented instances, the scope of interactions between birds and molluscs are much less surveyed than between birds and insects.

Avian ecological databases have been established in several regions (e.g. Europe³¹, North America [<https://www.audubon.org>], Australia [<https://birddata.birdlife.org.au>], New Zealand [<http://nzbirdsonline.org.nz>], the North Pacific³², and Antarctica³³) while others have global scope (e.g. BirdLife International [<http://www.birdlife.org>], Avibase [<https://avibase.bsc-eoc.org>], AS@S [<http://seabirds.saeon.ac.za>], and Birds of the World [<https://birdsoftheworld.org>]). Several platforms for citizen science also contribute to avian knowledge (e.g. eBird [<https://ebird.org>], iNaturalist [<https://www.inaturalist.org>] and GBIF [<https://www.gbif.org>]). However, most of these databases mainly focus attention on avian distributions, and often do not provide great detail on dietary preferences or predator-prey interactions^{24,27}. Moreover, very few databases recorded prey species in these categories instead providing general overviews of diet.

Among regions there tends to be strong observation and research biases dependent on population, economy, language, and resource availability. These area-based biases can skew understandings of real diversity on earth and prevent comprehensive insights from being made. In this study, we investigated the diet of all bird species in Japan, with a special focus on molluscan prey. The Japanese archipelago and associated islands are located from the subarctic to the subtropical zone, therefore this area has high biodiversity and the region is recognized as a hotspot for the planet³⁴. Similarly to other regions, ornithology in Japan has a long history and rich data to work from, but few researchers have reviewed the food preferences of all local bird species. Moreover, many publications reporting on the food composition of Japanese birds were written in Japanese, so it is difficult to disseminate this knowledge widely given the language barrier. There are some excellent works and databases showcasing the ecological traits of the entire avian fauna of Japan^{35–38}, but they are only available in Japanese.

Most databases fail to identify molluscan prey at the species level despite wide variability within the phylum. It is important to determine the exact interactions between predator and prey at the species level due to ecological differences and unique conservation considerations for each party to a predatory interaction. In this study, we attempted to classify the main diet of all Japanese bird species, list detailed information on molluscs in avian diets, and evaluate molluscs as a food resource for birds based on available literature. We also newly record five species of land snails in the crop and gizzard of two species of birds hunted on Hokkaido, the northernmost island of Japan. These data are also included in our up-to-date database.

Methods

Classification of dietary preferences and habitats for bird species in Japan via literature. We reviewed the food habits of 633 native avian species listed in the Check-list of Japanese Birds, 7th Revised Edition³⁹ in attempting to represent the whole avian fauna of Japan. Nine ecological traits related to distribution, habitat and diet are listed in our database along with references as shown below (Table 1): (1) the distribution and breeding status in each region of Japan (Fig. 1), (2) the endemism in Japan (Endemic, or —: not endemic to Japan)³⁹, (3) the species status in the Red List of Threatened Species of Japan, (4) main habitat (Terrestrial, Freshwater, and/or Marine, or Unknown), (5) dietary categories (I: carnivore, II: herbivore, IV: omnivore, or Unknown; Fig. 2), (6) main diet(s) (I: some animals, II: some plants, I-i: fishes, I-ii: vertebrates, I-iii: arthropods, I-iv: molluscs, I-v:




Fig. 1 Seven categories of distribution area in this study.

unknown or other animals, II-fr: plants [fruits and/or seeds], and/or III: scavenger, or Unknown; Fig. 2), (7) all recorded food habits (I-i, I-ii, I-iii, I-iv, I-v, II-fr, II-le: plants [leaves and/or others], or III; Fig. 2), (8) molluscs as avian food resources (iv-t: terrestrial molluscs, iv-f: freshwater molluscs, iv-mg: marine gastropods, iv-mb: marine bivalves, iv-mc: marine cephalopods, or iv-o: others or unknown molluscs; Fig. 2), (9) descriptions of molluscan prey in literature, and (10) referenced bibliographies.

To keep our findings relevant, we reviewed the validity of species binomial names listed in our database and provide updates reflective of current taxonomic knowledge in 2020. A review was conducted using the Birds of the World online research database³⁶ and apparent updates to binomials were cross-referenced using the




Fig. 2 The categories of preferred foods in this study. Food preferences were first categorized into four big groups (I. carnivore, II. herbivore, III. scavenger and IV. omnivore), and two of them (I and II) were further separated. In particular, molluscs were classified in detail.

International Union for Conservation of Nature's Red List of Threatened Species (<https://www.iucnredlist.org>). The updated binomial information is included in Online-only Table 1.

We roughly categorised seven regions for avian distribution in the Japanese archipelago (Hok: Hokkaido Island and/or surrounding islands, Hon: Honshu Island and/or surrounding islands, Shi: Shikoku Island and/or surrounding islands, Kyu: Kyushu Island and/or surrounding islands, Ryu: Ryukyu archipelago, Izu: Izu islands, Oga: Ogasawara islands; Fig. 1), and classified six categories for residency in each region of Japan (RB: resident breeder, MB: migrant breeder, WV: winter visitor, PV: passage visitor, FB: former breeder, or —: not distributed, rare, or unknown) based on the Check-list of Japanese Birds, 7th Revised Edition³⁹, and added seven categories for the species status in Japan based on the 2020, 4th Version of the Japanese Red Lists (EX: extinct, CR: critically endangered, EN: endangered, VU: vulnerable, NT: near threatened, DD: data deficient, or —: common species or not listed)⁵. To determine each species' main diet, we primarily focused on literature describing "preferred" or "main" food habits, although we also utilized information about the frequency of target foods in crop and gizzard contents. The taxonomies of molluscan prey written in the database were mainly based on MolluscaBase (<http://www.molluscabase.org>), the online database of world mollusc classifications. While our database does not contain perfect information on distribution, residency, and conservation status in terms of current knowledge, we believe it represents a high degree of accuracy and usefulness in pulling together comprehensive information from different sources.

The diet data in this study was collected from 165 scientific articles and books including dietary information on Japanese birds. We searched for the following two series of keywords in Google Scholar for each bird species: {"scientific name" AND ["food habits" OR "diet" OR "food habits (in Japanese)" OR "crop and gizzard contents (in Japanese)"]} and {"standard Japanese name (in Japanese)" AND ["food habits" OR "diet" OR "food habits (in Japanese)" OR "crop and gizzard contents (in Japanese)"]}. Keyword searching and browsing was conducted between 2nd May and 27th December in 2017, and the top one-hundred and all results for each series of keywords was checked, respectively. Moreover, we manually reviewed additional several literatures and books as possible. These included publications in English and Japanese and were published between 1913 and 2018. All 165 references citing the food habits for each bird species are recorded in the database, and listed on the reference list in Zenodo⁴⁰.

Land snails detected from the crop and gizzard of two bird species in Hokkaido, Japan. Crop and gizzard samples were obtained from two juvenile Oriental Turtle-Doves (*Streptopelia orientalis*; Columbidae,




Fig. 3 (A,B) Two bird species investigated in this study, *Streptopelia orientalis* (A), and *Tetrastes bonasia* (B). (C–H) The prey items detected from avian crops and gizzards of *S. orientalis*, (C) *Cochlicopa lubrica* (Cochlicopidae, Stylommatophora), (D) *Discus pauper* (Discidae, Stylommatophora), (E) *Karaftohelix gainesi* (Camaenidae, Stylommatophora), (F) *Parakaliella affinis* (Helicarionidae, Stylommatophora), (G) *Persicaria thunbergii* (Polygonaceae, Caryophyllales), and (H) *Schizopepon bryoniifolius* (Cucurbitaceae, Cucurbitales). I. The photograph of crop and gizzard contents of *T. bonasia*.

Columbiformes; Fig. 3A) and one juvenile Hazel Grouse (*Tetrastes bonasia*; Phasianidae, Galliformes; Fig. 3B). An individual *T. bonasia* was hunted at Ubaranai site no. 1 (Abashiri City, Hokkaido, Japan; N 43.9678°, E 144.0414°) on 7 November 2013, and two *S. orientalis* were shot at Ubaranai site no. 2 (Abashiri City, Hokkaido, Japan; N 43.9261°, E 144.0406°) on 28 October 2016. These birds were shot by a professional hunter for food and stored in a freezer; we then received them from the hunter and carefully extracted the crop and gizzard contents. Crop and gizzard contents of *T. bonasia* were identified from a photograph, while those of *S. orientalis* were identified directly from samples. In addition, the combined weight of crop and gizzard contents were measured for both *S. orientalis* individuals using an electronic scale (wet and dry weights for one, and dry weight only for the other; Online-only Table 2). The data collected from these samples is also included in our database.

Data Records

The database for the ecological traits of birds in Japan is available on Zenodo⁴⁰ with the original database in XLSX format (i.e. “Whole_Database.xlsx” on Zenodo), the reference list correspond to the database in PDF and CSV format (i.e. “Reference_List.pdf” and “Reference_List.csv” on Zenodo), and ten different data files for each trait category with described names provided in CSV format. For CSV format files on Zenodo, we used “0” and “1” instead of “–” and “+” in original database, respectively, “Unknown” for the missing data of mid three traits (i.e. variable no. 4, 5 and 6) like as in original database, and “NULL” for the non-existing values on the last four variables (i.e. variable no. 7, 8, 9 and 10). We also included the literary references as numbers in the data files as shown in “Reference_List.pdf” and “Reference_List.csv” on Zenodo. The first row of each CSV data file shows the description of each file, the second row includes a header indicating the names of variables, and the following rows present data for each bird species in Japan. The crop and gizzard contents from two individuals of *S. orientalis* used in this study were stored in Bihoro Museum (Bihoro, Hokkaido, Japan; Specimen IDs: BIHM0300372 and BIHM0300373).

Technical Validation

All records included in the database are based on articles in scientific journals and books; therefore we have confidence in their accuracy. We also listed the references cited in the database, making it possible for users to access the original sources. Moreover, some bird specialists, MK and T-Squires and mollusc specialists, Y-Morii, T-Saito and DY have carefully checked the database for possible errors and included information on outdated or updated avian binomials in Online-only Table 1.

Code availability

No code was used in this study.

Received: 16 July 2020; Accepted: 4 December 2020;

Published online: 20 January 2021

References

1. Jones, J. Habitat selection studies in avian ecology: A critical review. *Auk* **118**, 557–562 (2001).
2. Whelan, C. J., Sekercioglu, C. H. & Wenny, D. G. Why birds matter: from economic ornithology to ecosystem services. *J. Ornithol.* **156**, S227–S238 (2015).
3. Carignan, V. & Villard, M. Selecting indicator species to monitor ecological integrity: A Review. *Environ. Monit. Assess* **78**, 45–61 (2002).
4. Senzaki, M., Yamaura, Y. & Nakamura, F. The usefulness of top predators as biodiversity surrogates indicated by the relationship between the reproductive outputs of raptors and other bird species. *Biol. Conserv.* **191**, 460–468 (2015).
5. Ministry of the Environment Government of Japan. *The 4th Version of the Japanese Red Lists.* (2012).
6. Farber, P. L. The development of taxidermy and the history of ornithology. *ISIS* **68**, 550–566 (1977).
7. Mayr, E. Commentary: the contributions of ornithology to biology. *Bioscience* **34**, 250–255 (1984).
8. Sekercioglu, C. H. Increasing awareness of avian ecological function. *TRENDS Ecol. Evol.* **21**, 464–471 (2006).
9. Sekercioglu, C. H. Bird functional diversity and ecosystem services in tropical forests, agroforests and agricultural areas. *J. Ornithol.* **153**, S153–S161 (2012).
10. Anderson, S. H., Kelly, D., Ladley, J. J., Molloy, S. & Terry, J. Cascading effects of bird functional extinction reduce pollination and plant density. *Science* **331**, 1068–1071 (2011).
11. Bauer, S. & Hoyer, B. J. Migratory animals couple biodiversity and ecosystem functioning. *Science* **344**, 1242552 (2014).
12. Pauw, A. A bird's-eye view of pollination: biotic interactions as drivers of adaptation and community change. *Annu. Rev. Ecol. Evol. Syst.* **50**, 477–502 (2019).
13. Tews, J. *et al.* Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *J. Biogeogr.* **31**, 79–92 (2004).
14. Robb, G. N., McDonald, R. A., Chamberlain, D. E. & Bearhop, S. Food for thought: supplementary feeding as a driver of ecological change in avian populations. *Front. Ecol. Environ.* **6**, 476–484 (2008).
15. Sharam, G. J., Sinclair, A. R. E. & Turkington, R. Serengeti birds maintain forests by inhibiting seed predator. *Science* **325**, 51 (2009).
16. Wenny, D. G. *et al.* The need to quantify ecosystem services provided by birds. *Auk* **128**, 1–14 (2011).
17. Nyffeler, M., Şekercioglu, Ç. H. & Whelan, C. J. Insectivorous birds consume an estimated 400–500 million tons of prey annually. *Sci. Nat* **105**, 47 (2018).
18. Whelan, C. J., Wenny, D. G. & Marquis, R. J. Ecosystem services provided by birds. *Ann. N. Y. Acad. Sci* **1134**, 25–60 (2008).
19. Mäntylä, E., Klemola, T. & Laaksonen, T. Birds help plants: a meta-analysis of top-down trophic cascades caused by avian predators. *Oecologia* **165**, 143–151 (2011).
20. Pejchar, L. *et al.* Net effects of birds in agroecosystems. *Bioscience* **68**, 896–904 (2018).
21. Varennes, É., Hanssen, S. A., Bonardelli, J. & Guillemette, M. Sea duck predation in mussel farms: the best nets for excluding common eiders safely and efficiently. *Aquacult. Env. Interact* **4**, 31–39 (2013).
22. Sykes, P. W. Jr. The feeding habits of the snail kite in Florida, USA. *Colon. Waterbirds* **10**, 84–92 (1987).
23. Bourne, G. R. Differential snail-size predation by snail kites and limpkins. *Oikos* **68**, 217–223 (1993).
24. Reed, W. L. & Janzen, F. J. Natural selection by avian predators on size and colour of a freshwater snail (*Pomacea flagellata*). *Biol. J. Linn. Soc.* **67**, 331–342 (1999).
25. Tanaka, M. O., Souza, A. L. T. & Módena, É. S. Habitat structure effects on size selection of snail kites (*Rostrhamus sociabilis*) and limpkins (*Aramus guarauna*) when feeding on apple snails (*Pomacea* spp.). *Acta Oecologica* **30**, 88–96 (2006).
26. Cattau, C. E., Fletcher, R. J. Jr., Kimball, R. T., Miller, C. W. & Kitchens, W. M. Rapid morphological change of a top predator with the invasion of a novel prey. *Nat. Ecol. Evol* **2**, 108–115 (2018).
27. Graveland, J. Avian eggshell formation in calcium-rich and calcium-poor habitats: importance of snail shells and anthropogenic calcium sources. *Can. J. Zool.* **74**, 1035–1044 (1996).
28. Graveland, J. & van der Wal, R. Decline in snail abundance due to soil acidification causes eggshell defects in forest passerines. *Oecologia* **105**, 351–360 (1996).
29. Mänd, R., Tilgar, V. & Leivits, A. Calcium, snails, and birds: a case study. *Web Ecol* **1**, 63–69 (2000).
30. Barker, G. M. *Natural Enemies of Terrestrial Molluscs.* (Landcare Research Hamilton, 2004).
31. Storchová, L. & Hořák, D. Data from: Life-history characteristics of European birds. *Dataset* <https://doi.org/10.5061/dryad.n6k3n> (2018).
32. Drew, G. S. & Piatt, J. F. North Pacific Pelagic Seabird Database (NPPSD): U.S. geological survey data release. <https://doi.org/10.5066/F7WQ01T3> (2015).
33. Humphries, G. R. W. *et al.* Mapping Application for Penguin Populations and Projected Dynamics (MAPPPD): Data and tools for dynamic management and decision support. *Polar Rec. (Gr. Brit)* **53**, 160–166 (2017).
34. Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M. & Gascon, C. In *Biodiversity hotspots: Distribution and protection of conservation priority areas* (eds Zachos, F. E. & Habel, J. C.) Ch. 1 (Springer, 2011).
35. Yamashina, Y. *Birds in Japan and their ecology (in Japanese)*. (Iwanamishoten, 1941).
36. Yamashina, Y. Researches on food habits of Japanese birds, I. Food habits of the birds of the Izu Is. (in Japanese). *Tori* **11**, 1–46 (1941).
37. Kiyosu, Y. *The birds of Japan. I. (in Japanese)*. (Kodansha, 1952).
38. Takagawa, S. *et al.* JAVIAN Database: a species-level database of life history, ecology and morphology of bird species in Japan (in Japanese). *Bird Res* **7**, R9–R12 (2011).
39. The Ornithological Society of Japan. *Check-list of Japanese Birds 7th Revised Edition.* (2012).
40. Morii, Y. *et al.* A complete dietary review of Japanese birds with special focus on molluscs. *Zenodo* <https://doi.org/10.5281/zenodo.4293296> (2020).

Acknowledgements

We are grateful to Futoshi Nakamura, Craig Barnett, Masayuki Senzaki, Keita Kawajiri, Reiko Kurosawa, and Atsumi Morii for scientific discussions and data collections.

Author contributions

Y.-Morii designed the database and wrote the manuscript with T.-Squires. Y.-Morii, M.K., M.W., Y.W., A.U. and Y.-Machida collected the data, and Y.-Morii, M.K., T.-Squires, T.-Saito and D.Y. carefully reviewed and updated the database.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Y.M.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files associated with this article.

© The Author(s) 2021