

Proceeding Paper

# Exploring the Involvement of the *Alternative Respiratory Pathway* in *Pisum sativum* L. Seed Germination <sup>†</sup>

Lénia Rodrigues <sup>1</sup>, Ricardo Claudino <sup>2</sup>, Steven P.C. Groot <sup>3</sup>, Pierre Hohmann <sup>4</sup>, Amaia Nogales <sup>5</sup>, Lee Hansen <sup>6</sup> and Hélia Cardoso <sup>1,\*</sup>

<sup>1</sup> MED (Instituto Mediterrâneo para a Agricultura, Ambiente e Desenvolvimento), IIFA (Instituto de Investigação e Formação Avançada), Universidade de Évora, Pólo da Mitra, Ap. 94, 7002-554 Évora, Portugal; email1@gmail.com

<sup>2</sup> Escola de Ciências e Tecnologia, Universidade de Évora, Colégio Luís António Verney, Rua Romão Ramalho, 7000-671 Évora, Portugal; email2@gmail.com

<sup>3</sup> Wageningen Plant Research, Wageningen University & Research, Wageningen, The Netherlands; email3@gmail.com

<sup>4</sup> IRTA Institute of Agrifood Research and Technology, Sustainable Plant Protection Programme, Fruitcentre, 25003 Lleida, Spain; email4@gmail.com

<sup>5</sup> LEAF (Linking Landscape, Environment, Agriculture and Food) Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda, 1349-017 Lisboa, Portugal; email5@gmail.com

<sup>6</sup> Department of Chemistry and Biochemistry, Brigham Young University, Provo, UT 84602, USA; email6@gmail.com

\* Correspondence: liar@uevora.pt or hcardoso@uevora.pt

<sup>†</sup> Presented at the 1st International Electronic Conference on Horticulturae, 16–30 April 2022; Available online: <https://iecho2022.sciforum.net/>.

**Abstract:** Organic agriculture, recognized as a more sustainable agricultural system, strongly depends on the use of highly resilient genotypes. Resilient seeds, with increased tolerance to germinate and provide vigorous seedlings under environmental stresses, currently represent one of the most important agronomical traits. Seed germination involves the activation of several metabolic pathways, including cellular respiration. Alternative oxidase (AOX), a key enzyme in the alternative respiratory pathway, plays a crucial role in regulating cell reprogramming by controlling metabolic transitions related to the cellular redox state and the variable carbon balance. The involvement of the alternative respiratory pathway during germination was explored by analysis of *PsAOX* gene/protein expression. Seeds of four *Pisum sativum* L. cultivars ('Respect-1', 'S134', 'G78' and 'S91') were imbibed in sterile tap water for 16 h and metabolic parameters measured by calorimetry (heat and CO<sub>2</sub> emission rates) in a Multi-Cell Differential Scanning Calorimeter in isothermal mode at 25 °C. The involvement of *PsAOX* was evaluated by transcript quantification (*PsAOX1*, *PsAOX2a*, and *PsAOX2b*) through RT-qPCR, and by of analysing the *PsAOX* expression through Western blot. The results demonstrate that the cv. 'S91', characterized by a low germination rate, exhibited the lowest metabolic heat and CO<sub>2</sub> emission rate. However, contrary to expectations, *PsAOX* transcript accumulation and *PsAOX* protein expression were significantly higher for 'S91' than for the other cultivars. These results indicate that higher levels of AOX (transcript and protein) could be linked to lower metabolic rates for embryo growth when seed germination is compromised.

**Keywords:** LIVESEED; pea; AOX; alternative respiratory pathway; seed germination

**Citation:** Rodrigues, L.; Claudino, R.; Groot, S.; Hohmann, P.; Nogales, A.; Hansen, L. Exploring the Involvement of the *Alternative Respiratory Pathway* in *Pisum sativum* L. Seed Germination. *Biol. Life Sci. Forum* **2022**, *2*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Yuyang Zhang

Published: 16 April 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Legumes have been referred to as the basis of a healthy diet, representing the most prominent source of food protein [1]. Among the cultured legumes, pea (*Pisum sativum* L.) is one of the most widely spread crops in Europe, playing a very important role in human nutrition and animal feed [2]. Considering the importance of this grain legume

together with the selection of organic agriculture as a more sustainable agricultural system, the development of more resilient cultivars has been recognized as a key strategy. Seed germination and seedling establishment are among the most critical stage and key factors for successful crop production [3]. In order to support the imposed needs, several breeding programs have been implemented aiming to develop new cultivars characterized by seeds with greater plasticity toward environmental pressures, allowing them to more efficiently germinate. The European project LIVESEED [4] which integrates efforts by seed production companies and research/breeding Institutes, is one example of the high interest in seed research.

Considering that germination involves the activation of several metabolic pathways, including cellular respiration to provide the required energy, the involvement of the alternative respiratory pathway during germination was explored. Alternative oxidase (AOX) is a mitochondria inner membrane enzyme with a key role in the alternative respiratory pathway, in which this enzyme introduces a branch into the electron transport chain (ETC) at the ubiquinone pool. In this branch, AOX allows the transport of electrons to oxygen directly from the ubiquinone, preventing excessive reduction of the downstream complexes [5]. The involvement of this pathway in plant response to a diversity of environmental stresses and cell reprogramming events has been demonstrated [6,7], mainly associated with the ability to contribute to the control of reactive oxygen species (ROS), thereby reducing oxidative damage [8]. In addition, a link between AOX gene expression and metabolic and respiratory parameters was established in different plant species under biological systems/stress conditions [9]. Considering that seed germination involves the activation of several metabolic pathways, metabolic and respiratory changes were monitored by using calorimetry. Calorimetry simultaneously measures metabolic heat rates ( $R_q$ ) and  $\text{CO}_2$  emission rates ( $R_{\text{CO}_2}$ ) of biological samples and has been used as a screening tool to assess metabolic and respiratory changes associated with cell reprogramming events [10].

The present research investigates the involvement of PsAOX on efficient seed germination, and explores the link between AOX and metabolic/respiratory changes.

## 2. Methods

Seeds of four *P. sativum* L. cultivars ('Respect-1', 'S134', 'G78' and 'S91') provided by LIVESEED partners were selected for calorimetry measurements. Previously, seeds were imbibed in sterile tap water for 16 h. Calorimetric parameters (heat and  $\text{CO}_2$  emission rates) were measured at 25 °C using a Multi-Cell Differential Scanning Calorimeter in isothermal mode. A total of 16 biological replicates (16 seeds) were considered per cultivar. Heat rate ( $R_q$ ) and  $\text{CO}_2$  emission rate ( $R_{\text{CO}_2}$ ), were determined according to Rodrigues et al. [11]. Germination rates were recorded 6 days after calorimetric measurements.

The involvement of PsAOX in seed germination was evaluated by gene expression analysis of the three *P. sativum* AOX genes (*PsAOX1*, *PsAOX2a*, and *PsAOX2b*), and by the analysis of AOX protein levels through Western blot. Samples of the four pea cultivars were collected at 16 h post imbibition and further homogenized using liquid nitrogen. A total of 12 samples, each one consisting of a pull of 4 seeds, were used per cultivar.

For *PsAOX* gene expression studies, the cv. 'Respect-1' and cv. 'S91' were selected based on the results obtained by calorimetry. Total RNA was extracted from the harvested material using the Maxwell<sup>®</sup> 16 LEV simplyRNA Cells Kit (Promega) and integrity evaluated through 1.2% agarose gel electrophoresis. 1  $\mu\text{g}$  of total RNA was used for cDNA synthesis using the SensiFAST<sup>™</sup> cDNA Synthesis Kit (Bioline). Transcript level accumulation of the three pea AOX genes (*PsAOX1*, *PsAOX2a* and *PsAOX2b*) was assessed in an ABI 7500 system using SensiFAST SYBR Lo-ROX kit (Bioline). *PsPOB* and *PsSAR1* were used as reference genes for data normalization. Primer specificity was evaluated by visualization of a single peak at the dissociation curve and efficiency (E) was calculated by the


equation:  $E (\%) = 10^{-(1/\text{slope}) - 1} \times 100$  [12]. To obtain the slope value a standard curve of a 4-fold dilution series was generated for each primer pair.

For analysis of AOX protein levels, all four cvs. were considered: 'Respect-1', 'S134', 'G78' and 'S91'. Total protein content was extracted by phenol precipitation from 50 mg of plant material (adapted from [13]) and total protein quantification of 12 replicates of each cultivar were performed using the Pierce 660 nm Protein Assay Reagent (Thermo Scientific™). Western blotting was used for comparison of the levels of PsAOX protein between the cultivars. Briefly, after protein separation by SDS PAGE (25 µg total protein from each sample) in 14% polyacrylamide gels, using Laemmli buffer system [14], proteins were transferred to a PDVF membrane by electroblotting using a *Semi-Dry Trans-Blot Turbo Transfer* (Bio-Rad) system. After transferring, blocking was performed with 5% non-fat milk powder in TBS-Tween 20 and the membrane was incubated with primary antibody Anti-AOX1/2 (Agriseria AS04054; dilution: 1:1000), overnight at 4 °C. AOX bands were detected with an alkaline phosphatase-linked secondary antibody (anti-rabbit, Agriseria AS09607, 1:10,000 dilution), using a chemifluorescent substrate (ECF Plus Western Blotting Detection Reagents, GE, Healthcare). Membranes were placed in a transilluminator (Bio-Rad Gel-doc system) and a semi-quantitative analysis of band intensity was carried out using the software Bio-Rad Image Lab 5.2.1. For Western blot analysis, six replicates of each cultivar were used.

Statistical analyses were performed by SPSS version 22.0. Normality and homoscedasticity were checked in all data. One-way ANOVA followed by a Tukey HSD test was used for the comparison of calorespirometric parameters, germination rate and AOX band volume between the four cultivars. The comparison of transcript level of *PsAOX1*, *PsAOX2a* and *PsAOX2d* between the two cultivars was made using the T-Student. When the data did not meet the assumptions to perform parametric tests, the Kruskal-Wallis or Mann-Whitney nonparametric tests were conducted. Statistical significance was considered at  $p < 0.05$ .

### 3. Results and Discussion

At 25 °C, calorespirometric parameters  $R_q$  and  $R_{CO_2}$  (Figure 1) were significantly lower in cvs. 'S91' and 'G78', compared to cvs. 'Respect-1' and 'S134'. Later, at 6 days post imbibition, seeds of cvs. 'G78', 'S134' and 'Respect-1' presented significantly higher germination rate, compared to the cv. 'S91'. These results show that higher  $R_q$  and  $R_{CO_2}$  could be related with higher germination rates, as previously reported by Edelstein et al. [15], who observed that higher values of metabolic activity were associated with higher germination rates of melon seeds. With seed lots from onion and several Brassica species a positive correlation has been shown between single seed respiration and seed quality [16].



**Figure 1.** Calorespirometric parameters for the four pea (*Pisum sativum* L.) cultivars ('S91', 'G78', 'Respect-1' and 'S134') measured at 25 °C, 16 h post imbibition. (A) Respiratory heat rate— $R_q$ , (B)  $CO_2$  production rate— $R_{CO_2}$ . Data are the mean value of 16 measurements  $\pm$  standard error. Different letters indicate significant differences between the cultivars. Statistical significance was considered for  $p < 0.05$ .

The AOX gene family is nucleus encoded, composed of one to six members distributed in two subfamilies (AOX1 and AOX2) [17]. In pea, the AOX gene family is composed by one AOX1-subfamily member (*PsAOX1*) and two AOX2-subfamily members (*PsAOX2a* and *PsAOX2b*) [18]. Based on calorimetric results, for *PsAOX* gene expression studies, the cv. 'Respect-1' and cv. 'S91' were selected. Regarding the analysis of *PsAOX* gene expression performed 16 h post water imbibition, a higher transcript accumulation of *PsAOX2b* was observed in comparison with *PsAOX1* and *PsAOX2a*. Additionally, transcript accumulation was significantly higher in cvs. 'S91' when compared with cv. 'Respect-1' (data not shown).

Previous studies with tobacco and *Arabidopsis* have demonstrated the importance of AOX1 in photosynthesis-related pathways and respiration under light conditions [19]. Members of the AOX1-subfamily have also been associated with plant responses to abiotic stress factors [6,7]. At 16 h post imbibition, the photosynthetic apparatus is still not activated which could explain the lower levels of *PsAOX1* transcript. Additionally, besides AOX involvement in ROS control, at the timepoint considered, the ROS levels were already reduced through the enzymatic ROS scavenging system and the action of antioxidant metabolites.

On the other hand, members of the AOX2-subfamily have been mainly associated with developmental processes [6]. The higher level of transcript accumulation achieved for *PsAOX2a* lead us to suggest a higher involvement of this gene in the germination process, particularly in case of cv. 'S91'.

The analysis of total protein concentration in the four *P. sativum* cultivars showed significantly higher protein concentration in cv. 'Respect-1' compared to cvs. 'S134', 'G78' and 'S91'. No statistically significant differences in total protein concentration were observed among cvs. 'S134', 'G78' and 'S91' (data not shown). The *PsAOX* protein levels were evaluated 16 h post water imbibition by Western blot. The band corresponding to the *PsAOX* protein was detected in all cultivars and higher levels of protein were observed in cv. 'S91' when compared to cvs. 'Respect-1' (RS), 'S134' and 'G78'. Although we were not able to analyze the different isoforms of the AOX protein through the Western Blot analysis, the results at protein level confirm the results observed by transcript analysis, i.e., a higher AOX protein expression in the cultivar that presents the highest level of AOX gene expression.

The absence of a direct relationship between the total protein concentration and the level of the AOX protein could be expected, since the AOX is only a part of the total protein concentration [20]. In fact, the values of total protein concentration reflect essentially the concentration of storage proteins which is variable among cultivars [21]. Moreover, studies focused on the protein composition of pea seeds showed that a large percentage of the identified proteins correspond to storage proteins (mainly albumin, legumin and vicilin), with the rest being involved in the response to biotic and abiotic stress, energy production, metabolism and storage of essential non-protein compounds [21].

Overall, seeds from the cultivar characterized by lower  $R_q$  and  $R_{CO_2}$  (cv. 'S91'), coincident with lower vigour, exhibited significantly higher expression of the *PsAOX* genes (*PsAOX1*, *PsAOX2a* and *PsAOX2d*) and *PsAOX* protein. From these results, we hypothesize that higher levels of AOX (transcript and protein) in germinating seeds are linked to lower metabolic rates and germination. Further studies will be required to validate this hypothesis.

#### 4. Conclusions

To the best of our knowledge, the study presented hereby represents the first approach aiming to establish a link between respiratory parameters, monitored by calorimetry, and *PsAOX* transcript accumulation/*PsAOX* protein expression during germination of pea seed. The present work suggests the involvement of AOX and the alter-

native respiratory pathway during seed germination and the applicability of calorimetry to assess seed vigor, pointing to this method as a useful non-destructive phenotyping tool for selecting genotypes with superior germination capacity.

**Author Contributions:**

**Funding:**

**Institutional Review Board Statement:**

**Informed Consent Statement:**

**Data Availability Statement:**

**Acknowledgments:** This work was supported by the EU project LIVESEED—Improve performance of organic agriculture by boosting organic seed and plant breeding efforts across Europe funded by the European Union’s HORIZON 2020 research and innovation programme under the Grant Agreement no 727230, and by the Swiss State Secretariat for Education, Research and Innovation (SERI) under contract number 17.00090; and by National Funds through FCT under the Project UIDB/05183/2020. Authors are very thankful to IIFA for the fellowship given to Lénia Rodrigues. The opinions expressed and arguments employed herein do not necessarily reflect the official views of the EC and the Swiss government. Neither the European Commission/SERI nor any person acting behalf of the Commission/SERI is responsible for the use which might be made of the information provided on this document.

**Conflicts of Interest:**

**References**

1. Singh, N. Pulses: An overview. *J. Food Sci. Technol* **2017**, *54*, 853–857.
2. Dhall, R.K. *Pea Cultivation*; Centre for Communication and International Linkages, Punjab Agricultural, University Ludhiana: 2017.
3. McPhee, K.E. Pea. In *Pulses, Sugar and Tuber Crops*; Springer: Berlin/Heidelberg, Germany, 2007; pp. 33–47.
4. LIVESSED, IFOAM Organics Europe; Consulted in 12 July 2021. Available online: <https://www.liveseed.eu/tools-for-practitioners/eu-organic-seed-databases/> (accessed on).
5. Saha, B.; Borovskii, G.; Panda, S.K. Alternative oxidase and plant stress tolerance. *Plant Signal. Behav.* **2016**, *11*, e1256530.
6. Velada, I.; Ragonezi, C.; Arnholdt-Schmitt, B.; Cardoso, H. Reference genes selection and normalization of oxidative stress responsive genes upon different temperature stress conditions in *Hypericum perforatum* L. *PLoS ONE* **2014**, *9*, e115206.
7. Campos, M.D.; Campos, C.; Nogales, A.; Cardoso, H.G. Carrot alternative oxidase (*DcAOX*) sub-family differentially responds to growth and to chilling. *Plants—Mol. Breed. Hortic. Plants* **2021**, *10*, 2369.
8. Mohanapriya, G.; Bharadwaj, R.; Noceda, C.; Costa, J.H.; Kumar, S.R.; Sathishkumar, R.; Thiers, K.L.; Santos Macedo, E.; Silva, S.; Annicchiarico, P.; et al. Alternative oxidase (AOX) senses stress levels to coordinate auxin-induced reprogramming from seed germination to somatic embryogenesis—A role relevant for seed vigor prediction and plant robustness. *Front. Plant Sci.* **2019**, *10*, 1134.
9. Campos, M.D.; Nogales, A.; Cardoso, H.G.; Sarma, R.K.; Nobre, T.; Sathishkumar, R.; Arnholdt-Schmitt, B. Stress-induced accumulation of *DcAOX1* and *DcAOX2a* transcripts coincides with critical time point for structural biomass prediction in carrot primary cultures (*Daucus carota* L.). *Front. Genet.* **2016**, *7*, 1.
10. Arnholdt-Schmitt, B.; Mohanapriya, G.; Sathishkumar, R.; Macedo, E.S.; Costa, J.H. Predicting biomass production from plant robustness and germination efficiency by calorimetry. In *Biofuels: Greenhouse Gas Mitigation and Global Warming*; Springer: New Delhi, India, 2018; pp. 81–94.
11. Rodrigues, L.; Nogales, A.; Hansen, L.; Santos, F.; Rato, A.E.; Cardoso, H. Exploring the applicability of calorimetry to assess seed metabolic stability upon temperature stress conditions—*Pisum sativum* L. used as a case study. *Front. Plant Sci.* **2022**, 703.
12. Radonic, A.; Thulke, S.; Mackay, I.M.; Landt, O.; Siegert, W.; Nitsche, A. Guideline to reference gene selection for quantitative real-time PCR. *Biochem. Biophys. Res. Commun.* **2004**, *313*, 856–862.
13. Mamontova, T.; Lukasheva, E.; Mavropolo-Stolyarenko, G.; Proksch, C.; Bilova, T.; Kim, A.; Babakov, V.; Grishina, T.; Hoehenwarter, W.; Medvedev, S.; et al. Proteome map of pea (*Pisum sativum* L.) embryos containing different amounts of residual chlorophylls. *Int. J. Mol. Sci.* **2018**, *19*, 4066.
14. Laemmli, U.K. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **1970**, *227*, 680–685.
15. Edelstein, M.; Bradford, K.J.; Burger, D.W. Metabolic heat and CO<sub>2</sub> production rates during germination of melon (*Cucumis melo* L.) seeds measured by microcalorimetry. *Seed Sci. Res.* **2001**, *11*, 265–272.
16. Bradford, K.J.; Bello, P.; Fu, J.C.; Barros, M. Single-seed respiration: a new method to assess seed quality. *Seed Sci. Technol.* **2013**, *41*, 420–438.

17. Cardoso, H.G.; Nogales, A.; Frederico, A.M.; Svensson, J.T.; Santos Macedo, E.; Valadas, V.; Arnholdt-Schmitt, B. Exploring AOX gene diversity. In *Alternative Respiratory Pathways in Higher Plants*; Gupta, K.J., Mur, L.A.J., Neelwarne, B., Eds.; Wiley-Blackwell: Hoboken, NJ, USA, 2015; pp. 241–254.
18. Sweetman, C.; Soole, K.L.; Jenkins, C.L.; Day, D.A. Genomic structure and expression of alternative oxidase genes in legumes. *Plant Cell Environ.* **2019**, *42*, 71–84.
19. Dahal, K.; Vanlerberghe, G.C. Alternative oxidase respiration maintains both mitochondrial and chloroplast function during drought. *New Phytol.* **2017**, *213*, 560–571.
20. Barac, M.; Cabrilo, S.; Pesic, M.; Stanojevic, S.; Zilic, S.; Macej, O.; Ristic, N. Profile and functional properties of seed proteins from six pea (*Pisum sativum*) genotypes. *Int. J. Mol. Sci.* **2010**, *11*, 4973–4990.
21. Dhaliwal, S.K.; Salaria, P.; Kaushik, P. Pea seed proteins: A nutritional and nutraceutical update. In *Grain and Seed Proteins Functionality*; IntechOpen: London, UK, 2021.