



# Draft Genome Sequences of 10 Bacterial Strains Isolated from Root Nodules of *Alnus* Trees in New Hampshire

Ian Davis,<sup>a</sup> Joseph Sevigny,<sup>a,b</sup> Victoria Kleiner,<sup>a\*</sup> Kelsey Mercurio,<sup>a</sup> Céline Pesce,<sup>a</sup> Erik Swanson,<sup>a</sup> W. Kelley Thomas,<sup>a,b</sup> Louis S. Tisa<sup>a</sup>

<sup>a</sup>Department of Molecular, Cellular, and Biomedical Sciences, University of New Hampshire, Durham, New Hampshire, USA

<sup>b</sup>Hubbard Center for Genome Studies, University of New Hampshire, Durham, New Hampshire, USA

**ABSTRACT** Here, we report the draft genome sequences obtained for 10 bacterial strains isolated from root nodules of *Alnus* trees. These members of the nodule microbiome were sequenced to determine their potential functional roles in plant health. The selected strains belong to the genera *Rhodococcus*, *Kocuria*, *Rothia*, *Herbaspirillum*, *Streptomyces*, and *Thiopseudomonas*.

Actinorhizal plants form a symbiotic association with members of the nitrogen-fixing actinobacterial genus *Frankia* that allows these plants to colonize stressed environments (1–6). Besides containing *Frankia* spp., the actinorhizal root nodules contain large numbers of other actinobacteria occupying the same microniche as *Frankia* spp. (7–13). Many non-*Frankia* actinobacteria that have been isolated from actinorhizal root nodules might contribute to nodulation or aid in plant growth and health (8, 9, 14). As expected, this situation is not unique to actinorhizal nodules but is found in all plants that form root nodule structures, including legume nodules (15–17). There is a growing body of evidence suggesting that both wild and cultivated legume nodules are not exclusively inhabited by rhizobia but contain diverse assemblages of nonrhizobial bacteria (15–18).

Root nodule samples were collected from *Alnus* trees found by Adam's Pond at Jackson's Laboratory in Durham, NH. The root nodules were surface sterilized with hydrogen peroxide and rinsed several times with sterile distilled water. The last wash of the sterilized nodule was incubated in LB medium to ensure that all epiphytes associated with the plant were removed. The nodule was cut into a fine powder with a sterilized razor, and dilutions were plated onto Czapeck and R2A media containing cycloheximide and nalidixic acid. About 60 isolates were initially obtained, purified, and propagated on either Czapeck or R2A medium (Table 1). These isolates were incubated overnight in their respective isolation medium (Table 1), and genomic DNA (gDNA) was extracted by the cetyltrimethylammonium bromide (CTAB) DNA extraction protocol (19). RNA was removed by RNase treatment. The quality and quantity of the gDNA were verified using a Thermo Scientific NanoDrop instrument. These isolates were initially identified by amplifying and sequencing their 16S rRNA genes. Based on these results, 10 isolates were chosen for whole-genome sequencing analysis to provide insight into their plant-microbe interactions, including potential plant growth-promoting activity.

Whole-genome sequencing was performed at the Hubbard Center for Genome Studies (University of New Hampshire, Durham, NH) using Illumina technology techniques (20). A paired-end library was constructed using a Nextera DNA library preparation kit (Illumina, San Diego, CA) and sequenced on an Illumina HiSeq 2500 instrument to produce 250-bp paired-end reads. The total numbers of reads for all 10 strains are listed in Table 1. The Illumina sequence data were trimmed using Trimmomatic version 0.36 (21). TruSeq adapters were trimmed with an allowance of two mismatches.

**Citation** Davis I, Sevigny J, Kleiner V, Mercurio K, Pesce C, Swanson E, Thomas WK, Tisa LS. 2020. Draft genome sequences of 10 bacterial strains isolated from root nodules of *Alnus* trees in New Hampshire. *Microbiol Resour Announc* 9:e01440-19. <https://doi.org/10.1128/MRA.01440-19>.

**Editor** Irene L. G. Newton, Indiana University, Bloomington

**Copyright** © 2020 Davis et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Louis S. Tisa, [louis.tisa@unh.edu](mailto:louis.tisa@unh.edu).

\* Present address: Victoria Kleiner, Department of Microbiology, Boston University, Boston, Massachusetts, USA.

This article is New Hampshire Agricultural Experiment Station scientific contribution 2836.

**Received** 15 November 2019

**Accepted** 12 December 2019

**Published** 9 January 2020

**TABLE 1** Genome statistics

Bacterial species	Isolate <sup>a</sup>	GenBank accession no.	No. of reads	No. of contigs	Avg coverage (×)	Genome assembly size (bp)	N <sub>50</sub> contig size (kb)	No. of CDSs <sup>b</sup>	G+C content (%)	No. of rRNAs	No. of tRNAs
<i>Rothia dentocariosa</i>	1C11	SPNF000000000	4,631,207	97	210.5	2,536,666	1,708.8	2,212	55	10	52
<i>Streptomyces</i> sp.	4R-3d	SPNN000000000	8,912,300	93	429.3	8,495,698	213.7	7,601	72	8	65
<i>Rhodococcus</i> sp.	1R11	SPNG000000000	5,379,096	38	82.9	5,478,616	346.9	5,102	64	8	58
<i>Kocuria rhizophila</i>	4R-31	SPNK000000000	6,695,169	26	571.5	2,659,245	245.8	2,225	70	9	57
	4R-34	SPNL000000000	2,094,786	122	71.2	2,654,140	39.4	2,211	71	4	46
<i>Herbaspirillum</i> sp.	3C11	SPNH000000000	2,844,408	337	186.0	5,552,861	30.9	5,278	60	1	50
	3R-11	SPNI000000000	4,677,477	11	328.3	5,219,991	1,116.6	4,736	59	3	46
	3R11	SPNJ000000000	14,415,312	10	356.0	5,221,161	1,116.6	4,738	60	3	46
	3R-3a1	QUQN000000000	6,064,775	13	170.0	5,221,518	771.5	4,741	59	3	39
<i>Thiopsedomonas</i> sp.	4R-3cl	SPNM000000000	6,340,931	405	212.9	2,981,747	15.5	2,995	72	14	65

<sup>a</sup> C and R represent Czapecck and R2A media, respectively, that were used to culture each isolate.

<sup>b</sup> CDSs, coding DNA sequences.

Leading and trailing bases below a quality of three were trimmed. The reads were then scanned with a sliding window of 4 bp and trimmed if the average quality dropped below 30. Finally, reads were dropped if the length was less than 36 bp. Trimmed sequencing reads were assembled using SPAdes version 3.13 (22), with the default settings. The assembled genomes were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (23). The assembly metrics and annotation features are given in Table 1. The identities of the strains were determined by whole-genome-based taxonomic analysis via the Type (Strain) Genome Server (TYGS) platform (24) (<https://tygs.dsmz.de>), including digital DNA:DNA hybridization (dDDH) values (25). The type-based species clustering using a 70% dDDH radius around each of the type strains was used as previously described (26), while subspecies clustering was done using a 79% dDDH threshold, as previously introduced (27). Among the 10 strains, new species of the genera *Streptomyces*, *Rhodococcus*, *Herbaspirillum*, and *Thiopseudomonas* were identified, including one *Herbaspirillum* subspecies. Bioinformatic analysis of these genomes by the use of the antiSMASH 4.0 program (28) revealed the presence of high numbers of secondary metabolic biosynthetic gene clusters. Many of these potential natural products should be involved in plant-microbe interactions and aid in their plant growth-promoting activities.

**Data availability.** The draft genome sequences of these bacterial strains have been deposited in GenBank under the accession numbers listed in Table 1. Both the assembly and raw reads are available at DDBJ/ENA/GenBank under BioProject number [PRJNA480027](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA480027).

## ACKNOWLEDGMENTS

We thank the following T3 course participants for their efforts on this project: Chhetri Saroja, Tsunemi Yamashita, Devin Thomas, Mohammad Alam, Rami M. Alroobi, Eric C. Atkinson, Nick Baer, Kayla Bieser, Nicolas Blouin, Louise J. Brogan, Jack Chen, Nicholas P. Edgington, Olivia L. George, Ghanshyam D. Heda, Amber Howerton, Jenna Luek, Paula Mazzer, KellyAnn Miller, Daniel P. Moore, Shallee T. Page, Judith L. Roe, Kevin E. Shuman, and Kristy Townsend.

Partial funding was provided by the New Hampshire Agricultural Experiment Station. This work was supported by USDA National Institute of Food and Agriculture Hatch 02281 (to L.S.T.), National Institutes of Health grant 5R25GM125674, "Innovative Programs to Enhance Research Training (IPERT)" (to W.K.T.), National Institutes of Health grant P20GM103506 of the IDeA program (to W.K.T.), and the College of Life Sciences and Agriculture at the University of New Hampshire. Sequencing was performed on an Illumina HiSeq2500 instrument purchased with NSF MRI grant DBI-1229361 (to W.K.T.).

## REFERENCES

- Benson DR, Dawson JO. 2007. Recent advances in the biogeography and genealogy of symbiotic *Frankia* and its host plants. *Physiol Plant* 130: 318–330. <https://doi.org/10.1111/j.1399-3054.2007.00934.x>.
- Benson DR, Silvester WB. 1993. Biology of *Frankia* strains, actinomycete symbionts of actinorhizal plants. *Microbiol Rev* 57:293–319.
- Ngom M, Gray K, Diagne N, Oshone R, Fardoux J, Gherbi H, Hocher V, Svistoonoff S, Laplace L, Tisa LS, Sy MO, Champion A. 2016. Symbiotic performance of diverse *Frankia* strains on salt-stressed *Casuarina glauca* and *Casuarina equisetifolia* plants. *Front Plant Sci* 7:1331. <https://doi.org/10.3389/fpls.2016.01331>.
- Ngom M, Oshone R, Diagne N, Cissoko M, Svistoonoff S, Tisa LS, Laplace L, Sy MO, Champion A. 2016. Tolerance to environmental stress by the nitrogen-fixing actinobacterium *Frankia* and its role in actinorhizal plants adaptation. *Symbiosis* 70:17–29. <https://doi.org/10.1007/s13199-016-0396-9>.
- Diagne N, Ngom M, Djighaly PI, Ngom D, Ndour B, Cissokho M, Faye MN, Sarr A, Sy MO, Laplace L, Champion A. 2015. Remediation of heavy-metal-contaminated soils and enhancement of their fertility with actinorhizal plants, p 355–366. In Sherameti I, Varma A (ed), *Heavy metal contamination of soils, soil biology*, vol 44. Springer International Publishing, Basel, Switzerland.
- Diagne N, Arumugam K, Ngom M, Nambiar-Veetil M, Franche C, Narayanan KK, Laplace L. 2013. Use of *Frankia* and actinorhizal plants for degraded lands reclamation. *Biomed Res Int* 2013:948258. <https://doi.org/10.1155/2013/948258>.
- Valdés M, Pérez N-O, Estrada-de Los Santos P, Caballero-Mellado J, Peña-Cabrales JJ, Normand P, Hirsch AM. 2005. Non-*Frankia* actinomycetes isolated from surface-sterilized roots of *Casuarina equisetifolia* fix nitrogen. *Appl Environ Microbiol* 71:460–466. <https://doi.org/10.1128/AEM.71.1.460-466.2005>.
- Ghodhbane-Gtari F, Nouioui I, Hezbri K, Lundstedt E, D'Angelo T, McNutt Z, Laplace L, Gherbi H, Vaissayre V, Svistoonoff S, Ahmed HB, Boudabous A, Tisa LS. 2019. The plant-growth-promoting actinobacteria of the genus *Nocardia* induces root nodule formation in *Casuarina glauca*. *Antonie Van Leeuwenhoek* 112:75–90. <https://doi.org/10.1007/s10482-018-1147-0>.
- Ghodhbane-Gtari F, Tisa LS. 2014. Ecology and physiology of non-*Frankia* actinobacteria from actinorhizal plants, p 27–42. In Katsy El (ed), *Plasticity in plant-growth-promoting and phytopathogenic bacteria*. Springer, New York, NY.
- Ghodhbane-Gtari F, Nouioui I, Salem K, Ktari A, Montero-Calasanz M, d C, Tisa LS, Klenk H-P, Gtari M. 2014. *Nocardia casuarinae* sp. nov., an actinobacterial endophyte isolated from root nodules of *Casuarina glauca*. *Antonie Van Leeuwenhoek* 105:1099–1106. <https://doi.org/10.1007/s10482-014-0168-6>.

11. Ghodhbane-Gtari F, Essoussi I, Chattaoui M, Chouaia B, Jaouani A, Dafonchio D, Boudabous A, Gtari M. 2010. Isolation and characterization of non-*Frankia* actinobacteria from root nodules of *Alnus glutinosa*, *Casuarina glauca* and *Elaeagnus angustifolia*. *Symbiosis* 50:51–57. <https://doi.org/10.1007/s13199-009-0029-7>.
12. Carro L, Pujic P, Trujillo ME, Normand P. 2013. *Micromonospora* is a normal occupant of actinorhizal nodules. *J Biosci* 38:685–693. <https://doi.org/10.1007/s12038-013-9359-y>.
13. Malishkaite YB, Nechaeva GA, Kuimova TF, Evtushenko LI, Agre NS. 1987. *Nocardia* strains from actinorhizal plant nodules. *Microbiology* 56:100–106.
14. Echbab H, Prin Y, Ducouso M, Nourissier-Mountou S, Lahlou H, Araho M. 2004. Helper bacteria associated with *Casuarina cunninghamiana*-*Frankia* symbiosis: selection of isolates for their effects on plant growth in axenic conditions. *Acta Bot Gall* 151:429–440. <https://doi.org/10.1080/12538078.2004.10515445>.
15. Dudeja SS, Giri R, Saini R, Suneja-Madan P, Kothe E. 2012. Interaction of endophytic microbes with legumes. *J Basic Microbiol* 52:248–260. <https://doi.org/10.1002/jobm.201100063>.
16. Aserse AA, Rasanen LA, Aseffa F, Hailemariam A, Lindstrom K. 2013. Diversity of sporadic symbionts and nonsymbiotic endophytic bacteria isolated from nodules of woody, shrub, and food legumes in Ethiopia. *Appl Microbiol Biotechnol* 97:10117–10134. <https://doi.org/10.1007/s00253-013-5248-4>.
17. Leite J, Fischer D, Rouws LFM, Fernandes PI, Hofmann A, Kublik S, Schlöter M, Xavier GR, Radl V. 2016. Cowpea nodules harbor non-rhizobial bacterial communities that are shaped by soil type rather than plant genotype. *Front Plant Sci* 7:2064. <https://doi.org/10.3389/fpls.2016.02064>.
18. Martínez-Hidalgo P, Hirsch AM. 2017. The nodule microbiome: N<sub>2</sub>-fixing rhizobia do not live alone. *Phytobiomes J* 1:70–82. <https://doi.org/10.1094/PBIOMES-12-16-0019-RVW>.
19. Murray MG, Thompson WF. 1980. Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res* 8:4321–4325. <https://doi.org/10.1093/nar/8.19.4321>.
20. Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <https://doi.org/10.1517/14622416.5.4.433>.
21. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
22. Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clinngenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol* 20:714–737. <https://doi.org/10.1089/cmb.2013.0084>.
23. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
24. Meier-Kolthoff JP, Göker M. 2019. TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nat Commun* 10:2182. <https://doi.org/10.1038/s41467-019-10210-3>.
25. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <https://doi.org/10.1186/1471-2105-14-60>.
26. Liu Y, Lai Q, Göker M, Meier-Kolthoff JP, Wang M, Sun Y, Wang L, Shao Z. 2015. Genomic insights into the taxonomic status of the *Bacillus cereus* group. *Sci Rep* 5:14082. <https://doi.org/10.1038/srep14082>.
27. Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, Fiebig A, Rohde C, Rohde M, Fartmann B, Goodwin LA, Chertkov O, Reddy TBK, Pati A, Ivanova NN, Markowitz V, Kyrpidis NC, Woyke T, Göker M, Klenk H-P. 2014. Complete genome sequence of DSM 30083<sup>T</sup>, the type strain (U5/41<sup>T</sup>) of *Escherichia coli*, and a proposal for delineating subspecies in microbial taxonomy. *Stand Genomic Sci* 9:2. <https://doi.org/10.1186/1944-3277-9-2>.
28. Blin K, Wolf T, Chevrette MG, Lu XW, Schwalen CJ, Kautsar SA, Duran HGS, Santos E, Kim HU, Nave M, Dickschat JS, Mitchell DA, Shelest E, Breitling R, Takano E, Lee SY, Weber T, Medema MH. 2017. antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res* 45:W36–W41. <https://doi.org/10.1093/nar/gkx319>.