


BMJ Open Efficacy of soft palatal augmentation prosthesis for oral functional rehabilitation in patients with dysarthria and dysphagia: a protocol for a randomised controlled trial

Tomonori Yokoyama,^{1,2} Takao Mukai,³ Naoki Kodama ,¹ Kana Takao,³ Takashi Hiraoka,⁴ Nobuyuki Arai,⁴ Jitsuro Yano,⁵ Hiroaki Nagatsuka,^{1,6} Yousuke Manda,¹ Kozo Hanayama,⁴ Shogo Minagi¹

To cite: Yokoyama T, Mukai T, Kodama N, *et al.* Efficacy of soft palatal augmentation prosthesis for oral functional rehabilitation in patients with dysarthria and dysphagia: a protocol for a randomised controlled trial. *BMJ Open* 2022;**12**:e060040. doi:10.1136/bmjopen-2021-060040

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-060040>).

Received 01 February 2022
Accepted 28 June 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Naoki Kodama;
naoki-k@md.okayama-u.ac.jp

ABSTRACT

Introduction Palatal augmentation prosthesis (PAP) is used in patients with articulation and swallowing disorders caused by postoperative loss of tongue tissue due to tongue cancer, cerebrovascular disease sequelae and age-related hypofunction. We have previously reported a newly designed soft PAP fabricated using an thermoplastic material that is particularly appropriate for early intervention. However, the effect of soft PAP on oral function improvement remains to be elucidated. The aim of this study is to investigate whether soft PAP can improve dysarthria and dysphagia occurring as cerebrovascular disease sequelae.

Methods and analysis This prospective, randomised, controlled trial will compare the immediate and training effects of rehabilitation using soft PAP with those of rehabilitation without using it. Primary outcomes are the single-word intelligibility test score and pharyngeal transit time (PTT). Secondary outcomes are tongue function (evaluated based on maximum tongue pressure, repetitions of tongue pressure and endurance of tongue pressure), articulation function (evaluated based on speech intelligibility, oral diadochokinesis, Voice-Related Quality of Life (V-RQOL)) and swallowing function (evaluated using Eating Assessment Tool-10). The study results will help determine the efficacy of Soft PAP in improving functional outcomes of word intelligibility and PTT. We hypothesised that early rehabilitation using Soft PAP would more effectively improve articulation and swallowing function compared with conventional rehabilitation without using soft PAP.

Ethics and dissemination Ethical approval was obtained from the Okayama University Certified Review Board. The study findings will be published in an open access, peer-reviewed journal and presented at relevant conferences and research meetings.

Trial registration number jRCTs062200054.

INTRODUCTION

Palatal augmentation prosthesis (PAP) is used in patients with articulation and swallowing

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ A randomised controlled design will minimise bias and allow for a direct comparison between the groups.
- ⇒ The intervention group will undertake rehabilitation using soft palatal augmentation prosthesis, although the control group will receive normal training.
- ⇒ The study sample size was calculated based on previous study regarding the improvement in pharyngeal transit time.
- ⇒ The limitation of this study is that only the evaluator is blinded to allocation results, not the therapist or participants.

disorders caused by postoperative loss of tongue tissue due to tongue cancer, cerebrovascular disease sequelae and age-related hypofunction.¹ The Japan Council for Quality Health Care guideline for PAP states that it enhances the rehabilitation effect when fabricated and used appropriately.¹ PAP use has been recommended in the PAP guidelines issued by the Japan Prosthodontic Society.¹ PAP can be used to improve dysphagia or dysarthria caused not only by tissue loss after glossectomy but also by posterior movement disorders occurring as cerebrovascular disease sequelae.² In patients requiring upper and/or lower limb rehabilitation, early initiation of rehabilitation effectively improves the prognosis of movement disorders.³ However, an early intervention protocol involving PAP use for rehabilitation therapy has not yet been established. This might be partly attributable to the practical difficulties faced by general dental practitioners in fabricating and adjusting PAP during the acute stage of the primary disease. Therefore, medical



Figure 1 Soft palatal augmentation prosthesis.

and dental interprofessional collaboration in acute care-oriented hospitals in this field is required for PAP use in early rehabilitation therapy.

Most studies reporting the effect of PAP on dysphagia and dysarthria occurring due to stroke and neuromuscular diseases are either case reports or retrospective studies.⁴ Moreover, the methods used for evaluating clinical efficacy in these studies were not well structured; therefore, a meta-analysis could not be performed.⁴ Thus, prospective cohort studies examining the effect of rehabilitation using PAP on patients with stroke and neuromuscular disease are needed.

Conventionally, PAP has been fabricated using acrylic resin.^{5–8} Acrylic resin is a denture base material and exhibits stable mechanical properties that are appropriate for long-term use, even after repeated clinical adjustment. To achieve the best effect on patients' oral function, frequent adjustments by a skilled dentist are usually required, even when PAP is fabricated at a rather stable or chronic stage of the primary disease. However, given the atmosphere of acute care units providing early intervention in hospitals, a device that is easy to fit and requires less adjustment might be easily accepted. We have previously reported a newly designed Soft PAP fabricated using an elastic thermoplastic material that is particularly appropriate for early intervention (figure 1).^{9 10} Soft PAP is a simplified PAP that has a flat palatal surface. Although a major disadvantage of Soft PAP is that it may not be fully adjustable to exactly reproduce patient's oral function, it provides a mechanical target point as a terminus ad quem for tongue movement during rehabilitation. Its inherent properties of ease of fabrication and no requirement of adjustment to provide a mechanical target point are expected to benefit patients undergoing rehabilitation.¹¹ Early rehabilitation interventions using this simplified type of PAP may improve dysarthria and dysphagia occurring as cerebrovascular disease sequelae. We conducted a preliminary study using a small sample size to ascertain the effect of early

intervention using soft PAP.¹² The results of this preliminary study suggest that early rehabilitation using soft PAP improves the functional speed of the tongue tip during articulation movements. We hypothesised that early rehabilitation using soft PAP would more effectively improve articulation and swallowing function compared with conventional rehabilitation without using it. Therefore, our null hypothesis is that the effect of rehabilitation using soft PAP on articulation and swallowing function is not different from that of conventional rehabilitation without using soft PAP.

METHODS AND ANALYSIS

Trial design

This study will be a prospective, randomised, controlled trial comparing the immediate and training effects of rehabilitation using soft PAP (SP group) with those of conventional rehabilitation without using soft PAP (NSP group) on patients with dysarthria and dysphagia. Figure 2 shows the flowchart of this clinical trial. Two PAPs with different palatal plate thicknesses will be fabricated for each participant. One will extend up to the height of the flat palatal surface at the cervical line of the upper posterior tooth, whereas the other one would be half as thick as the previous one (figure 3). The participants will select the PAP they feel most comfortable with. They will be instructed to wear soft PAP for as long as possible, except during meals and sleep, with each participant maintaining a record of the actual duration of wearing the PAP. Each group will be provided the same articulation training during the study period. The patients will be assessed at baseline and at 2 and 8 weeks postintervention. Enrolment in the study will be performed by a medical doctor or dentist. Soft PAP will be fabricated by a dentist, and video fluorography (VF) assessment will be performed by a medical doctor, radiologist and speech–language pathologist. Other evaluations will be performed by medical doctors, dentists and speech–language pathologists. The criteria for stopping the trial include deterioration of patients' condition and a request to stop.

Patient and public involvement

Patients admitted to Kawasaki Medical School Hospital who meet the inclusion criteria receive an explanation for this study. They are not involved in the design or conducting of the study. Results of the study will be disseminated to the participants who want to know the results. In addition, they are not asked to assess the burden of the intervention and time required to participate in the research.

Study sites

This study is funded by the Department of Occlusal and Oral Functional Rehabilitation, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University.




Figure 2 Study flow chart. PAP, palatal augmentation prosthesis.

Study participant recruitment will take place at the Kawasaki Medical School Hospital.

Ethical approval

The study protocol was approved by the Okayama University Certified Review Board and the ethics review of board of Kawasaki Medical School.

Study population

Study participants will be selected from among patients admitted to the Kawasaki Medical School Hospital with a primary diagnosis of stroke, neuromuscular disease or head trauma and with a speech articulation disorder score of ≥ 2 .¹³ Speech intelligibility will be evaluated by recording the patient reading a text aloud and assessed

by three speech–language pathologists with at least 3 years of experience. The following paragraphs provide a detailed description of the inclusion and exclusion criteria.

Inclusion criteria

Patients fulfilling the following five conditions will be enrolled in the study:

1. Those who are acutely hospitalised for stroke, neuromuscular disease, head injury or other related diseases in the Kawasaki Medical School Hospital.
2. Those suffering from dysarthria with a speech intelligibility score of ≥ 2 or dysphagia with problems in the preparatory, oral or pharyngeal phase.




Figure 3 Soft palatal augmentation prosthesis (soft PAP) used in the SP group: type 1 and type 2.




Figure 5 The tongue pressure measurement device (TPM-01, JMS, Hiroshima) is connected to a personal computer. Measurements are performed with real-time visual feedback on the screen.

P_{max} .¹⁶ Two measurements, with a 1 min break between them, will be performed. For the offline measurement of ETP from the waveform, the starting point of the duration will be defined as the moment at which tongue pressure exceeds 50% of P_{max} . The endpoint of the duration will be defined according to the following three settings (figure 7).

A: Tongue pressure decreases between 40% and 50% of P_{max} for 2 s with subsequent pressure recovery to >50% of P_{max} .

B: Tongue pressure decreases below 40% of P_{max} for 0.5 s with subsequent pressure recovery to >50% of P_{max} .

C: Tongue pressure decreases below 50% of P_{max} without any subsequent recovery to >50% of P_{max} .

The earliest time point for the above three settings will be regarded as the endpoint.

Articulation function

Oral diadochokinesis (Oral DDK)

Patients will be instructed to repeat each of the three target sounds (/ta/, /ka/, /taka/) as quickly as possible for 5 s, and the sounds will be recorded using a digital voice (integrated chip; IC) recorder (Sony ICD-SX850, Tokyo, Japan). Two measurements will be performed for




Figure 6 An example of measurements obtained when 50% of P_{max} is considered the target tongue pressure.




Figure 7 Measurement of the endurance of tongue pressure (ETP): A, B and C show a schematic sample of the endpoint of ETP. A: Tongue pressure decreased between 40% and 50% of Pmax for 2 s with subsequent pressure recovery to >50% of Pmax. B: Tongue pressure decreased below 40% of Pmax for 0.5 s with subsequent pressure recovery to >50% of Pmax. C: Tongue pressure decreased below 50% of Pmax without any subsequent recovery to >50% of Pmax.

each sound. The average of the two measurements will be considered the oral DDK for each sound.

Forty-word intelligibility test

Ishihara's 40-word list will be used for a 40-word intelligibility test, with the words listed in a randomised order. Participants will read out the words from the list, and their voices will be recorded using an IC recorder. The assessment will be completed by three healthy adults without hearing impairment whose native language is Japanese. The average percentage of correct responses by the three examiners will be calculated.

Speech intelligibility

Patients' speech will be recorded in two conditions: (1) reading aloud a fairy tale, 'Jack and the Beanstalk' and (2) 'routine conversation about patients' events of the day' in 1 min. The assessment will be completed by three healthy adults without hearing impairment whose native language is Japanese. Speech intelligibility will be rated on a five-point scale for each recording.¹⁷ Conventional criteria will be used for this scale in all cases, as in Taguchi's method. The conventional criteria and scores are as follows: 1=intelligible, 2=partially intelligible, 3=intelligible when the topic is known, 4=mostly unintelligible and 5=unintelligible. The average of the three examiners' ratings will be calculated.

Voice-Related Quality of Life

Voice-Related Quality of Life (V-RQOL) is a questionnaire used for examining the V-RQOL.¹⁸ Patients will fill in a self-assessment V-RQOL sheet using the following five-point scale for each of the 10 questions about their voice: 1=none, not a problem; 2=a small amount; 3=a moderate (medium) amount; 4=a lot; and 5=problem is 'as bad as it can be'.

Swallowing function

Swallowing VF

The patient will sit in a chair and swallow 5 mL of thickened barium twice. During this task, VF will be recorded in the lateral and frontal views and assessed using the following parameters:

Oral transit duration: time elapsed between the start of swallowing motion of the tongue and the arrival of the bolus head at the ramus of the mandible.

Pharyngeal transit time (PTT): time elapsed between the bolus head crossing the ramus of the mandible and its passage through the upper oesophageal sphincter.

Stage transition duration: time elapsed between the bolus head crossing the ramus of the mandible and the onset of hyoid bone elevation.

Hyoid movement distance: the maximum elevation of the hyoid bone from the resting position to an anterior and superior position during the swallowing reflex, with the lower border of the spinous process of the fourth cervical vertebra used as the reference point.

Videofluoroscopic dysphagia scale: assessment of VF recordings obtained for evaluating dysphagia according to the report by Han *et al.*¹⁹

Questionnaire on swallowing

Patients will answer a questionnaire on swallowing function (Eating Assessment Tool-10 (EAT-10)) that comprises 10 items on swallowing-related quality of life.²⁰

Sample size calculation

Sample size was determined using the results of a previous study on PTT, which is the primary endpoint, as reference Matsubara *et al.*²¹ Matsubara *et al.*²¹ reported an improvement in PTT after 4 weeks of a high-speed jaw-opening exercise in elderly patients with mealtime malaise

($p=0.01$; Cohen's $d=0.57$). Using an unpaired two-tailed t -test with Cohen's $d=0.57$, $\alpha=0.05$, power=0.8, and allocation ratio=1, the sample size will be 50 patients per group. Thus, the target study participants are estimated to be 100. We will be collaborating with our research partners for participant recruitment.

Data processing and analysis

We will follow the intention-to-treat (ITT) principle and perform data collection and analysis based on the treatment assigned to patients by stratified random allocation. ITT analysis is an analytical method based on the allocation determined before the start of the intervention. Missing data will be statistically analysed using the last observation carried forward method. The obtained data will be entered into a dedicated database for collection and analysis. The background and functional test results of participants as well as adverse events experienced by them will be compared between the SP and NSP groups. The primary outcomes are the single-word intelligibility test score and PTT. The secondary outcomes are tongue function (evaluated based on maximum tongue pressure, RTP and ETP), articulation function (evaluated based on speech intelligibility, oral diadochokinesis, V-RQOL) and swallowing function (evaluated using EAT-10). The immediate effect will be assessed under paired conditions. Training effects will be assessed using two-way analysis of variance. All statistical analyses will be performed using SPSS V.22 (IBM). A p value <0.05 will be considered significant.

Data management

The principal investigator or subinvestigator will explain the study in writing to participants and obtain their consent. Furthermore, a case report form (CRF) has been prepared. The data collected in this study will be entered into and analysed using the researcher's computer (not connected to the internet) to protect participants' personal information; moreover, the data will be stored in a digital versatile disc recordable (DVD-R) as a password-protected file. The data will remain in a locked cabinet in the Division of Oral Surgery, Kawasaki Medical School Hospital, for 5 years after study completion. Subsequently, the DVD-Rs will be destroyed and the consent forms shredded.

Data monitoring

On-site monitoring of this study is conducted by a person designated by the principal investigator in accordance with the protocol. The monitoring manager has no conflicts of interest to declare.

All adverse events will be recorded on the CRF that will include information on the nature and timing of onset and resolution, severity, treatment and outcome of the adverse event as well as assessment of its severity. Follow-up investigations will be performed if deemed necessary.

Compensation

The study will be covered by clinical research insurance in case of liability for compensation in the event that study participants experience health problems.

DISCUSSION

Acute rehabilitation is usually primarily achieved through functional training and compensatory methods; therefore, prosthesis use has rarely been incorporated into acute rehabilitation. Studies reporting the starting point of rehabilitation interventions for cerebrovascular diseases have shown that patients starting rehabilitation within 72 hours of hospitalisation had a shorter hospital stay and better gait at discharge than those starting rehabilitation later than 72 hours.²² Furthermore, Takashima and Abe have reported that early construction of a knee-ankle-foot orthosis during the acute phase of stroke improves gait independence at an early stage.²³ Accordingly, using prosthetic orthotics to support functional movement may contribute to early rehabilitation of patients with central nervous system-related locomotor disability. Moreover, as soft PAP is easy to fabricate and use, it may become a widely used training device for early intervention. Future research will help identify the optimal timing of initiating a soft PAP intervention and its duration and other impacts.

ETHICS AND DISSEMINATION

Ethical approval was obtained from the Okayama University Certified Review Board. Study findings will be published in an open access, peer-reviewed journal and presented at relevant conferences and research meetings. Any changes or revisions to the research protocol or consent explanatory document will be approved in advance by an accredited clinical research review committee.

Author affiliations

¹Department of Occlusal & Oral Functional Rehabilitation, Okayama University Graduate School of Medicine Dentistry and Pharmaceutical Sciences, Okayama, Japan

²Division of Speech-Language-Hearing Therapy, Rehabilitation Center, Kawasaki Medical School Hospital, Kurashiki, Japan

³Division of Oral Surgery, Kawasaki Medical School Hospital, Kurashiki, Japan

⁴Department of Rehabilitation Medicine, Kawasaki Medical School, Kurashiki, Japan

⁵Department of Speech-Language Pathology and Audiology, Faculty of Rehabilitation, Kawasaki University of Medical Welfare, Kurashiki, Japan

⁶Sadamitsu Dental Clinic, Gakuenasahicho, Japan

Acknowledgements The authors would like to express their sincere gratitude to Mr. Katsuya Nakamura and Dr. Shinsuke Nagami for their help in conducting this study.

Contributors TY and SM conceptualised the original study and drafted the manuscript. KT, TM, TH, NA, JY, HN, KH and NK contributed to refining the study design. YM is the monitoring manager. NK and SM critically revised the manuscript. TM and SM are the principal investigators. SM is the lead researcher. All authors have approved the final draft of the manuscript.

Funding This work is funded by the Department of Occlusal and Oral Functional Rehabilitation, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s)

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Naoki Kodama <http://orcid.org/0000-0003-2489-0186>

REFERENCES

- Practice Guidelines for palatal augmentation prosthesis (PAP) for Dysphagia and Dysarthria in Japan 2020. Available: https://hotetsu.com/files/files_536.pdf
- Ueda K, Mukai Y, Morita M. The effectiveness of palatal augmentation prosthesis for dysphagia. *Jap J Dysphag Rehabil* 2012;16:32–41.
- Bernhardt J, Dewey H, Thrift A, *et al*. A very early rehabilitation trial for stroke (AVERT): phase II safety and feasibility. *Stroke* 2008;39:390–6.
- Marunick M, Tselios N. The efficacy of palatal augmentation prostheses for speech and swallowing in patients undergoing glossectomy: a review of the literature. *J Prosthet Dent* 2004;91:67–74.
- Wheeler RL, Logemann JA, Rosen MS. Maxillary reshaping prostheses: effectiveness in improving speech and swallowing of postsurgical oral cancer patients. *J Prosthet Dent* 1980;43:313–9.
- Aramany MA, Downs JA, Beery QC, *et al*. Prosthodontic rehabilitation for glossectomy patients. *J Prosthet Dent* 1982;48:78–81.
- Arioka K, Ishida R, Mori T. The applications of palatal augmentation plate (PAP) to five patients with dysphagia after glossectomy caused by oral cancer. *Jap J Dysphagia Rehabil* 2005;9:76–82.
- Ono T, Hamamura M, Honda K, *et al*. Collaboration of a dentist and speech-language pathologist in the rehabilitation of a stroke patient with dysarthria: a case study. *Gerodontology* 2005;22:116–9.
- Kozaki K-I, Kawakami S, Konishi T, *et al*. Structure of a new palatal plate and the artificial tongue for articulation disorder in a patient with subtotal glossectomy. *Acta Med Okayama* 2016;70:205–11.
- Minagi S, Abe M, Gofuku A, *et al*. Dream of speaking project and medical-dental-engineering collaboration. *Ann Jpn Prosthodont Soc* 2017;9:186–90.
- Ohta K, Miyazaki F, Kawakami S. Fabrication procedure of soft PAP for speech improvement of glossectomy patient. *Maxillofac Prosthet* 2016;39:50–5.
- Yokoyama T, Nagatsuka H, Yano J. The effects of early intervention using PAP on oral function recovery: a randomized controlled trial. *Ann Jpn Prosthodont Soc* 2020;12:81.
- Hattori M, Sumita YI, Kimura S, *et al*. Application of an automatic conversation intelligibility test system using computerized speech recognition technique. *J Prosthodont Res* 2010;54:7–13.
- Nagashima K. A computerized number generator through the stratified block randomization [online] (in Japanese). Available: <https://nshi.jp/contents/js/randblock/> [Accessed 16 Jan 2021].
- Yokoyama T, Yano J, Kumakura I. Investigation of the relationship between tongue motor function and speech mechanism in healthy young adults [in Japanese]. *J Speech Lang Hear Res* 2017;14:47–55.
- Solomon NP, Robin DA, Luschei ES. Strength, endurance, and stability of the tongue and hand in Parkinson disease. *J Speech Lang Hear Res* 2000;43:256–67.
- Gengoshougaichiryogaku TT [in Japanese]. Tokyo: Igaku Shoin 1966:37–8.
- Hogikyan ND, Sethuraman G. Validation of an instrument to measure voice-related quality of life (V-RQOL). *J Voice* 1999;13:557–69.
- Han TR, Paik N-J, Park J-W, *et al*. The prediction of persistent dysphagia beyond six months after stroke. *Dysphagia* 2008;23:59–64.
- Belafsky PC, Mouadeb DA, Rees CJ, *et al*. Validity and reliability of the eating assessment tool (EAT-10). *Ann Otol Rhinol Laryngol* 2008;117:919–24.
- Matsubara M, Tohara H, Hara K, *et al*. High-speed jaw-opening exercise in training suprahyoid fast-twitch muscle fibers. *Clin Interv Aging* 2018;13:125–31.
- Hayes SH, Carroll SR. Early intervention care in the acute stroke patient. *Arch Phys Med Rehabil* 1986;67:319–21.
- Takashima Y, Abe H. The effect of early construction of a knee-ankle-foot-orthosis during the acute phase in severe hemiplegic patients for functional independence measure of walking and stair climbing. *Boll Jap Soc Prosthet Orthot* 2018;34:52–9.