

Plasma miR-151-3p as a candidate diagnostic biomarker for head and neck cancer: a cross-sectional study within the INHANCE consortium

Roberta Pastorino,^{1,2} Michele Sassano,^{2,3} Danilo Francesco Tiziano,^{4,5} Luca Giraldi,² Rosarita Amore,² Dario Arzani,² Emanuela Abiusi,^{4,5} Wolfgang Ahrens,⁶ Laia Alemany Vilches,^{7,8} Cristina Canova,⁹ Claire Mary Healy,¹⁰ Ivana Holcatova,¹¹ Pagona Lagiou,¹² Jerry Polesel,¹³ Maja Popovic,¹⁴ Ståle Nygård,¹⁵ Gabriella Cadoni,^{16,17} Ariana Znaor,¹⁸ Paolo Boffetta,^{3,19} Keitaro Matsuo,^{20,21} Isao Oze,²⁰ Paul Brennan,¹⁸ Stefania Boccia^{1,2}

¹ Department of Woman and Child Health and Public Health—Public Health Area, Fondazione Policlinico Universitario A. Gemelli IRCCS, Roma, Italia

² Section of Hygiene, University Department of Life Sciences and Public Health, Università Cattolica del Sacro Cuore, Roma, Italia

³ Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy

⁴ Department of Life Sciences and Public Health, Section of Genomic Medicine, Università Cattolica del Sacro Cuore, Roma, Italia

⁵ Unit of Medical Genetics, Department of Laboratory science and Infectious Diseases, Fondazione Policlinico Universitario A. Gemelli IRCCS, Roma, Italia

⁶ Leibniz Institute for Prevention Research and Epidemiology, BIPS, and University of Bremen, Faculty of Mathematics and Computer Science, Institute of Statistics, Bremen, Germany

⁷ Cancer Epidemiology Research Programme, IDIBELL, Catalan Institute of Oncology, L'Hospitalet de Llobregat, Spain.

⁸ Consortium for Biomedical Research in Epidemiology and Public Health (CIBERESP), Madrid, Spain.

⁹ Department of Cardiac Thoracic Vascular Sciences and Public Health, Università di Padova, Padova, Italia

¹⁰ Trinity College Dublin School of Dental Science, Dublin, Ireland

¹¹ Institute of Hygiene & Epidemiology, 1st Faculty of Medicine, Charles University, Prague, Czech Republic.

¹² Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece.

¹³ Cancer Epidemiology Unit, Centro di Riferimento Oncologico (CRO) Aviano, IRCCS, Aviano, Italia.

¹⁴ Cancer Epidemiology Unit, Department of Medical Sciences, Università di Torino, Torino, Italia.

¹⁵ Cancer Registry of Norway, Norway

¹⁶ Dipartimento Scienze dell'Invecchiamento, Neurologiche, Ortopediche e della Testa-Collo, Fondazione Policlinico Universitario A. Gemelli IRCCS, Roma, Italia.

¹⁷ Dipartimento Patologia Testa Collo e Organi di Senso, Facoltà Medicina e Chirurgia Università Cattolica Sacro Cuore, Roma, Italia.

¹⁸ International Agency for Research on Cancer, world Health Organization, Lyon, France.

¹⁹ Stony Brook Cancer Center, Stony Brook University, Stony Brook, NY, USA

²⁰ Division of Cancer Epidemiology and Prevention, Aichi Cancer Center, Nagoya, Japan.

²¹ Division of Cancer Epidemiology, Nagoya University Graduate School of Medicine, Nagoya, Japan.

Running title. Plasma miR-151-3p as candidate diagnostic biomarker for HNC

Keywords. head and neck cancer; microRNAs; Plasma; Diagnostic biomarker; INHANCE Consortium

Financial support. This study has received funding from AIRC (Associazione Italiana per la lotta contro il Cancro), contract no. 14220 to SB, Fondazione Veronesi, n. CUP: J54G13000430007 to SB and Università Cattolica del Sacro Cuore (funds line D.3.1) to SB to cover the journal fee of the

publication. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript

Corresponding author.

Michele Sassano, Department of Medical and Surgical Sciences, Università di Bologna, Bologna, Italy. E-mail: michele.sassano02@icatt.it; Ph: +39 (6) 30154396

Conflict of Interest

The authors have no conflicts of interest to declare.

Author Disclaimer

Where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer /World Health Organization.

Word count, figures and tables

Abstract: 204 words

Manuscript: 2232 words

n.1 figure and n.5 tables

Supplementary materials: n.2 tables

Abstract

Background

Identification of screening tests for the detection of head and neck cancer at an early stage is an important strategy to improving prognosis. Our objective was to identify plasma circulating microRNAs for the diagnosis of head and neck cancer (oral and laryngeal subsites), within a multicenter International Head and Neck Cancer Epidemiology (INHANCE) consortium.

Methods

A high-throughput screening phase with 754 miRNAs was performed in plasma samples of 88 cases and 88 controls, followed by a validation phase of the differentially expressed miRNAs, identified in the screening, in samples of 396 cases and 396 controls. Comparison of the Fold Changes were carried out using the Wilcoxon rank-sum test and the Dunn's multiple comparison test.

Results

We identified miR-151-3p (FC=1.73, p=0.007) as differentially expressed miRNAs in the screening and validation phase. The miR-151-3p was the only overexpressed miRNA in validation sample of head and neck (HNC) patients with early stage at diagnosis (FC=1.81, p=0.008) and it was confirmed upregulated both in smoker early-stage cases (FC=3.52, p=0.024) and in non-smoker early-stage cases (FC=1.60, p=0.025) compared with controls.

Conclusions

We identified miR-151-3p as an early marker of HNC. This miRNA was the only upregulated in patients at early stages of the disease, independently of the smoking status.

Impact

The prognosis for head and neck is still poor. The discovery of a new diagnostic biomarker could lead to an earlier tumor discovery and therefore to an improvement in patient prognosis.

Introduction

Head and Neck Cancer (HNC) represents the sixth most common type of cancer worldwide, with more than 930,000 cases and 460,000 deaths annually (1). These cancers originate from several sites of the head and neck region: pharynx (oropharynx, nasopharynx, and hypopharynx), larynx or the oral cavity, and are strongly associated with environmental and lifestyle risk factors such as tobacco and alcohol consumption (2–4). Other known risk factors are human papillomavirus infection (HPV), diet, lack of physical activity and/or family history (5–9).

The prognosis of HNCs is still poor, despite the combined treatment involving surgery, radiotherapy, and chemotherapy (10). The 5-year disease-specific survival rate in HNC patients in Europe ranges between 25.5% for hypopharyngeal cancer and 63.1% for laryngeal cancer (11). The main reasons for the poor prognosis are frequent loco-regional recurrence and metastasis (12,13). Fifty percent of HNC patients, in fact, develop recurrent disease within the first two years of treatment (14), and it is currently well established that HNC survivors also have an increased risk of developing second primary cancers (SPC) compared to the overall population, with frequent SPC of the head and neck, esophagus, and lung (15–18).

As in all cancer types, also for HNC an early detection is expected to improve the prognosis: clinical/instrumental tools coupled with the identification of novel sensitive and reliable biomarkers would be crucial to this purpose. Although several studies have been performed to depict the molecular landscape of HNC over the last years, the identification of reliable genomic biomarkers is still lacking (19); subclassifications based on genomic profiling might be useful to better understand the biological mechanisms responsible for carcinogenesis (20). The level of molecular markers, including microRNAs (miRNAs), in biofluids is considered a promising non-invasive approach for the early detection of cancer, including HNC (21).

miRNAs are short noncoding RNA molecules of 19-25 nucleotides which play a crucial role in the regulation of gene expression, including oncogenes or tumor suppressor genes, and are emerging as a new molecular tool for non-invasive cancer diagnosis and prognosis (22). Several miRNAs involved in the pathogenesis of HNC have now been identified, and some are currently being evaluated for their diagnostic and/or prognostic performance (23). A recently published meta-analysis revealed that miR-21 and miR-93 are mostly upregulated in HNC, whereas miR-9, miR-203, miR-218 and miR-375 are downregulated (24). Although the meta-analysis demonstrated that these molecules are promising diagnostic tool with moderate accuracy, to our knowledge, none of

these has been broadly used as a HNC diagnostic biomarker in a clinical setting due to the limited accuracy in discriminating cases and controls.

The aim of the present study was to identify specific diagnostic miRNA signatures for oral and laryngeal HNC, within a multicenter International Head and Neck Cancer Epidemiology (INHANCE) consortium (25).

Materials and Methods

Study participants

Participants were selected from the three studies of the INHANCE Consortium (25): Rome (Italy), ARCAGE study (with 10 Western European centers) and Aichi (Japan). We included, as study cases, patients ≥ 18 years of age with histologically confirmed primary HNC arising in the anatomical sites of the oral cavity or larynx.

The controls were selected among hospital visitors without a history of cancer, with a case-control ratio of 1:1. Participants completed a questionnaire, including information about age, gender, tobacco smoking at the diagnosis, and donated a sample of peripheral blood. Cases and controls were matched by age, ethnicity, gender and cigarette smoke exposure. A total of 692 Caucasian and 276 Japanese subjects were included. The characteristics of all subjects are listed in Supplementary Table 1.

The study was approved by the local Ethical Committee of each participating center and written informed consents were obtained from all study subjects.

Variables

Different study centres used different cancer codes, which were converted into International Classification of Diseases for Oncology (ICD-O-2) when included into this study. HNCs were classified according to the following anatomic sites using the ICD-10 codes: oral cavity (C00.3–C00.9, C02.0–C02.3, C03.0, C03.1, C03.9, C04.0, C04.1, C04.8, C04.9, C05.0, C06.0–C06.2, C06.8, and C06.9) and larynx (codes C10.1, C32.0–C32.3 and C32.8–C32.9). Cancers were staged according to the TNM Staging System, 7th edition, and categorized as early (stage I and II) and

advanced (stage III and IV) stage at diagnosis (26). Individuals were categorized according to smoking status as current smokers or non-current smokers.

Blood Samples

Peripheral blood was collected in EDTA tubes at the time of interview and processed as rapidly as possible (generally within 2 hours). For cases, blood draw was performed at the diagnosis before surgery or any other treatment. Plasma samples were isolated by centrifugation of whole blood at 2000xg for 10 minutes at room temperature. Samples were stored at -80°C .

Molecular analyses and study design

Cell-free RNA (cfRNA) was isolated from 300 μL of plasma using the NucleoSpin miRNA Plasma kit (Macherey-Nagel, Duren, Germany). All plasma samples were spiked-in with 10 pmol of *Arabidopsis thaliana* synthetic miR-159a (synthesized by Eurofins MWG Operon, Ebersberg, Germany) as a control of RNA extraction.

In order to identify HNC diagnostic miRNA signatures, a two-stage approach was chosen: a *high-throughput screening phase* performed in plasma samples of 88 HNC cases and 88 controls, followed by a *validation phase* of the differentially expressed miRNAs, identified in the screening phase, in samples of 396 HNC cases and 396 controls (Supplementary Table 2).

For the screening phase, the expression levels of 754 miRNAs (those annotated in miRBase v14 at the time of the study design) were quantified using the TaqMan Human MicroRNA Array A + B Card Set v3.0 and the ABI 7900HT instrument (Applied Biosystems, Foster City, CA), according to manufacturer's instructions (including the pre-amplification step). qPCR data were collected by the SDS software v2.4 (Applied Biosystems, Foster City, CA). Threshold cycle (Ct) values were determined using the Expression Suite software (Applied Biosystems, Foster City, CA), setting automatic baseline and threshold. Only miRNAs with Ct values <35 in at least one group were evaluated in comparative analyses. The relative levels were calculated according to the Ct method and expressed as fold change (FC) compared to the median of controls. Only miRNAs with multiple testing corrected $p < 0.05$ were considered as differentially expressed and selected for the validation phase.

For the second stage of the study (validation phase), custom Low-density Arrays (Life Technologies, Foster City, CA) were designed with spotted primers and probes, specific for the

significant miRNAs selected in the screening phase. Amplification was performed using the TaqMan™ Universal Master Mix II, no UNG (Life Technologies, Foster City, CA) and the ABI 7900HT instrument (Applied Biosystems, Foster City, CA), according to manufacturer's instructions (including the pre-amplification step). The Ct-method was used to determine the relative expression levels, compared to the median level of controls. miRNA 159 was used as the endogenous control.

Statistical Analysis

For each miRNA, Ct or Δ Ct values (CT target-CT reference gene), and FC were reported as median and interquartile range (IQR). Subgroup analyses were performed by considering current smokers compared with those who were not current smokers, and by restricting the main analyses to the cases with an early stage of disease at diagnosis (stage I and II).

Comparison of the FCs were carried out using the Wilcoxon rank-sum test and the Dunn's multiple comparison test. Results with a $p < 0.05$ after multiple comparison correction were considered significant.

In addition, we investigated the significance of the interaction between miRNA expression levels and smoking status by including an interaction term in logistic regression models.

Analysis were performed using R 3.5.1 (<https://www.r-project.org/>) and Stata softwares (StataCorp. 2016. Stata Statistical Software: Release 16. College Station, TX: StataCorp LP).

Data Availability

The data generated in this study are available upon request from the corresponding author.

Results

Screening phase: miR-145, -151-3p, -191, -409-3p, -425, -628-3p and -766 are differentially expressed in HNC

In the screening phase, we detected a total of seven differentially expressed miRNAs between cases and controls, including six upregulated (miR-425, miR-151-3p, miR-191, miR-409-3p, miR-628-3p, miR-766) and one downregulated miRNA (miR-145), that were selected for validation (Table 1).

Our data indicate that about 1% of the total miRNAs were deregulated in cases. Most miRNAs were not detectable in plasma.

Validation phase: miR-145 and -151-3p are upregulated in HNC

In the validation phase, two out of the seven miRNAs from the screening were confirmed to be differentially expressed between cases and controls (Table 2).

While miR-151-3p was confirmed to be upregulated (Replication sample: FC=1.73, p=0.007; Discovery sample: FC=1.72, p=0.003), the behavior of miR-145 was opposite in the two phases of our study (Replication sample: FC=2.35, p=0.008; Discovery sample: 0.62, p=0.0003). The other miRNAs (miR-191, -409-3p, -425, -628-3p, and miR-766) were not differentially expressed, although the direction of the change was the same in the discovery and replication sample. In the analysis restricted to the early-stage cases, miR-151-3p was the only overexpressed miRNA in HNC patients (FC=1.81, p=0.008, Table 2).

Smoker patients display a wider miRNA dysregulation

When comparing the expression of the selected miRNAs between smoker (n=199) and non-smoker (n=197) cases (Table 3), miR-191 was not differentially expressed at the significance level (p=0.17), although the fold-change indicates overexpression in smokers. All other tested miRNAs showed overexpression in current smokers compared to non-current smokers (miR-145: FC=1.80, p=0.045; miR-151-3p: FC=2.41, p<0.001; miR-409-3p: FC=2.67, p=0.007; miR-628-3p: FC=2.49, p=0.002; miR-766: FC=3.01, p=0.008). These differences were not related to cigarette-smoke per se, but rather to the HNC, since there were no significant differences according to smoking status among controls (smokers n=199; non-smokers n=197, Table 3). Indeed, no significant interactions between miRNA expression levels and smoking status were found (Table 4). Additionally, among early-stage cases, no differences were observed between current smokers and non-smokers (n=38 and 85 for smokers and non-smokers, respectively; Table 3).

Besides the differences between smoker and non-smoker cases, the differential expression of all miRNAs was particularly evident when comparing current smokers with controls (Table 5).

None of the seven miRNAs was differentially expressed in the analysis of non-current smokers vs. controls. Finally, only miR-151-3p was upregulated both in smoker early-stage cases (FC=3.52, p=0.024) and non-smoker early-stage cases (FC=1.60, p=0.025) compared with controls (Table 6).

Discussion

In the present study, we identified miR-151-3p as a candidate diagnostic biomarker for HNC. We firstly used a high-throughput approach to identify candidate deregulated biomarkers in plasma of patients compared to controls. In this first phase of the study we found that less than 1% of miRNA are differentially expressed in HNC, suggesting that the effect of the background noise of global miRNA release in cfRNA is limited, and confirming the robustness of miRNA dosage as disease biomarkers in general, as widely demonstrated also in other studies. Our study has several strengths, including the availability of a large validation sample that has been analyzed independently from the first screening group, the different geographic origin of cases and controls enrolled for the validation phase, the availability of both early and advanced stages of HNC. The main limitations are the paucity of available genomic data of patients and, more importantly, the cross-sectional nature of the study with the consequent lack of longitudinal samples, especially after surgical removal of the HNC. Furthermore, at the time of the study design NGS miRNome analysis was not available and we adopted the microarray method in the screening phase. However in the validation phase we performed a targeted analysis of the differentially expressed miRNAs identified during the screening. Lastly, our conclusions refer to tumors of the oral cavity and larynx and we don't know if they are generalizable to all sub-sites of head and neck cancer.

As schematized in fig. 1, we identified miR-151-3p as an early marker of HNC. This miRNA was the only upregulated in patients at early stages of the disease, independently to the smoking status. With disease progression, the increase in miR-151-3p is more prominent in smokers compared to non-smokers. Differently from other miRNAs, miR-151-3p regulation has never been related with smoking per se in previous studies. Also, in our group of controls, no differences in miRNA levels were found on the basis of the smoking status. This indicates that the upregulation of this miRNA is HNC-related. MiR-151-3p has been involved also in other neoplasms, playing a pivotal role in colorectal cancer (27), osteosarcoma (28), cholangiocarcinoma (29), and some others. Thus, miR-151-3p has a potential clinical application as an early diagnostic marker of HNC; conceivably,

miRNA levels could be monitored over time, as a marker of progression, especially in smoker patients.

With the progression of the disease, the differences between smokers and non-smokers are more pronounced (Figure 1): only miR-151-3p and miR-145 are upregulated in both groups.

This latter miRNA provided different results in the screening compared with the validation phase: it was downregulated in the first part of the study, and then upregulated. The explanation of this discrepancy is unknown but it could be related to the different representation of smokers in the two cohorts, which is statistically significant ($X^2=7.13$; $p=0.008$). Indeed, in the validation cohort, miR-145 is significantly upregulated in smoker patients, compared to non-smokers. Based on the data of the literature, miR-145 acts as tumor suppressor miRNA via different signaling pathways in some cancer models (30,31). Additionally, in a recent meta-analysis, this miRNA has been reported as downregulated in saliva of patients with squamous HNC (32). Further studies are thus necessary to better elucidate our findings.

The other differentially expressed miRNAs were specific of smoker patients. To our knowledge, very few information is available regarding the role of miR-409-3p in cancer (33), while, no association has been identified with HNC. Several studies have reported the upregulation of miR-425 in different cancer types, such as non-small-cell lung, colorectal, or breast cancers: to our knowledge, this is instead the first report for HNC (34,35). The role of miR-628 is still debated in the literature: while acting as a tumor suppressor in some models (36), the overexpression of this miRNA has been linked with higher aggressivity in some cancer types (37). Similarly, also miR-766 promotes cancer progression in several models, likely by the accumulation of P53 (38). To the best of our knowledge, for the last two miRNAs there is no evidence on their association with HNC. In conclusion, we report novel miRNAs as potential biomarkers for HNC. In particular, the most promising finding is miR-151-3p, which appears to be an early marker of disease; additionally, the longitudinal evaluation for prognostic purposes of the other miRNAs could be more advisable for smoker patients (miR-145, -409-3p, -425, 628, -766).

References

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin.* 2021;71(3):209-249.
2. Wyss A, Hashibe M, Chuang S-C, Lee Y-CA, Zhang Z-F, Yu G-P, *et al.* Cigarette, Cigar, and Pipe Smoking and the Risk of Head and Neck Cancers: Pooled Analysis in the International Head and Neck Cancer Epidemiology Consortium. *Am J Epidemiol* 2013;178(5):679–90.
3. Hashibe M, Brennan P, Benhamou S, Castellsague X, Chen C, Curado MP, *et al.* Alcohol Drinking in Never Users of Tobacco, Cigarette Smoking in Never Drinkers, and the Risk of Head and Neck Cancer: Pooled Analysis in the International Head and Neck Cancer Epidemiology Consortium. *J Natl Cancer Inst* 2007;99(10):777–89.
4. Pezzuto F, Buonaguro L, Caponigro F, Ionna F, Starita N, Annunziata C, *et al.* Update on Head and Neck Cancer: Current Knowledge on Epidemiology, Risk Factors, Molecular Features and Novel Therapies. *Oncology* 2015;89(3):125–36.
5. Gaudet MM, Olshan AF, Chuang S-C, Berthiller J, Zhang Z-F, Lissowska J, *et al.* Body mass index and risk of head and neck cancer in a pooled analysis of case-control studies in the International Head and Neck Cancer Epidemiology (INHANCE) Consortium. *Int J Epidemiol* 2010;39(4):1091–102.
6. Kawakita D, Lee Y-CA, Turati F, Parpinel M, Decarli A, Serraino D, *et al.* Dietary fiber intake and head and neck cancer risk: A pooled analysis in the International Head and Neck Cancer Epidemiology consortium. *Int J Cancer* 2017;141(9):1811–21.
7. Nicolotti N, Chuang S-C, Cadoni G, Arzani D, Petrelli L, Bosetti C, *et al.* Recreational physical activity and risk of head and neck cancer: a pooled analysis within the international head and neck cancer epidemiology (INHANCE) Consortium. *Eur J Epidemiol* 2011;26(8):619–28.
8. Gillison ML, Lowy DR. A causal role for human papillomavirus in head and neck cancer. *Lancet* 2004;363(9420):1488–9.
9. Negri E, Boffetta P, Berthiller J, Castellsague X, Curado MP, Dal Maso L, *et al.* Family history of cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Int J cancer* 2009;124(2):394–401.
10. Budach V, Tinhofer I. Novel prognostic clinical factors and biomarkers for outcome prediction in head and neck cancer: a systematic review. *Lancet Oncol* 2019;20(6):e313–26.

11. Gatta G, Botta L, Sánchez MJ, Anderson LA, Pierannunzio D, Licitra L; EUROCARE Working Group: Prognoses and improvement for head and neck cancers diagnosed in Europe in early 2000s: The EUROCARE-5 population-based study. *Eur J Cancer* 2015;51(15):2130-2143.
12. Denaro N, Merlano MC, Russi EG. Follow-up in Head and Neck Cancer: Do More Does It Mean Do Better? A Systematic Review and Our Proposal Based on Our Experience. *Clin Exp Otorhinolaryngol* 2016;9(4):287-97.
13. Leemans CR, Braakhuis BJM, Brakenhoff RH. The molecular biology of head and neck cancer. *Nat Rev Cancer* 2011;11(1):9-22.
14. Argiris A, Karamouzis M V, Raben D, Ferris RL. Head and neck cancer. *Lancet* 2008;371(9625):1695-709.
15. Leoncini E, Vukovic V, Cadoni G, Pastorino R, Arzani D, Bosetti C, *et al.* Clinical features and prognostic factors in patients with head and neck cancer: Results from a multicentric study. *Cancer Epidemiol* 2015;39(3):367-74.
16. Jones AS, Morar P, Phillips DE, Field JK, Husband D, Helliwell TR. Second primary tumors in patients with head and neck squamous cell carcinoma. *Cancer* 1995;75(6):1343-53.
17. Lubin JH, Purdue M, Kelsey K, Zhang Z-F, Winn D, Wei Q, *et al.* Total Exposure and Exposure Rate Effects for Alcohol and Smoking and Risk of Head and Neck Cancer: A Pooled Analysis of Case-Control Studies. *Am J Epidemiol* 2009;170(8):937-47.
18. Liu W-S, Chang Y-J, Lin C-L, Liang J-A, Sung F-C, Hwang I-M, *et al.* Secondary primary cancer in patients with head and neck carcinoma: the differences among hypopharyngeal, laryngeal, and other sites of head and neck cancer. *Eur J Cancer Care* 2014;23(1):36-42.
19. Kasradze D, Juodzbaly G, Guobis Z, Gervickas A, Cicciù M. Genetic and proteomic biomarkers of head-and-neck cancer: A systematic review. *J Cancer Res Ther* 2020;16(3):410.
20. Leemans CR, Snijders PJF, Brakenhoff RH. The molecular landscape of head and neck cancer. *Nat Rev Cancer* 2018;18(5):269-82.
21. Arantes LMRB, De Carvalho AC, Melendez ME, Carvalho AL. Serum, plasma and saliva biomarkers for head and neck cancer. *Expert Rev Mol Diagn* 2018 Jan;18(1):85-112.
22. Filipów S, Łaczmański Ł. Blood circulating miRNAs as cancer biomarkers for diagnosis and surgical treatment response. *Front Genet* 2019;10:169
23. Liu C, Yu Z, Huang S, Zhao Q, Sun Z, Fletcher C, *et al.* Combined identification of three miRNAs in serum as effective diagnostic biomarkers for HNSCC. *EBioMedicine* 2019;50:135-143

24. Lubov J, Maschietto M, Ibrahim I, Mlynarek A, Hier M, Kowalski LP, et al. Meta-analysis of micrnas expression in head and neck cancer: Uncovering association with outcome and mechanisms. *Oncotarget* 2017;8(33):55511-55524
25. Winn D, Lee Y-C, Hashibe M, Boffetta P, INHANCE consortium. The INHANCE consortium: toward a better understanding of the causes and mechanisms of head and neck cancer. *Oral Dis* 2015;21(6):685–93.
26. Sobin LH, Gospodarowicz MK, Wittekind C. TNM classification of malignant tumours. Hoboken: Wiley; 2011.
27. Yue C, Chen X, Li J, Yang X, Li Y, Wen Y. miR-151-3p Inhibits Proliferation and Invasion of Colon Cancer Cell by Targeting Close Homolog of L1. *J Biomed Nanotechnol* 2020;16(6):876–84.
28. Li X, Liu Y, Zhang X, Shen J, Xu R, Liu Y, et al. Circular RNA hsa_circ_0000073 contributes to osteosarcoma cell proliferation, migration, invasion and methotrexate resistance by sponging miR-145-5p and miR-151-3p and upregulating NRAS. *Aging (Albany NY)* 2020;12(14):14157–73.
29. McNally ME, Collins A, Wojcik SE, Liu J, Henry JC, Jiang J, et al. Concomitant dysregulation of microRNAs miR-151-3p and miR-126 correlates with improved survival in resected cholangiocarcinoma. *HPB* 2013;15(4):260–4.
30. He JH, Han ZP, Zhou J Bin, Chen WM, Lv YB, He ML, et al. MiR-145 affected the circular RNA expression in prostate cancer LNCaP cells. *J Cell Biochem* 2018;119(11):9168–77.
31. Hang W, Feng Y, Sang Z, Yang Y, Zhu Y, Huang Q, et al. Downregulation of miR-145-5p in cancer cells and their derived exosomes may contribute to the development of ovarian cancer by targeting CT. *Int J Mol Med* 2019;43(1):256–66.
32. Kang JW, Eun YG, Lee YC. Diagnostic Value of Salivary miRNA in Head and Neck Squamous Cell Cancer: Systematic Review and Meta-Analysis. *Int J Mol Sci* 2021;22(13):7026.
33. Sommerova L, Anton M, Bouchalova P, Jasickova H, Rak V, Jandakova E, et al. The role of miR-409-3p in regulation of HPV16/18-E6 mRNA in human cervical high-grade squamous intraepithelial lesions. *Antiviral Res* 2019;163:185–92.
34. Liu D, Zhang H, Cui M, Chen C, Feng Y. Hsa-miR-425-5p promotes tumor growth and metastasis by activating the CTNND1-mediated β -catenin pathway and EMT in colorectal cancer. *Cell Cycle* 2020;19(15):1917–27.
35. Yan YF, Gong FM, Wang BS, Zheng W. MiR-425-5p promotes tumor progression via modulation of CYLD in gastric cancer. *Eur Rev Med Pharmacol Sci* 2017;21(9):2130–6.

36. Li M, Qian Z, Ma X, Lin X, You Y, Li Y, *et al.* MiR-628-5p decreases the tumorigenicity of epithelial ovarian cancer cells by targeting at FGFR2. *Biochem Biophys Res Commun* 2018;495(2):2085–91.
37. Wang JY, Wang JQ, Lu SB. miR-628-5p promotes growth and migration of osteosarcoma by targeting IFI44L. *Biochem Cell Biol* 2020 Apr;98(2):99-105
38. Wang Q, Selth LA, Callen DF, Wang Q, Selth LA, Callen DF. MiR-766 induces p53 accumulation and G2/M arrest by directly targeting MDM4. *Oncotarget* 2017;8(18):29914–24.

TABLES

Table 1. miRNAs differentially expressed between cases and controls in the screening phase.

miRNA	Mean Ct, cases	Mean Ct, controls	FC	p-value
miR-145	27.2	26.5	0.62	0.0003
miR-151-3p	25.2	26.0	1.72	0.0030
miR-191	32.7	34.5	3.48	0.0031
miR-409-3p	27.3	28.6	2.60	0.0080
miR-425	29.2	31.1	3.64	0.0002
miR-628-3p	30.0	31.0	2.04	0.0026
miR-766	26.0	27.3	2.28	0.0052

p-values refer to multiple testing-corrected p-values.

CT: Threshold cycle; FC: Fold change

Table 2. Comparison of miRNA expression between cases and controls in the validation phase.

miRNA	Δ Ct* cases, median [IQR]	Δ ct early stage cases, median [IQR]	Δ Ct controls, median [IQR]	FC, cases vs controls	FC, early stage cases vs controls	p-value, cases vs controls	p-value, early stage cases vs controls
miR-145	9.60 [7.75, 11.76]	10.18 [8.91-12.66]	10.83 [8.54, 13.24]	2.35	1.58	0.008	0.517
miR-151-3p	6.89 [4.90, 8.38]	6.82 [4.59-8.26]	7.68 [5.75, 9.64]	1.73	1.81	0.007	0.008
miR-191	16.54 [11.64, 18.52]	16.01 [11.67-18.24]	17.43 [11.68, 18.56]	1.86	2.68	0.400	0.352
miR-409-3p	8.74 [6.56, 10.82]	8.98 [7.42-11.67]	9.30 [6.99, 11.82]	1.48	1.25	0.117	0.887
miR-425	12.15 [9.66, 17.81]	12.65 [9.17-17.94]	13.26 [9.68, 17.98]	2.16	1.52	0.182	0.550
miR-628-3p	11.43 [9.14, 16.20]	11.79 [9.18-16.01]	11.46 [9.26, 16.94]	1.02	0.79	0.799	0.964
miR-766	9.37 [6.98, 11.90]	9.93 [7.41-15.00]	9.98 [7.38, 13.28]	1.53	1.04	0.151	0.901

CT: Threshold cycle; Δ Ct values (CT target-CT reference gene); FC: Fold change; IQR: Interquartile Range

Bold values denote statistical significance at the $p < 0.05$ level.

*: difference between Ct value of each miRNA and that of miR-159

Table 3. Comparison of miRNA expression between smokers and non-smokers among all cases, early stage cases, and controls separately.

Group	miRNA	n smokers	Δ ct smokers, median [IQR]	n nonsmokers	Δ ct nonsmokers, median [IQR]	FC	p-value
Cases	miR-145	199	9.46 [6.94, 11.41]	197	10.31 [8.86, 12.37]	1.80	0.045
	miR-151-3p		6.14 [4.44, 8.00]		7.40 [6.00, 8.82]	2.41	<0.001
	miR-191		13.87 [10.98, 18.85]		17.03 [12.37, 18.52]	8.91	0.178
	miR-409-3p		7.70 [5.89, 10.36]		9.12 [7.66, 11.68]	2.67	0.007
	miR-425		11.61 [9.01, 17.80]		13.14 [10.05, 17.94]	2.89	0.224
	miR-628-3p		10.77 [8.01, 13.37]		12.09 [10.13, 17.03]	2.49	0.002
	miR-766		8.65 [6.11, 11.04]		10.23 [7.87, 13.37]	3.01	0.008
Early stage cases	miR-145	38	10.16 [9.45, 11.68]	85	10.46 [8.65, 13.37]	1.23	0.763
	miR-151-3p		5.86 [5.03, 6.86]		7.00 [4.52, 8.34]	2.19	0.248
	miR-191		17.80 [12.62, 19.25]		15.51 [11.47, 18.00]	0.20	0.451
	miR-409-3p		7.99 [6.20, 9.47]		9.54 [7.59, 12.14]	2.93	0.183
	miR-425		11.90 [9.01, 17.80]		14.12 [9.47, 18.26]	4.68	0.393
	miR-628-3p		10.99 [9.18, 14.22]		12.36 [9.03, 16.30]	2.58	0.436
	miR-766		10.77 [7.66, 17.94]		9.92 [7.18, 14.82]	0.55	0.688
Controls	miR-145	199	10.16 [8.10-11.99]	197	10.69 [8.36-12.52]	1.45	0.652
	miR-151-3p		7.58 [5.78-9.16]		8.14 [5.76-10.36]	1.47	0.560
	miR-191		17.04 [11.96-18.48]		17.50 [11.21-18.73]	1.38	0.802
	miR-409-3p		8.64 [6.91-10.68]		9.35 [6.83-11.28]	1.64	0.919
	miR-425		12.23 [9.75-17.72]		13.81 [9.55-18.44]	2.99	0.655
	miR-628-3p		11.33 [8.74-15.53]		12.23 [9.89-17.90]	1.86	0.350
	miR-766		9.62 [7.08-11.14]		10.38 [7.54-14.80]	1.69	0.430

CT: Threshold cycle; Δ Ct values (CT target-CT reference gene); FC: Fold change; IQR: Interquartile Range

Bold values denote statistical significance at the $p < 0.05$ level.

Table 4. Results of the interaction analysis between miRNA levels and smoking status

Group	miRNA	P _{interaction}
All cases vs controls	miR-145	0.139
	miR-151-3p	0.082
	miR-191	0.142
	miR-409-3p	0.106
	miR-425	0.087
	miR-628-3p	0.386
	miR-766	0.061
Early stage cases vs controls	miR-145	0.787

	miR-151-3p	0.657
	miR-191	0.467
	miR-409-3p	0.777
	miR-425	0.665
	miR-628-3p	0.577
	miR-766	0.284

Table 5. Post-hoc analysis comparing miRNA expression between cases (according to smoking status) and controls.

miRNA	Δ ct smoker cases, median [IQR]	Δ ct non-smoker cases, median [IQR]	Δ ct controls, median [IQR]	FC, smoker cases vs controls	p-value, smoker cases vs controls	FC, non-smoker cases vs controls	p-value, non-smoker cases vs controls
miR-145	9.46 [6.94-11.41]	10.31 [8.86-12.37]	10.83 [8.54, 13.24]	2.59	<0.001	1.44	0.100
miR-151-3p	6.14 [4.44-8.00]	7.40 [6.00-8.82]	7.68 [5.75, 9.64]	2.91	<0.001	1.21	0.100
miR-191	13.87 [10.98-18.85]	17.03 [12.37-18.52]	17.43 [11.68, 18.56]	11.80	0.227	1.32	0.279
miR-409-3p	7.70 [5.89-10.36]	9.12 [7.66-11.68]	9.30 [6.99, 11.82]	3.03	0.003	1.13	0.484
miR-425	11.61 [9.01-17.80]	13.14 [10.05-17.94]	13.26 [9.68, 17.98]	3.13	0.048	1.08	0.246
miR-628-3p	10.77 [8.01-13.37]	12.09 [10.13-17.03]	11.46 [9.26, 16.94]	1.61	0.046	0.65	0.181
miR-766	8.65 [6.11-11.04]	10.23 [7.87-13.37]	9.98 [7.38, 13.28]	2.53	0.004	0.84	0.487

CT: Threshold cycle; Δ Ct values (CT target-CT reference gene); FC: Fold change; IQR: Interquartile Range

Bold values denote statistical significance at the $p < 0.05$ level.

Table 6. Post-hoc analysis comparing miRNA expression between early stage cases (according to smoking status) and controls.

miRNA	Δ ct smoker cases, median [IQR]	Δ ct non-smoker cases, median [IQR]	Δ ct controls, median [IQR]	FC, smoker cases vs controls	p-value, smoker cases vs controls	FC, non-smoker cases vs controls	p-value, non-smoker cases vs controls
miR-145	10.16 [9.45-11.68]	10.46 [8.65-13.37]	10.83 [8.54-13.24]	1.59	0.117	1.29	0.282
miR-151-3p	5.86 [5.03-6.86]	7.00 [4.52-8.34]	7.68 [5.75-9.64]	3.52	0.024	1.60	0.025
miR-191	17.80 [12.62-19.25]	15.51 [11.47-18.00]	17.43 [11.68-18.56]	0.77	0.344	3.80	0.160
miR-409-3p	7.99 [6.20-9.47]	9.54 [7.59-12.14]	9.30 [6.99-11.82]	2.48	0.088	0.85	0.372
miR-425	11.90 [9.01-17.80]	14.12 [9.47-18.26]	13.26 [9.68-17.98]	2.58	0.195	0.55	0.375
miR-628-3p	10.99 [9.18-14.22]	12.36 [9.03-16.30]	11.46 [9.26-16.94]	1.38	0.361	0.54	0.411
miR-766	10.77 [7.66-17.94]	9.92 [7.18-14.82]	9.98 [7.38-13.28]	0.58	0.423	1.05	0.392

CT: Threshold cycle; Δ Ct values (CT target-CT reference gene); FC: Fold change; IQR: Interquartile Range

Bold values denote statistical significance at the $p < 0.05$ level.

FIGURE LEGEND

Figure 1. miRNAs and disease progression. Distribution of identified miRNAs among early-stage and advanced cancer cases.

* Significant in the screening cohort only.




Figure 1