

<https://helda.helsinki.fi>

Evidence and consensus on technical aspects of embryo transfer

D'Angelo, Arianna

2022-09-10

D'Angelo , A , Panayotidis , C , Alteri , A , Mcheik , S & Veleva , Z 2022 , ' Evidence and consensus on technical aspects of embryo transfer ' , Human reproduction open , vol. 2022 , no. 4 , 038 . <https://doi.org/10.1093/hropen/hoac038>

<http://hdl.handle.net/10138/350048>

<https://doi.org/10.1093/hropen/hoac038>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.


This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Evidence and consensus on technical aspects of embryo transfer

Arianna D'Angelo ^{1,*}, Costas Panayotidis², Alessandra Alteri ³,
Saria Mcheik ⁴, and Zdravka Veleva ⁵

¹Wales Fertility Institute, Swansea Bay Health Board, University Hospital of Wales, Cardiff University, Cardiff, UK ²Attiki Iatriki Advanced Gynaecological Ultrasound and Hysteroscopic Centre Private Practice, Pallini, Athens, Greece ³IRCCS San Raffaele Scientific Institute, Milan, Italy ⁴European Society of Human Reproduction and Embryology (ESHRE) Central Office, Strombeek-Bever, Belgium ⁵Helsinki University Central Hospital, Helsinki, Finland

*Correspondence address. Wales Fertility Institute, Swansea Bay Health Board, University Hospital of Wales, Cardiff University, Cardiff, UK. E-mail: dangelo@cardiff.ac.uk  <https://orcid.org/0000-0002-1505-8489>

Submitted on March 15, 2022; resubmitted on August 18, 2022; editorial decision on September 1, 2022

BACKGROUND: Ultrasound-guided embryo transfer (US-GET) is a widely performed procedure, but standards for the best practice are not available.

OBJECTIVE AND RATIONALE: This document aims to provide an overview of technical aspects of US-GET after considering the published data and including the preparation for the embryo transfer (ET) procedure, the actual procedure, the post-procedure care, associated pathologies, complications and risks, quality assurance and practitioners' performance.

SEARCH METHODS: A literature search for evidence on key aspects of the ET procedure was carried out from database inception to November 2021. Selected papers ($n = 359$) relevant to the topic were analysed by the authors. The following key points were considered in the papers: whether ultrasound (US) practice standards were explained, to what extent the ET technique was described and whether complications or incidents and how to prevent such events were reported. In the end, 89 papers could be used to support the recommendations in this document, which focused on transabdominal US-GET.

OUTCOMES: The relevant papers found in the literature search were included in the current document and described according to the topic in three main sections: requirements and preparations prior to ET, the ET procedure and training and competence for ET. Recommendations are provided on preparations prior to ET, equipment and materials, ET technique, possible risks and complications, training and competence. Specific aspects of the laboratory procedures are covered, in particular the different loading techniques and their potential impact on the final outcomes. Potential future developments and research priorities regarding the ET technique are also outlined.

LIMITATIONS, REASONS FOR CAUTION: Many topics were not covered in the literature review and some recommendations were based on expert opinions and are not necessarily evidence based.

WIDER IMPLICATIONS: ET is the last procedural step in an ART treatment and is a crucial step towards achieving a pregnancy and live birth. The current paper set out to bring together the recent developments considering all aspects of ET, especially emphasizing US quality imaging. There are still many questions needing answers, and these can be subject of future research.

STUDY FUNDING/COMPETING INTEREST(S): No funding. A.D.A. has received royalties from CRC Press and personal honorarium from Cook, Ferring and Cooper Surgical. The other co-authors have no conflicts of interest to declare that are relevant to the content of this article.

Key words: ART / embryo transfer / good practice / IVF / pregnancy / recommendations / ultrasound

WHAT DOES THIS MEAN FOR PATIENTS?

In this review, we aimed to advise on the best practice for the ultrasound-guided embryo transfer (US-GET) procedure based on the available evidence (both published articles and expert opinion). As the final step in assisted reproduction, expertise and precision in use of this technique is critical for success of the treatment, i.e. achieving a live birth. The authors attempted to establish a standardized protocol for US-GET to improve pregnancy and live birth rates, and to minimize exposure of women to unnecessary or harmful interventions. In addition, an optimal US-GET is needed to avoid multiple treatment cycles, the high expenses of which make IVF inaccessible to many, especially in developing countries. A more effective embryo transfer procedure also helps reduce the physical and psychological burden after failed cycles and would lead to fewer couples giving up before achieving a pregnancy. We have also identified the gaps in research and the need for new trials in order to optimize and standardize the US-GET technique for clinical practice, as some interventions have not been shown to be beneficial for patients or lack sufficient evidence in support of their effectiveness and safety.

Introduction

Embryo transfer (ET) is the procedure in which one or several embryos are placed into the uterine cavity. It is the final and one of the most critical steps within ART for both patient and doctor.

The effectiveness of the ET procedure is evaluated by the success rate of a commencing pregnancy using three parameters: the positive pregnancy test (urine or blood), the ultrasound (US) verification around 6–7 weeks confirming a gestational sac and/or embryo cardiac activity, and ultimately the live birth rate.

A plethora of published papers on ET technique (Tiras and Cenksoy, 2014; De los Santos et al., 2016; Practice Committee of the American Society for Reproductive Medicine, 2017; Saravelos and Li, 2019) demonstrates that different protocols, different approaches and system set-ups are followed within clinics, some of which are reported to be associated with improved outcomes after ART (Tiras and Cenksoy, 2014; Saravelos and Li, 2019). However, to date, no international consensus on ET standards of practice has been established. One possible explanation is that success of the ET procedure depends on many factors, several of which are difficult to standardize and hence to investigate. Among these are operator experience, difficulty in catheterization, embryo catheter loading technique (air bubbles, culture medium characteristics, volume of fluid), pressure and speed of injection, duration of the ET and US settings.

This article summarizes the recent evidence on ET technique through a review of the literature. It further provides some practical recommendations for practitioners and formulates standards for future ET practice, based on the collected evidence and expert opinion. The recommendations focus on ultrasound-guided embryo transfer (US-GET), in which US guidance is transabdominal.

Transabdominal US guidance is the gold standard procedure performed for ET. Transmyometrial US-GET is performed in extremely rare cases, e.g. a resistant cervix or the presence of other anatomical obstacles (e.g. severe stenosis on the cervical part, repetitive failed mock ET) (Sharif et al., 1996; Groutz et al., 1997). A transvaginal US technique for ET has also been described (Porat et al., 2010; Bodri et al., 2011; Larue et al., 2017). Because of their rare use, transmyometrial US-GET, transvaginal US-GET and clinical touch ET (where the clinician tactilely judges the correct catheter position without technical assistance) were considered to be outside the scope of this paper.

Methods

PUBMED, Embase and the Cochrane database were searched from inception to November 2021 for papers on the topic of ET technique. References (titles and abstracts) were screened and considered against the defined inclusion criteria (English language, human studies) and exclusion criteria [publication type (editorial, letter, book, studies on commercial kits, basic science studies), reviews of which a more recent version is available, not on topic of US-GET] (Fig. 1). Only papers focusing on US-GET were selected and papers on other techniques for ET were excluded. For references considered to be relevant, full-text papers were collected, assessed and summarized by the appointed author, depending on the topic (Fig. 1). Further information from guidelines and regulatory papers was added based on the experience and research of the authors.

Recommendations


The literature search resulted in 89 papers being included in the current review and described according to topic in three main sections: requirements and preparations prior to ET, the ET procedure and training and competence for ET. A summary of all the recommendations is listed in Figs 2–4.

Requirements and preparations prior to ET

Requirements prior to ET

Preparation for ET follows roughly the same steps as the preparation for oocyte retrieval in terms of infection screening and disinfection requirements (D'Angelo et al., 2019). However, there are several important differences. There is usually no need for sedation prior to ET, while filling the urinary bladder and ruling out ovarian hyperstimulation syndrome (OHSS) are additional steps that are required and are specific to ET.

Patients should be provided with preparatory information about the ET procedures (Gameiro et al., 2015). It is also important to review the patient's file before planning for ET and, in case of a concern, to consider performing a mock ET in advance. Figure 5 summarizes the list of items to be checked prior to ET and Fig. 6 summarizes the information required on the patient's record before ET is carried out.



Preparations prior to ET

Pelvic US. A pelvic US evaluation should be performed before starting ART, to decide on the ovarian stimulation protocol and to determine whether there is an anatomical abnormality of the uterus or malposition of the ovaries (Grimbizis et al., 2016). This baseline diagnostic US examination also allows for the detection of recent lesions, such as endometrial abnormalities or ovarian cysts, in a timely manner and is helpful to visualize the uterus and predict potential difficulties during ET. The time frame to perform the US is at the discretion of the clinician. The working group (WVG) recommends a baseline US, if the previous baseline US was performed more than 4–6 months earlier to

highlight any difficulties or reconfirm previous findings (D’Angelo et al., 2019). This time frame should be shortened in cases of significant conditions (endometriosis, surgery, specific symptoms).

Uterine measurement prior to ET. A large, randomized controlled trial (RCT) comparing transabdominal US-GET with a technique based on uterine length measurement before ET found no difference between the two techniques in terms of success, but the ET based on uterine length was better tolerated and easier to perform as a single operator was needed (Revelli et al., 2016).

Similarly, a prospective study found that US-GET does not enhance pregnancy rates compared with ET based on previous uterine length

- Verbal and written information should be provided to patients, according to local templates, explaining the procedure, risks, and their incidence. Confirmation of written informed consent for treatment should be obtained.
- Taking accurate patient history before ET is essential to highlight potential complications that can occur during the procedure.
- The WG recommends a baseline US if the previous baseline US was performed more than 4-6 months earlier to highlight any difficulties or reconfirm previous findings.
- Ideally, ET is performed in a room in the close proximity to the lab. The preferred temperature of the room is about 22-23°C, and if necessary, a warming blanket and socks can be used by the patient.
- The US system should be fit for ET in order to have the best image in clarity and the settings should be optimized to focus on two elements: the cervical-uterine cavity and the catheter itself.
- Soft tip catheter or rigid stylet catheter can be used. Different size speculums should be available. Other gynaecological instruments for vaginal examination and cervical manipulation in case of difficult catheterisation should be available.
- The WG recommends the use of specialised gynaecological chairs with full ability to modify the patient position.
- During the ET, the practitioner should wear clean/sterile clothes, shoes, gloves, and mask.
- The patient's records should be up to date with cervical smear/Pap test results.
- Antibiotic prophylaxis prior to ET is not recommended.
- A mock ET can be performed in order to choose the best ET catheter in cases of cervical stenosis, pelvic adhesions, or tortuous cervical canal.

Figure 2. Recommendations regarding the requirements and preparations prior to embryo transfer. ET, embryo transfer; US, ultrasound; WG, working group.

measurement (Lambers et al., 2006). However, there seems to be a relation between uterine length and successful implantation (Chun et al., 2010).

Uterine cavity measurement has been proposed as a strategy to improve ET outcomes, however, studies in the literature failed to show any cut-off value.

Doppler study. Variable practices are noted with regards to the Doppler study. It could be of value in cases of adenomyosis and for evaluating the maturity of the endometrium. Three-dimensional (3D)-US and power Doppler angiography can offer a comprehensive assessment of endometrial and sub-endometrial vascularization by using three indices: vascularization index, flow index and vascularization-flow index, which are important factors underlying endometrial receptivity.

Extensive work suggests that Doppler pulsatility index evaluation may be predictive of ET success and that better vascularity gives

better intra-myometrial and sub-endometrial blood flow (Cacciatore and Tiitinen, 1996). Chien et al. (2002) observed better ET results in patients with the presence of both endometrial and sub-endometrial flow in comparison with patients with sub-endometrial flow only or no detectable endometrial-sub-endometrial flow. In their meta-analysis, Wang et al. (2018) concluded that endometrial and sub-endometrial vasculature may be associated with the ET outcome.

Uterine 3D-US examination. 3D-US and Doppler investigations are considered helpful for the operator to become familiar with the patient's anatomy, the shape of the uterine cavity, the myometrium, the presence of possible cervical stenosis and to measure the cervical canal (D'Angelo et al., 2019). Some authors have performed 3D-US just before, during and after ET and reported the latter provided them with information about the location of the pregnancy in relation to where

the embryo was deposited (Letterie, 2005; Fang *et al.*, 2009). Yet, an RCT comparing 3D versus 2D US-GET has demonstrated that 3D US-GET is an imaging technology that does not improve clinical outcomes compared with 2D US-GET (Saravolos *et al.*, 2016). The ongoing pregnancy rates between 3D- and 2D-US groups were not significantly different (35.4% versus 37.1%, $P=0.070$, risk ratio 0.96, 95% CI 0.75–1.21). 3D-US prior to ART could be considered as a

complementary imaging study benefiting certain individual patients (e.g. in case of suspected or previous difficult ET).

Operating room, equipment and consumables

Ideally, ET should be performed in a room in close proximity to the laboratory to minimize exposure of the embryos to temperature drops. If the laboratory is some distance from the ET room,

- For practical reasons and because of individual variations in bladder volume and personal tolerance, the WG suggests that the bladder should be full enough so that the patient feels the urge to urinate
- Speculum examination just prior to ET is the final opportunity to rule out vaginal or cervical infection
- Cervical mucus can be removed with a cotton swab or alternatively with a cervical brush
- Cleansing of the vagina/cervix using sterile water or saline should be done prior to ET to minimize bacterial vagina/cervical contamination.
- Endocervical canal and/or endometrial flushing cannot be recommended as a routine procedure.
- Contraindications for ET (such as OHSS) could be ruled out by an ultrasonographic assessment of the patient immediately before ET. The ultrasonographic characteristics of OHSS such as enlarged ovaries and evidence of ascites can also be helpful in guiding towards elective freeze all embryos.
- No particular US machine is better. Individual image quality and variable preferences on US machine quality parameters have been reported
- The tubes should be assessed in order to exclude hydrosalpinx or pelvic inflammatory disease (PID). In case of tubal disease, FET should be considered until therapy (antibiotics or surgery) is completed.
- No need for sedation before ET.
- The use of myorelaxant in conjunction with ET is not recommended.
- The double step catheterisation (outer and inner catheter technique) is more reassuring than the single step catheterisation to accomplish the embryo transfer.
- There is insufficient evidence to suggest the superiority of the air-fluid or fluid-only methods during embryo loading.

Figure 3. Recommendations regarding the embryo transfer procedure. ET, embryo transfer; WG, working group; OHSS, ovarian hyperstimulation syndrome; FET, frozen embryo transfer.

(continued)

- The placement of the air bubble depends on many factors such as the pressure during the ET, the positioning of the catheter tip, the shape of the uterine cavity, the rotation of the uterine body or any uterine malformations.
- the use of small volume of medium (10-30µl) is highly recommended.
- The optimal embryo distance from fundus during ET is <10mm from the fundal endometrium
- The uterine myometrial contractility is a factor to consider during ET. ET could be more successful with less irritated uterus.
- The use of an automated syringe system where the outflow is always constant could avoid sudden expulsion of the embryo(s)
- The duration of the ET seems to be inversely related to the ET success rates.
- When the ET is completed, the embryologist has to check the catheter for possible retained embryo(s)
- Bed rest should not be recommended

Figure 3. Continued

- ET should be performed by practitioners who are competently trained (75 ETs over 2 years) in reproductive medicine. Competence should be assessed regularly through peer-to-peer observational audits
- The embryologist must be trained and have acquired embryo transfer experience under supervision before performing transfers independently. The embryologists who perform embryo transfers must pass a documented training process, confirming their high proficiency level (50 cases). Staff competence should be assessed by monitoring KPIs

Figure 4. Recommendations regarding quality assurance and performance. ET, embryo transfer; KPIs, key performance indicators.

arrangements should be made to maintain temperature and pH while transporting embryos (De los Santos et al., 2016). The preferred temperature of the ET room is 22–23°C (D'Angelo et al., 2019).

All essential items, equipment and supplies required for ET should be available. During the ET, the practitioner should wear clean/sterile clothes, shoes, gloves and a mask in accordance with European standards and local regulations.

There are no studies investigating the association between the US system quality used for ET and ET outcomes. Yet, the US system should have the ability to: adjust the field of view, depth and zoom; adjust the focal zone to the region of interest; adjust the acoustic power, colour and power Doppler capabilities; display the mechanical

and thermal indices on screen; display the catheter guide superimposed on the field of view; and to print or save images/cine loops in the system's hard drive or a central picture archiving and communication system, including image gain adjustment controls (D'Angelo et al., 2019). Settings that can increase the contrast of the different tissues or increase the black for fluid can be used to avoid artefacts.

A soft tip catheter or rigid stylet catheter can be used for the ET. The choice of transfer catheter can be based on the following characteristics: a design that is less traumatic, length, tip diameter, lumen opening at the tip for the embryo(s) passage, cost, echogenicity of the body and tip, and, overall, their consistency and whether they are easy to manipulate. A meta-analysis of two RCTs and two cohort studies

- Consent forms, clinical data and serological examinations undertaken by patients/donors prior to admission to the treatment should be available to the laboratory staff (De los Santos et al. 2016).
- US scan documents for the assessment of the endometrial lining and uterine cavity: in particular anatomical observation on the cervix – assessment of ET difficulty, angle, shortness, glands - the shape of the uterine cavity and the presence of possible internal lesions (e.g., polyps submucosal fibroids, homogeneous endometrium) as well on the adnexa, and exclusion of ovarian pathology and/or presence of hydrosalpinx.
- Information about the previous ET procedures and their evaluation (whether they were easy, feasible with stylet or difficult). The operator should be vigilant for:
 - Women who previously received a cervical treatment, cone biopsy or trachelectomy,
 - Women who have difficulties in tolerating vaginal examination (vaginismus)
 - Women with cervical canal ultrasound measures suggesting stenosis from cervical glands,
 - Women who had hysteroscopic myomectomy - particularly for those treated for myomas near the cervix
 - Women with complicated caesarean section (isthmoceles or niche)
 - Women who had permanent cervical cerclage
- Possible previous traumatic experiences (difficult examination, sexual assaults, severe anxiety)

Figure 5. List of items to be checked before embryo transfer. ET, embryo transfer.

Based on the revised guidelines for good practice in IVF laboratories(De los Santos et al. 2016), the patient records required prior to ET procedure should include:

1. Batch number and type of media used for transfer.
2. Time from oocyte retrieval to transfer.
3. Time from oocyte insemination to transfer.
4. Date and time of embryo transfer.
5. Name of the operator loading the catheter.
6. Name of the practitioner performing the transfer.
7. Number, developmental stage and quality of embryo(s) at the time of transfer.
8. Type of catheter used for transfer.
9. Fate of supernumerary embryos.
10. Patient medical history: previous cervical interventions, previous ET, previous difficulties in cannulation, up to date cervical smear test.

A double identity check of the patient, the patient file, and the culture dish(es) are mandatory immediately before the transfer.

The patient records post- ET procedure should include details about the procedure, e.g., presence of blood, retained embryo(s) and justification when more than one embryo is transferred.

Figure 6. Patient information required on the patient records. ET, embryo transfer.

showed that the pregnancy rates were improved when soft catheters were used compared to rigid catheters ([Practice Committee of the American Society for Reproductive Medicine, 2017](#)).

Gynaecological instruments for vaginal examination and cervical manipulation in case of a difficult catheterization should be available. These include speculum, vorcellum/tenaculum, dilators, sponge

holder, cotton swabs, cotton buds for cervical mucus, forceps and a cleaning set including saline/cleaning solution.

A standard medium-size Cusco-type speculum is the most commonly used, but different sizes of speculum should also be available. However, in specific patients (e.g. obese patients, or those with previous vaginal surgeries) other types of speculum can be used, such as Colin's or Grave's.

Patient and practitioner positioning

Gynaecological positioning of the patient in the semi-lithotomy or lithotomy can facilitate the ET performance. The position of the patient may need to be adapted to patient mobility.

The gynaecological examination chair should provide optimal comfort for the patient and ergonomic positioning for the doctor and assistant. The WG recommends the use of specialized gynaecological chairs with full ability to modify the patient position.

Mock ET

A mock ET (or a dummy ET) can be performed in a preceding cycle (Ali et al., 2008), at the time of oocyte retrieval (Mirkin et al., 2003) or immediately before ET (Prapas et al., 1995). It has been shown not to increase the frequency of uterine contractions (Torre et al., 2010). A mock ET can diminish the incidence of difficult transfers by allowing the physician to choose the most suitable catheter. Other potential benefits, such as an accurate measurement of the uterine cavity length, are most beneficial in settings where US is not available at ET (Shamonki et al., 2005). However, most patients who proceed to ET might have already had mapping of the cervical canal, through previous IUIs or sonohysterography. Moreover, US guidance during ET is widely available nowadays. US can be used for moulding the ET catheter to the uterocervical angle (Sallam et al., 2002). It should also be noted that a retroverted uterus will often change position, even to anteversion, thus challenging the effectiveness of a mock ET (Henne and Milki, 2004).

A mock ET seems to be less valuable in case of US-GET and in the presence of a thorough documentation of observations of previous transcervical procedures. Nonetheless, it can be performed in order to choose the best ET catheter in cases of cervical stenosis, pelvic adhesions or tortuous cervical canal.

Prevention of infection

The patient's records should be up to date with cervical smear/PAP test results. A vaginal infection screening (including, if necessary, a vaginal swab for bacteriological examination) should be performed during diagnostic work-up according to local guidelines and regulations.

The evidence from the literature does not support the administration of prophylactic antibiotics in association with ET. A systematic review of the literature in 2012 concluded that prophylaxis with amoxicillin and clavulanic acid did not improve IVF success rates (Kroon et al., 2012). Similarly, a retrospective study of 876 fresh and frozen-thawed transfers, with or without oral doxycycline and methylprednisolone, found no independent effect of antibiotic prophylaxis at ET on the success of treatment (Kaye et al., 2017).

A survey of IVF clinics in the USA reported that 40% of them use antibiotic prophylaxis (Beshar et al., 2021); the authors of this study conducted a retrospective analysis of the transfer of 250 single euploid

blastocysts in frozen cycles and showed that doxycycline prophylaxis did not result in higher live birth and ongoing pregnancy rates.

In women with symptoms of infection, it is recommended to perform specific microbiological testing and take appropriate actions (D'Angelo et al., 2019). In general, antibiotic prophylaxis should be used only when supported by evidence, since it can induce resistance and can have negative side effects, including *Clostridium difficile* and fungal infection (Shirlow et al., 2017).

Regarding vaginal screening, the literature search did not provide information on this topic.

Associated pathologies and factors affecting success

The successful implantation requires a receptive intrauterine environment for the embryo(s) and the presence of uterine pathologies can negatively affect the success rate of the ET.

Preparation for a difficult ET is required in cases of the following associated pathologies: uterine malformation, fibroids, obesity, endometriosis and post-surgery pelvic inflammatory disease (PID). Ideally, these pathologies are detected in the pre-ET US and either treated or considered in the preparation of a difficult ET.

The use of screening hysteroscopy may reveal intrauterine pathologies that may not be diagnosed by transvaginal US and the use of hysteroscopic surgery to optimize the uterine cavity (e.g. septum resection) may be of value.

On the one hand, a Cochrane systematic review including 11 RCTs showed that there is insufficient data to decide whether routine screening hysteroscopy increases live birth and clinical pregnancy, be it for all women or those with two or more failed IVF attempts (Kamath et al., 2019). A recent RCT assessed the role of office hysteroscopy prior to the first ART cycle and the authors reported that the office hysteroscopy did not improve ART results. Minimal intrauterine anomalies not diagnosed by transvaginal US or hysterosalpingography do not seem to negatively affect ART outcomes (Ben Abid et al., 2021). On the other hand, there is low-quality evidence that operative hysteroscopy increases the pregnancy rate in infertile women with previously diagnosed polyps (Di Spiezio Sardo et al., 2016).

More robust and high-quality RCTs are needed to demonstrate the benefit of diagnostic and/or operative hysteroscopy before ET in the general population. Thus, this intervention could not be recommended based on the existing evidence.

Patient preparation for ET

Before carrying out the ET, the identity of the patient should be checked and the World Health Organization surgical safety checklist applied (World Alliance for Patient Safety, 2008).

A full urinary bladder

The patient should attend the ET procedure with a full bladder. This straightens the angle between the uterine cervix and uterine body (Sundström et al., 1984; Lewin et al., 1997) and facilitates visualization using the transabdominal US scan. A straighter cervical canal and smaller inclination of the uterine body facilitate the effortless insertion of an ET catheter into the correct spot in the uterine cavity (Abou-Setta, 2007). It was suggested in a large study by Lewin et al. (1997) that performing ET with a full bladder increases the clinical pregnancy rate. Two smaller RCTs failed to show such an effect (Mitchell et al., 1989; Lorusso et al., 2005). A difference in fluid intake instructions

may account for the difference in results: [Lewin et al. \(1997\)](#) instructed patients to take 1000 ml of liquids, whereas [Mitchell et al. \(1989\)](#) required only 250 ml, and [Lorusso et al. \(2005\)](#) required a 'moderately filled' bladder.

For practical reasons and because of individual variations in bladder volume and personal tolerance, the WG suggests that the bladder should be full enough so that the patient feels the urge to urinate. If necessary, after the initial assessment, the patient can be instructed to drink an additional one to two cups of water in order to achieve optimal distension of cervico-uterine angle. Excessive distension of the bladder can cause significant discomfort to the patient, and the assistant may not be able to produce the best US images despite the posterior bladder enhancement of the image. However, by optimal positioning of the patient, the anteverted uterus can change its position by force of gravity, even with smaller bladder volumes. Therefore, patient positioning and use of gynaecological couches that feature bottom and back tilting with fully optimizable leg support, can avoid significant technical difficulties in ET and improve the patient experience with the procedure.

In the case of a retroverted uterus, a full bladder makes the uterine-cervical angle more pronounced and therefore more difficult to catheterize.

Speculum examination

Speculum examination just prior to ET is the final opportunity to rule out vaginal or cervical infection. The practitioner should also evaluate the external appearance of the cervix in order to rule out potential signs of difficult ET such as cervical polyps or cervical ectropion/inflammation.

Removing mucus from the cervical canal. Removing mucus from the cervical canal can facilitate the insertion of an ET catheter into the uterine cavity and it can potentially avoid a clogged catheter tip, or relocation of mucus within the uterine cavity, which may affect implantation. On the other hand, removing cervical mucus might stimulate uterine contractility or cervical bleeding, which can have a negative effect on the ET outcome.

Cervical mucus can be removed with a cotton swab or a cervical brush, although the latter is considered to be slightly more traumatic with a higher risk of provoking uterine contractions. Careful catheter aspiration of mucus is another option. In an early trial, [Mansour et al. \(1994\)](#) injected methylene blue dye into the uterine cavity in a mock ET and concluded that expulsion of the dye was significantly reduced after the removal of cervical mucus. One RCT in which cervical mucus was removed with sterile cotton swabs ([Moini et al., 2011](#)) and a prospective cohort study of catheter aspiration ([Eskandar et al., 2007](#)) demonstrated that removing mucus improved clinical outcomes. However, a meta-analysis of eight RCTs (including the previously cited RCTs) involving 1715 women reported very little evidence of an overall benefit of cervical mucus removal before ET ([Craciunas et al., 2014](#)). A similar conclusion was reported by a Cochrane meta-analysis, even if the methods of mucus removal and studies included are questionable ([Derks et al., 2009](#)).

Disinfection

Cleansing of the vagina/cervix should be carried out prior to ET to minimize bacterial vaginal/cervical contamination. Currently, most practitioners achieve this by using sterile water or saline. Cleaning prior

to ET should be performed delicately in order to avoid bleeding. This is important not only for the success of ET catheterization but also for diminishing the subsequent stress of the patient if she detects spotting after ET.

Flushing the endocervical canal and endometrial cavity prior to ET

A Cochrane meta-analysis including studies on the effect of flushing the endocervical canal or the endometrial cavity on pregnancy rates found no evidence of any substantial benefit ([Derks et al., 2009](#)). Owing to the lack of benefit, endocervical canal and/or endometrial flushing cannot be recommended as a routine procedure.

Pelvic US immediately prior to ET

The aim of the US assessment of the patient immediately before ET is to rule out contraindications for ET. Among contraindications for ET, OHSS is the most common and potentially life-threatening. While the ultimate decision on cancelling ET because of OHSS relies also on laboratory findings and subjective symptoms, the US characteristics of OHSS, such as enlarged ovaries and evidence of ascites, can also help guide towards elective freeze-all of embryos ([D'Angelo and Amso, 2002](#)).

US prior to ET additionally aims to confirm a beneficial uterine environment and endometrium, i.e. an endometrial thickness of preferably >7 mm ([Kasius et al., 2014](#)). Although optimal results from ART can be achieved in patients with regular uterine cavities with no deformities (e.g. septa, fibroids or polyps), a small study on patients with uterine polyps up to 15 mm in length, some of which were treated through hysteroscopic resection, reported that the presence of small polyps was not associated with poorer pregnancy and implantation rates ([Isikoglu et al., 2006](#)).

Occasionally, fluid in the uterine cavity at the time of ET can be observed in patients with hydrosalpinx whose tubes communicate freely with the uterine cavity (i.e. those who have not undergone salpingectomy or tubal obliteration) ([Melo et al., 2020](#)).

The presence of intrauterine fluid prior to ET seems to be an unfavourable prognostic factor; the tubes should be assessed in order to exclude hydrosalpinx or PID. Elective freeze-all of the embryos should be considered until therapy of tubal disease (antibiotics or surgery) is completed ([Melo et al., 2020](#)).

Pain relief and uterine myorelaxant

It is believed that patient feedback during ET is important in ensuring an atraumatic procedure with minimal uterine contractions and minimal disruption to the endometrium.

A trial evaluated the use of phenazopyridine, a bladder analgesic, for reducing discomfort during ET ([Frishman et al., 2007](#)). A single dose of the medicine, administered 1 h prior to ET, failed to reduce discomfort, as measured with a visual analogue pain scale. In the American Society for Reproductive Medicine guidelines on ET, acupuncture, analgesics, massage, general anaesthesia and traditional Chinese medicine were listed as having no beneficial effect on pregnancy ([Practice Committee of the American Society for Reproductive Medicine, 2017](#)). While not necessarily required for ET, variable sedation techniques have been reported, including the use of sedative drugs, such as Propofol, Rapifen, Fentanyl and Diazepam. Verbal analgesia by the seditionist/assistant is another important anxiolysis form ([D'Angelo et al., 2019](#)).

The ET procedure

There are two potential ET practices: single step and double step ET.

The single-step option involves the use of a prepared soft catheter. These catheters have a very smooth and flexible inner part and a second external tube to protect the inner part as well to give more stability during the insertion. With this concept, internal cervical/uterine trauma is avoided and the embryo(s) can be passed through the cervical canal. There is a risk, however, that difficulties are encountered in passing through the cervical canal, or even that it is impossible to pass the soft catheter.

In the double-step option, a rigid double catheter is used and passed through the cervical canal up to the top end of the cervix. The second step is to remove the inner part and replace it with the softer catheter containing the embryo(s). Although the double-step option may increase the risk of cervical/internal trauma or increase patient discomfort, it is more reassuring for the operator to successfully pass and complete the ET.

Catheter loading

The embryo loading technique represents a critical aspect of the procedure and might affect ART outcomes. The choice of the syringe, type of catheter, type and volume of transfer medium, presence of air bubble, catheter loading speed and embryo(s) placement in the catheter may be variables involved in the success of the procedure.

Two main catheter loading methods have been described: the air-fluid method (air-embryo-air or medium-air-embryo-air-medium) (Fig. 7A and B, respectively) and the fluid-only method (Fig. 7C).

In the air-fluid method, the loading of the syringe-catheter complex with the transferred volume consists of the transfer media (which contains the embryo(s)) separated by air spaces on both sides. In the fluid-only method, the embryo(s) is placed in a complete column of fluid, without any air brackets or bubbles. A systematic review and meta-analysis of two prospective randomized trials concluded that there was insufficient evidence to suggest the superiority of the air-fluid or fluid-only methods during embryo loading (Abou-Setta et al., 2007).

Moreover, the effect of medium volume and the presence of air bubbles on clinical outcomes appear to be controversial. Generally, the use of a small volume of medium (10–30 μ l) is highly recommended. Indeed, a large volume of transfer medium (>60 μ l) may increase the chance of dislocation of the transferred embryo(s) from the uterus into the cervix or Fallopian tubes, predisposing to ectopic pregnancy. Likewise, a very small medium volume (<10 μ l) along with air bubbles seemed to have a negative effect on implantation and pregnancy rates (Ebner et al., 2001). A comparison between 40–50 and 15–20 μ l showed that a higher volume is associated with increased implantation and pregnancy rates (Montag et al., 2002). However, some studies found no difference in terms of clinical outcomes between low (15–25 μ l) and high (35–45 μ l) transfer volume (Omid et al., 2015; Sigalos et al., 2018).

Of note is that loading the catheter directly from the culture micro drop under the oil versus loading from the transfer dish without an oil layer leads to similar pregnancy rates (Halvaei et al., 2013).

Air bubbles might help with US visualization of the ET catheter and proper placement of the embryo(s) (Schoolcraft, 2016). The use of air bubbles in the catheter might also protect the embryos from the cervical mucus and accidental discharge before entering the endometrial

cavity (Tiras et al., 2012). The embryo is within the surface of this bubble and follows the flow of the liquid used as propulsion, usually to the top part of the uterine cavity (floating).

ET technique and procedure

Once the resolution of the cervical image on the US is optimized, the practitioner attempts to pass the flexible tip of the catheter directly through the cervical canal under US guidance. If an angled uterine body in relation to the cervix is detected, this can be corrected by manipulating the speculum holder ascending or descending the initial portion of the cervix situated between the speculum valves. The possibility to adjust the pelvic part of the gynaecological couch aiming to descend the bottom of the patient and the back simultaneously can be very helpful and makes the uterine body lower (due to gravity) and aligned with the cervical canal.

The practitioner should try to pass the catheter as smoothly as possible in an axial trajectory without bending it or irritating the patient's cervix and/or endometrium. Bending, repeated attempts to catheterize or difficulties to progress within the uterine cavity can result in irritation of the myometrium, creating micro-contractions (Fanchin et al., 1998). Two studies concluded that during the ET those who had better IVF outcome had a less irritated uterus (Sammali et al., 2018; Blank et al., 2020). However, the use of atosiban, an oxytocin receptor antagonist, to reduce the uterine contraction during ET is unlikely to improve the clinical pregnancy rate or the live birth rate in the general IVF patient population (Buddhabunyan et al., 2021). The clinical pregnancy rate in older women (>35 years old) in the atosiban group was twice that of the placebo group, but the result was not statistically significant (Buddhabunyan et al., 2021).

3D-US and the four-dimensional-US with abdominal probes can demonstrate the positioning of the embryo(s) or show more realistically the catheter in relation to the uterine cavity. Reports showed 80% accuracy of the embryo(s) positioning with 3D-US imaging just after ET with a subsequent scan confirming an intrauterine pregnancy (Baba et al., 2000).

To improve the US image, the assistant can apply gentle pressure on the patient's abdomen, although a similar effect can be reached by adjusting the US settings on more recent US systems (Fig. 8).

Extensive work on identifying the optimal place to release the embryo has taken place because the movement of the final position of the embryo is unlikely to be a factor predicting the success of ET (Allahbadia et al., 2008; Ozcan et al., 2016). The middle upper area gives better results in terms of implantation and pregnancy rates (Oliveira et al., 2004). Cavagna et al. (2006) similarly suggested avoiding ET in the lower regions of the uterine cavity, as this may result in higher miscarriage rates.

Higher pregnancy rates were obtained when the position of the air bubble from the fundal endometrial surface was <10 mm. When the inner catheter tip is placed 1.5–2 cm from the fundal endometrium, a best performance is expected (Cenksoy et al., 2014). In a more recent retrospective study, it was shown that the probability of pregnancy, clinical pregnancy and ongoing pregnancy decreases as the distance from the fundus (DFF) to the air bubble (SD: 10.27 ± 3.0 mm) increases (Bayram et al., 2021). When all variables remained constant, an increase of 1 mm of DFF changed the odds of pregnancy by 0.882; of clinical pregnancy by 0.891 and of ongoing pregnancy by 0.925.




Figure 7. Catheter loading methods used in human embryo transfer. (A and B) Air–fluid method: the transfer media containing the embryo(s) separated by air spaces on both sides. (C) fluid-only method: the embryo(s) is placed in a complete column of fluid without any air brackets or bubbles.

There are no studies to evaluate what is the best way to withdraw the catheter, i.e. as inner and outer together, or separately.

Pressure in the piston. Some authors have proposed an automated syringe system where the outflow is always constant (Caanen *et al.*, 2016) with the potential benefit to avoid sudden expulsion of the embryo(s).

The injection pressure during the ET is related to the catheter length and therefore if the catheter end is in the proximity of the fundal end, one should use the lowest pressure possible in order to complete the ET with a gentle injection and maintain the embryo(s) as close to the upper fundal area as possible. In cases of larger uterine cavities, the ET catheter end may not achieve ideal positioning towards the fundal end. Chun *et al.* (2010) reported a possible higher miscarriage rate when the embryo bubble did not reach the highest point.

Duration of the ET procedure. The duration of the ET procedure is the time taken for: the embryologist to prepare the loaded catheter; catheter transport to operator hands; the catheterization (one step or two steps), passing through the cervix and moving the catheter tip to the targeted point within the endometrial cavity; the injection; and catheter withdrawal. The duration of the ET procedure has been shown to have a significant influence on pregnancy success rates, with a duration of the transfer of more than 120 s having a negative effect (Matorras *et al.*, 2004).

A plausible explanation for the association between duration of the procedure and ET outcomes could be the time during which embryo(s) is(are) outside the incubator as well as the difficulty of ET. The duration of the ET seems to be inversely related to the ET success rates (Abdelmassih *et al.*, 2007). Cetin *et al.* (2010) observed that the

highest ET success rates were noted when the time from when the embryo(s) was loaded on the catheter to the time when the embryo(s) was released was <44 s. They showed a 35% success rate for young women (<35 years old). However, for the difficult ETs, the time was detrimental for the older women (>35 years old) and if the ET time is more than 60 s, the difference between the two groups was significant, with 30% success for women <35 years old and 13% success for women >35 years old.

End of procedure and post-procedure care

When the ET is completed, the embryologist has to check the catheter for possible retained embryo(s). This is a crucial quality control procedure. The embryo should be reinjected immediately; a retrospective analysis of data from 12 studies showed that the implantation rate, the clinical pregnancy rate and the pregnancy loss rates were not decreased for patients undergoing immediate re-transfer after embryo retention (Practice Committee of the American Society for Reproductive Medicine, 2017).

A Cochrane review (Abou-Setta *et al.*, 2014) and two systematic reviews (Craciunas and Tsampras, 2016; Cozzolino *et al.*, 2019) looked at bed rest after ET and found that immediate mobilization after ET does not influence success rates. Therefore, bed rest should not be recommended.

Complications and troubleshooting

Complications and risks associated with the ET procedure

Complications during ET (e.g. the presence of cervical trauma/bleeding, retained embryo in the catheter, rare expulsion of the embryo




Figure 8. Ultrasound images showing various factors related to a transabdominal ultrasound-guided embryo transfer. (A) Zoom window focused on the uterine cavity. (B) Deep field view of embryo transfer (ET) during catheterization. (C) Retroverted uterus ET. (D) ET catheter approaching the fundal part of the endometrium. (E) Measurement of distances between tip of catheter and fundal endometrium and between embryo bubble and endometrium. (F) Artefacts during ET: example of mirror image artefact (uterus is anteverted but there is a false image showing a retroverted uterus).

from the cervix and short-term post-ET infection) are very rare. Long-term complications in ART are failure to achieve pregnancy, ectopic pregnancy, miscarriage and multiple pregnancies.

Although the majority of ETs are straightforward, some degree of difficulty can be encountered, even if there is no consensus on what qualifies as a difficult ET. Generally, ETs have been defined as difficult when they cause discomfort to the patient or when there is the presence of blood at the end of the catheter. There may be some anatomical difficulty to access the uterine cavity and, in this case, the ET requires the use of specific tools such as catheters with sheaths and rigid mandrels (stylet). In addition, the embryo may be retained in the

transfer catheter, which is usually detected by the embryologist and requires a repeated transfer.

The effect of difficult transfers on clinical outcomes is debated. While some studies reported no harmful effect of difficult transfers, others reported detrimental effects on clinical outcomes (Arora and Mishra, 2018).

Troubleshooting during ET

In cases of cervical catheterization difficulties, where the catheter does not pass the cervical canal, only partially passes or is bent, a more forced catheterization can be attempted using a tenaculum to stabilize

Table 1 Staff training and competence in performing embryo transfer.

	Ovarian stimulation and trigger	Oocyte retrieval	Embryo transfer
Training		Number of procedures to complete	
	100 cycles*	75*	75*
Competence		Monitor PIs to check competence and skills Take appropriate action when there is a gap between actual and expected performance	

*The numbers are those proposed by the ESHRE clinic PI working group.

Table adapted from [Vlaisavljevic et al. \(2021\)](#).

PI, performance indicators.

the cervix (traction), a thicker stylet catheter or external cervical dilators. In case of a complete cervical catheterization failure, it is advised to proceed to freeze all of the embryos and hysteroscopic assessment.

An important point to consider in troubleshooting is when to abandon the procedure and when to try again, for example, with another catheter (e.g. change to a rigid one) or resort to gentle dilatation of the cervix. There are very limited data informing such a decision. [Tur-Kaspa et al. \(1998\)](#) showed that in case of a difficult ET, uterine manipulation or cervical dilatation or repeated attempts at the time of the ET could be performed without adversely affecting the pregnancy outcomes. Until further conclusive data are available, it can be recommended that the practitioner decides in a case-by-case approach on the most appropriate course of action.

Quality assurance and performance

Training and competence

Clinicians. Two RCTs confirmed that US-GET performed by either midwives or experienced nurses does not impact negatively on the outcomes ([Bjuresten et al., 2003](#); [Rinaldi et al., 2014](#)).

The assistant's experience does not seem to impact outcomes following US-GET ([Harris et al., 2009](#)). Thus, assistance during ET for someone without formal US training is a reasonable option.

ET should be performed by practitioners who are competently trained in reproductive medicine. In some countries, fertility specialists or nurses can be trained to perform ET procedures, but there are currently no generally accepted minimal requirements for training. For safety reasons, and wherever feasible, a simulator could be the initial part of structured training for novices who want to perform this procedure, enabling them to acquire basic skills and reach a predefined level of performance in a safe and controlled environment before applying the procedure to patients ([Soave et al., 2019](#)).

The number of procedures to be completed for training (within 2 years) is 75 according to the recently published paper based on the Maribor Consensus ([Vlaisavljevic et al., 2021](#)).

In addition to training, competence in a certain procedure should be maintained. Criteria for assessing proficiency/competency on the technical aspects of ET have not been described, but suggested criteria are pregnancy rates, number of ETs performed relative to the size of the clinic and ectopic pregnancy rates.

Competence should be assessed regularly through peer-to-peer observational audits, the frequency of which should be decided within the team.

Embryologists. The embryologist must be trained and have acquired ET experience under supervision before performing transfers independently. The embryologists who perform ET must pass a documented training process, confirming a high proficiency level. Each laboratory has its own training programme that includes: reading and understanding the standard operating procedures; observing the ET procedures performed by qualified and experienced embryologists; loading of discarding material without any loss; and performing a minimum of 50 ET procedures under supervision. After reaching pregnancy rates within two standard deviations of the average pregnancy rate in the laboratory/clinic, the trainee can be authorized to perform the ET procedure without supervision ([Montag and Morbeck, 2017](#)). A good training programme includes training on how to deal with difficulties and problems such as embryo return, re-loading of embryos and options for difficult transfers.

Maintaining embryologists' competence is critical as well. To demonstrate each embryologist's expertise, a certain procedure number should be recorded in a logbook ([Alpha Scientists in Reproductive Medicine, 2015](#)). For example, ESHRE requires the completion of 50 cases in 3 years for the ET procedure in order to evaluate the competency of a Clinical Embryologist ([Kovačič et al., 2020](#)).

Staff competence should be assessed by monitoring key performance indicators (ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine, 2017) (Table 1). The maintenance of achieved competence should be monitored annually, even for senior embryologists and if necessary, re-training is recommended.

Future developments

Transvaginal US-GET has some benefits such as no need for a full bladder avoiding bladder discomfort, the practitioner is doing both transvaginal sonography (TVS) and ET with no need for assistance, and the TVS has a far better image. However, the process to catheterize prior to the transvaginal US-GET takes more time for preparation than a standard ET under the transabdominal US-GET and it is unclear what could be done in case of a difficult ET ([Bodri et al., 2011](#)). [Bodri et al. \(2011\)](#) did not show superior success rates between transvaginal US-GET and standard transabdominal US-GET.

[Cozzolino et al. \(2018\)](#) concluded that in three recent RCTs the quality of evidence supporting the equivalence of the transvaginal versus transabdominal approach in clinical pregnancy or live birth rates is low and they identified the need for larger RCTs.

Future research should focus on factors and methods that could increase the ET success rate. An association between the US system

quality used for ET and the outcomes has not yet been investigated; whether optimizing the image quality can help practitioners avoid cervical catheterization difficulties, manipulate the ET catheter more gently and complete the ET procedure in an atraumatic and precise way is not known.

More evidence-based knowledge is needed regarding the catheter loading techniques to compare the air-fluid or fluid-only methods during embryo loading and also regarding the best way to withdraw the catheter (as inner and outer together, or separately).

Conclusion

ET is the last procedural step in ART and is crucial for achieving a pregnancy and live birth. The current paper set out to bring together recent developments concerning all aspects of ET, especially emphasizing US quality imaging. There are still many questions needing answers, and these can be the subject of future research.

What is clear is that the performance of ET is not researched in-depth and objective data that are based on US criteria are not routinely recorded or checked. The sequence of what steps are needed to perform ET is similar in most fertility clinics around the world.

Although ETs are performed frequently, clear standards and quality criteria to improve their effectiveness are needed.

The authors' opinion is that US quality combined with a gentle tactile technique can make a difference in pregnancy rate improvement at ET.

Data availability

This article conducts a literature review of existing research records, and no new data were generated or analysed in support of this manuscript.

Acknowledgements

The authors would like to thank the ESHRE central office for its support.

Authors' roles

A.D.A. and C.P. proposed the topic of the paper. S.M. performed the literature search and contributed to the coordination of the WG. All authors contributed equally to drafting the article or revising it critically. All authors approved the final version.

Funding

The authors declare no funding was given to this work.

Conflict of interest

A.D.A. has received, during the last 36 past months, royalties from CRC press and personal honorarium from Cook, Ferring and Cooper Surgical. The other co-authors have no conflicts of interest to declare that are relevant to the content of this article.

References

- Abdelmassih VG, Neme RM, Dozortsev D, Abdelmassih S, Diamond MP, Abdelmassih R. Location of the embryo-transfer catheter guide before the internal uterine os improves the outcome of in vitro fertilization. *Fertil Steril* 2007;**88**:499–503.
- Abou-Setta AM. Effect of passive uterine straightening during embryo transfer: a systematic review and meta-analysis. *Acta Obstet Gynecol Scand* 2007;**86**:516–522.
- Abou-Setta AM, Mansour RT, Al-Inany HG, Aboulghar MM, Aboulghar MA, Serour GI. Among women undergoing embryo transfer, is the probability of pregnancy and live birth improved with ultrasound guidance over clinical touch alone? A systemic review and meta-analysis of prospective randomized trials. *Fertil Steril* 2007;**88**:333–341.
- Abou-Setta AM, Peters LR, D'Angelo A, Sallam HN, Hart RJ, Al-Inany HG; Cochrane Gynaecology and Fertility Group. Post-embryo transfer interventions for assisted reproduction technology cycles. *Cochrane Database Syst Rev* 2014;**8**:CD006567.
- Ali CR, Khashan AS, Horne G, Fitzgerald CT, Nardo LG. Implantation, clinical pregnancy and miscarriage rates after introduction of ultrasound-guided embryo transfer. *Reprod Biomed Online* 2008;**17**:88–93.
- Allahbadia GN, Gandhi G, Kadam K, Arora S, Awasthi A, Nagwekar A, Allahbadia S, Wolman I. Antibubble trajectory during embryo transfers in donor egg IVF does not predict success. *Reprod Biomed Online* 2008;**16**:881–885.
- Alpha Scientists in Reproductive Medicine. The Alpha Consensus Meeting on the professional status of the clinical embryologist: proceedings of an expert meeting. *Reprod Biomed Online* 2015;**30**:451–461.
- Arora P, Mishra V. Difficult embryo transfer: a systematic review. *J Hum Reprod Sci* 2018;**11**:229–235.
- Baba K, Ishihara O, Hayashi N, Saitoh M, Taya J, Kinoshita K. Three-dimensional ultrasound in embryo transfer. *Ultrasound Obstet Gynecol* 2000;**16**:372–373.
- Bayram A, De Munck N, Elkhatib I, Aranz A, El-Damen A, Abdala A, Coughlan C, Garrido N, Vidales LM, Lawrenz B et al. The position of the euploid blastocyst in the uterine cavity influences implantation. *Reprod Biomed Online* 2021;**43**:880–889.
- Ben Abid H, Fekih M, Fathallah K, Chachia S, Bibi M, Khairi H. Office hysteroscopy before first in vitro fertilization. A randomized controlled trial. *J Gynecol Obstet Hum Reprod* 2021;**50**:102109.
- Beshar I, Johal JK, Bavan B, Milki AA. Withholding antibiotics does not reduce clinical pregnancy outcomes of natural cycle frozen embryo transfers. *Fertil Steril* 2021;**115**:1225–1231.
- Bjuresten K, Hreinsson JG, Fridström M, Rosenlund B, Ek I, Hovatta O. Embryo transfer by midwife or gynecologist: a prospective randomized study. *Acta Obstet Gynecol Scand* 2003;**82**:462–466.
- Blank C, Sammalı F, Kuijsters N, Huang Y, Rabotti C, de Sutter P, Mischi M, Schoot B. Assessment of uterine activity during IVF by quantitative ultrasound imaging: a pilot study. *Reprod Biomed Online* 2020;**41**:1045–1053.
- Bodri D, Colodrón M, García D, Obradors A, Vernaev V, Coll O. Transvaginal versus transabdominal ultrasound guidance for embryo transfer in donor oocyte recipients: a randomized clinical trial. *Fertil Steril* 2011;**95**:2263–2268, 2268.e1.

- Buddhabunyan N, Sothornwit J, Seejorn K, Buppasiri P, Salang L. Effects of atosiban on uterine peristalsis following frozen embryo transfer: a randomized controlled trial. *Eur J Obstet Gynecol Reprod Biol* 2021;**265**:96–101.
- Caanen MR, van der Houwen LE, Schats R, Vergouw CG, de Leeuw B, Lambers MJ, Groeneveld E, Lambalk CB, Hompes PG. Embryo transfer with controlled injection speed to increase pregnancy rates: a randomized controlled trial. *Gynecol Obstet Invest* 2016;**81**: 394–404.
- Cacciatore B, Tiitinen A. Does ovarian stimulation affect uterine artery impedance? *J Assist Reprod Genet* 1996;**13**:15–18.
- Cavagna M, Contart P, Petersen CG, Mauri AL, Martins AM, Baruffi RL, Oliveira JB, Franco JG Jr. Implantation sites after embryo transfer into the central area of the uterine cavity. *Reprod Biomed Online* 2006;**13**:541–546.
- Cenksoy PO, Ficicioglu C, Yesiladali M, Akcin OA, Kaspar C. The importance of the length of uterine cavity, the position of the tip of the inner catheter and the distance between the fundal endometrial surface and the air bubbles as determinants of the pregnancy rate in IVF cycles. *Eur J Obstet Gynecol Reprod Biol* 2014;**172**: 46–50.
- Cetin MT, Kumtepe Y, Kiran H, Seydaoglu G. Factors affecting pregnancy in IVF: age and duration of embryo transfer. *Reprod Biomed Online* 2010;**20**:380–386.
- Chien LW, Au HK, Chen PL, Xiao J, Tzeng CR. Assessment of uterine receptivity by the endometrial-subendometrial blood flow distribution pattern in women undergoing in vitro fertilization-embryo transfer. *Fertil Steril* 2002;**78**:245–251.
- Chun SS, Chung MJ, Chong GO, Park KS, Lee TH. Relationship between the length of the uterine cavity and clinical pregnancy rates after in vitro fertilization or intracytoplasmic sperm injection. *Fertil Steril* 2010;**93**:663–665.
- Cozzolino M, Troiano G, Esencan E. Bed rest after an embryo transfer: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2019;**300**:1121–1130.
- Cozzolino M, Vitagliano A, Di Giovanni MV, Laganà AS, Vitale SG, Blaganje M, Drusany Starić K, Borut K, Patrelli TS, Noventa M. Ultrasound-guided embryo transfer: summary of the evidence and new perspectives. A systematic review and meta-analysis. *Reprod Biomed Online* 2018;**36**:524–542.
- Craciunas L, Tsampras N. Bed rest following embryo transfer might negatively affect the outcome of IVF/ICSI: a systematic review and meta-analysis. *Hum Fertil (Camb)* 2016;**19**:16–22.
- Craciunas L, Tsampras N, Fitzgerald C. Cervical mucus removal before embryo transfer in women undergoing in vitro fertilization/intracytoplasmic sperm injection: a systematic review and meta-analysis of randomized controlled trials. *Fertil Steril* 2014;**101**: 1302–1307.
- D'Angelo A, Amso NN. Embryo freezing for preventing ovarian hyperstimulation syndrome: a Cochrane review. *Hum Reprod* 2002;**17**:2787–2794.
- D'Angelo A, Panayotidis C, Amso N, Marci R, Matorras R, Onofriescu M, Turp AB, Vandekerckhove F, Veleva Z, Vermeulen N et al.; ESHRE Working Group on Ultrasound in ART. Recommendations for good practice in ultrasound: oocyte pick up. *Hum Reprod Open* 2019;**2019**:hoz025.
- De los Santos MJ, Apter S, Coticchio G, Debrock S, Lundin K, Plancha CE, Prados F, Rienzi L, Verheyen G, Woodward B et al.; ESHRE Guideline Group on Good Practice in IVF Labs. Revised guidelines for good practice in IVF laboratories (2015). *Hum Reprod* 2016;**31**:685–686.
- Derks RS, Farquhar C, Mol BWJ, Buckingham K, Heineman MJ. Techniques for preparation prior to embryo transfer. *Cochrane Database of Systematic Reviews* 2009;**4**:CD007682.
- Di Spiezio Sardo A, Di Carlo C, Minozzi S, Spinelli M, Pistotti V, Alviggi C, De Placido G, Nappi C, Bifulco G. Efficacy of hysteroscopy in improving reproductive outcomes of infertile couples: a systematic review and meta-analysis. *Hum Reprod Update* 2016;**22**: 479–496.
- Ebner T, Yaman C, Moser M, Sommergruber M, Pölz W, Tews G. The ineffective loading process of the embryo transfer catheter alters implantation and pregnancy rates. *Fertil Steril* 2001;**76**: 630–632.
- ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine. The Vienna consensus: report of an expert meeting on the development of ART laboratory performance indicators. *Reprod Biomed Online* 2017;**35**:494–510.
- Eskandar MA, Abou-Setta AM, El-Amin M, Almushait MA, Sobande AA. Removal of cervical mucus prior to embryo transfer improves pregnancy rates in women undergoing assisted reproduction. *Reprod Biomed Online* 2007;**14**:308–313.
- Fanchin R, Righini C, Olivennes F, Taylor S, de Ziegler D, Frydman R. Uterine contractions at the time of embryo transfer alter pregnancy rates after in-vitro fertilization. *Hum Reprod* 1998;**13**: 1968–1974.
- Fang L, Sun Y, Su Y, Guo Y. Advantages of 3-dimensional sonography in embryo transfer. *J Ultrasound Med* 2009;**28**:573–578.
- Frishman GN, Allsworth JE, Gannon JB, Wright KP. Use of phenazopyridine for reducing discomfort during embryo transfer. *Fertil Steril* 2007;**87**:1010–1014.
- Gameiro S, Boivin J, Dancet E, de Klerk C, Emery M, Lewis-Jones C, Thorn P, Van den Broeck U, Venetis C, Verhaak CM et al. ESHRE guideline: routine psychosocial care in infertility and medically assisted reproduction—a guide for fertility staff. *Hum Reprod* 2015;**30**: 2476–2485.
- Grimbizis GF, Di Spiezio Sardo A, Saravelos SH, Gordts S, Exacoustos C, Van Schoubroeck D, Bermejo C, Amso NN, Nargund G, Timmerman D et al. The Thessaloniki ESHRE/ESGE consensus on diagnosis of female genital anomalies. *Hum Reprod* 2016;**31**:2–7.
- Groutz A, Lessing JB, Wolf Y, Azem F, Yovel I, Amit A. Comparison of transmyometrial and transcervical embryo transfer in patients with previously failed in vitro fertilization-embryo transfer cycles and/or cervical stenosis. *Fertil Steril* 1997;**67**:1073–1076.
- Halvaei I, Khalili MA, Razi MH, Agha-Rahimi A, Nottola SA. Impact of different embryo loading techniques on pregnancy rates in in vitro fertilization/embryo transfer cycles. *J Hum Reprod Sci* 2013;**6**:65–69.
- Harris ID, Styer AK, Petrozza JC. Ultrasonographer experience does not impact outcomes following ultrasound-guided embryo transfer. *Fertil Steril* 2009;**92**:918–922.
- Henne MB, Milki AA. Uterine position at real embryo transfer compared with mock embryo transfer. *Hum Reprod* 2004;**19**:570–572.

- Isikoglu M, Berkkanoglu M, Senturk Z, Coetzee K, Ozgur K. Endometrial polyps smaller than 1.5 cm do not affect ICSI outcome. *Reprod Biomed Online* 2006;**12**:199–204.
- Kamath MS, Bosteels J, D'Hooghe TM, Seshadri S, Weyers S, Mol BWJ, Broekmans FJ, Sunkara SK. Screening hysteroscopy in subfertile women and women undergoing assisted reproduction. *Cochrane Database Syst Rev* 2019;**4**:CD012856.
- Kasius A, Smit JG, Torrance HL, Eijkemans MJ, Mol BW, Opmeer BC, Broekmans FJ. Endometrial thickness and pregnancy rates after IVF: a systematic review and meta-analysis. *Hum Reprod Update* 2014;**20**:530–541.
- Kaye L, Will EA, Bartolucci A, Nulsen J, Benadiva C, Engmann L. Pregnancy rates for single embryo transfer (SET) of day 5 and day 6 blastocysts after cryopreservation by vitrification and slow freeze. *J Assist Reprod Genet* 2017;**34**:913–919.
- Kovačić B, Prados FJ, Plas C, Woodward BJ, Verheyen G, Ramos L, Mäkinen S, Apter SJ, Vidal F, Ziebe S et al. ESHRE Clinical Embryologist certification: the first 10 years. *Hum Reprod Open* 2020;**2020**:hoaa026.
- Kroon B, Hart RJ, Wong BMS, Ford E, Yazdani A; Cochrane Gynaecology and Fertility Group. Antibiotics prior to embryo transfer in ART. *Cochrane Database Syst Rev* 2012;**3**:CD008995.
- Lambers MJ, Dogan E, Kosteljik H, Lens JW, Schats R, Hompes PG. Ultrasonographic-guided embryo transfer does not enhance pregnancy rates compared with embryo transfer based on previous uterine length measurement. *Fertil Steril* 2006;**86**:867–872.
- Larue L, Keromnes G, Massari A, Roche C, Moulin J, Gronier H, Bouret D, Cassuto NG, Ayel JP. Transvaginal ultrasound-guided embryo transfer in IVF. *J Gynecol Obstet Hum Reprod* 2017;**46**:411–416.
- Letterie GS. Three-dimensional ultrasound-guided embryo transfer: a preliminary study. *Am J Obstet Gynecol* 2005;**192**:1983–1987; discussion 1987–1988.
- Lewin A, Schenker JG, Avrech O, Shapira S, Safran A, Friedler S. The role of uterine straightening by passive bladder distension before embryo transfer in IVF cycles. *J Assist Reprod Genet* 1997;**14**:32–34.
- Lorusso F, Depalo R, Bettocchi S, Vacca M, Vimercati A, Selvaggi L. Outcome of in vitro fertilization after transabdominal ultrasound-assisted embryo transfer with a full or empty bladder. *Fertil Steril* 2005;**84**:1046–1048.
- Mansour RT, Aboulghar MA, Serour GI, Amin YM. Dummy embryo transfer using methylene blue dye. *Hum Reprod* 1994;**9**:1257–1259.
- Matorras R, Mendoza R, Expósito A, Rodriguez-Escudero FJ. Influence of the time interval between embryo catheter loading and discharging on the success of IVF. *Hum Reprod* 2004;**19**:2027–2030.
- Melo P, Georgiou EX, Johnson N, van Voorst SF, Strandell A, Mol BWJ, Becker C, Granne IE. Surgical treatment for tubal disease in women due to undergo in vitro fertilisation. *Cochrane Database Syst Rev* 2020;**10**:CD002125.
- Mirkin S, Jones EL, Mayer JF, Stadtmauer L, Gibbons WE, Oehninger S. Impact of transabdominal ultrasound guidance on performance and outcome of transcervical uterine embryo transfer. *J Assist Reprod Genet* 2003;**20**:318–322.
- Mitchell JD, Wardle PG, Foster PA, Hull MG. Effect of bladder filling on embryo transfer. *J In Vitro Fert Embryo Transf* 1989;**6**:263–265.
- Moini A, Kiani K, Bahmanabadi A, Akhoond M, Akhlaghi A. Improvement in pregnancy rate by removal of cervical discharge prior to embryo transfer in ICSI cycles: a randomised clinical trial. *Aust N Z J Obstet Gynaecol* 2011;**51**:315–320.
- Montag M, Kupka M, van der Ven K, van der Ven H. Embryo transfer on day 3 using low versus high fluid volume. *Eur J Obstet Gynecol Reprod Biol* 2002;**102**:57–60.
- Montag M, Morbeck D. *Principles of IVF Laboratory Practice*. Cambridge, UK: Cambridge University Press, 2017.
- Oliveira JBA, Martins AMVC, Baruffi RLR, Mauri AL, Petersen CG, Felipe V, Contart P, Pontes A, Franco JG. Increased implantation and pregnancy rates obtained by placing the tip of the transfer catheter in the central area of the endometrial cavity. *Reprod Biomed Online* 2004;**9**:435–441.
- Omid M, Halvaei I, Mangoli E, Khalili MA, Razi MH. The effect of embryo catheter loading technique on the live birth rate. *Clin Exp Reprod Med* 2015;**42**:175–180.
- Ozcan P, Ficioglu C, Kokulu MB, Alagoz O, Yesiladali M. Can the movement of the air bubbles after embryo transfer predict the success of IVF treatment? *J Clin Anal Med* 2016;**7**:371–374.
- Porat N, Boehnlein LM, Schouweiler CM, Kang J, Lindheim SR. Interim analysis of a randomized clinical trial comparing abdominal versus transvaginal ultrasound-guided embryo transfer. *J Obstet Gynaecol Res* 2010;**36**:384–392.
- Practice Committee of the American Society for Reproductive Medicine. Performing the embryo transfer: a guideline. *Fertil Steril* 2017;**107**:882–896.
- Prapas Y, Prapas N, Hatziparasidou A, Prapa S, Nijs M, Vanderzwalmen P, Vlassis G, Jones EE. The echoguide embryo transfer maximizes the IVF results. *Acta Eur Fertil* 1995;**26**:113–115.
- Revelli A, Rovei V, Dalmasso P, Gennarelli G, Racca C, Evangelista F, Benedetto C. Large randomized trial comparing transabdominal ultrasound-guided embryo transfer with a technique based on uterine length measurement before embryo transfer. *Ultrasound Obstet Gynecol* 2016;**48**:289–295.
- Rinaldi L, Floccari A, Selman H. Ultrasound guidance of embryo transfer: a role for midwife. *Sex Reprod Healthc* 2014;**5**:47–49.
- Sallam HN, Agameya AF, Rahman AF, Ezzeldin F, Sallam AN. Ultrasound measurement of the uterocervical angle before embryo transfer: a prospective controlled study. *Hum Reprod* 2002;**17**:1767–1772.
- Sammali F, Kuijsters NPM, Schoot BC, Mischi M, Rabotti C. Feasibility of transabdominal electrohysterography for analysis of uterine activity in nonpregnant women. *Reprod Sci* 2018;**25**:1124–1133.
- Saravelos SH, Kong GW, Chung JP, Mak JS, Chung CH, Cheung LP, Li TC. A prospective randomized controlled trial of 3D versus 2D ultrasound-guided embryo transfer in women undergoing ART treatment. *Hum Reprod* 2016;**31**:2255–2260.
- Saravelos SH, Li TC. Embryo transfer techniques. *Best Pract Res Clin Obstet Gynaecol* 2019;**59**:77–88.
- Schoolcraft WB. Importance of embryo transfer technique in maximizing assisted reproductive outcomes. *Fertil Steril* 2016;**105**:855–860.
- Shamonki MI, Spandorfer SD, Rosenwaks Z. Ultrasound-guided embryo transfer and the accuracy of trial embryo transfer. *Hum Reprod* 2005;**20**:709–716.

- Sharif K, Afnan M, Lenton W, Bilalis D, Hunjan M, Khalaf Y. Transmyometrial embryo transfer after difficult immediate mock transcervical transfer. *Fertil Steril* 1996;**65**:1071–1074.
- Shirlow R, Healey M, Volovsky M, MacLachlan V, Vollenhoven B. The effects of adjuvant therapies on embryo transfer success. *J Reprod Infertil* 2017;**18**:368–378.
- Sigalos G, Michalopoulos Y, Kastoras AG, Triantafyllidou O, Vlahos NF. Low versus high volume of culture medium during embryo transfer: a randomized clinical trial. *J Assist Reprod Genet* 2018;**35**:693–699.
- Soave I, D'Angelo A, Piva I, Marci R. A pilot study on oocyte retrieval simulator: a new tool for training? *J Med Syst* 2019;**43**:202.
- Sundström P, Wramsby H, Persson PH, Liedholm P. Filled bladder simplifies human embryo transfer. *Br J Obstet Gynaecol* 1984;**91**:506–507.
- Tiras B, Cenksoy PO. Practice of embryo transfer: recommendations during and after. *Semin Reprod Med* 2014;**32**:291–296.
- Tiras B, Korucuoglu U, Polat M, Saltik A, Zeyneloglu HB, Yarali H. Effect of air bubble localization after transfer on embryo transfer outcomes. *Eur J Obstet Gynecol Reprod Biol* 2012;**164**:52–54.
- Torre A, Scheffer JB, Schönauer LM, Frydman N, Fanchin R. Mock embryo transfer does not affect uterine contractility. *Fertil Steril* 2010;**93**:1343–1346.
- Tur-Kaspa I, Yuval Y, Bider D, Levron J, Shulman A, Dor J. Difficult or repeated sequential embryo transfers do not adversely affect in-vitro fertilization pregnancy rates or outcome. *Hum Reprod* 1998;**13**:2452–2455.
- Vlaisavljevic V, Apter S, Capalbo A, D'Angelo A, Gianaroli L, Griesinger G, Kolibianakis EM, Lainas G, Mardesic T, Motrenko T et al.; ESHRE Clinic PI Working Group. The Maribor consensus: report of an expert meeting on the development of performance indicators for clinical practice in ART. *Hum Reprod Open* 2021;**2021**:hoab022.
- Wang J, Xia F, Zhou Y, Wei X, Zhuang Y, Huang Y. Association between endometrial/subendometrial vasculature and embryo transfer outcome: a meta-analysis and subgroup analysis. *J Ultrasound Med* 2018;**37**:149–163.
- World Alliance for Patient Safety. *World Alliance for Patient Safety: Forward Programme 2008-2009*. Geneva: World Health Organization, 2008.