

1 ***Pseudomonas aeruginosa* utilises the host-derived polyamine spermidine to**
2 **facilitate antimicrobial tolerance**

3

4

5 Chowdhury M. Hasan¹, Sian Pottenger¹, Angharad E. Green¹, Adrienne A. Cox^{1,2}, Jack
6 White,² Trevor Jones¹, Craig Winstanley¹, Aras Kadioglu¹, Megan Wright², Daniel R. Neill^{1,*},
7 and Joanne L. Fothergill^{1,*}

8

9 ¹ Department of Clinical Infection, Microbiology and Immunology, Institute of Infection,
10 Veterinary and Ecological Sciences, University of Liverpool, United Kingdom

11 ² School of Chemistry and Astbury Centre for Structural Molecular Biology, University of
12 Leeds, United Kingdom

13

14

15 **Correspondence:** Joanne L Fothergill (j.fothergill@liverpool.ac.uk, +44(0)151 79 59646) or
16 Daniel R Neill (d.neill@liverpool.ac.uk, +44(0)151 79 59622), Department of Clinical
17 Infection, Microbiology and Immunology, Institute of Infection, Veterinary and Ecological
18 Sciences, Ronald Ross Building, 8 West Derby Street, Liverpool, UK, L69 7BE

19

20 * Equal contributions

21

22 The authors have declared that no conflict of interest exists.

23

24

25

26

27 **Abstract**

28 *Pseudomonas aeruginosa* undergoes diversification during infection of the cystic fibrosis (CF)
29 lung. Understanding these changes requires model systems that capture the complexity of
30 the CF lung environment. We previously identified loss-of-function mutations in the two-
31 component regulatory system sensor kinase gene *pmrB*, in *P. aeruginosa* from CF and from
32 experimental infection of mice. Here, we demonstrate that whilst such mutations lower in
33 vitro MICs for multiple antimicrobial classes, this is not reflected in increased antibiotic
34 susceptibility in vivo. Loss of PmrB impairs aminoarabinose modification of
35 lipopolysaccharide, increasing the negative charge of the outer membrane and promoting
36 uptake of cationic antimicrobials. However, in vivo, this can be offset by increased membrane
37 binding of other positively charged molecules present in lungs. The polyamine spermidine
38 readily coats the surface of PmrB-deficient *P. aeruginosa*, reducing susceptibility to
39 antibiotics that rely on charge differences to bind the outer membrane and increasing biofilm
40 formation. Spermidine is elevated in lungs during *P. aeruginosa* infection in mice and during
41 episodes of antimicrobial treatment in people with CF. These findings highlight the need to
42 study antimicrobial resistance under clinically relevant environmental conditions. Microbial
43 mutations carrying fitness costs in vitro may be advantageous during infection, where host
44 resources can be utilised.

45 **Introduction**

46 *Pseudomonas aeruginosa* is a ubiquitous environmental bacterium and a metabolically
47 versatile opportunistic pathogen, responsible for severe, acute nosocomial infections (1, 2).
48 It is also the most frequently recovered pathogen of the cystic fibrosis (CF) lung (3), where it
49 causes chronic infection that is associated with pulmonary exacerbations and declining lung
50 function. Such infections are difficult to treat, due in part to intrinsic antimicrobial resistance
51 of *P. aeruginosa*, together with adaptive resistance mechanisms induced by the presence of
52 antimicrobial agents or other environmental factors (4). The chronic nature of *P. aeruginosa*
53 infection of the CF lung necessitates long-term, high dose antimicrobial therapy, creating
54 conditions conducive to the emergence and selection of acquired resistance mechanisms
55 (5). Phenotypic flexibility and a large genome encoding complex regulatory machinery makes
56 chronically colonised *P. aeruginosa* a challenge to eradicate and necessitates frequent
57 review of treatment regimens for people with CF.

58

59 Lipopolysaccharide (LPS) is a key component of the Gram-negative outer membrane and
60 can be stabilized by the addition of divalent cations, including Mg^{2+} and Ca^{2+} . Cationic
61 antimicrobials, including polymyxin B, colistin and host-derived peptides such as LL37, exert
62 their effects via disruption of cell membrane integrity, but rely on charge differentials with the
63 outer membrane in order to bind (6). Modifications of LPS that reduce cationic antimicrobial
64 binding affinity and penetration can result in resistance (7). One such modification is the
65 addition of positively charged 4-amino-4-deoxy-L-arabinose (L-Ara4N) to the lipid A
66 component of LPS (8), mediated in *P. aeruginosa* by the proteins encoded by the
67 *arnBCADTEF-ugd* operon (9). Expression of operon genes is regulated by two component
68 signalling systems, such as PhoPQ and PmrAB (10, 11), which are activated under conditions
69 of locally decreased divalent cation concentrations. This ensures that the charge of the
70 membrane can be maintained when Mg^{2+} and Ca^{2+} are limited. PhoPQ- and PmrAB-induced

71 expression of the *arn* operon results in high level resistance to both cationic peptides and
72 aminoglycosides (12). Specific mutations in *pmrAB* have been implicated in polymyxin
73 resistance, via upregulation of both the lipid A deacylase *pagL* and the *arn* operon (13-15).

74

75 Expression of genes *PA4775*, *speE2* (*PA4774*) and *speD2* (*PA4773*) (16), located adjacent
76 to *pmrAB* on the chromosome, is also induced by Mg²⁺ limiting conditions, due to the
77 presence of a PmrA binding site close to *speD2* (6). These genes are involved in the
78 synthesis of the polyamine spermidine. During environmental stress or periods of low cation
79 availability, PmrAB stimulates polyamine synthesis and these coat the bacterial surface,
80 increasing the outer membrane charge and providing protection against both antimicrobial
81 agents and oxidative stress (6).

82

83 Polyamines are polycationic hydrocarbons, containing two or more amine groups, and are
84 abundant across all three kingdoms of life. Polyamine utilisation and uptake genes are found
85 in all bacteria (17) and species including *P. aeruginosa* can utilise polyamines as a sole
86 carbon source for growth (18). Putrescine, spermidine and cadaverine, the principal
87 polyamines of prokaryotes, have been implicated in iron and free radical scavenging, acid
88 resistance, biofilm formation, protection from the phagolysosome, interaction with
89 components of cell envelopes and antimicrobial resistance (17). However, there is
90 uncertainty regarding the effect, if any, that polyamines have on antimicrobial susceptibility
91 in *P. aeruginosa*. Increased susceptibility to multiple classes of antibiotics was observed
92 when PAO1 was cultured with polyamines, (19) but addition of exogenous polyamines to a
93 PAO1 lacking a functional spermidine synthase (*speE2*) partially protected the outer
94 membrane from polymyxin B (6). Whilst the extent of the role played by polyamines in *P.*
95 *aeruginosa* growth, virulence and antimicrobial resistance has not been fully determined, it is
96 notable that spermine was found to be elevated in the airways of those with CF and that

97 levels have been reported to decrease during treatment of pulmonary exacerbations (20),
98 whilst those of putrescine have been found to decrease(21).

99

100 In a previous study, we identified loss of function mutations in *pmrB* in *P. aeruginosa* isolated
101 from the airways of mice, following experimental infection, and in isolates taken from people
102 with CF (22, 23). These isolates showed enhanced susceptibility to multiple classes of
103 antibiotics. Here, we sought to understand why loss of function *pmrB* mutations might be
104 retained in *P. aeruginosa*, in an environment of prolonged antimicrobial exposure, such as
105 the CF lung. As we had observed altered LPS structure in *pmrB* mutants (23), we
106 hypothesised that host-derived molecules might play a role in stabilising the outer membrane
107 of *P. aeruginosa* in vivo, thereby overcoming the lack of PmrAB-driven modifications of lipid
108 A. Here, we propose that the host cationic polyamine spermidine acts in this way, negating
109 the antimicrobial susceptibility phenotype of *P. aeruginosa* lacking functional PmrB. These
110 findings highlight the need to conduct antimicrobial susceptibility testing under
111 environmentally-relevant conditions.

112

113 **Results**

114 ***P. aeruginosa* lacking PmrB show enhanced antimicrobial susceptibility in vitro but** 115 **not in vivo**

116 We previously described susceptibility to multiple classes of antibiotics in *P. aeruginosa* with
117 naturally acquired loss of function mutations in *pmrB* and in a *pmrB*-deletion strain of LESB65
118 (23). To determine whether this susceptibility would result in improved infection outcomes
119 following onset of antimicrobial therapy, we infected mice with LESB65 or a LESB65 mutant
120 lacking *pmrB* (LESB65 Δ *pmrB*) and then treated with intra-nasal colistin at 6 and 24 h post-
121 infection. In LESB65-infected mice, colistin treatment led to significant reductions in the
122 number of *P. aeruginosa* recovered from both the upper airways (nasopharynx and sinuses)

123 (Figure 1A) and the lungs (Figure 1B), with four out of eight mice clearing the infection
124 completely. By contrast, colistin treatment did not significantly alter the bacterial burdens
125 recovered from LESB65 Δ *pmrB*-infected animals (Figure 1A and B). Consistent with our
126 previous findings, the LESB65 Δ *pmrB* strain colonised lungs at a higher density than its wild
127 type parent strain (Figure 1B). We subsequently performed in vitro antimicrobial susceptibility
128 testing with bacteria recovered from the infections and confirmed that LESB65 Δ *pmrB*
129 retained its susceptibility to colistin in vitro (Supplementary Figure 1).

130

131 **Proteomics analysis suggests a switch from polyamine synthesis to uptake and** 132 **utilisation in LESB65 Δ *pmrB***

133 To further explore the environment-dependent antimicrobial susceptibility profile of LESB65
134 and LESB65 Δ *pmrB*, we revisited proteomics data obtained from late-exponential bacterial
135 cultures grown in LB (23). We identified inter-strain abundance differences in a group of
136 functionally related proteins involved in polyamine transport, biosynthesis and metabolism.
137 Bacteria can synthesise polyamines or acquire them via environmental uptake. Genes
138 involved in polyamine synthesis are co-transcribed with *pmrA* and *pmrB* (24) and the proteins
139 encoded by those genes (*SpeD2*, *SpeE2* and PA4775) were found at greatly reduced
140 abundance both in LESB65 Δ *pmrB* and in an LESB65-derived isolate with a naturally-
141 acquired missense mutation in *pmrB* (LESB65*pmrB*^{L255Q}) (Figure 2A). There is a *pmrA*
142 binding sequence close to the start codon of *speD2* (6) and these data suggest that in the
143 absence of a functional PmrAB system, expression of the operon is significantly reduced.
144 However, the reduced abundance of polyamine synthesis proteins in these strains appears
145 to be offset by a corresponding increased abundance of the polyamine binding, uptake and
146 utilisation proteins of the *SpuABCDEFGH* operon (Figure 2B). Of the six proteins of this
147 operon that were detected in proteomics analysis, four were significantly more abundant in
148 both LESB65 Δ *pmrB* and LESB65*pmrB*^{L255Q}, as compared to LESB65.

149

150 **Spermidine is abundant in both the airways and increases during infection and**
151 **antimicrobial treatment**

152 The apparent increase in polyamine binding and acquisition proteins in the *pmrB* mutant
153 strains may be advantageous in environments that are rich in free polyamines. Others have
154 reported polyamine abundance in CF sputum and changes in bioavailability associated with
155 pulmonary exacerbations(20, 21). As the spermidine synthesis proteins were significantly
156 decreased in abundance in PmrB-deficient *P. aeruginosa*, we sought to determine whether
157 the polyamine could instead be scavenged from the environment. We measured spermidine
158 levels in the sinuses and lungs of both uninfected mice and those with chronic *P. aeruginosa*
159 LESB65 infection (Figure 3). Spermidine was detectable at comparable concentrations in
160 sinuses and lungs and was found to increase in the context of infection. This increase is
161 unlikely to result from polyamine production in *P. aeruginosa*, as the levels of spermidine
162 produced by high density cultures of LESB65 or LESB65 Δ *pmrB* were ~1000-fold lower than
163 those detected in respiratory tissues (Supplementary Figure 2).

164

165 We also detected free polyamines in sputum from people with CF with chronic *P. aeruginosa*
166 infection (Figure 4). Spermidine was quantifiable in samples from 18 of 19 people tested
167 (Figure 4A). There was considerable inter- (Figure 4A) and intra- (Figure 4B) participant
168 variability in sputum spermidine levels, but levels were higher during periods of antimicrobial
169 treatment (Figure 4C), suggesting potential for polyamines to influence treatment efficacy.

170

171 Another polyamine, spermine, was detected in all sputum samples and was found to be more
172 abundant, relative to spermidine. Like spermidine, there was considerable interparticipant
173 variation (Supplementary Figure 3A), however there were no significant differences between
174 baseline, exacerbation and treatment samples (Supplementary Figure 3B). Spermine levels
175 were also investigated in the lungs and sinuses of uninfected and *P. aeruginosa* infected

176 mice (Supplementary Figure 3C). Spermine levels were below the detection limit of the ELISA
177 in all lung samples tested and in 7/10 sinus samples. Thus, whilst both spermidine and
178 spermine are produced in both human and mouse respiratory tissue, their relative abundance
179 appears to be species specific.

180

181 **PmrB genotype influences *P. aeruginosa* surface interactions with spermidine**

182 Having demonstrated that spermidine is available within the airways, we next sought to
183 characterise whether *P. aeruginosa* might interact with this cationic molecule. Using purified
184 spermidine tagged with fluorescent nitrobenzoxadiazole (NBD), we first investigated whether
185 the polyamine could interact with the bacterial surface and if such interactions were transient
186 or prolonged. For these assays, we used non-toxic concentrations of spermidine, determined
187 by broth microdilution (Supplementary Figure 4). LESB65 and LESB65 Δ *pmrB* were co-
188 incubated with 4 mM unlabelled spermidine or spermidine-NBD for 30 minutes before the
189 bacteria were pelleted by centrifugation and resuspended in PBS. We then determined the
190 extent of spermidine binding to the bacterial surface by flow cytometry. Spermidine-NBD
191 bound both LESB65 and LESB65 Δ *pmrB*, as evidenced by increasing median fluorescence
192 intensity [MFI] for *P. aeruginosa* co-cultured with labelled vs unlabelled spermidine (MFI 10.4
193 vs 3.00 for LESB65 with spermidine-NBD vs unlabelled spermidine, MFI 73.3 vs 3.13 for
194 LESB65 Δ *pmrB*, $p < 0.0001$ for both strains) (Figure 5A and B). The NBD fluorescence of the
195 LESB65 Δ *pmrB* population was significantly higher than that of LESB65 (MFI 73.3
196 LESB65 Δ *pmrB* vs 10.4 LESB65, $p < 0.001$). Furthermore, fluorescence was retained longer
197 in LESB65 Δ *pmrB* cultures (MFI 73.3 at 0 mins, 67.5 at 30 mins, 62.3 at 60 mins, 56.9 at 120
198 mins, 54.6 at 240 mins) than in LESB65 cultures (MFI 10.4 at 0 mins, 7.30 at 30 mins, 5.49
199 at 60 mins, 4.60 at 120 mins, 4.59 at 240 mins) (Figure 5A and B), suggesting prolonged
200 binding or uptake in the LESB65 Δ *pmrB* strain. Binding, and surface coating of NBD-

201 spermidine to the *P. aeruginosa* membrane was confirmed by fluorescence microscopy
202 (Supplementary Figure 5).

203

204 We next investigated whether the same interactions between *P. aeruginosa* and spermidine
205 might take place in more chemically-complex environments, more reflective of CF lung
206 conditions. To this end, we repeated the flow cytometry assay using irradiated CF sputum as
207 the assay buffer. Binding to spermidine was reduced for both LESB65 and LESB65 Δ *pmrB* in
208 CF sputum, as compared to in PBS, although significant interaction was still apparent for both
209 strains (Figure 5C). Furthermore, as was observed in PBS, LESB65 Δ *pmrB* demonstrated a
210 significantly enhanced spermidine binding capacity in CF sputum, relative to the wild type
211 control (Figure 5C). The staining pattern of both strains was altered in CF sputum, with
212 evidence of brightly and dimly stained sub-populations of bacteria (Figure 5D). These
213 subpopulations may result from dynamic modification of surface charge in response to
214 PmrAB-independent environmental sensing mechanisms activated in CF sputum.

215

216 **Surface-coated spermidine protects *P. aeruginosa* from antimicrobials and offsets the**
217 **susceptibility associated with loss of PmrB function**

218 Polyamines have been implicated in resistance to several classes of antibiotics, with binding
219 of these positively charged molecules to the *P. aeruginosa* membrane reducing charge
220 interactions with cationic antimicrobials (6, 24). The decreased abundance of SpeD2, SpeE2
221 and PA4775 in loss-of-function *pmrB* mutants may, therefore, contribute to the observed
222 increases in antimicrobial susceptibility, under conditions where polyamines cannot be readily
223 scavenged from the environment. However, where polyamines are abundant, the increased
224 polyamine-binding potential of LESB65 Δ *pmrB* might offset the inherent susceptibility to
225 antimicrobials associated with loss of PmrB function. To explore this idea, we performed MIC
226 assays with *P. aeruginosa* without spermidine, in the presence of spermidine, or with *P.*

227 *aeruginosa* that had been pre-incubated with spermidine and then pelleted and washed
228 before addition of antibiotics. (Table 1). Assays were performed with PmrB-deficient strains
229 on both the LESB65 and PAO1 backgrounds. In both cases, spermidine increased the colistin
230 MIC50 of the *pmrB*-deficient strain, but not the wild type ancestor, by 4-8 fold. This was the
231 case both when the spermidine was present throughout the assay and when strains were
232 pre-incubated with the polyamine. In polyamine-rich environments, such as the respiratory
233 tract, surface-coating of *P. aeruginosa* with cationic polyamines may achieve a comparable
234 outcome to PmrAB-driven L-Ara-4N addition to LPS lipid A, by increasing the positive charge
235 of the outer membrane. This finding may explain why loss-of-function *pmrB* mutations are
236 retained in *P. aeruginosa* causing chronic infection of the CF lung, despite prolonged, high
237 dose antimicrobial treatment.

238

239 **Exogenous spermidine influences *P. aeruginosa* biofilm formation**

240 Polyamines contribute to biofilm formation in both Gram positive and Gram negative bacterial
241 species (25). They serve as structural components of the extracellular matrix, but also
242 stimulate signalling that can promote or inhibit biofilm formation (26, 27). However, the role
243 of polyamines in *P. aeruginosa* biofilm formation is not well described. To determine whether
244 spermidine-induced reductions in antimicrobial susceptibility in LESB65 Δ *pmrB* might be
245 augmented by enhanced biofilm formation, we quantified surface attached biofilm by crystal
246 violet staining. LESB65 formed biofilm more readily than LESB65 Δ *pmrB* (Figure 6A).
247 However, whilst the addition of spermidine to cultures led to a modest reduction in LESB65
248 biofilm mass, it induced a significant increase in LESB65 Δ *pmrB* biofilm mass over a 48 hour
249 period (Figure 6B). LESB65 Δ *pmrB* biofilms were more susceptible to colistin treatment than
250 those formed by LESB65, but this susceptibility was lost with the addition of spermidine to
251 cultures (Figure 6C). Thus, spermidine can enhance biofilm formation and promote biofilm
252 tolerance to colistin via a mechanism dependent on *pmrB* genotype.

253

254 **Discussion**

255 Despite their ubiquity and essentiality across all kingdoms of life, understanding of polyamine
256 biology is limited. *P. aeruginosa* can synthesise polyamines from methionine, arginine or
257 ornithine, but also scavenge them from their environment. In the context of infection,
258 spermidine levels have been reported to be reduced in lungs in idiopathic pulmonary fibrosis,
259 relative to healthy controls (28), whilst levels are increased in chronic obstructive pulmonary
260 disease, particularly in those who smoke (29). It has been suggested that spermidine has a
261 synergistic effect when co-administered with β -lactams for treatment of MRSA keratitis (19),
262 but that polyamines also limit host-derived antimicrobial killing (30) and enhance *S. aureus*
263 growth and virulence in this context (31). The expression of *P. aeruginosa* polyamine
264 synthesis, uptake and utilisation genes is under environmental control, including by the
265 PmrAB two-component regulatory system.

266

267 The PmrAB locus is present in pathogenic Gram negative species, including *P. aeruginosa*,
268 *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Escherichia coli* and *Salmonella enterica*
269 (32). The PmrAB system plays a key role in environmental sensing, stimulating production of
270 polyamines and modification of LPS to buffer the outer membrane charge in conditions of low
271 divalent cation availability (14). Mutations in *pmrB* are frequently identified in *P. aeruginosa*
272 infecting the CF lung (33), and both activating and loss-of-function mutations have been
273 described (14, 23). This apparent dichotomy may reflect environmental differences between
274 individuals or between niches within the host. Where divalent cations or other positively
275 charged molecules, including polyamines, can be co-opted to buffer membrane charge, loss
276 of function *pmrB* mutations may be retained due to the advantages they confer in terms of
277 lysozyme resistance and enhanced adherence to host surfaces (23). However, under

278 conditions where cationic molecules are scarce, activating mutations in *pmrB* would likely
279 confer a greater advantage, by promoting the LPS modifications that reduce binding of
280 antimicrobials to the bacterial outer membrane. Of note, *P. aeruginosa* has the capacity to
281 actively promote polyamine bioavailability during infection. Contact with host cells induces a
282 characteristic gene expression pattern in *P. aeruginosa* that leads to spermidine production
283 and localisation to the cell surface. This process is dependent on flagellar-mediated motility,
284 with non-motile bacteria failing to produce spermidine upon host cell contact (34). However,
285 in the present study, the observed effects do not appear to be flagella-dependent, as our
286 findings were comparable with *pmrB* deletion mutants generated in both LESB65 and PAO1.
287 PAO1 carries a single polar flagellum, whilst LESB65 lacks one.

288

289 The differences in the kinetics of interaction with spermidine in wild type and PmrB-deficient
290 LESB65 suggest that surface charge may influence the capacity of *P. aeruginosa* to utilise
291 exogenous polyamines. Whilst it is clear that detection of divalent cations through PhoPQ
292 and PmrAB can modulate polyamine synthesis and uptake pathways, as well as inducing
293 surface charge modifications, it remains to be determined whether direct sensing of
294 polyamine abundance can modulate those same processes. An ability to alter surface
295 charge, and thus change the efficiency of polyamine binding, in response to the local
296 availability of those molecules, would offer advantages in metabolic resource management.

297

298 Similarly, whilst the findings presented here demonstrate spermidine binding to the surface
299 of *P. aeruginosa*, it is unclear whether the changes in antimicrobial susceptibility and biofilm
300 formation that we observed are a direct result of that physical interaction or whether they are
301 a consequence of polyamine-induced signalling. The increase in positive charge associated
302 with spermidine coating of the outer membrane may be sufficient to explain the increased
303 resistance to cationic antimicrobials such as colistin, and spermidine might act as a substrate
304 for biofilm formation, encourage greater surface interactions via the change in membrane

305 charge or aid in chelation of negatively charged biofilm DNA. However, we can't rule out
306 further contributions from spermidine-induced signalling and others have reported an effect
307 of exogenous polyamines on bacterial pathogen gene expression, including in *P. aeruginosa*
308 (35, 36).

309

310 The composition and charge of the outer membrane of Gram-negative bacteria are key
311 determinants of antimicrobial resistance and major barriers to antibiotic uptake (37). The
312 findings presented here go some way towards explaining the dichotomy of retention of loss-
313 of-function *pmrB* mutations in *P. aeruginosa*, in the face of antimicrobial pressure. The
314 susceptibility of PmrB-deficient LESB65 to antimicrobials in vitro did not translate to
315 susceptibility within the lung environment and this may be explained by buffering of the
316 negatively-charged outer membrane with host-derived cationic molecules, including
317 spermidine, together with the phenotypic advantages in vivo, including lysozyme resistance
318 and increased attachment to host surfaces, that derive from loss of PmrB (23). This highlights
319 the need to perform antimicrobial susceptibility testing under conditions that are relevant to
320 infection and that capture key environmental cues that are sensed by pathogens or with which
321 they interact. This is a particular challenge for those interested in pathogens of the CF lung,
322 given the complexity of that environment and the difficulty in replicating its physical, chemical
323 and microbiological features in the laboratory. However, substantial progress has been made
324 in this area (38), through use of metabolic profiling of CF pathogens (39) and analytical
325 methods that utilise next-generation sequencing data to aid in the benchmarking of new
326 models that aim to replicate conditions of the CF lung (40). As models continue to be refined,
327 consideration should be given to the inclusion of host-derived factors, including polyamines,
328 that might influence pathogen membrane charge and antimicrobial susceptibility.

329

330 **Materials and methods**

331 **Bacteria and culture conditions**

332 *P. aeruginosa* Liverpool Epidemic Strain (LES)B65, LESB65 Δ *pmrB*, PAO1 and PAO1 Δ *pmrB*
333 were used throughout. Deletion of *pmrB* in LESB65 and PAO1 was performed as part of a
334 previous study (23). Bacterial stocks were stored at -80°C in 15% (v/v) glycerol. Prior to
335 experiments, isolates were streaked onto Mueller Hinton (MH) agar and then liquid cultures
336 were prepared in MH broth, unless otherwise stated, from a single colony, and incubated at
337 37°C in a shaking incubator (180 rpm).

338

339 **Sputum samples**

340 Sputum samples were collected from people with CF at the Adult CF centre (Liverpool Heart
341 and Chest Hospital) during periods of stable infection and periods of exacerbation in
342 accordance with ethical approval (IRAS:216408, ethics reference no: 17/NW/0091). Samples
343 were expectorated between 2017 and 2020, and stored at -80°C within 2 h of production.
344 Exacerbation definitions were physician-based, using a clear set of criteria (drop in FEV₁ of
345 generally 10%, increased sputum production and discoloration, increased temperature
346 measured on more than one occasion (>38°C), increased cough and dyspnoea, malaise,
347 lethargy, fatigue, poor appetite, and poor exercise tolerance). Treatment was dual therapy
348 intravenous antibiotics, prescribed by specialist physicians on an individual patient basis.

349

350 **Chemicals and reagents**

351 Antibiotics and spermidine were purchased from Sigma-Aldrich (Sigma, UK). Stock solutions
352 were prepared using DEPC water and filtered through 0.22 μ m syringe filter.

353

354 **LESB65 infection of mice**

355 All infections were performed at the University of Liverpool with prior approval by the UK
356 Home Office and the University of Liverpool Ethics Committee. Female BALB/c mice of 6-8

357 weeks of age (Charles River, UK) were used for infection experiments and housed in
358 individually ventilated cages. Mice were acclimatised for one week prior to infection. Mice
359 were randomly assigned to an experimental group on arrival at the unit by staff with no role
360 in study design. For infection, 2×10^6 colony forming units of mid-exponential growth *P.*
361 *aeruginosa* were instilled into the nares of mice that had been lightly anaesthetised with a
362 mixture of isoflurane and oxygen. At 6 and 24 hours post-infection, mice were intranasally
363 administered a 50 μ l dose of 400 μ g/ml colistin in PBS or else PBS only for control animals,
364 under light anaesthesia. Following this, cage labels were reversed to blind researchers to
365 experimental groups. Mice were culled at 48 hours post-infection and upper airway (sinus
366 and nasopharynx) tissue and lungs were removed post-mortem and homogenised in 3 ml
367 PBS using an IKA T10 handheld tissue homogeniser (IKA, USA). Homogenates were serially
368 dilution onto *Pseudomonas* selective agar (Oxoid, UK) for enumeration of infectious burden.
369 Following enumeration, researchers were unblinded. No animals were excluded from
370 analysis.

371

372 **Spermidine and Spermine ELISA**

373 Spermidine was quantified from lysates of overnight *P. aeruginosa* cultures, from mouse
374 upper airway (sinus and nasopharynx) and lower airway (lung) tissue homogenates, and from
375 CF sputum. Bacterial lysates were prepared by sonication. Overnight cultures were pelleted
376 by centrifugation and resuspended in 1 ml PBS prior to sonication. Mouse tissues were
377 dissected at 48 hours post intranasal administration of 2×10^6 colony forming units of mid-
378 log phase LESB65 (infected group) or PBS (control group). Competitive ELISA for spermidine
379 (Abbexa) or spermine (MyBiosource) detection was performed in precoated 96-well plates,
380 according to manufacturer's instructions.

381

382 **Minimum inhibitory concentration assays**

383 MIC assays were performed by broth microdilution. Isolates were first streaked onto fresh
384 MH agar, and then a single colony from each plate was further grown overnight in 5 ml MH
385 broth on an orbital shaker (180 r.p.m) at 37 °C. A fresh dilution in MH broth was made by
386 incubating 200 µl of the overnight culture in 5 ml MH media. A hundred microliters of this
387 culture was incubated in 96 well-plates with 100 µl of 1:2 serially diluted antibiotic in MH broth.
388 After a 24 h static incubation at 37 °C, the OD₆₀₀ was determined to assess bacterial growth.

389

390 **MIC assays with spermidine**


391 MIC assays were performed as above, with the addition of 4 mM spermidine to the assay
392 throughout or using *P. aeruginosa* that had been pre-incubated with 4 mM spermidine. In the
393 latter case, overnight cultures of *P. aeruginosa*, prepared as above, were pelleted by
394 centrifugation and resuspended in 5 ml PBS containing 4 mM spermidine. Tubes were
395 incubated at 37 °C, 180 r.p.m. for 30 minutes and then bacteria were again pelleted and
396 resuspended in MH broth for use in MIC assays.

397

398 **Synthesis of fluorescent spermidine-NBD (nitrobenzoxadiazole)**

399 7-Nitro-2,1,3-benzoxadiazol-4-N⁸-spermidine hydrogen chloride

400 (NBD-N⁸-Spermidine)



401

402 A solution of N¹-N⁴-Bis-Boc-spermidine (50 mg, 0.29 mmol) in acetonitrile (2.0 ml) was added
403 to 4-chloro-7-nitrobenzofurazan (58 mg, 0.29 mmol) and caesium carbonate (94 mg, 0.29
404 mmol) and heated to 80 °C under reflux conditions for 40 minutes. The reaction mixture was
405 concentrated *in vacuo*, and purified through a silica plug, flushed with EtOAc to elute the boc

406 protected fluorophore. The compound was then dissolved in a solution of 4M HCl in dioxane
407 (5 mL) and stirred for 3 hours at room temperature, before removing the solvent *in vacuo*.
408 The crude mixture was dissolved in DCM was and extracted with water. The aqueous layer
409 was lyophilised to give NBD-N⁸-Spermidine as a hydrochloride salt (41) as a red, amorphous
410 solid and a global yield of 74%, after both the addition of the fluorophore and the subsequent
411 removal of the boc protection group. δ_H (500 MHz, D₂O) 8.48 (1H, d, J 9.0, Ar 6-H), 6.35 (1H,
412 d, J 9.0, Ar 5-H), 3.66 (2H, broad s, 2-H₂), 3.17 (4H, m, 5- and 7-H₂), 3.11 (2H, appt t, J 7.9,
413 9-H₂), 2.08 (2H, apt tt, J 8, 2.4, 8-H₂), 1.88 (4H, m, 3 and 4-H₂). δ_C (125 MHz, D₂O) 147.2
414 (Ar 4-C), 145.0 (Ar 7a-C), 144.7 (Ar 3a-C), 139.8 (Ar 5-C), 100.6 (Ar 6-C), 48.0 (5-C), 45.1
415 (7-C), 43.6 (2-C), 37.2 (9-C), 25.3 (3-C), 24.4 (8-C), 23.8 (4-C). HRMS (ESI): C₁₃H₂₀N₆O₃
416 requires [M+H]⁺, calculated 309.1675 , found 309.1673.

417

418 **Treatment of CF sputum for use in flow cytometry**

419 CF sputum was treated as follows, prior to use in spermidine binding assays; 1 ml of sputum
420 was irradiated with Ultra-Violet light for 20 mins before minimal dilution in polyamine-free PBS
421 and vigorous vortexing, as described by Devereux *et al.*, (2015) (42). Treated CF-sputum
422 was split into 2 ml aliquots and stored at -80 °C until required.

423

424 **Flow cytometry analysis of spermidine-*P. aeruginosa* interactions**

425 *P. aeruginosa* were pre-incubated with 4 mM spermidine-NBD or unlabelled spermidine,
426 following the same protocol as that used for MIC assays. Following incubation, bacteria were
427 pelleted by centrifugation and resuspended in either PBS or CF sputum. Immediately, and at
428 30, 60, 120,180 and 240 minutes, 200 μ l samples were removed from the cultures and
429 analysed for NBD fluorescence on a FACS Aria II flow cytometer (BD Biosciences). Twenty
430 thousand individual bacteria were recorded. Side-scatter and forward-scatter limits for
431 bacterial flow cytometry were pre-determined using *P. aeruginosa* stained with the DNA dye
432 thiazole orange (Sigma Aldrich).

433

434 **Biofilm assay**

435 Biofilm experiments were set up using overnight cultures. Each culture was first diluted 1:100
436 in fresh broth, then added to 96 well plates and incubated for 48 hours at 37° C under static
437 conditions. Plates were washed twice with PBS and stained with 200 µL of a 0.25% solution
438 of crystal violet in water. After incubating at room temperature for 15 minutes, the plates were
439 rinsed twice with water and allowed to dry for 24 hours. The stain was then dissolved in 1 mL
440 of 25% acetic acid in water and incubated at room temperature for 2 minutes. Biofilm
441 formation was quantified by measuring the optical density of this final solution at 590 nm. To
442 test the effect of spermidine on biofilm formation in the presence of colistin, cultures were set
443 up as follows; a 1:100 dilution of each strain was added to 96 well plates with half of the wells
444 supplemented with 4 mM of spermidine. Plates were then incubated for 24 hours. After 24
445 hours, 1:2 serially diluted suspensions of colistin were added to both spermidine (+) and
446 spermidine (-) biofilms. Plates were incubated for a further 24 hours before being processed
447 as above.

448

449 **Statistics**

450 Data analysis was carried out using GraphPad Prism v.8.02 and JMP version 14.0. Data were
451 tested for normality. One-way or two-way ANOVA was used for comparison between groups,
452 and post-hoc analysis included correction for multiple comparisons (as stated in the figure
453 legend). Significance was determined from label free proteomics data using Progenesis Q1.
454 A P value less than 0.05 was considered significant.

455

456 **Study Approval**

457 Ethical approval for collection of CF sputum was obtained from the North West Research
458 Ethics Committee (IRAS:216408, ethics reference no: 17/NW/0091). Written informed
459 consent was obtained from all study participants, prior to enrolment. Ethical approval for

460 animal studies was obtained from the UK Home Office (project licence PP2072053) and the
461 University of Liverpool Animal Welfare Ethical Review Board.

462

463 **Author Contributions**

464 CW, AK, MW, DN and JF designed the study, contributed resources and reagents and
465 supervised staff. CH, SP, AG, AC, TJ and DN performed experiments. CH, SP, AG, AC, MW,
466 DN, JF analysed data. DN, JF wrote the manuscript, with input from all authors.

467

468 **Acknowledgements**

469 This work was funded by an Action Medical Research and Cystic Fibrosis Trust grant
470 awarded to JF, DN, CW and AK (Grant number GN2444). AC was supported by an MRC
471 Doctoral Training Partnership studentship. SP, AG and DN were supported by a Wellcome
472 and Royal Society Sir Henry Dale Fellowship awarded to DN (Grant Number 204457/Z/16/Z).
473 JF was supported by a Medical Research Foundation Fellowship (Grant number MRF-091-
474 0006-RG-FOTHE). JW was supported by a Leverhulme Trust Research Grant (Grant number
475 RPG-2018-030). JF, DN, CW and AK received additional support from the PIPE-CF Strategic
476 Research Centre Grant (SRC022), jointly funded by the Cystic Fibrosis Trust and the Cystic
477 Fibrosis Foundation.

478

479 **References**


- 480 1. Rajan S, and Saiman L. Pulmonary infections in patients with cystic fibrosis. *Semin*
481 *Respir Infect.* 2002;17(1):47-56.
- 482 2. Sadikot RT, Blackwell TS, Christman JW, and Prince AS. Pathogen-host interactions
483 in *Pseudomonas aeruginosa* pneumonia. *Am J Respir Crit Care Med.*
484 2005;171(11):1209-23.
- 485 3. Lyczak JB, Cannon CL, and Pier GB. Lung infections associated with cystic fibrosis.
486 *Clin Microbiol Rev.* 2002;15(2):194-222.
- 487 4. Langendonk RF, Neill DR, and Fothergill JL. The Building Blocks of Antimicrobial
488 Resistance in *Pseudomonas aeruginosa*: Implications for Current Resistance-
489 Breaking Therapies. *Front Cell Infect Microbiol.* 2021;11:665759.

- 490 5. Govan JR, and Deretic V. Microbial pathogenesis in cystic fibrosis: mucoid
491 *Pseudomonas aeruginosa* and *Burkholderia cepacia*. *Microbiol Rev.* 1996;60(3):539-
492 74.
- 493 6. Johnson L, Mulcahy H, Kanevets U, Shi Y, and Lewenza S. Surface-localized
494 spermidine protects the *Pseudomonas aeruginosa* outer membrane from antibiotic
495 treatment and oxidative stress. *J Bacteriol.* 2012;194(4):813-26.
- 496 7. Gunn JS. Bacterial modification of LPS and resistance to antimicrobial peptides. *J*
497 *Endotoxin Res.* 2001;7(1):57-62.
- 498 8. Ernst RK, Moskowitz SM, Emerson JC, Kraig GM, Adams KN, Harvey MD, et al.
499 Unique lipid A modifications in *Pseudomonas aeruginosa* isolated from the airways of
500 patients with cystic fibrosis. *J Infect Dis.* 2007;196(7):1088-92.
- 501 9. Raetz CR, Reynolds CM, Trent MS, and Bishop RE. Lipid A modification systems in
502 gram-negative bacteria. *Annu Rev Biochem.* 2007;76:295-329.
- 503 10. McPhee JB, Lewenza S, and Hancock RE. Cationic antimicrobial peptides activate a
504 two-component regulatory system, PmrA-PmrB, that regulates resistance to
505 polymyxin B and cationic antimicrobial peptides in *Pseudomonas aeruginosa*. *Mol*
506 *Microbiol.* 2003;50(1):205-17.
- 507 11. McPhee JB, Bains M, Winsor G, Lewenza S, Kwasnicka A, Brazas MD, et al.
508 Contribution of the PhoP-PhoQ and PmrA-PmrB two-component regulatory systems
509 to Mg²⁺-induced gene regulation in *Pseudomonas aeruginosa*. *J Bacteriol.*
510 2006;188(11):3995-4006.
- 511 12. Mulcahy H, Charron-Mazenod L, and Lewenza S. Extracellular DNA chelates cations
512 and induces antibiotic resistance in *Pseudomonas aeruginosa* biofilms. *PLoS*
513 *Pathog.* 2008;4(11):e1000213.
- 514 13. Moskowitz SM, Brannon MK, Dasgupta N, Pier M, Sgambati N, Miller AK, et al.
515 PmrB mutations promote polymyxin resistance of *Pseudomonas aeruginosa* isolated
516 from colistin-treated cystic fibrosis patients. *Antimicrob Agents Chemother.*
517 2012;56(2):1019-30.
- 518 14. Moskowitz SM, Ernst RK, and Miller SI. PmrAB, a two-component regulatory system
519 of *Pseudomonas aeruginosa* that modulates resistance to cationic antimicrobial
520 peptides and addition of aminoarabinose to lipid A. *J Bacteriol.* 2004;186(2):575-9.
- 521 15. Han ML, Zhu Y, Creek DJ, Lin YW, Anderson D, Shen HH, et al. Alterations of
522 Metabolic and Lipid Profiles in Polymyxin-Resistant *Pseudomonas aeruginosa*.
523 *Antimicrob Agents Chemother.* 2018;62(6).
- 524 16. Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, and Brinkman FS. Enhanced
525 annotations and features for comparing thousands of *Pseudomonas* genomes in the
526 *Pseudomonas* genome database. *Nucleic Acids Res.* 2016;44(D1):D646-53.
- 527 17. Shah P, and Swiatlo E. A multifaceted role for polyamines in bacterial pathogens.
528 *Mol Microbiol.* 2008;68(1):4-16.
- 529 18. Chou HT, Li JY, Peng YC, and Lu CD. Molecular characterization of PauR and its
530 role in control of putrescine and cadaverine catabolism through the gamma-
531 glutamylation pathway in *Pseudomonas aeruginosa* PAO1. *J Bacteriol.*
532 2013;195(17):3906-13.
- 533 19. Kwon DH, and Lu CD. Polyamine effects on antibiotic susceptibility in bacteria.
534 *Antimicrob Agents Chemother.* 2007;51(6):2070-7.
- 535 20. Grasemann H, Shehnaz D, Enomoto M, Leadley M, Belik J, and Ratjen F. L-
536 ornithine derived polyamines in cystic fibrosis airways. *PLoS One.*
537 2012;7(10):e46618.
- 538 21. Twomey KB, Alston M, An SQ, O'Connell OJ, McCarthy Y, Swarbreck D, et al.
539 Microbiota and metabolite profiling reveal specific alterations in bacterial community
540 structure and environment in the cystic fibrosis airway during exacerbation. *PLoS*
541 *One.* 2013;8(12):e82432.

- 542 22. Fothergill JL, Neill DR, Loman N, Winstanley C, and Kadioglu A. *Pseudomonas*
543 *aeruginosa* adaptation in the nasopharyngeal reservoir leads to migration and
544 persistence in the lungs. *Nat Commun.* 2014;5:4780.
- 545 23. Bricio-Moreno L, Sheridan VH, Goodhead I, Armstrong S, Wong JKL, Waters EM, et
546 al. Evolutionary trade-offs associated with loss of PmrB function in host-adapted
547 *Pseudomonas aeruginosa*. *Nat Commun.* 2018;9(1):2635.
- 548 24. Bolard A, Schniederjans M, Haussler S, Triponney P, Valot B, Plesiat P, et al.
549 Production of Norspermidine Contributes to Aminoglycoside Resistance in pmrAB
550 Mutants of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother.* 2019;63(10).
- 551 25. Karatan E, and Michael AJ. A wider role for polyamines in biofilm formation.
552 *Biotechnol Lett.* 2013;35(11):1715-7.
- 553 26. McGinnis MW, Parker ZM, Walter NE, Rutkovsky AC, Cartaya-Marin C, and Karatan
554 E. Spermidine regulates *Vibrio cholerae* biofilm formation via transport and signaling
555 pathways. *FEMS Microbiol Lett.* 2009;299(2):166-74.
- 556 27. Goytia M, Dhulipala VL, and Shafer WM. Spermine impairs biofilm formation by
557 *Neisseria gonorrhoeae*. *FEMS Microbiol Lett.* 2013;343(1):64-9.
- 558 28. Kang YP, Lee SB, Lee JM, Kim HM, Hong JY, Lee WJ, et al. Metabolic Profiling
559 Regarding Pathogenesis of Idiopathic Pulmonary Fibrosis. *J Proteome Res.*
560 2016;15(5):1717-24.
- 561 29. Holz O, DeLuca DS, Roepcke S, Illig T, Weinberger KM, Schudt C, et al. Smokers
562 with COPD Show a Shift in Energy and Nitrogen Metabolism at Rest and During
563 Exercise. *Int J Chron Obstruct Pulmon Dis.* 2020;15:1-13.
- 564 30. O'Callaghan RJ, Girgis DO, Dajcs JJ, and Sloop GD. Host defense against bacterial
565 keratitis. *Ocul Immunol Inflamm.* 2003;11(3):171-81.
- 566 31. Hume EB, Dajcs JJ, Moreau JM, Sloop GD, Willcox MD, and O'Callaghan RJ.
567 *Staphylococcus corneal* virulence in a new topical model of infection. *Invest*
568 *Ophthalmol Vis Sci.* 2001;42(12):2904-8.
- 569 32. Huang J, Li C, Song J, Velkov T, Wang L, Zhu Y, et al. Regulating polymyxin
570 resistance in Gram-negative bacteria: roles of two-component systems PhoPQ and
571 PmrAB. *Future Microbiol.* 2020;15:445-59.
- 572 33. Moore MP, Lamont IL, Williams D, Paterson S, Kukavica-Ibrulj I, Tucker NP, et al.
573 Transmission, adaptation and geographical spread of the *Pseudomonas aeruginosa*
574 Liverpool epidemic strain. *Microb Genom.* 2021;7(3).
- 575 34. Felgner S, Preusse M, Beutling U, Stahnke S, Pawar V, Rohde M, et al. Host-
576 induced spermidine production in motile *Pseudomonas aeruginosa* triggers
577 phagocytic uptake. *Elife.* 2020;9.
- 578 35. Durand JM, and Bjork GR. Putrescine or a combination of methionine and arginine
579 restores virulence gene expression in a tRNA modification-deficient mutant of
580 *Shigella flexneri*: a possible role in adaptation of virulence. *Mol Microbiol.*
581 2003;47(2):519-27.
- 582 36. Zhou L, Wang J, and Zhang LH. Modulation of bacterial Type III secretion system by
583 a spermidine transporter dependent signaling pathway. *PLoS One.*
584 2007;2(12):e1291.
- 585 37. Delcour AH. Outer membrane permeability and antibiotic resistance. *Biochim*
586 *Biophys Acta.* 2009;1794(5):808-16.
- 587 38. Bonfield TL. Preclinical Modeling for Therapeutic Development in Cystic Fibrosis. *Am*
588 *J Respir Crit Care Med.* 2020;201(3):267-8.
- 589 39. Moyne O, Castelli F, Bicout DJ, Bocard J, Camara B, Cournoyer B, et al.
590 Metatypes of *Pseudomonas aeruginosa* Correlate with Antibiotic Resistance,
591 Virulence and Clinical Outcome in Cystic Fibrosis Chronic Infections. *Metabolites.*
592 2021;11(2).

- 593 40. Cornforth DM, Diggle FL, Melvin JA, Bomberger JM, and Whiteley M. Quantitative
594 Framework for Model Evaluation in Microbiology Research Using *Pseudomonas*
595 *aeruginosa* and Cystic Fibrosis Infection as a Test Case. *mBio*. 2020;11(1).
- 596 41. Jagu E, Pomel S, Pethe S, Cintrat JC, Loiseau PM, and Labruere R. Spermine-NBD
597 as fluorescent probe for studies of the polyamine transport system in *Leishmania*
598 *donovani*. *Bioorg Med Chem Lett*. 2019;29(14):1710-3.
- 599 42. Devereux G, Fraser-Pitt D, Robertson J, Devlin E, Mercer D, and O'Neil D.
600 Cysteamine as a Future Intervention in Cystic Fibrosis Against Current and
601 Emerging Pathogens: A Patient-based ex vivo Study Confirming its Antimicrobial and
602 Mucoactive Potential in Sputum. *EBioMedicine*. 2015;2(10):1507-12.
603

604




605

606 **Figure 1. Difference in in vitro and in vivo antimicrobial susceptibility in PmrB-deficient**
 607 ***P. aeruginosa*. LESB65 or LESB65 Δ pmrB colony forming units (CFU) in (A) sinuses and (B)**
 608 **lungs at 48 hours post-infection. Mice were intranasally infected with 2×10^6 CFU *P.***
 609 ***aeruginosa*, at 6 and 24 hours post-infection, mice were intranasally administered a 50 μ l**
 610 **dose of 20 μ g colistin or PBS control. Each circle represents an individual mouse and p values**
 611 **were determined by two-way ANOVA with Bonferroni correction. Data are representative of**
 612 **two independent experiments, n=8 for each treatment group.**

613

614



615

616 **Figure 2. Loss of PmrB influences the relative abundance of polyamine synthesis and**

617 **utilisation proteins in *P. aeruginosa*.** Abundance of proteins of **(A)** the spermidine

618 synthesis operon and **(B)** the polyamine binding, uptake and utilisation operon in

619 LESB65ΔpmrB and LESB65pmrB^{L255Q} relative to LESB65. Values are fold change vs

620 LESB65 and a negative value denotes decreased abundance. Bars show mean +/- SEM (n


621 = 5 per group) and significance was determined from label free proteomics data using

622 Progenesis QI. NS = adjusted p value > 0.05, * = p < 0.05, ** = p < 0.01, *** = p < 0.001, ****

623 = p < 0.0001, ND = not detected. Block arrows denote gene position and orientation, line

624 arrows show transcriptional start site and directionality. Source data can be found in Bricio-

625 Moreno *et al.* (23), Supplementary Dataset 2.



626

627 **Figure 3. Spermidine is abundant in the murine respiratory tract and bioavailability**

628 **increases during *P. aeruginosa* infection.** Concentration of spermidine in the sinuses and


629 lungs of mice at 48 hours post intranasal administration of PBS (-) or LESB65 (+). Spermidine

630 was measured by ELISA and each square represents a tissue sample from an individual

631 mouse. Significance was determined by two-way ANOVA with Bonferroni correction. Data

632 are from a single experiment, n=5 for each treatment group.

633



634

635

Figure 4. Spermidine is detectable in CF sputum. Concentration of spermidine in sputum

636

from people with CF, chronically infected with *P. aeruginosa*, determined by competition

637

ELISA. **(A)** Spermidine levels in 19 study participants. Between 1 and 13 samples were

638

available per participant. No two samples from the same participant were collected at the

639

same visit (n=65). **(B)** Changes in spermidine abundance in sputum from a single participant

640

over time. This participant received meropenem during the treatment phases and ceftazidime

641

during exacerbation and recovery/maintenance. The individual was co-infected with MRSA

642

and so received teicoplanin throughout. **(C)** Collected samples were defined as baseline,

643

treatment or pulmonary exacerbation, defined by participant clinical data. Participants


644

received two or more of aztreonam, fosfomycin, colistimethate, meropenem, tobramycin,

645 colomycin, ceftazidime, teicoplanin and tazocin, with individual patient antibiotic
646 combinations chosen by their specialist clinician. Whiskers show 10-90 percentile.
647 Significance was determined by one-way ANOVA with Dunnett's multiple comparison testing.
648 Data are from a single experiment.

649

650



652

653 **Figure 5. Prolonged interaction with environmental spermidine in PmrB-deficient *P.***654 ***aeruginosa*. LESB65 (A) and LESB65 Δ pmrB (B) from mid-log cultures were incubated for**655 30 minutes with unlabelled spermidine (blue histogram) or spermidine-NBD (all other
656 histograms), then pelleted, washed in saline and resuspended in polyamine-free PBS.657 Spermidine-NBD binding to *P. aeruginosa* was determined by flow cytometry at 0, 30, 60,

658 120, 180 and 240 minutes after co-incubation. Data are representative of two independent


659 experiments with n = 5 samples per group. **(C)** Spermidine was co-incubated with LESB65660 (red bars) or LESB65 Δ pmrB (blue bars) for 30 minutes in PBS or in CF sputum. Bacteria

661 were the pelleted, washed in saline and resuspended in PBS or CF sputum. Fluorescence

662 was determined after 30 minutes by flow cytometry. P-values are from two-way ANOVA with
663 Šidák's multiple comparison test. Data are representative of two independent experiments
664 with n = 4 samples per group. **(D)** Representative flow cytometry histograms of LESB65 and
665 LESB65 Δ *pmrB* following 30 minutes in the presence (solid lines) or absence (dashed lines)
666 of spermidine-NBD.

667

668



669

670 **Figure 6. Spermidine promotes biofilm production in PmrB-deficient *P. aeruginosa*.**

671 Surface-attached biofilm production was quantified by crystal violet staining. **(A)** Biofilm mass

672 after 48 hours culture of LESB65 and LESB65 $\Delta pmrB$. **(B)** Fold-change in biofilm production

673 vs the no spermidine control for LESB65 and LESB65 $\Delta pmrB$. **(C)** Biofilm formation, relative

674 to untreated controls, in presence of 4 mM spermidine, 8 g/ml colistin, or both spermidine

675 and colistin. Lines represent the mean and error bars are standard deviation. Data are

676 representative of four **(A, B)** or three **(C)** independent experiments, with $n=15$ **(A)** or $n=18$ **(B)**

677 for each treatment group. In **(C)** each data point represents a biological replicate that is the

678 mean of five technical replicates. P values are from two-tailed paired t tests **(A, B)** or two-

679 way ANOVA with Tukey's multiple comparison test **(C)**.

680

681 **Table 1. Presence of spermidine reduces antimicrobial susceptibility of PmrB-deficient**
682 **but not wild type *P. aeruginosa*.** The concentration of colistin required to inhibit 50 percent
683 of growth (minimum inhibitory concentration [MIC]₅₀) was determined for LESB65, PAO1
684 and their PmrB-deficient derivatives. Assays were conducted without spermidine, in the
685 presence of 4 mM spermidine, or with bacteria that had been pre-incubated for 30 minutes
686 with 4 mM spermidine and then washed in PBS before addition of antibiotics. Data shown
687 are the median and range of MIC₅₀ values from 5 independent experiments.

Strain	Spermidine throughout	Preincubation with spermidine	MIC ₅₀ µg/ml [Median (range)]
LESB65	-	-	2 (2-4)
	+	-	2 (2-4)
	-	+	2 (1-4)
LESB65Δ <i>pmrB</i>	-	-	0.25 (0.125-0.5)
	+	-	2 (0.5-4)
	-	+	1 (1)
PAO1	-	-	0.5 (0.5)
	+	-	1 (0.5-1)
	-	+	0.5 (0.25-0.5)
PAO1Δ <i>pmrB</i>	-	-	0.125 (0.125-0.25)
	+	-	2 (0.5-2)
	-	+	1 (0.5-2)

688

689