









Assessment of physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health – a systematic review

Lidia Romero-Gallardo ^{1,2} Olga Roldan Reoyo ^{3,4} Jose Castro-Piñero ^{5,6}
Linda E May ^{7,8} Olga Ocón-Hernández ^{9,10} Michelle F Mottola ¹¹
Virginia A Aparicio ^{2,12} Alberto Soriano-Maldonado ^{13,14}

To cite: Romero-Gallardo L, Roldan Reoyo O, Castro-Piñero J, *et al.* Assessment of physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health – a systematic review. *BMJ Open Sport & Exercise Medicine* 2022;**8**:e001318. doi:10.1136/bmjsem-2022-001318

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2022-001318>).

ORR and JC-P contributed equally.

Accepted 2 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Lidia Romero-Gallardo;
lidiaromerogallardo@gmail.com

ABSTRACT

Objectives To systematically review studies evaluating one or more components of physical fitness (PF) in pregnant women, to answer two research questions: (1) What tests have been employed to assess PF in pregnant women? and (2) What is the validity and reliability of these tests and their relationship with maternal and neonatal health?

Design A systematic review.

Data sources PubMed and Web of Science.

Eligibility criteria Original English or Spanish full-text articles in a group of healthy pregnant women which at least one component of PF was assessed (field based or laboratory tests).

Results A total of 149 articles containing a sum of 191 fitness tests were included. Among the 191 fitness tests, 99 (ie, 52%) assessed cardiorespiratory fitness through 75 different protocols, 28 (15%) assessed muscular fitness through 16 different protocols, 14 (7%) assessed flexibility through 13 different protocols, 45 (24%) assessed balance through 40 different protocols, 2 assessed speed with the same protocol and 3 were multidimensional tests using one protocol. A total of 19 articles with 23 tests (13%) assessed either validity (n=4), reliability (n=6) or the relationship of PF with maternal and neonatal health (n=16).

Conclusion Physical fitness has been assessed through a wide variety of protocols, mostly lacking validity and reliability data, and no consensus exists on the most suitable fitness tests to be performed during pregnancy.

PROSPERO registration number CRD42018117554.

BACKGROUND

Physical fitness (PF) has been defined as the ability to carry out daily tasks with vigour and alertness, without undue fatigue and with ample energy to enjoy leisure-time pursuits and meet unforeseen emergencies.^{1,2} PF is considered a powerful marker of health that is associated with a lower risk of cardiovascular events, cancer and all-cause mortality in all

WHAT IS ALREADY KNOWNWHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ The assessment of physical fitness during pregnancy requires special considerations to preserve fetal and maternal health.
- ⇒ Although physical fitness during pregnancy has been assessed inconsistently across studies, these tests have not been systematically compiled to date.
- ⇒ The validity and reliability of the variety of tests used to assess physical fitness during pregnancy has not been comprehensively reviewed.

WHAT THIS STUDY ADDS

- ⇒ During pregnancy, physical fitness including cardiorespiratory fitness, muscular strength, flexibility and balance have been assessed inconsistently, using a wide variety of protocols.
- ⇒ Most of the tests used to assess physical fitness during pregnancy lack validity and reliability data.
- ⇒ Higher physical fitness might be associated with better maternal and neonatal health, although further research is needed.

HOW THIS STUDY MAY AFFECT RESEARCH, PRACTICE AND POLICY

- ⇒ The extent to which the data derived from current physical fitness tests during pregnancy is valid and reliable is still unclear and, therefore, should be interpreted with caution.
- ⇒ Developing a battery of fitness tests to assess the different fitness components during pregnancy must be set as a priority for relevant institutions.
- ⇒ An expert consensus to develop a battery of physical fitness tests is recommended.

ages.^{3–7} In pregnant individuals, some studies have recently highlighted the potential impact of PF on maternal and fetal health.^{8–15} Low PF levels are associated with low infant birth weight,⁸ increased risk of gestational diabetes mellitus,^{9,10} poor postpartum recovery¹¹ and



worse delivery outcomes.^{12 13} Moreover, the anatomical, biomechanical, physiological and psychological changes during the pregnancy might compromise PF levels.^{16–18} Consequently, it is of clinical and public health interest to assess PF during the pregnancy, and to understand which available tests are best to assess PF during this critical period of life.

Two categories of PF components have been defined as follows: (1) health-related components (cardiorespiratory fitness (CRF), muscular fitness, muscular endurance and flexibility) and (2) skill-related components (ability, coordination, balance, power, reaction time and speed).¹² These PF components can be assessed subjectively through questionnaires,¹⁵ objectively and accurately through laboratory tests and efficiently, economically and easily through field-based tests. During the pregnancy, a wide variety of fitness tests have been used to assess PF, although a compilation of these tests has not been published to date. Compiling all fitness tests performed in pregnant women would help practitioners to select the most useful test according to their purpose. It is also important to note that, although laboratory tests are generally the gold standard for assessing PF, these tests are not accessible to everyone because they need sophisticated and expensive equipment, and it is not possible to evaluate a relatively large sample in a short period of time. As an alternative, a number of field tests exist that provide an opportunity to assess PF in a more accessible way.² However, there is no consensus on which fitness tests should be used to assess PF in pregnant individuals, and the validity and reliability of many of the tests used to assess PF during the pregnancy are unknown.¹⁹

Since the assessment of PF in pregnancy requires special consideration to preserve fetal and maternal health,^{18 20 21} understanding which fitness tests are valid, reliable, and associated with maternal and neonatal health outcomes, would provide a framework for improving PF assessment during pregnancy and also for improving exercise prescription in this population.

The aims of this systematic review were to: (1) describe which fitness tests have been used to evaluate PF in pregnant individuals; and (2) to evaluate the validity and reliability of the fitness tests, and their relationship with maternal and neonatal health.

METHODS

Registration and review guidelines and checklist

This systematic review was prospectively registered at PROSPERO (CRD42018117554; available at <http://www.t.ly/fS6a>). In addition, the review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines²² and the PRISMA checklist²³ is included as online supplemental material 1, table S1. (1) .

Search strategy

Articles were searched by two independent reviewers from two major databases, MEDLINE (PubMed) and the

Web of Science (WOS) from inception to January 2021. For the search strategy undertaken in PubMed Medical Subject Heading, (MeSH) terms were used. All terms were combined using the connector OR for similar criteria the connector 'AND' was used to combine population group (ie, pregnant women), to delimit date of publication ('0001/01/01'(PDat): '2021/01/15'(PDat)), to include full text papers, and to include studies performed in humans.

A similar search strategy and term combination was undertaken in the WoS (online supplemental material 2, table S2), although MeSH terms and its appropriate terms connection were not used as they are exclusive for PubMed. The complete search strategy and further details are presented in online supplemental material 2, tables S1 and S2.

Inclusion criteria

The inclusion criteria were as follows: (1) healthy pregnant individuals (no restriction regarding gestational week); (2) at least one component of PF assessed either through field based or laboratory tests; (3) access to full text; (4) only one original article from the same study/project using the same test were included and (5) text in English or Spanish.

Quality assessment of the articles

To assess quality of the articles included in aim 2, three quality scores were applied. To assess validity and reliability, authors adapted two quality scores ad hoc previously used in two different systematic reviews following the same goal as the present review, however, undertaken in different populations.^{24 25} To assess the association of PF with health-related outcomes the Effective Public Health Practice Project was used.²⁶ All procedures are comprehensively described in online supplemental material 3, tables S3–S5.

Process and data extraction

After checking title and abstract, only the studies meeting all inclusion criteria were introduced in a reference manager software (Mendeley). In the event of disagreement between the two independent reviewers concerning the inclusion/exclusion of an article, a consensus was reached (there was no need of a third person). The snowball strategy was also used. Information including reference, age, sample size and fitness test description are summarised in online supplemental material 5, table S6.

RESULTS

A comprehensive PRISMA flow diagram is presented in [figure 1](#).

Overall results, quality assessment and gestational week

The search identified 2617 studies, of which 149 were included ([figure 1](#)). These articles contained 191 fitness tests, using 149 different protocols that were included for Aim 1. A summary of the number of articles that




Figure 1 Flow chart of the literature search and paper selection process.

assessed PF during the pregnancy and the protocols used for its assessment is presented in [figure 2](#). This has been organised based on each of the different PF components assessed in those articles. Moreover, a comprehensive diagram of the fitness tests and the different protocols performed to date, organised by PF component, is presented in [figure 3](#).

Regarding aim 1, 99 tests (including 75 different protocols) were used to assess CRF,^{8 12 13 18 27–108 28} (including 16 different protocols) to assess muscular fitness,^{8 12 13 61 86 109–122} 14 (including 13 different protocols) to assess flexibility,^{12 13 110 114 123–127} 45 tests (including 40 different protocols) to assess balance,^{110 116 128–167} 2 tests

using the same protocol to assess speed^{168 169} and 3 tests using the same protocol were multidimensional.^{168–170} No results were found for other PF components such as agility or coordination.

Regarding aim 2, a total of 19 articles (13% of the total number of articles included) assessed at least validity (n=3) and reliability (n=4) of fitness tests. These articles are summarised in [table 1](#). Of the three articles^{74 75 169} that assessed validity, two articles were classified as low quality^{74 169} and one as high quality.⁷⁵ Of the four articles that assessed reliability criteria, three were considered high quality^{74 117 168} and one low quality.¹²¹ The relationship of PF with maternal and neonatal health outcomes (n=16 tests) are summarised in [table 2](#). Of these 16 tests, 11 were classified as very low quality^{13 57 68 95 108 111 126 157 158} and 5 were classified as low quality.^{8 63 115 128 170}

The gestational week at PF assessment ranged from 8 to 41 across articles. Some articles assessed PF at different time points throughout pregnancy; therefore, we divided pregnancy into two stages. Early pregnancy (ie, from week 0 to week 20 of gestation) and late pregnancy (ie, from week 21 to week 40). Using this approach, 11 articles (7%) were performed in early pregnancy; 57 articles (38%) were performed in late pregnancy; 55 articles (37%) were performed several times (ie, range 2–5 times) throughout pregnancy; 7 (5%) articles specified a range of weeks that included early to late pregnancy; 14 articles (9%) reported only the trimester without specifying gestational week; 4 articles (3%) provided no information and 1 article (1%) assessed PF on the day of labour.

Aim 1: fitness tests used to evaluate PF in pregnant women

Cardiorespiratory fitness

We identified 99 tests assessing CRF, of which 61 (62%) were performed on a cycle ergometer, 25 (25%) on a treadmill, 10 (10%) on a track and 3 (3%) used step




Figure 2 Number of tests and protocols that assessed the different components of physical fitness during pregnancy.




Figure 3 Diagram of the fitness tests and the different protocols organised by PF component. PF, physical fitness.

protocols (figure 3). Of the 99 tests, a total of 75 corresponded to different protocols. For instance, there were 56 different protocols using a cycle ergometer, distributed as follows: only one article used the Arstila test⁶⁸; one used the Bruce Protocol at 75% HR_{max}²⁷; one applied the Modified Bruce ramp protocol at anaerobic threshold¹⁰⁴; two employed the Modified Balke protocol at 70% HR_{max}^{34 41}; one used a YMCA protocol;¹⁰⁶ The remaining of articles (n=55) used ad hoc tests (ie, specifically designed for the purpose of the investigation); 11 of which^{32 37 38 41 45 57 64 79 107} used steady-state tests and 44^{28-31 33 35 36 39-41 43 44 46-56 59-63 65-69 90 100-106 108 171-173} used incremental tests. When analysing the type of test based on intensity, we found that 13 tests were maximal tests,^{31 43 44 47-49 59 60 67 103-105 171} 37 submaximal tests^{29 30 35-40 42 45 46 50-52 54-57 62-66 68 69 79 90 100-102 106 108 172} and 3 used mixed tests^{28 33 41} containing submaximal and maximal stages within the same protocol.

There were 25 treadmill tests that used 14 different protocols, distributed as follows: the Modified Balke protocol was used in 10 articles^{8 31 71 73 75-78 82 96}; the Modified Bruce protocol in 2 articles^{13 97} and the traditional Balke protocol used twice in the same article⁷⁰; the traditional Bruce protocol,⁹⁸ the Cornell protocol,⁷⁴ the SWET protocol and the Ebbelling single-stage protocol¹⁸ were each used in one article. There were seven ad hoc tests of which two were steady-state,^{38 81} and five were incremental tests.^{72 73 80 83 90} According to intensity, three were maximal tests^{73 80 81} and four submaximal tests.^{38 72 83 90}

Of the 10 tests performed on a track, 6 articles used the 6 min walk test protocol,^{84 85 87-89 92} and 4 were ad hoc tests (ie, maximal and 4 were submaximal). With regard

to the three step tests, one Canadian Home Fitness test⁹³ was used and two ad hoc incremental submaximal tests were used.^{94 95}

Muscular fitness

A total of 28 tests (ie, 14% of all included articles) that included 16 different protocols assessed muscular fitness, of which 10 performed maximal hand-grip strength tests,^{8 12 13 86 109-115} performed endurance hand-grip tests, 2 for 3 min^{118 120} and 1 for a 2 min period¹¹⁹ (figure 3). In two of the articles conducting an endurance hand-grip test,^{118 119} a hand-grip sphygmomanometer was used instead of dynamometry. On the other hand, one used a hand-held dynamometer fixed to a chair to assess quadriceps strength¹¹⁶ and one used a toe-grip dynamometer.¹¹⁶ Moreover, two ad hoc isometric tests were used to assess maximal voluntary hip extension and back flexors endurance in the same article.¹⁷⁴ Finally, 13 dynamic endurance tests were found, 9 were listed as ad hoc tests^{12 112 122} and another 3 (30s Chair Stand Test, 5 Times Sit to Stand test, Trendelenburg's test) were classified as 'other' dynamic tests.^{13 112 117}

Flexibility

Our search identified 14 (7%) tests that assessed flexibility using 13 different protocols, including the side bending test,¹²⁵ the sit-and-reach test,¹² the back-scratch test (twice),^{13 110} the motion analysis (ie, including three different tests such as the seated and standing forward flexion, seated and standing side to side flexion and seated axial rotation)¹²³ and an optoelectrical system (ie, performing four different tests).¹²⁷ Goniometry

Table 1 Overview of studies that assessed the validity and/or reliability of fitness tests during pregnancy

Reference (authors, year)	Validity	Reliability	Capacity evaluated, short test description and maternal and neonatal health outcomes or statistical results	Quality score
Cardio-respiratory fitness				
Yeo <i>et al</i> (2005) ⁷⁴	Yes	Yes	Cornell Protocol on treadmill platform. Validity: Bland-Altman plots. The mean difference was 4.4±3.6 mL/kg/min. Data indicated that VO ₂₀₀₀ overestimates VO ₂ by an average of 4.4 mL/kg/min compared with CPX/D. Pearson correlation coefficient between the average and difference of paired measurements was close but not significant (r=0.48; p>0.01). Reliability: Paired t test (t (45)=3.9, p<0.001). Linear regression: y=0.96X-1.6; 95% CI for the slope: 0.94 to 1.1; R ² =0.91, p<0.001	4–8
Mottola <i>et al</i> (2006) ⁷⁵	Yes	No	Modified Balke protocol on treadmill platform. Validity: Pearson Correlation: R ² =0.72, R ² adjusted=0.71 and SEE=2.7 (The prediction equation was compared with cross validation (n=39; p=0.78).	5
Muscular fitness				
Gutke <i>et al</i> (2008) ¹²¹	No	Yes	Maximal voluntary isometric hip extension Reliability: Spearman's r and Intercorrelation coefficient (ICC). Right leg: r=0.82; ICC=0.87. Left leg: r=0.88; ICC=0.85 (both p value no reported).	3
Yenişehir <i>et al</i> (2020) ¹¹⁷	No	Yes	Five Times Sit to Stand Test (5TSS) Reliability: Inter-rater reliability of 5TSS was excellent for subjects with and without pregnancy-related pelvic girdle pain (ICC ¼ 0.999, 95% CI ¼ 0.999 to 1.000; ICC ¼ 0.999, 95% CI ¼ 0.999 to 0.999, respectively). Test-retest reliability of 5TSS was also very high for subjects with and without PGP (ICC ¼ 0.986, 95% CI ¼ 0.959 to 0.995; ICC ¼ 0.828, 95% CI ¼ 0.632 to 0.920, respectively).	5–7
Flexibility				
Lindgren and Kristiansson (2014) ¹²⁶	No	Yes	Ad hoc passive abduction of the left fourth finger. Reliability: Intraindividual coefficient of variance. (1) Between the first and second measurement=0.077; (2) Between the second and third=0.070 and between the third and fourth=0.071.	2
Speed				
Evensen <i>et al</i> (2015) ¹⁶⁸	No	Yes	Ten metres Timed walk Test Reliability: ICC from a one-way random effects model and reporting the 95% CI. Coefficients for test-retest reliability for 10mTWT: (ICC=0.74; 95% CI=0.42 to 0.90; SEM=0.17 m/s; MDC ₉₅ =0.47 m/s) Coefficients for intertester reliability 10mTWT: (ICC=0.94; 95% CI=0.82 to 0.98; SEM=0.09 m/s; MDC ₉₅ =0.25 m/s).	8
Evensen <i>et al</i> (2016) ¹⁶⁹	Yes	No	Ten metres Timed walk Test Validity: Spearman correlation coefficient. Between the 10mTWT and ASLR (r=-0.65, p=0.003). Between the 10mTWT and PGQ (r=-0.25 to -0.56).	3
Multidimensional				
Evensen <i>et al</i> (2015) ¹⁶⁸	No	Yes	Timed Up and Go Test (TUG) Reliability: ICC from a one-way random effects model and reporting the 95% CI. Coefficients for test-retest reliability TUG: (ICC=0.88; 95% CI 0.70 to 0.95; SEM=0.42 s; MDC ₉₅ =1.16 s.) Coefficients for intertester reliability TUG: (ICC=0.95; 95% CI 0.84 to 0.98; SEM=0.36 m/s; MDC ₉₅ =1.00 m/s).	8
Evensen <i>et al</i> (2016) ¹⁶⁹	Yes	No	TUG Validity: Spearman correlation coefficient. Between the TUG and ASLR (r _s =0.73, p=0.001). Between the TUG and ASLR (r _s =0.73, p=0.001). Between the TUG and PGQ (r _s =0.41 to 0.52).	3
ASLR, Active Straight Leg Raised; MDC, minimal detectable change; 10mTWT, Ten-metre Timed Walk Test; PGP, Pelvic girdle pain; SEM, SE of measurement.				

was used in two different articles to measure hamstring flexibility,¹¹⁴ wrist flexion-extension and medial lateral deviation.¹²⁴ Only one article used an ad hoc machine to test passive abduction of the left fourth finger.¹²⁶

Balance

We identified 45 (24%) articles assessing balance of which 19 analysed static balance and 26 used dynamic balance with 40 different protocols. With regard to static balance, 18 were laboratory tests of which

12 assessed balance through stabilometry tests on a force platform,^{129 131 132 138 149 158–160 162–164 167} one on a pressure platform¹⁶³ and another on an Equitest platform.¹⁶⁵ Four articles did not mention the type of platform used.^{117 132 133 175} Regarding protocols, all articles conducted the tests with participants standing with bipedal support. However, standing position varied between articles. Ten articles maintained a standing posture with feet separated,^{116 131 132 147 158 159 162 165–167}

**Table 2** Summary of studies assessing PF and its relationship with maternal and neonatal health outcomes

Health-related outcome	Related to PF					Assoc (-/+)	Statistics	Quality score (0-5)	Unrelated to PF	
	Biblio no								Biblio no	Quality score (0-5)
	CRF	MF	Flexibility	Balance	Multidimensional					
Maternal Health										
Prepregnancy weight	57					-	r=-0.63, P=0.001	2		
Maternal HR at submaximal exercise	108					-	NR, P<0.05	1		
Duration of gestation	63					+	r=0.12, P=0.01	3		
Physical activity practice	95					+	P=0.01	2	108	1
Back pain			126			+	OR=1.09, 95% CI 1.01 to 1.17, P=0.022	2		
Anxiety				158		-	r=0.559, P=0.02	2		
Fall risk				128		-	P<0.0001	3		
				157		-	P<0.001	2		
Pelvic girdle pain					170	+	P<0.001	4		
Birth										
Length of labour in nulliparas	57					-	r=-0.65, P=0.05	2		
Second stage of labour	108					-	NR	1		
Caesarean	13					-	NR, P<0.001	2		
Pain during contractions	111					+	r=0.67, P<0.001	2		
Fetal and neonatal health										
Fetal umbilical artery pH	68					+	NR, P<0.001	2		
			13			+	r=0.220, P<0.05	2		
Asphyxiated babies	108					-	NR, P<0.05	2		
Arterial umbilical cord PO2	13					+	r=0.267, P<0.05	2		
			13			+	r=0.237, P<0.05	2		
Arterial umbilical cord PCO2			13			-	R=0.331, P<0.01	2		
Neonatal birth weight		8				+	r=0.27, P=0.048	3		
		115				+	F (2182)=3.15, P=0.004	4		
		13				+	r=0.191, P<0.05	2	93, 26	2,1
New-born length									93	
New-born head circumference									93	
Apgar Score									93, 26	2,1

Related to PF refers to those variables where authors found either a positive or negative association of the variable with PF levels. Unrelated to PF refers to those variables where authors could not find any association between the variable and PF.
 +, direct association of the variable with PF; -, inverse association of the variable with PF; CRF, cardiorespiratory fitness; HR, heart rate; MF, muscular fitness; NR, not reported; PCO2, pressure of CO2; PF, physical fitness; PO2, pressure of O2.

one with feet together,¹²⁹ two used mixed protocols,^{128 160} one with medial malleoli separated¹³⁰ and four did not mention the standing posture.^{138 149 163 164} Moreover, three articles used protocols with eyes open^{132 149 162} exclusively, eight articles used mixed protocols with eyes open and closed, one used visual target and visual tasks¹⁶⁴ and six did not specify whether participants kept their eyes closed or opened. Only one article used a field test, the one-legged standing protocol.¹¹⁰ On the other hand, one test was a field-test without a platform.

In relation to the 26 articles measuring dynamic balance, 9 assessed balance using platforms. Each of these articles used a different testing tool such as a balance master platform,¹³³ pressure platform,¹⁶³ force platform,¹³⁵ Equitest platform¹³⁴ and a movable platform, which was used in two articles.^{136 137} Two of these articles were walking protocols,^{135 163} one with translational perturbations,¹⁵⁷ one was standing with one knee flexed and arms across the chest.^{136 137} Another 15 articles used three-dimensional (3-D) camera motion capture systems

using 13 different protocols. Twelve of the 15 articles were walking protocols^{139 140 142–144 148 150 152–156 161} and 2 used a stand to sit motion protocol.^{141 151} Moreover, one article used a triaxial accelerometer¹⁴⁶; another article assessed balance through recording (without specification of camera type)¹⁴⁵ and another using instrumented insoles.¹⁷⁶ All three of these articles used walking protocols.

Speed

The only protocol that was used to assess speed during pregnancy was the 10 m timed walk test (10mTWT). However, the same test was identified in two different articles.^{168 169} In the 10mTWT, the participants commenced standing at a chair. When told to start, subjects walked as fast as possible along 14 m marked with white tape placed at 0 m, 2 m, 12 m and 14 m. The time (100th of a second) required to walk between the 2 m and 12 m markers was recorded and converted into speed in metres per second (m/sec).

Agility and coordination

No articles of agility and coordination were identified.

Multidimensional

Our search identified a walking multidimensional test that was used in three studies.^{168–170} In the Timed Up and Go Test (TUG), the participant began seated in a chair with their arms on armrests and their toes against a start line. The purpose was to cross the front white line at 3 m away, turn around and walk back to the chair and sit down as fast as possible. The performance is measured in time (100th of a second).

Aim 2: evaluation of the validity and reliability of the fitness tests, and their relationship with maternal and neonatal health

Articles assessing validity and reliability are summarised in table 1. Articles assessing PF and its relationship with maternal and neonatal health outcomes are presented in table 2 and follows a similar format as Sallis *et al.*¹⁷⁷

Cardiorespiratory fitness

We identified two articles examining validity.^{74 75} Yeo *et al.*⁷⁴ aimed to validate a portable metabolic testing system (VO2000) on healthy sedentary pregnant individuals. The VO2000 consistently overestimated VO₂ measurements, compared with the same manufacturer's reference system, by 4.4±3.6 SD mL/kg/min although the Pearson correlation was significant (r=0.48; p=0.01). When the VO2000 was used twice, the mean difference was statistically significant (1.0±1.8 mL/kg/min; t(45)=3.9, p<0.001). Mottola *et al.*⁷⁵ provided a prediction equation for VO_{2peak} in pregnant individuals between 16 and 22 weeks of gestation, using a modified Balke protocol. The results of this equation revealed an adjusted R² of 0.71 and differences between actual and predicted VO₂ of 2.7 mL/kg/min. When the authors used this equation to predict VO_{2peak} in a cross-validation group (n=39), they

found a predicted value of 23.38±4.03 mL/kg/min, while the actual value was 23.54±5.9 mL/kg/min (p=0.78).

A total of six articles analysed the association of CRF with maternal and neonatal health outcomes. Pomerance *et al.*⁵⁷ observed that VO_{2max} was inversely associated with the length of labour in multiparas (r=-0.65; p=0.001) and prepregnancy weight (r=-0.63; p=0.001). However, VO_{2max} was not correlated with newborn weight, length or head circumference, or with the 1 min Apgar scores (all p>0.05). In the same line, Wong and McKenzie¹⁰⁸ observed that fit mothers showed lower HR at submaximal exercise intensity (p<0.05) and the second stage of labour was shorter (no statistics reported) compared with unfit pregnant mothers. However, there was no difference between fit and unfit in the length of gestation or weight gained (no statistics reported). In the same article, the authors showed neither positive nor negative effects of maternal fitness on newborn weight or Apgar scores.

In addition, Erkkola and Rauramo⁶⁸ found that newborns from fit pregnant individuals had higher pH than fetuses of less physically fit women (p<0.01). In this article, participants with low physical performance were more likely to have asphyxiated neonates than neonates of physically fit women (p<0.05). In the same line, Baena-García *et al.*¹³ observed that maternal CRF at the 16th gestational week was related to higher arterial umbilical cord PO₂ (r=0.267, p<0.05), and those who had caesarean sections had significantly lower CRF compared with those who had vaginal births (p<0.001).

Moreover, Bisson *et al.*⁸ studied the association of CRF in early pregnancy with physical activity before and during early pregnancy. The authors found that a higher VO_{2peak} in early pregnancy was positively associated with physical activity spent at sports and exercise before and during early pregnancy (p<0.001).

Muscular fitness

Only two muscular fitness tests assessed reliability.^{117 121} Yenişehir *et al.*¹¹⁷ analysed reliability and validity of Five Times Sit-to-Stand. Inter-rater reliability was excellent for subjects with and without pelvic girdle pain (PGP) (intra-class correlation coefficient, ICC=0.999, 95% CI 0.999 to 1.000; ICC=0.999, 95% CI 0.999 to 0.999, respectively). Test-retest reliability was also very high for subjects with and without PGP (ICC=0.986, 95% CI 0.959 to 0.995; ICC=0.828, 95% CI 0.632 to 0.920, respectively).

Gutke *et al.*¹²¹ analysed the reliability for an ad hoc test. This test consisted of a maximal voluntary isometric hip extension with a fixed sensor holding a sling around the thigh and pulling for 5 s during 3 reps with 5–10 s of rest (r=0.82 for the right leg and r=0.88 for the left leg; ICC=0.87 for the right leg and 0.85 for the left leg; with p value not reported).

Bisson *et al.*⁸ observed that hand-grip strength was positively associated with infant birth weight (r=0.34, p=0.0068) even after adjustment for confounders (r=0.27, p=0.0480). Żelazniewicz and Pawłowski *et al.*¹¹⁵ observed that hand-grip strength was associated with offspring

birth weight when controlled for the newborn sex and gestational age at delivery ($F(2,182)=3.15$; $p=0.04$). Baena-García *et al*¹³ found greater hand-grip strength weakly associated with greater neonatal birth weight ($r=0.191$, $p<0.05$). Wickboldt *et al*¹¹¹ found that hand-grip strength was moderately correlated with pain scores, where the mean hand-grip strength during contractions had the highest correlation coefficient ($r=0.67$; $p<0.001$) compared with peak hand-grip strength ($r=0.56$; $p<0.001$) and the area under the curve of hand-grip force ($r=0.55$; $p<0.001$).

Flexibility

Lindgren and Kristiansson¹²⁶ designed an ad hoc machine to test passive abduction of the left fourth finger and its relationship with low-back pain during pregnancy and early postpartum. Abduction angle was measured at three different times throughout the pregnancy and once in the postnatal period. Reliability of the abduction angle was analysed by the intraindividual coefficient of variance. The coefficients of variance between the first and second measurement was 0.077, between the second and third 0.070 and between the third and fourth 0.071.

Only two flexibility tests evaluated associations with maternal and neonatal health outcomes. Lindgren and Kristiansson¹²⁶ found that women with greater passive abduction angle of the left fourth finger was associated with the highest back pain incidence (OR 1.09; 95% CI 1.01 to 1.17; $p=0.022$) and the highest number of previous pregnancies (OR 3.24; 95% CI 1.57 to 6.68; $p=0.002$). Baena-García *et al*¹³ found increased flexibility associated with a more alkaline arterial pH ($r=0.220$, $p<0.05$), higher arterial PO_2 ($r=0.237$, $p<0.05$) and lower arterial PCO_2 ($r=-0.331$, $p<0.01$) in the umbilical cord blood.

Balance

No validity or reliability assessments were performed regarding balance tests.

Three articles associated balance with neonatal and maternal health-related outcomes. Öztürk *et al*¹²⁸ observed that static balance decreased and fall risk increased in pregnant individuals with lower back pain (49.90 ± 24.47 vs 28.47 ± 19.60 ; $p<0.0001$). In relation to exercise, McCrory *et al*¹⁵⁷ showed that exercise may play a role in fall prevention in pregnancy ($p=0.005$) and they also found that dynamic balance is altered in pregnant individuals who have fallen compared with non-fallers and non-pregnant individuals ($p<0.001$). Nagai *et al*¹⁵⁸ studied the relationship between anxiety and balance. They concluded that when anxiety increases during pregnancy, the standing posture is destabilised ($r=0.559$, $p=0.020$), which may increase the chance of falling.

Speed

Validity and reliability for 10mTWT was studied by Evensen *et al* in two different articles.^{168 169} In 2015, Evensen *et al*¹⁶⁸ analysed the test-retest reliability of 10mTWT showing an ICC of (0.74). Intertester reliability was determined in the

first 13 participants with strong correlation (ICC=0.94). In 2016,¹⁶⁹ the same authors analysed the convergent validity of 10mTWT by comparing performances with scores achieved on the Active Straight Leg Raise (ASLR) test and observed moderate positive correlations between 10mTWT and ASLR ($r=0.65$, $p=0.003$).

This systematic review did not find any articles that analysed the association of speed with maternal and neonatal health outcomes.

Agility and coordination

No articles were identified.

Multidimensional

Validity and reliability for TUG was analysed by Evensen *et al* in two different studies.^{168 169} The TUG showed good test-retest reliability (ICC=0.88) and intertester reliability (ICC=0.95). Regarding reliability, strong correlations were found between the TUG and ASLR ($r=0.73$, $p=0.001$).

The time on TUG among pregnant individuals with PGP was significantly higher (mean (95% CI) 6.9 (6.5 to 7.3) seconds) than for asymptomatic pregnant (5.8 (5.5 to 6.0), $p<0.001$) and non-pregnant (5.5 (5.4 to 5.6), $p<0.001$) individuals.

DISCUSSION

Summary of the evidence

This systematic review revealed that PF has been assessed through a wide variety of tests during pregnancy. However, little is known on the validity and reliability of the tests performed, and the large variety of tests makes it challenging to compare results from different studies. Until a battery of specific fitness tests for pregnant women is developed and validated, the confidence of PF data obtained during pregnancy is limited and should be interpreted with caution. Consequently, the appropriateness of using this PF data to prescribe exercise during pregnancy could be questioned and is a matter that requires special attention. In this context, it is also difficult to evaluate the association of PF with maternal and neonatal health which, in fact, is of wide clinical and public health interest. However, some studies observed associations of PF with maternal and neonatal health outcomes, which needs to be replicated once a PF test battery is released. We strongly suggest that extensive research must be performed to validate such battery of PF tests.

Cardiorespiratory fitness

This systematic review identified that a cycle ergometer has been the equipment most frequently used to assess CRF followed by treadmill and field tests, although step tests have also been conducted. There is a large disparity of protocols and wide variety of ad hoc tests used, which makes comparing results between studies difficult. However, the Modified Balke treadmill Protocol validated by Mottola *et al*⁷⁵ for pregnant women has been the most

frequently used test. There have been more incremental tests used for CRF tests during the pregnancy compared with steady-state tests and more submaximal compared with maximal tests. There is no consensus regarding test termination criteria for submaximal tests, which undoubtedly needs further research. Some articles used relative intensity using physiological variables such as %HR_{max} or %VO_{2max}, and others used absolute intensity, such as specific HR (beats per minute). Among the studies that used %HR_{max} as a test termination criterion, there was a variety of percentages such as 70%,^{34 35 90 100} 75%^{27 29 69 97} or 85%.^{13 54 74} Among the studies that used %VO_{2max}, there were different percentages such as 40%,³⁸ 50%,^{37 101} 60%^{32 38} or 70%.³⁰ Among the studies that used absolute HR as a test termination criterion, the HR for finalising the tests were set either at 125,⁶¹ 150,^{36 45 62 108} 155,⁹⁴ 160⁶⁵ or 170^{50 53 55 56} beats per minute. Some studies even used the rate of perceived exertion as complementary criteria^{46 50 106} or peak aerobic power.³⁹ These complementary criteria have been recommended and studied in pregnant women by authors like Hesse *et al*⁸ since the physical and emotional changes during pregnancy limit performance. It must be noted that the same equation was not used to estimate HR_{max}. Some articles used the traditional 220-age formula^{29 35 54 69 97} while others used the Karvonen⁷⁴ or Tanaka¹⁰⁰ formulas. Some articles did not specify how HR_{max} was estimated.^{27 34 90} This heterogeneity could be due to the physiological complexity of pregnancy, in terms of cardiac changes and response to exercise and the lack of scientific information in this regard. Moreover, the gestational week could be a determinant for physiological responses since Bijl *et al*¹⁰⁰ observed a slower haemodynamic recovery and an increased ventilatory response to exercise in early pregnancy compared with non-pregnant women. With regard to maximal tests, different terms have been used for maximal criteria such as volitional fatigue,^{30 33 43 44 47 48 98 103 105} exhaustion,³¹ anaerobic threshold^{73 80 104 171} and point of symptom limitation.^{59 60 102}

This lack of consensus has many drawbacks that should be resolved in view of the need to accurately assess CRF during the pregnancy. We advocate for an expert consensus to be developed in the following years to achieve the goal of appropriate and effective CRF assessment during the pregnancy. In particular, it seems essential to develop a treadmill and a cycle ergometer submaximal test that reveals sufficient validity to confidently estimate VO_{2max} throughout gestation.

Muscular fitness

Muscular fitness tests included muscular strength, endurance and power.² The studies included in this systematic review show that muscular strength was the most frequently assessed component of muscular fitness, since only six studies^{12 13 112 117 122 178 179} assessed endurance and none of them assessed power in pregnancy. In most studies, muscular strength was evaluated through hand-grip maximal strength using a dynamometer. However,

two studies used a hand-grip sphygmomanometer test.^{118 119} Some of the hand-grip tests were performed in a standing position,^{8 109} while others used a sitting position¹¹⁰ or supine position,¹¹³ and others did not reveal the position used for the assessment.^{86 112 114 115} Some tests were completed three times,¹¹² others twice^{8 86 115} and others only once.^{110 113 114} This clearly reveals a large methodological variability that might influence the results and make comparing results between studies difficult. Another limitation is the fact that the main strength outcome was hand-grip strength. While hand-grip strength is a good marker of health,¹⁸⁰ it is unclear whether hand-grip responds to changes following exercise interventions. Therefore, validating other muscular strength tests, including lower limb strength tests, is needed for researchers and practitioners to confidently assess muscular strength during the pregnancy.

There were no validity studies and the reliability was assessed only in one maximal isometric hip extension test.¹²¹ This test has limitations since the pregnant abdomen must be on a bed and, as acknowledged by the authors, it cannot be performed during the third trimester. It must be noted that higher hand-grip strength was associated with higher birth weight.^{8 115} Moreover, increased hand-grip strength was produced during uterine contraction.¹¹¹ The advantage of using hand-grip is that it represents an inexpensive, rapid and easy-to-use assessment with minimal training needed to appropriately administer. However, assessing the performance of pregnant athletes with this test seems clearly insufficient. More quality in tests employed is necessary since the association of muscular strength with maternal and neonatal health outcomes is of clinical importance. Moreover, other studies are needed to understand the extent to which preserving strength throughout pregnancy and post partum relates to clinical outcomes.

Flexibility

Although there were seven studies assessing flexibility, none of them used the same protocol. Once again, this reflects a lack of agreement when assessing the same component of PF. Moreover, Lindgren and Kristiansson¹²⁶ found that higher flexibility showed higher low back pain. Despite the limitation of a finger laxity test, we considered these findings an interesting association that warrants further investigation since passive stretching is one of the most common practical prescriptions for exercise professionals instead of mobility and breathing exercises. On the other hand, the results of Baena-García *et al*¹³ are very relevant to fetal health since flexibility was associated with a better pH, PO₂ and PCO₂ in umbilical cord blood. Hence, more research about flexibility tests, their outcomes and their prescription are needed.

Balance

We identified that balance was the second PF component most frequently evaluated during pregnancy, following CRF. This makes sense since the centre of gravity changes during pregnancy as a result of expansion of the



uterus and the risk of falls increases. However, there is high heterogeneity between the protocols employed in different studies. For static balance, the protocol most frequently used was stabilometry on a force platform with bipedal support and eyes open and eyes closed within the same test. For dynamic balance, there was a greater heterogeneity across protocols both in the platform used and in the movements over the platforms. Regarding the assessment tool, the 3-D camera was the device most frequently used.^{139-142 144 165} Likewise, we observed differences between the number of platform pieces, trials and Hz used. Some protocols were performed on two piece platforms,^{130 131 149} others on one piece platforms^{129 132 138 158 160 166 167} and others did not specify the type of platform.¹⁶³⁻¹⁶⁵ Although the number of trials and the frequency of recording (ie, Hz) are important protocol parameters that should be carefully documented, only 5 (out of 13) articles described the number of trials^{131 138 166 167 175} and 1 described frequency of recording.¹⁴⁹ The usefulness of these tests is restricted to the research area and all of them use expensive technological tools; therefore, it is difficult to extrapolate these tests to fitness centres or clinical settings. Falls during pregnancy could be prevented if balance was easily assessed. For this reason, it is necessary to develop an inexpensive and easy-to-use balance field test.

Validity and reliability of PF tests, and association with maternal and neonatal health

Unfortunately, studies that examine validity and reliability of PF tests are scarce. The PF component most frequently studied was CRF. However, we only found two studies that analysed the validity of the CRF tests, and no studies examined the reliability of these tests. On a treadmill platform, Mottola *et al*,⁷⁵ validated a special equation for modified Balke protocol that has been used by numerous other authors. In contrast, Yeo *et al*⁷⁴ aimed to validate a portable metabolic testing system (mod. VO2000) but it overestimated VO₂ measurements for pregnant individuals compared with non-pregnant females and males.

Regarding muscular fitness, the hand-grip test was most commonly used; this test was used as the gold standard for muscular fitness during pregnancy. Only Gutke *et al*¹²¹ studied the reliability of a test for hip extension. However, the *p* value was not reported, and the position adopted in the test could be uncomfortable for pregnant participants. Finally, the studies evaluating validity and reliability of speed and multidimensional tests of PF have been researched by Evensen *et al*.^{168 169} They demonstrated that TUG and 10mTWT are reliable and valid tests for use during the pregnancy.

The validity and reliability of balance (without tests), agility and/or coordination tests has not been investigated to date.

We suggest that specific tests to be performed in pregnancy are needed and their validity and reliability must be assessed to understand the extent to which one might

rely on such measures when prescribing exercise, or making clinical recommendations.

Regarding the association of PF with maternal and neonatal health outcomes, we conclude that more research is also necessary. Nevertheless, from this review we can highlight some interesting associations with different fitness components. A better CRF was associated with a shorter labour^{57 108} and a lower risk of caesarean section.¹³ However, no association was found regarding other fetal outcomes such as Apgar scores or the newborn anthropometrics.^{57 108} By contrast, muscular strength was associated with optimum infant birth weight.^{8 13 115} Other neonatal outcomes like fetal umbilical cord pH were positively associated with maternal CRF.⁶⁸ On the other hand, better balance scores were associated with lower risk of falls,^{128 158 181} which is of particular interest for exercise professionals, who might include balance as a component of exercise programs for pregnant women. Finally, Evensen *et al*¹⁶⁹ found that PGP could be a limiting factor to assess PF in pregnant individuals since the time of TUG was significantly higher in those with pain than in asymptomatic pregnant and non-pregnant individuals.

None of the studies reviewed in this article have described adverse events during PF assessment. Moreover, official bodies such as the American College of Obstetricians and Gynecologists, the Canadian Society of Exercise Physiology and the Society of Obstetricians and Gynaecologists of Canada have highlighted the benefits of an adequate PF assessment, and assert the need of consensus in PF assessment during the pregnancy.¹⁸² Consequently, the findings from this study have important research and clinical implications.

Limitations and strengths

A limitation of this article is that, although PubMed and WOS are among the most relevant databases in the medical literature, the possibility that a small number of studies have been overlooked cannot be discarded. Nevertheless, these two databases are the biggest databases in sports medicine and sports sciences and, therefore, include the vast majority of studies.

A strength of this systematic review is the fact that, to the best of our knowledge, this is the first article to comprehensively analyse PF assessments, the validity and reliability of fitness tests, and their relationship with maternal and neonatal health outcomes during the pregnancy. The results from this systematic review provide an overall picture of how PF is being assessed in this population, what type of tests are being performed, their specific characteristics, whether these tests have been tested for validity and/or reliability; and whether PF is associated with maternal and neonatal health outcomes. All this information is of wide and undoubted clinical interest.

CONCLUSIONS

The main finding of this systematic review is that PF has been assessed through a wide variety of protocols, mostly lacking validity and reliability data, and that no

consensus exists on the most suitable fitness tests to be performed during pregnancy. In addition, the available evidence regarding the association of PF with maternal and neonatal health outcomes is scarce and is a matter of further investigation. Provided the need to assess PF during the pregnancy and the importance not only to understand the physical state of the pregnant individual but also to precisely prescribe exercise in this population, extensive research is needed to design and validate a battery of fitness tests to be used for the safe and effective assessment of PF during pregnancy. We advocate for an expert consensus panel to develop a battery of PF tests to assess the different PF components during pregnancy.

Author affiliations

¹Department of Physical Education and Sport, Universidad de Granada, Granada, Spain

²Sport and Health University Research Centre, Universidad de Granada, Granada, Spain

³Applied Sports Technology Exercise and Medicine Research Centre, Swansea University, Swansea, UK

⁴Sport Science Department, Swansea University, Swansea, UK

⁵GALENO Research Group, Department of Physical Education, Faculty of Education Sciences, Universidad de Cadiz, Cadiz, Spain

⁶The Biomedical Research and Innovation Institute of Cadiz (INIBICA), Cádiz, España

⁷Kinesiology, East Carolina University College of Health and Human Performance, Greenville, North Carolina, USA

⁸Department of Obstetrics & Gynecology, Brody School of Medicine, East Carolina University, Greenville, North Carolina, USA

⁹Gynaecology and Obstetrics Unit, 'San Cecilio' University Hospital, Universidad de Granada, Granada, Spain

¹⁰The Biosanitary Research Institute of Granada.ibs, Granada, Spain

¹¹R. Samuel McLaughlin Foundation- Exercise and Pregnancy Lab, School of Kinesiology, University of Western Ontario, London, Ontario, Canada

¹²Department of Physiology, Institute of Nutrition and Food Technology and Biomedical Research Centre, Universidad de Granada, Granada, Spain

¹³Department of Education, Faculty of Education Sciences, University of Almería, Almería, Spain

¹⁴SPORT Research Group (CTS-1024), CERNEP Research Center, University of Almería, Almería, Spain

Twitter Lidia Romero-Gallardo @lidiaromero_owa and Alberto Soriano-Maldonado @AlbertoSoriano_

Contributors Conceptualisation: LR-G, VAA, JC-P and AS-M. Literature search and data analysis: LR-G and ORR. Methodology: LR-G, ORR, JC-P, MM, VAA and AS-M. Formal analysis and investigation: LR-G, ORR and AS-M. Writing-original draft preparation: LR-G. Writing-review and editing: LR-G, ORR, JC-P, LEM, OO, LM, MM, VAA and AS-M. Resources: VAA, JC-P and AS-M. Supervision: VAA, OO, JC-P and AS-M.

Funding This study has been partially funded by the University of Granada, Plan Propio de Investigación 2016, Excellence actions: Units of Excellence: Unit of Excellence on Exercise and Health (UCEES), and by the Junta de Andalucía, Consejería de Conocimiento, Investigación y Universidades and European Regional Development Fund (ERDF), ref. SOMM17/6107/UGR. This study is included in the thesis of LRG enrolled in the Doctoral Program in Biomedicine of the University of Granada.

Competing interests Yes.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability

of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

Author note Please note that this work was posted as a pre-print prior to submission. The pre-print version can be accessed here <https://doi.org/10.1101/2021.06.26.21259584>.

ORCID iDs

Lidia Romero-Gallardo <http://orcid.org/0000-0001-6341-855X>

Olga Roldan Reoyo <http://orcid.org/0000-0003-2625-4060>

Jose Castro-Piñero <http://orcid.org/0000-0002-7353-0382>

Linda E May <http://orcid.org/0000-0002-8231-2280>

Olga Ocón-Hernández <http://orcid.org/0000-0002-8113-3651>

Michelle F Mottola <http://orcid.org/0000-0002-8707-4656>

Virginia A Aparicio <http://orcid.org/0000-0002-2867-378X>

Alberto Soriano-Maldonado <http://orcid.org/0000-0002-4626-420X>

REFERENCES

- Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985;100:126–31.
- Riebe D, Ehrman JK, Liguori G, eds. *ACSM's Guidelines for Exercise Testing and Prescription*. 10th ed. American College of Sports Medicine, 2018.
- Blair SN, Kohl HW, Barlow CE. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA* 1995;273:1093–8.
- Kodama S, Saito K, Tanaka S. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women. *JAMA* 2009;301:2024–35.
- Gibbons LW, Blair SN, Cooper KH, et al. Association between coronary heart disease risk factors and physical fitness in healthy adult women. *Circulation* 1983;67:977–83.
- Ortega FB, Cadenas-Sanchez C, Lee D-C, et al. Fitness and fatness as health markers through the lifespan: an overview of current knowledge. *Prog Prev Med* 2018;3:e0013.
- Ortega FB, Ruiz JR, Castillo MJ, et al. Physical fitness in childhood and adolescence: a powerful marker of health. *Int J Obes* 2008;32:1–11.
- Bisson M, Almérás N, Plaisance J, et al. Maternal fitness at the onset of the second trimester of pregnancy: correlates and relationship with infant birth weight. *Pediatr Obes* 2013;8:464–74.
- Gar C, Rottenkolber M, Grallert H, et al. Physical fitness and plasma leptin in women with recent gestational diabetes. *PLoS One* 2017;12:e0179128.
- Engberg E, Tikkanen HO, Koponen A, et al. Cardiorespiratory fitness and health-related quality of life in women at risk for gestational diabetes. *Scand J Med Sci Sports* 2018;28:1–9.
- Weissgerber TL, Wolfe LA, Davies GAL. Exercise in the prevention and treatment of maternal – fetal disease : a review of the literature. *Appl Physiol Nutr Metab* 2006;674:661–74.
- Price BB, Amini SB, Kappeler K. Exercise in pregnancy: effect on fitness and obstetric outcomes—a randomized trial. *Med Sci Sports Exerc* 2012;44:2263–9.
- Baena-García L, Coll-Risco I, Ocón-Hernández O, et al. Association of objectively measured physical fitness during pregnancy with maternal and neonatal outcomes. The GESTAFIT project. *PLoS One* 2020;15:e0229079.
- Marín-Jiménez N, Acosta-Manzano P, Borges-Cosic M, et al. Association of self-reported physical fitness with pain during pregnancy: the GESTAFIT project. *Scand J Med Sci Sports* 2019;29:1022–30.
- Romero-Gallardo L, Soriano-Maldonado A, Ocón-Hernández O, et al. International fitness Scale—IFIS: validity and association with health-related quality of life in pregnant women. *Scand J Med Sci Sport*. 2020;30:505–14.
- Miller MJ, Kutcher J, Adams KL. Effect of pregnancy on performance of a standardized physical fitness test. *Mil Med* 2017;182:e1859–63.

- 17 Treuth MS, Butte NF, Puyau M. Pregnancy-Related changes in physical activity, fitness, and strength. *Med Sci Sports Exer* 2005;37:832–7.
- 18 LeMoyné EL, Curnier D, Ellemberg D. Pregnancy and cognition: deficits in inhibition are unrelated to changes in fitness. *J Clin Exp Neuropsychol* 2014;36:178–85.
- 19 Meah VL, Backx K, Davenport MH. International Working group on maternal hemodynamics. functional hemodynamic testing in pregnancy: recommendations of the International Working group on maternal hemodynamics. *Ultrasound Obstet Gynecol* 2018;51:331–40.
- 20 May LE, Allen JJB, Gustafson KM. Fetal and maternal cardiac responses to physical activity and exercise during pregnancy. *Early Hum Dev* 2016;94:49–52.
- 21 Wolfe LA, Weissgerber TL. Clinical physiology of exercise in pregnancy: a literature review. *Journal of Obstetrics and Gynaecology Canada* 2003;25:473–83.
- 22 Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009;6:e1000100.
- 23 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 24 Castro-Piñero J, Artero EG, España-Romero V, et al. Criterion-related validity of field-based fitness tests in youth: a systematic review. *Br J Sports Med* 2010;44:934–43.
- 25 Artero EG, España-Romero V, Castro-Piñero J, et al. Reliability of field-based fitness tests in youth. *Int J Sports Med* 2011;32:159–69.
- 26 Armijo-Olivo S, Stiles CR, Hagen NA, et al. Assessment of study quality for systematic reviews: a comparison of the Cochrane collaboration risk of bias tool and the effective public health practice project quality assessment tool: methodological research. *J Eval Clin Pract* 2012;18:12–18.
- 27 Kulpa PJ, White BM, Visscher R. Aerobic exercise in pregnancy. *Am J Obstet Gynecol* 1987;156:1395–403.
- 28 Carpenter MW, Sady SP, Hoegsberg B. Fetal heart rate response to maternal exertion. *JAMA* 1988;259:3006–9.
- 29 Moore D, Jarrett J, Bendick P. Exercise-induced changes in uterine artery blood flow, as measured by Doppler ultrasound, in pregnant subjects. *Am J Perinatol* 1988;5:94–7.
- 30 Sady SP, Carpenter MW, Sady MA, et al. Prediction of VO₂max during cycle exercise in pregnant women. *J Appl Physiol* 1988;65:657–61.
- 31 Artal R, Masaki DI, Khodiguian N, et al. Exercise prescription in pregnancy: weight-bearing versus non-weight-bearing exercise. *Am J Obstet Gynecol* 1989;161:1464–9.
- 32 Hume RF, Bowie JD, McCoy C, et al. Fetal umbilical artery Doppler response to graded maternal aerobic exercise and subsequent maternal mean arterial blood pressure: predictive value for pregnancy-induced hypertension. *Am J Obstet Gynecol* 1990;163:826–9.
- 33 Sady MA, Haydon BB, Sady SP, et al. Cardiovascular response to maximal cycle exercise during pregnancy and at two and seven months postpartum. *Am J Obstet Gynecol* 1990;162:1181–5.
- 34 Field SK, Bell SG, Cenaiko DF, et al. Relationship between inspiratory effort and breathlessness in pregnancy. *J Appl Physiol* 1991;71:1897–902.
- 35 Rafla NM, Beazely JM. The effect of maternal exercise on fetal umbilical artery waveforms. *Eur J Obstet Gynecol Reprod Biol* 1991;40:119–22.
- 36 Bung P, Huch R, Huch A. Maternal and fetal heart rate patterns: a pregnant athlete during training and laboratory exercise tests; a case report. *Eur J Obstet Gynecol Reprod Biol* 1991;39:59–62 [https://www.ejog.org/article/0028-2243\(91\)90143-9/pdf](https://www.ejog.org/article/0028-2243(91)90143-9/pdf)
- 37 Young JC, Treadway JL. The effect of prior exercise on oral glucose tolerance in late gestational women. *Eur J Appl Physiol Occup Physiol* 1992;64:430–3.
- 38 Clapp JF, Little KD, Capeless EL. Fetal heart rate response to sustained recreational exercise. *Am J Obstet Gynecol* 1993;168:198–206.
- 39 Lotgering FK, Struijk PC, van Doorn MB, et al. Anaerobic threshold and respiratory compensation in pregnant women. *J Appl Physiol* 1995;78:1772–7 <http://www.ncbi.nlm.nih.gov/pubmed/7649911>
- 40 Artal R, Fortunato V, Welton A, et al. A comparison of cardiopulmonary adaptations to exercise in pregnancy at sea level and altitude. *Am J Obstet Gynecol* 1995;172:1170–80.
- 41 Souttanakis HN, Artal R, Wiswell RA. Prolonged exercise in pregnancy: glucose homeostasis, ventilatory and cardiovascular responses. *Semin Perinatol* 1996;20:315–27.
- 42 O'Neill ME. Maternal rectal temperature and fetal heart rate responses to upright cycling in late pregnancy. *Br J Sports Med* 1996;30:32–5.
- 43 Manders MA, Sonder GJ, Mulder EJ, et al. The effects of maternal exercise on fetal heart rate and movement patterns. *Early Hum Dev* 1997;48:237–47.
- 44 Kemp JG, Greer FA, Wolfe LA. Acid-Base regulation after maximal exercise testing in late gestation. *J Appl Physiol* 1997;83:644–51.
- 45 Wolfe LA, Preston RJ, Burggraf GW, et al. Effects of pregnancy and chronic exercise on maternal cardiac structure and function. *Can J Physiol Pharmacol* 1999;77:909–17.
- 46 Brenner IK, Wolfe LA, Monga M, et al. Physical conditioning effects on fetal heart rate responses to graded maternal exercise. *Med Sci Sports Exerc* 1999;31:792–9.
- 47 MacPhail A, Davies GA, Victory R, et al. Maximal exercise testing in late gestation: fetal responses. *Obstet Gynecol* 2000;96:565–70.
- 48 Heenan AP, Wolfe LA, Davies GA. Maximal exercise testing in late gestation: maternal responses. *Obstet Gynecol* 2001;97:127–34.
- 49 Kennelly MM, Geary M, McCaffrey N, et al. Exercise-related changes in umbilical and uterine artery waveforms as assessed by Doppler ultrasound scans. *Am J Obstet Gynecol* 2002;187:661–6.
- 50 Wolfe LA, Heenan AP, Bonen A. Aerobic conditioning effects on substrate responses during graded cycling in pregnancy. *Can J Physiol Pharmacol* 2003;81:696–703.
- 51 Lindqvist PG, Marsal K, Merlo J, et al. Thermal response to submaximal exercise before, during and after pregnancy: a longitudinal study. *J Matern Fetal Neonatal Med* 2003;13:152–6.
- 52 Lynch A-M, McDonald S, Magann EF, et al. Effectiveness and safety of a structured swimming program in previously sedentary women during pregnancy. *J Matern Fetal Neonatal Med* 2003;14:163–9.
- 53 Heenan AP, Wolfe LA. Plasma osmolality and the strong ion difference predict respiratory adaptations in pregnant and nonpregnant women. *Can J Physiol Pharmacol* 2003;81:839–47.
- 54 Pirhonen JP, Lindqvist PG, Marsal K. A longitudinal study of maternal oxygen saturation during short-term submaximal exercise. *Clin Physiol Funct Imaging* 2003;23:37–41.
- 55 McAuley SE, Jensen D, McGrath MJ, et al. Effects of human pregnancy and aerobic conditioning on alveolar gas exchange during exercise. *Can J Physiol Pharmacol* 2005;83:625–33.
- 56 Weissgerber TL, Wolfe LA, Hopkins WG, et al. Serial respiratory adaptations and an alternate hypothesis of respiratory control in human pregnancy. *Respir Physiol Neurobiol* 2006;153:39–53.
- 57 Pomerance JJ, Gluck L, Lynch VA. Physical fitness in pregnancy: its effect on pregnancy outcome. *Am J Obstet Gynecol* 1974;119:867–76.
- 58 Jensen D, Webb KA, Wolfe LA, et al. Effects of human pregnancy and advancing gestation on respiratory discomfort during exercise. *Respir Physiol Neurobiol* 2007;156:85–93.
- 59 Jensen D, Webb KA, Davies GAL, et al. Mechanical ventilatory constraints during incremental cycle exercise in human pregnancy: implications for respiratory sensation. *J Physiol* 2008;586:4735–50.
- 60 Kardel KR, Johansen B, Voldner N, et al. Association between aerobic fitness in late pregnancy and duration of labor in nulliparous women. *Acta Obstet Gynecol Scand* 2009;88:948–52.
- 61 Thorell E, Svärdsudd K, Andersson K, et al. Moderate impact of full-term pregnancy on estimated peak oxygen uptake, physical activity and perceived health. *Acta Obstet Gynecol Scand* 2010;89:1140–8.
- 62 Vega SR, Kleinert J, Sulprizio M, et al. Responses of serum neurotrophic factors to exercise in pregnant and postpartum women. *Psychoneuroendocrinology* 2011;36:220–7.
- 63 Thorell E, Goldsmith L, Weiss G, et al. Physical fitness, serum relaxin and duration of gestation. *BMC Pregnancy Childbirth* 2015;15:1–7.
- 64 Kim J-H, Roberge RJ, Powell JB. Effect of external airflow resistive load on postural and exercise-associated cardiovascular and pulmonary responses in pregnancy: a case control study. *BMC Pregnancy Childbirth* 2015;15:45.
- 65 Nakagaki A, Inami T, Minoura T, et al. Differences in autonomic neural activity during exercise between the second and third trimesters of pregnancy. *J Obstet Gynaecol Res* 2016;42:951–9.
- 66 Jędrzejko M, Nowosielski K, Poręba R, et al. Physical efficiency and activity energy expenditure in term pregnancy females measured during cardiopulmonary exercise tests with a supine cycle ergometer. *J Matern Fetal Neonatal Med* 2016;29:3800–5.
- 67 Kardel KR. Effects of intense training during and after pregnancy in top-level athletes. *Scand J Med Sci Sports* 2005;15:79–86.
- 68 Erkkola R, Rauramo L. Correlation of maternal physical fitness during pregnancy with maternal and fetal pH and lactic acid at delivery. *Acta Obstet Gynecol Scand* 1976;55:441–6.

- 69 Ong MJ, Guelfi KJ, Hunter T, *et al.* Supervised home-based exercise may attenuate the decline of glucose tolerance in obese pregnant women. *Diabetes Metab* 2009;35:418–21.
- 70 Sibley L, Ruhling RO, Cameron-Foster J, *et al.* Swimming and physical fitness during pregnancy. *J Nurse Midwifery* 1981;26:3–12.
- 71 Lewis RD, Yates CY, Driskell JA. Riboflavin and thiamin status and birth outcome as a function of maternal aerobic exercise. *Am J Clin Nutr* 1988;48:110–6.
- 72 Marquez-Sterling S, Perry AC, Kaplan TEDA, *et al.* Physical and psychological changes with vigorous exercise in sedentary primigravidae. *Med Sci Sports Exerc* 2000;32:58–62.
- 73 Santos IA, Stein R, Fuchs SC, *et al.* Aerobic exercise and submaximal functional capacity in overweight pregnant women. *Obstetrics & Gynecology* 2005;106:243–9.
- 74 Yeo S, Ronis DL, Antonakos CL, *et al.* Need for population specific validation of a portable metabolic testing system: a case of sedentary pregnant women. *J Nurs Meas* 2005;13:207–18.
- 75 Mottola MF, Davenport MH, Brun CR, *et al.* VO₂peak prediction and exercise prescription for pregnant women. *Med Sci Sports Exerc* 2006;38:1389–95.
- 76 Davenport MH, Charlesworth S, Vanderspank D, *et al.* Development and validation of exercise target heart rate zones for overweight and obese pregnant women. *Appl Physiol Nutr Metab* 2008;33:984–9.
- 77 de Oliveria Melo AS, Silva JLP, Tavares JS, *et al.* Effect of a physical exercise program during pregnancy on Uteroplacental and fetal blood flow and fetal growth. *Obstet Gynecol* 2012;120:302–10.
- 78 Ruchat S-M, Davenport M, Giroux I, *et al.* Walking program of low or vigorous intensity during pregnancy confers an aerobic benefit. *Int J Sports Med* 2012;33:661–6.
- 79 Morton MJ, Paul MS, Campos GR, *et al.* Exercise dynamics in late gestation: effects of physical training. *Am J Obstet Gynecol* 1985;152:91–7.
- 80 Salvesen Kjell Å, Hem E, Sundgot-Borgen J. Fetal wellbeing may be compromised during strenuous exercise among pregnant elite athletes. *Br J Sports Med* 2012;46:279–83.
- 81 Mottola MF, Inglis S, Brun CR, *et al.* Physiological and metabolic responses of late pregnant women to 40 min of steady-state exercise followed by an oral glucose tolerance perturbation. *J Appl Physiol* 2013;115:597–604.
- 82 Bisson M, Rhéaume C, Bujold E, *et al.* Modulation of blood pressure response to exercise by physical activity and relationship with resting blood pressure during pregnancy. *J Hypertens* 2014;32:1450–7.
- 83 Marshall MR, Pivarnik JM. Perceived exertion of physical activity during pregnancy. *J Phys Act Heal* 2015;12:1039–43.
- 84 da Silva EG, de Godoy I, de Oliveira Antunes LC, *et al.* Respiratory parameters and exercise functional capacity in preeclampsia. *Hypertens Pregnancy* 2010;29:301–9.
- 85 Ramírez-Vélez R, Aguilar de Plata AC, Escudero MM, *et al.* Influence of regular aerobic exercise on endothelium-dependent vasodilation and cardiorespiratory fitness in pregnant women. *J Obstet Gynaecol Res* 2011;37:1601–8.
- 86 Hjorth MF, Kloster S, Girma T, *et al.* Level and intensity of objectively assessed physical activity among pregnant women from urban Ethiopia. *BMC Pregnancy Childbirth* 2012;12:154.
- 87 Radzikowska E, Wiater E, Franczuk M. Lung function in pregnancy in Langerhans cell histiocytosis. *Adv Exp Med Biol* 2018;1023:73–83.
- 88 Oviedo-Caro MA, Bueno-Antequera J, Munguía-Izquierdo D. Explanatory factors and levels of health-related quality of life among healthy pregnant women at midpregnancy: a cross-sectional study of the PregnActive project. *J Adv Nurs* 2018;74:2766–76.
- 89 Dennis AT, Salman M, Paxton E, *et al.* Resting hemodynamics and response to exercise using the 6-minute walk test in late pregnancy: an international prospective multicentre study. *Anesth Analg* 2019;129:450–7.
- 90 Veille JC, Hohimer AR, Burry K, *et al.* The effect of exercise on uterine activity in the last eight weeks of pregnancy. *Am J Obstet Gynecol* 1985;151:727–30.
- 91 Birnbaumer P, Dietz P, Watson ED, *et al.* Absolute Accelerometer-Based intensity prescription compared to physiological variables in pregnant and nonpregnant women. *Int J Environ Res Public Health* 2020;17:5651.
- 92 Amola M, Pawara S, Kalra S. Effect of inspiratory muscle training and diaphragmatic breathing exercises on dyspnea, pulmonary functions, fatigue and functional capacity in pregnancy during third trimester. *J Clin Diagnostic Res* 2019;13:YC1–4.
- 93 Dibblee L, Graham TE. A longitudinal study of changes in aerobic fitness, body composition, and energy intake in primigravid patients. *Am J Obstet Gynecol* 1983;147:908–14.
- 94 Williams A, Reilly T, Campbell I, *et al.* Investigation of changes in responses to exercise and in mood during pregnancy. *Ergonomics* 1988;31:1539–49.
- 95 Melzer K, Schutz Y, Soehnchen N, *et al.* Effects of recommended levels of physical activity on pregnancy outcomes. *Am J Obstet Gynecol* 2010;202:266.e1–266.e6.
- 96 Szymanski LM, Satin AJ. Strenuous exercise during pregnancy: is there a limit? *Am J Obstet Gynecol* 2012;207:179.e1–179.e6.
- 97 Winn H, Hess O, Goldstein I, *et al.* Fetal responses to maternal exercise: effect on fetal breathing and body movement. *Am J Perinatol* 1994;11:263–6.
- 98 Hesse CM, Tinius RA, Pitts BC, *et al.* Assessment of endpoint criteria and perceived barriers during maximal cardiorespiratory fitness testing among pregnant women. *J Sports Med Phys Fitness* 2018;58:1844–51.
- 99 Santos CMD, Santos WMD, Gallarreta FMP, *et al.* Effect of maternal exercises on biophysical fetal and maternal parameters: a transversal study. *Einstein* 2016;14:455–60.
- 100 Bijl RC, Cornette JMJ, van der Ham K, *et al.* The physiological effect of early pregnancy on a woman's response to a submaximal cardiopulmonary exercise test. *Physiol Rep* 2020;8:e14624.
- 101 Jovanovic L, Kessler A, Peterson CM. Human maternal and fetal response to graded exercise. *J Appl Physiol* 1985;58:1719–22.
- 102 da Silva Corrêa M, Catai AM, Milan-Mattos JC, *et al.* Is pelvic floor muscle training able to alter the response of cardiovascular autonomic modulation and provide a possible cardiovascular benefit to pregnant women? *NeuroUrol Urodyn* 2020;39:2272–83.
- 103 Matenchuk BA, James M, Skow RJ, *et al.* Longitudinal study of cerebral blood flow regulation during exercise in pregnancy. *J Cereb Blood Flow Metab* 2020;40:2278–88.
- 104 Bilodeau J-F, Bisson M, Larose J, *et al.* Physical fitness is associated with prostaglandin F_{2α} isomers during pregnancy. *Prostaglandins Leukot Essent Fatty Acids* 2019;145:7–14.
- 105 Purdy GM, James MA, Wakefield PK, *et al.* Maternal cardioautonomic responses during and following exercise throughout pregnancy. *Appl Physiol Nutr Metab* 2019;44:263–70.
- 106 Sussman D, Saini BS, Schneiderman JE, *et al.* Uterine artery and umbilical vein blood flow are unaffected by moderate habitual physical activity during pregnancy. *Prenat Diagn* 2019;39:976–85.
- 107 O'Neill ME, Cooper KA, Hunyor SN, *et al.* Cardiorespiratory response to walking in trained and sedentary pregnant women. *J Sports Med Phys Fitness* 1993;33:40–3 <http://www.ncbi.nlm.nih.gov/pubmed/8350606>
- 108 Wong S, McKenzie D. Cardiorespiratory fitness during pregnancy and its effect on outcome. *Int J Sports Med* 1987;08:79–83.
- 109 Petrov Fieril K, Glantz A, Fagevik Olsen M. The efficacy of moderate-to-vigorous resistance exercise during pregnancy: a randomized controlled trial. *Acta Obstet Gynecol Scand* 2015;94:35–42.
- 110 Atay E, Başalan Iz F. Investigation of the effect of changes in muscle strength in gestational age upon fear of falling and quality of life. *Turk J Med Sci* 2015;45:977–83.
- 111 Wickboldt N, Savoldelli G, Rehberg-Klug B. Continuous assessment of labour pain using handgrip force. *Pain Res Manag* 2015;20:159–63.
- 112 Kalliokoski P, Rodhe N, Bergqvist Y, *et al.* Long-Term adherence and effects on grip strength and upper leg performance of prescribed supplemental vitamin D in pregnant and recently pregnant women of Somali and Swedish birth with 25-hydroxyvitamin D deficiency: a before-and-after treatment study. *BMC Pregnancy Childbirth* 2016;16:353.
- 113 Ngaka TC, Coetsee JF, Dyer RA. The influence of body mass index on sensorimotor block and vasopressor requirement during spinal anesthesia for elective cesarean delivery. *Anesth Analg* 2016;123:1527–34.
- 114 Rodríguez-Díaz L, Ruiz-Frutos C, Vázquez-Lara JM. Efectividad de un programa de actividad física mediante El método Pilates en El embarazo Y en El proceso del parto. *Enfermería Clínica* 2017;27:271–7.
- 115 Żelazniewicz A, Pawłowski B. Maternal hand grip strength in pregnancy, newborn sex and birth weight. *Early Hum Dev* 2018;119:51–5.
- 116 Takeda K, Yoshikata H, Imura M. Do squat exercises with weight shift during pregnancy improve postural control? *Int J of Women's Health and Reproduction Sciences* 2019;7:10–16.
- 117 Yenişehir S, Çıtak Karakaya Ilkim, Sivaslıoğlu AA, *et al.* Reliability and validity of five times sit to stand test in pregnancy-related pelvic girdle pain. *Musculoskelet Sci Pract* 2020;48:102157.
- 118 Baker PN, Johnson IR. The use of the hand-grip test for predicting pregnancy-induced hypertension. *Eur J Obstet Gynecol Reprod Biol* 1994;56:169–72.

- 119 Rogers MS, Todinson B. Change in cardiovascular indices with position and isometric exercise throughout pregnancy: assessment by impedance cardiography and oscillometric sphygmomanometry. *Hypertens Pregnancy* 1998;17:191–202.
- 120 Feiner B, Weksler R, Ohel G, et al. The influence of maternal exercise on placental blood flow measured by simultaneous Multigate spectral Doppler imaging (SM-SDI). *Ultrasound Obstet Gynecol* 2000;15:498–501.
- 121 Gutke A, Östgaard HC, Öberg B. Association between muscle function and low back pain in relation to pregnancy. *J Rehabil Med* 2008;40:304–11.
- 122 O'Connor PJ, Poudevigne MS, Cress ME, et al. Safety and efficacy of supervised strength training adopted in pregnancy. *J Phys Act Health* 2011;8:309–20 <http://www.ncbi.nlm.nih.gov/pubmed/21487130>
- 123 Gilleard W, Crosbie J, Smith R. Effect of pregnancy on trunk range of motion when sitting and standing. *Acta Obstet Gynecol Scand* 2002;81:1011–20.
- 124 Marnach ML, Ramin KD, Ramsey PS, et al. Characterization of the relationship between joint laxity and maternal hormones in pregnancy. *Obstet Gynecol* 2003;101:331–5.
- 125 Garshasbi A, Faghih Zadeh S, et al. The effect of exercise on the intensity of low back pain in pregnant women. *Int J Gynecol Obstet* 2005;88:271–5.
- 126 Lindgren A, Kristiansson P. Finger joint laxity, number of previous pregnancies and pregnancy induced back pain in a cohort study. *BMC Pregnancy Childbirth* 2014;14:61.
- 127 Cherni Y, Desseauve D, Decatoire A, et al. Evaluation of ligament laxity during pregnancy. *J Gynecol Obstet Hum Reprod* 2019;48:351–7.
- 128 Öztürk G, Geler Külcü D, Aydoğ E, et al. Effects of lower back pain on postural equilibrium and fall risk during the third trimester of pregnancy. *J Matern Neonatal Med* 2016;29:1358–62.
- 129 Shibayama Y, Kuwata T, Yamaguchi J, et al. Changes in standing body sway of pregnant women after long-term bed rest. *J Obstet Gynaecol* 2016;36:479–82.
- 130 Takeda K, Yoshikata H, Imura M. Changes in posture control of women that fall during pregnancy. *International Journal of Women's Health and Reproduction Sciences* 2018;6:255–62.
- 131 Moreira LS, Elias LA, Gomide AB. A longitudinal assessment of myoelectric activity, postural sway, and low-back pain during pregnancy. *Acta Bioeng Biomech* 2017;19:77–83.
- 132 Opala-Berdzik A, Błaszczyk JW, Świder D, et al. Trunk forward flexion mobility in reference to postural sway in women after delivery: a prospective longitudinal comparison between early pregnancy and 2- and 6-month postpartum follow-ups. *Clin Biomech* 2018;56:70–4.
- 133 Davies J, Fernando R, McLeod A, et al. Postural stability following ambulatory regional analgesia for labor. *Anesthesiology* 2002;97:1576–81.
- 134 McCrory JL, Chambers AJ, Daftary A, et al. Dynamic postural stability during advancing pregnancy. *J Biomech* 2010;43:2434–9.
- 135 Branco M, Santos-Rocha R, Aguiar L, et al. Kinematic analysis of gait in the second and third trimesters of pregnancy. *J Pregnancy* 2013;2013:1–9.
- 136 Cakmak B, Inanir A, Nacar MC. The effect of maternity support belts on postural balance in pregnancy. *PM&R* 2014;6:624–8.
- 137 Inanir A, Cakmak B, Hisim Y, et al. Evaluation of postural equilibrium and fall risk during pregnancy. *Gait Posture* 2014;39:1122–5.
- 138 Butler EE, Colón I, Druzin ML, et al. Postural equilibrium during pregnancy: decreased stability with an increased reliance on visual cues. *Am J Obstet Gynecol* 2006;195:1104–8.
- 139 Wu W, Meijer OG, Lamoth CJC, et al. Gait coordination in pregnancy: transverse pelvic and thoracic rotations and their relative phase. *Clin Biomech* 2004;19:480–8.
- 140 Forczek W, Staszkievicz R. Changes of kinematic gait parameters due to pregnancy. *Acta Bioeng Biomech* 2012;14:113–9.
- 141 Takeda K. A Kinesiological analysis of the Stand-to-Sit during the third trimester. *J Phys Ther Sci* 2012;24:621–4.
- 142 Gottschall JS, Sheehan RC, Downs DS. Pregnant women exaggerate cautious gait patterns during the transition between level and Hill surfaces. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 2013;23:1237–42.
- 143 McCrory JL, Chambers AJ, Daftary A, et al. The pregnant “waddle”: An evaluation of torso kinematics in pregnancy. *J Biomech* 2014;47:2964–8.
- 144 Krkeljas Z. Changes in gait and posture as factors of dynamic stability during walking in pregnancy. *Hum Mov Sci* 2018;58:315–20.
- 145 Błaszczyk JW, Opala-Berdzik A, Plewa M. Adaptive changes in spatiotemporal gait characteristics in women during pregnancy. *Gait Posture* 2016;43:160–4.
- 146 Sawa R, Doi T, Asai T, et al. Differences in trunk control between early and late pregnancy during gait. *Gait Posture* 2015;42:455–9.
- 147 Valerio PM, Gonçalves VE, Zordão CC, et al. Influence of type 1 diabetes on the postural control of women in the third gestational trimester. *Clin Biomech* 2020;77:105062.
- 148 McCrory JL, Chambers AJ, Daftary A, et al. Torso kinematics during gait and trunk anthropometry in pregnant fallers and non-fallers. *Gait Posture* 2020;76:204–9.
- 149 Ribas IS, Guirro ECO. Analysis of plantar pressure and postural balance during different phases of pregnancy. *BRAZILIAN J Phys Ther* 2007;11:391–6.
- 150 Rothwell SA, Eckland CB, Campbell N, et al. An analysis of postpartum walking balance and the correlations to anthropometry. *Gait Posture* 2020;76:270–6.
- 151 Catena RD, Bailey JP, Campbell N, et al. Stand-to-sit kinematic changes during pregnancy correspond with reduced sagittal plane hip motion. *Clin Biomech* 2019;67:107–14.
- 152 Catena RD, Campbell N, Werner AL, et al. Anthropometric changes during pregnancy provide little explanation of dynamic balance changes. *J Appl Biomech* 2019;35:232–9.
- 153 Forczek W, Ivanenko Y, Curyło M, et al. Progressive changes in walking kinematics throughout pregnancy—A follow up study. *Gait Posture* 2019;68:518–24.
- 154 Forczek W, Masłoń A, Frączek B, et al. Does the first trimester of pregnancy induce alterations in the walking pattern? *PLoS One* 2019;14:e0209766.
- 155 Gimunová M, Zvonar M, Sebera M, et al. Special footwear designed for pregnant women and its effect on kinematic gait parameters during pregnancy and postpartum period. *PLoS One* 2020;15:e0232901.
- 156 Forczek W, Ivanenko Y, Salamaga M, et al. Pelvic movements during walking throughout gestation - the relationship between morphology and kinematic parameters. *Clin Biomech* 2020;71:146–51.
- 157 McCrory JL, Chambers AJ, Daftary A, et al. Dynamic postural stability in pregnant fallers and non-fallers. *BJOG* 2010;117:954–62.
- 158 Nagai M, Isida M, Saitoh J, et al. Characteristics of the control of standing posture during pregnancy. *Neurosci Lett* 2009;462:130–4.
- 159 Catena RD, Campbell N, Wolcott WC, et al. Anthropometry, standing posture, and body center of mass changes up to 28 weeks postpartum in Caucasians in the United States. *Gait Posture* 2019;70:196–202.
- 160 Oliveira LF, Vieira TMM, Macedo AR, et al. Postural sway changes during pregnancy: a descriptive study using stabilometry. *Eur J Obstet Gynecol Reprod Biol* 2009;147:25–8.
- 161 Catena RD, Bailey JP, Campbell N, et al. Correlations between joint kinematics and dynamic balance control during gait in pregnancy. *Gait Posture* 2020;80:106–12.
- 162 Fontana Carvalho AP, Dufresne SS, Rogério De Oliveira M, et al. Effects of lumbar stabilization and muscular stretching on pain, disabilities, postural control and muscle activation in pregnant woman with low back pain. *Eur J Phys Rehabil Med* 2020;56:297–306.
- 163 Karadag-Saygi E, Unlu-Ozkan F, Basgul A. Plantar pressure and foot pain in the last trimester of pregnancy. *Foot Ankle Int* 2010;31:153–7.
- 164 Yu Y, Chung HC, Hemingway L, et al. Standing body sway in women with and without morning sickness in pregnancy. *Gait Posture* 2013;37:103–7.
- 165 Ersal T, McCrory JL, Sienko KH. Theoretical and experimental indicators of falls during pregnancy as assessed by postural perturbations. *Gait Posture* 2014;39:218–23.
- 166 Opala-Berdzik A, Bacik B, Markiewicz A, et al. Comparison of static postural stability in exercising and non-exercising women during the perinatal period. *Med Sci Monit* 2014;20:1865–70.
- 167 Opala-Berdzik A, Błaszczyk JW, Bacik B, et al. Static postural stability in women during and after pregnancy: a prospective longitudinal study. *PLoS One* 2015;10:e0124207.
- 168 Evensen NM, Kvåle A, Brækken IH. Reliability of the timed up and go test and Ten-Metre timed walk test in pregnant women with pelvic girdle pain. *Physiother Res Int* 2015;20:158–65.
- 169 Evensen NM, Kvåle A, Brækken IH. Convergent validity of the timed up and go test and Ten-metre timed walk test in pregnant women with pelvic girdle pain. *Man Ther* 2016;21:94–9.
- 170 Christensen L, Vøllestad NK, Veierød MB, et al. The Timed Up & Go test in pregnant women with pelvic girdle pain compared to

- asymptomatic pregnant and non-pregnant women. *Musculoskelet Sci Pract* 2019;43:110–6.
- 171 Kennelly MM, McCaffrey N, McLoughlin P, *et al.* Fetal heart rate response to strenuous maternal exercise: not a predictor of fetal distress. *Am J Obstet Gynecol* 2002;187:811–6.
- 172 Heenan AP, Wolfe LA, Davies GAL, *et al.* Effects of human pregnancy on fluid regulation responses to short-term exercise. *J Appl Physiol* 2003;95:2321–7.
- 173 Jensen D, Webb KA, O'Donnell DE. Chemical and mechanical adaptations of the respiratory system at rest and during exercise in human pregnancy. *Appl Physiol Nutr Metab* 2007;32:1239–50.
- 174 Gutke A, Östgaard HC, Öberg B. Predicting persistent pregnancy-related low back pain. *Spine* 2008;33:E386–93.
- 175 Cakmak B, Ribeiro AP, Inanir A. Postural balance and the risk of falling during pregnancy. *J Matern Fetal Neonatal Med* 2016;29:1623–5.
- 176 Martínez-Martí F, Martínez-García MS, Carvajal M Á, *et al.* Fractal behavior of the trajectories of the foot centers of pressure during pregnancy. *Biomed Phys Eng Express* 2019;5:025007.
- 177 Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 2000;18:963–75.
- 178 Gutke A, Kjellby-Wendt G, Öberg B. The inter-rater reliability of a standardised classification system for pregnancy-related lumbopelvic pain. *Man Ther* 2010;15:13–18.
- 179 Wade M, Prime H, Hoffmann TJ, *et al.* Birth weight interacts with a functional variant of the oxytocin receptor gene (OXTR) to predict executive functioning in children. *Dev Psychopathol* 2018;30:203–11.
- 180 García-Hermoso A, Cavero-Redondo I, Ramírez-Vélez R, *et al.* Muscular strength as a predictor of all-cause mortality in an apparently healthy population: a systematic review and meta-analysis of data from approximately 2 million men and women. *Arch Phys Med Rehabil* 2018;99:2100–13.
- 181 McCrory JL, Chambers AJ, Daftary A, *et al.* Ground reaction forces during gait in pregnant fallers and non-fallers. *Gait Posture* 2011;34:524–8.
- 182 Birsner ML, Gyamfi-Bannerman C. Physical activity and exercise during pregnancy and the postpartum period. *Obstet Gynecol* 2020;135:e178–88.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Online Supplemental Material 1, table S1. PRISMA Checklist 2020

Section and Topic	Item #	Checklist item	Location where item is reported
TITLE			
Title	1	Identify the report as a systematic review.	1
ABSTRACT			
Abstract	2	See the PRISMA 2020 for Abstracts checklist.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	3
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	3-4
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	4
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	4
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	4
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	4
Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	5
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	5
	10b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	5
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	5
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results.	-
Synthesis methods	13a	Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)).	-

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Section and Topic	Item #	Checklist item	Location where item is reported
	13b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	-
	13c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	-
	13d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	4
	13e	Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression).	-
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	-
Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	ESM2-3
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	ESM2-3
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram.	5
	16b	Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded.	5
Study characteristics	17	Cite each included study and present its characteristics.	5-12
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	5-12
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	5-12
Results of syntheses	20a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	5-12
	20b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	5-12
	20c	Present results of all investigations of possible causes of heterogeneity among study results.	5-12
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	5-12
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	-
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	-

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Section and Topic	Item #	Checklist item	Location where item is reported
DISCUSSION			
Discussion	23a	Provide a general interpretation of the results in the context of other evidence.	12-16
	23b	Discuss any limitations of the evidence included in the review.	17
	23c	Discuss any limitations of the review processes used.	17
	23d	Discuss implications of the results for practice, policy, and future research.	17
OTHER INFORMATION			
Registration and protocol	24a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	2
	24b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	4
	24c	Describe and explain any amendments to information provided at registration or in the protocol.	4
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	17
Competing interests	26	Declare any competing interests of review authors.	36
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	ESM5

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71

ESM=Electronic Material Supplement

For more information, visit: <http://www.prisma-statement.org/>

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Online Supplemental Material 2. Search strategy in PubMed and Web of Science.

For PubMed, we used Medical Subject Heading (MeSH) terms. This is a powerful method to enhance the quality of the search. In addition, all MeSH terms were included without the command MeSH attached, to consolidate our results and avoid losing those papers not included in MeSH database. This is because some MeSH terms were introduced in a specific date (e.g., ‘physical fitness’ was included in 1996). Hence papers published in a previous date would be lost. The same process was developed with terms not available in the MeSH database such as agility, aerobic capacity, etc. (see ESM 2-Table-S1) for search criteria and related terms.

All terms were combined using the connector OR for similar criteria. The connector ‘AND’ was used to combine population group (i.e., pregnant women), to delimit date of publication ("0001/01/01"[PDat]: "2021/01/15"[PDat]), to include full text papers, and to include studies performed in humans. A similar search strategy and terms combination was undertaken in WoS (ESM 2-Table-S2), although MeSH terms and its appropriate terms connection were not used as they are exclusive for PubMed.

The first step of the search was to look for systematic reviews and meta-analysis within the field of this systematic review. Since there was no such article published regarding our topic, the research team agreed on starting the search with no limit on the publication date. Then, an initial search was undertaken in both databases following the strategy explained in ESM 2-Table-S1 and ESM 2-Table-S2 for PubMed and WoS database respectively. The results from both, were merged.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Online Supplemental Material 2, Table S2. Search strategy used and number of articles found in **Pubmed**.

<i>Search Strategy</i>			
("Pregnant Women"[Mesh] OR "Pregnant Women" OR "Pregnancy"[Mesh] OR "Pregnancy") AND ("Physical Fitness"[Mesh] OR "Physical Fitness" OR "Physical Conditioning" OR "Exercise Test"[Mesh] OR "Exercise Test" OR "Fitness Trackers"[Mesh] OR "Fitness Trackers" OR "Muscle Strength"[MeSH] OR "Muscle Strength" OR "Muscular fitness" OR "Range of motion, articular"[Mesh] OR "Range of motion, articular" OR "Postural Balance"[MeSH] OR "Postural Balance" OR "Walk Test"[Mesh] OR "Walk Test" OR "Cardiorespiratory Fitness"[Mesh] OR "Cardiorespiratory Fitness" OR "Agility" OR "running speed" OR "aerobic fitness" OR "aerobic capacity" OR "maximal oxygen consumption" OR "V02max" OR "Physical function") AND full text[<i>sb</i>] AND ("0001/01/01"[<i>PDat</i>] : "2021/01/15"[<i>PDat</i>]) AND Humans[Mesh]			
Search criteria 1	MeSH Entry Terms for Criteria 1	Search criteria 2	MeSH Entry Terms for Criteria 2
Pregnant Women (MeSH)	Women, Pregnant Pregnant Woman Woman, Pregnant	Physical fitness (MeSH)	Fitness, Physical
Pregnancy (MeSH)		Exercise Test (MeSH)	Exercise Tests Test, Exercise Tests, Exercise Arm Ergometry Test Arm Ergometry Tests Ergometry Test, Arm Ergometry Tests, Arm Test, Arm Ergometry Tests, Arm Ergometry Bicycle Ergometry Test Bicycle Ergometry Tests

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Ergometry Test, Bicycle
Ergometry Tests, Bicycle
Test, Bicycle Ergometry
Tests, Bicycle Ergometry
Fitness Testing
Fitness Testings
Testing, Fitness
Testings, Fitness
Step Test
Step Tests
Test, Step
Tests, Step
Stress Test
Stress Tests
Test, Stress
Tests, Stress
Treadmill Test
Test, Treadmill
Tests, Treadmill
Treadmill Tests
Physical Fitness Testing
Fitness Testing, Physical
Fitness Testings, Physical
Physical Fitness Testings
Testing, Physical Fitness
Testings, Physical Fitness
Cardiopulmonary Exercise Test
Cardiopulmonary Exercise Tests
Exercise Test, Cardiopulmonary
Exercise Tests, Cardiopulmonary

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

	Test, Cardiopulmonary Exercise
Fitness Trackers (MeSH)	Fitness Tracker Tracker, Fitness Trackers, Fitness Physical Fitness Trackers Fitness Tracker, Physical Fitness Trackers, Physical Physical Fitness Tracker Tracker, Physical Fitness Trackers, Physical Fitness Activity Trackers Activity Tracker Tracker, Activity Trackers, Activity Personal Fitness Trackers Fitness Tracker, Personal Fitness Trackers, Personal Personal Fitness Tracker Tracker, Personal Fitness Trackers, Personal Fitness
Muscle Strength (MeSH)	Strength, Muscle
Muscle strength dynamometer (MeSH)	Dynamometer, Muscle Strength Dynamometers, Muscle Strength Muscle Strength Dynamometers
Range of motion, articular (MeSH)	Joint Range of Motion Joint Flexibility Flexibility, Joint Range of Motion Passive Range of Motion
Postural Balance (MeSH)	Musculoskeletal Equilibrium

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

		Equilibrium, Musculoskeletal Postural Equilibrium Equilibrium, Postural Balance, Postural
	Walk Test (MeSH)	Test, Walk Tests, Walk Walk Tests 6-Minute Walk Test 6 Minute Walk Test 6-Minute Walk Tests Test, 6-Minute Walk Tests, 6-Minute Walk Walk Test, 6-Minute Walk Tests, 6-Minute Incremental Shuttle Walk Test Endurance Shuttle Walk Test
		Cardiorespiratory fitness (MeSH) Fitness, Cardiorespiratory
Total items found	Without filters:	1657
	With Humans filter:	1135
	With Full Text filter:	1388
	With Humans & Full Text Filter:	930

The search recruited articles published until 15.01.21: no starting date limit was set for the search.

MeSH (Medical Subject Headings) is the National Library of Medicine controlled vocabulary thesaurus used for indexing articles for PubMed

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health— a systematic review

Online Supplemental Material 2, Table S3. Search strategy used and number of articles found in **Web of Science**

Search Strategy

TS=("Pregnant women" OR "pregnancy" OR "pregnan*") AND (("Physical Conditioning" OR "Physical fitness" OR "Exercise Test*" OR "Arm Ergometry Test*" OR "Bicycle Ergometry Test*" OR "Step Test*" OR "Treadmill Test*" OR "Physical Fitness Test*" OR "Cardiopulmonary Exercise Test*" OR "Fitness Tracker*" OR "Physical Fitness Tracker*" OR "Activity Tracker*" OR "Personal Fitness Tracker*") OR ("Muscle Strength" OR "Muscular Fitness" OR "Muscle strength dynamometer*") OR ("Joint Range of motion" OR "Joint flexibility" OR "Flexibility" OR "Range of motion" OR "Passive Range of Motion") OR ("Postural Balance" OR "Musculoskeletal Equilibrium" OR "Equilibrium" OR "Postural Equilibrium") OR ("Walk Test*" OR "6-Minute Walk Test*" OR "Incremental Shuttle Walk Test*" OR "Endurance Shuttle Walk Test") OR ("Cardiorespiratory Fitness" OR "Cardiovascular Fitness OR “Aerobic Fitness” OR “Aerobic Capacity” OR “Maximal Oxygen Consumption” OR “V02max”) OR (“Agility” OR “running speed” OR “aerobic fitness”))

Total items found 1687

The search recruited articles published until 15.01.21 no starting date limit was set for the search.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health— a systematic review

Online Supplemental Material 3. Comprehensive description of the three quality assessment scores used in the present systematic review.

The first quality score [24], was used to evaluate the quality of the articles that assessed validity. This list included three items based on sample size, description of the article population and statistical analysis to assess validity of each article. The validity quality score ranged from 0 to 6 (Table-S3). A score of 0-2 defined a very low-quality article; a score of 3-4 defined a low-quality article; and a score of 5-6 defined a high-quality article.

The second quality score [25] was employed to rate the studies that measured reliability (ESM 4 – Table-S4). This ranking was formed by four items based on description of the participants, the time interval, the results and appropriateness of statistical analyses. Each item in both, was rated from 0 (the lowest quality) to 2 (the highest quality). The reliability quality score ranged from 0 to 8 (ESM 4– Table-S4). A score of 0-1 defined a very low-quality article; a score of 2-5 defined a low-quality article; and a score of 6-8 defined a high-quality article.

The third quality score (ESM 4– Table-S5) was created to evaluate those studies that assessed association of PF with health-related outcomes. We adapted a score previously used in the Effective Public Health Practice Project (EPHPP) [26] which has been used in similar reviews [27]. The health-related outcomes quality score ranged from 0 to 5 (ESM 4– Table S5). A score of 0-2 defined a very low-quality article, a score of 3-4 defined a low-quality article, and a score of 5 defined a high-quality score. Three quality scores were calculated by counting the number of positive items.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Online Supplemental Material 3 Table S4. Quality assessment criteria to evaluate validity studies.

Grading system parameter	Grade	Criterion
Number of study subjects	0	n < 10
	1	n= 11-50
	2	n>51
Description of the study population regarding to age, sex, health status, fitness levels, etc	0	Less items than required for grade 1
	1	At least age and week of gestation.
	2	Age, week of gestation, health status and fitness levels and more.
Statistical analysis included in the study	0	Those not included in grade 1
	1	Error indexes or regression analysis
	2	≥3 items of Bland-Altamn plot and or ANOVA for repeated measurements

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health– a systematic review

Electronic Supplementary Material 4 Table S5. Quality assessment criteria to evaluate reliability studies.

Grading system parameter	Grade	Criterion
Description of the participants	0	Less items than required for grade 1.
	1	At least age and week of gestation.
	2	Age, week of gestation, health status and fitness levels and more.
Description of the time interval	0	Interval unknown.
	1	Vague and imprecise information about interval.
	2	Precise and complete description about interval.
Description of the results	0	Less results presented than required for grade.
	1	Description of test-retest results or description of the differences.
	2	Description of test-retest results and description of the differences.
Appropriateness of statistic	0	Only coefficient of variation
	1	Everything between grades 0 and 2 (normally – but not always – correlation plus an additional statistic).
	2	At least paired statistics, ANOVA for repeated measures (or non-parametrical corresponding tests) or Bland-Altman method.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with maternal and neonatal health— a systematic review

Electronic Supplementary Material 4 Table S6. Quality assessment criteria to evaluate health-related outcomes studies.

Grading system parameter	Grade	Criterion
Description of the study sample regarding to number of participants, age, sex, health status, fitness levels, etc	0	$n \leq 25$ and including less item than required for grade 1.
	1	$N \geq 26$ and at least age and gestational week.
Adequate assessment and report of physical fitness test.	0	Items for grade 1 are not included within the article.
	1	Validity and/or reliability reported of test and detailed description of testing protocol.
Adequate assessment of health-related outcomes	0	Items for grade 1 are not included within the article.
	1	Validity or reliability of the outcome measure reported and/or measurement procedure adequately described.
Adequate adjustment of confounders	0	No adjustment was done.
	1	Adjustment of confounders such as age and sex were done.
Description of both number and reasons to withdrawal and dropout.	0	No description included.
	1	Description included.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Electronic Supplementary Material 5 Table S7. Overview of studies included in the systematic review and description of physical fitness tests.

Reference (authors, year)	Sample Size (n)	Gestation Weeks (SD) or range in weeks	Mean age (SD), or range, in years	Fitness Test and Short Description
<i>Cardiorespiratory Fitness</i>				
<i>Cycle-ergometer protocol</i>				
Pomerance et al., (1974) ¹	54	17.5-27	35-37	Ad hoc, steady-state test at 60 rpm at 450, 600 and 300 kpm.
Erkkola, (1976) ²	120	(2 weeks before term)	20-26	1) Ad hoc, incremental submaximal test at 150, 300 and 450 kpm/min. 2) Arstila ECG test.
Morton et al, (1985) ³	23	40.15 (1.5)	28.5 (2.1)	Ad hoc, steady-state test at 40 to 50 rpm and at 300 kpm . min ⁻¹ for 6 min.
Veille et al., (1985) ⁴	17	35 (2)	31 (1)	Ad hoc, incremental submaximal test at 50 and 60 rpm at 50W for 10-15 min to 70% HR max (no formula).
Jovanovic et al., (1985) ⁵	6	37.1 (0.9)	28.5 (1.7)	Ad hoc, incremental submaximal self-administered test to 50% VO ₂ max or exertion equivalent to usual training.
Wong & McKenzie, (1987) ⁶	20	3 time-points, (10-14; 22-24; 34-36)	29.13	Ad hoc, incremental submaximal test at 50 rpm at 25, 50, 75 and 100 W for 5-6 min to 150 bpm.
Kulpa et al., (1987) ⁷	141	First trimester	18-34	Bruce protocol to 75% of HR max.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Carpenter et al., (1988) ⁸	45	29 (3.7)	25.2 (3)	Ad hoc, incremental test. 2 phases: a) <i>submaximal</i> : at 0, 30 and 60W for 6 min. b) <i>maximal</i> : at 60W to volitional fatigue.
Moore et al., (1988) ⁹	11	21.3	26.6	Ad hoc, incremental submaximal test at free pace for 20 min to 60 to 75% HR max. (220-age).
Sady & Carpenter, (1988) ¹⁰	40	29.2 (3.9)	25.9 (3.3)	2 incremental tests: 1) <i>Submaximal test</i> at 0 W, 30 W and 60 W at 30%, 50% and 70% of VO ₂ max. 2) <i>Maximal test</i> increasing 10 W every 2-min stage to volitional fatigue.
Artal et al., (1989) ¹¹	37	29.8 (0.5)	28.3 (1.8)	Ad hoc, incremental maximal test at 25, 50, 75W and increments of 25W every 2-min stage until exhaustion.
Hume et al., (1990) ¹²	30		28	Ad hoc, steady-state submaximal test at 60% VO ₂ max for 20min.
Sady et al., (1990) ¹³	9	25.6 (3.0);	29 (4.9)	Ad hoc, incremental test. 2 phases: a) <i>submaximal</i> : at 0, 30 and 60 W for 18 min. 6-min each stage. b) <i>maximal</i> : incremental continuous to volitional fatigue.
Field et al., (1991) ¹⁴	13	33 ± 2	30 (4)	Modified Balke protocol to 70% HR max (no formula)
Rafla & Beazely, (1991) ¹⁵	21	28-37	-	Ad hoc, incremental submaximal test from 60 rpm to 70% HR max (220-age)
Bung et al., (1991) ¹⁶	1	3 time-points, (24, 28, 37)	25	Ad hoc, incremental submaximal test from 15 W to 150 bpm.
Young & Treadway, (1992) ¹⁷	5	33 (1)	29 (1)	Ad hoc, steady-state submaximal test at 50% VO ₂ max for 30 min.
Clapp et al., (1993) ¹⁸	120	16-39	-	Ad hoc, steady-state submaximal test at 60% ± 3% VO ₂ max for 30 min.
Lotgering et al., (1995) ¹⁹	33	3 time-points, 16.1 (1); 25 (0.7); 35 (0.6)	30.9 (0.7)	Ad hoc, incremental submaximal test. After 3 min at 15W, to increase 10 W every 30 sec until peak aerobic power.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Artal et al., (1995) ²⁰	7	33.86±1.46	24.9 (2.18)	Ad hoc, incremental submaximal test. After 5 min per stage at 25, 50 and 75W, to increase 25W every 2 min to volitional fatigue.
O'Neill, (1996) ²¹	11	35.8 (1.1)	30.3 (3.3)	1) Ad hoc, steady-state test at 62.5 W for 15 min. 2) Ad hoc, steady-state test at 87.5 W for 15 min. 3) Ad hoc, steady-state test at 62.5 W for 30 min.
Soultanakis et al., (1996) ²²	20	27.1 (1.3)	31.4 (1.5)	1) Incremental maximal with modified Balke protocol, increasing 25 W every 2-min at 60 rpm to VO ₂ max 2) Ad hoc, steady-state submaximal test during 1 hour at 50%-60% VO ₂ max at 60 rpm.
Manders et al., (1997) ²³	12	29-32	20-36	Ad hoc, incremental maximal test. After 5-min per stage at 50W, to increase 25 W/min to volitional fatigue.
Kemp et al., (1997) ²⁴	23	33 (1)	-	Ad hoc, incremental maximal test at 20 W for 4 min. Then, increasing 20 W/min until exhaustion.
McGrath et al., (1999) ²⁵	41	3 time-points: 17.45 (0.45); 26.5 (0.2) and 37.15 (0.15)	29.4 (0.85)	Ad hoc, steady-state test with three 6-min stages and exercise brief (<5-min) between them. 1) 20 W to 110 bpm, 2) 45 W to 130 bpm 3) 70 W to 150 bpm.
Brenner et al., (1999) ²⁶	20	27.0 (1.0) and 37.0 (1.0)	29 (3.35)	Ad hoc incremental submaximal test for 3 min without resistance, then, increased 30 W/min to 170 bpm or RPE of 18.
MacPhail et al., (2000) ²⁷	23	32 (4)	20-40	Idem Kemp et al., (1997)
Heenan et al., (2001) ²⁸	28	34.7 (0.4)	30.8 (1.5)	Idem Kemp et al., (1997)
Kennelly et al., (2002) ²⁹	22	32.1 (1.4)	25.9 (4.9)	Ad hoc incremental maximal test. After 2-min at 30 W, increasing 10 W/min at 50-60 rpm to achieve AT.
Heenan & Wolfe, (2003) ³⁰	22	37.0 (0.2)	29 (1.1)	1) Ad hoc, incremental submaximal test at 20 W for 4 min. Then, to increase 20 W/min until 170 bpm. 2) Ad hoc, incremental ramp test from 0 W increasing work rate in 30-sec periods to 70 or 110% of VT.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Wolfe et al., (2003) ³¹	18	3 time-points: 19.2 (0.8) 27.8 (0.3) 37.0 (0.3)	28.3 (0.25)	Ad hoc incremental submaximal test for 3 min of no resistance. Then, to increase 30 W/min to 170 bpm or RPE of 18.
Lindqvist et al., (2003) ³²	14	5 time-points: 8, 15, 22, 29 and 36.	29 (5)	Ad hoc incremental submaximal test for 2 min of no resistances. Then, to increase 20 W every 2 min to HR max or pulse oximetry below 95%.
Lynch et al., (2003) ³³	23	16, 20, 24, 28, 32, 36	28.7(4)	Ad hoc incremental submaximal test at 60 rpm no resistance. Then, to increase 0.5 or 1 kP during two 3-min stages to 130 ± 5 bpm and 1 stage more to 145 ± 5 beats/min.
Heenan et al., (2003) ³⁴	39	37.0 (0.2)	28.5 (1.4)	1) Ad hoc incremental submaximal test at 20 W for 4 min. Then, to increase 20 W/min until 170 bpm. 2) Ad hoc, incremental ramp test from 0 W increasing work rate in 30-sec period. (70 or 110% of VT)
Pirhonen et al., (2003) ³⁵	14	5 time-points: (8, 15, 22, 29, 36)	29.2 (4.6)	Ad hoc incremental submaximal test at 0 W and 20 W for 2 min. Then, to increase at 40 W and thereafter 30 W/min to 85% HR max (220-age) or pulse oximetry below 95%.
Kardel, (2005) ³⁶	41	17, 30, 36	27.7 (1.95)	Ad hoc, incremental maximal test for 3-min stages at 50 W, 100 W and 150 W. After a rest, (no longer than 3-min) work maximally (200-280 W) for the first 30 seconds of 3-min stages.
McAuley et al., (2005) ³⁷	14	17.05 (2.05)	29.9 (0.85)	Ad hoc incremental submaximal and maximal test for 4 min at 20 W at 60-80 rpm. Then, to increase 20 W/min to 170 bpm or volitional fatigue.
Weissgerber et al., (2006) ³⁸	11	7 - 22	25-40	Ad hoc incremental submaximal test at 20 W for 4 min. Then, increasing 5 W/min until volitional fatigue or 170 bpm.
Jensen et al., (2007) ³⁹	22	3 time-points: 19.7 (1.2), 28.2 (0.3), 36.3 (0.3)	30.9 (0.9)	Idem test 1 of Heenan & Wolfe (2003).
Jensen et al., (2008) ⁴⁰	15	34-38	30.6 (1.0)	Ad hoc incremental maximal test from 6-min resting period. After 25 W/2 min at cadence of 60 and 70 rpm to the point of volitional fatigue.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Kardel et al., (2009) ⁴¹	40	35-37	20-40	Ad hoc incremental maximal test at 20 W for 2 min. Then, to increase (8-12 min) to ramp up 10% of the predicted maximal load.
Ong et al., (2009) ⁴²	12	2 time-points: 18 and 28.	30 (4)	Ad hoc, incremental submaximal test increasing 25 W/min to 75 % HR Max (220-age).
Thorell et al., (2010) ⁴³	520	4 time-points: 10.9, 24.0, 29.7, 36.5.	29.0 (4.4)	Ad hoc incremental submaximal test at 50 or 75 W (based on previous level) increasing 25 W/min to ≥ 125 bpm.
Rojas-Vega et al., (2011) ⁴⁴	20	34 \pm 1.6	35.2 (3.6)	Ad hoc incremental submaximal test free of cadence and speed for 2 min. Then, to increase 25 W/ 2 min at 60 rpm to 150 bpm.
Thorell et al., (2015) ⁴⁵	520	10.9	29.6	Idem Thorell et al. (2010).
Kim et al., (2015) ⁴⁶	32	13-35	24.8 (2.5)	Ad hoc, steady-state test with three 20-min phases:1) standing 2) pedalling at 50 W for 20 min 3) sitting.
Nakagaki et al., (2016) ⁴⁷	20	25.1(6.3)	33.7(4.2)	Ad hoc, incremental submaximal test at 50 rpm to 160 bpm or impossibility to maintain the pedalling rate.
Jedrzejko,et al., (2016) ⁴⁸	22	37-41	24.4 (3.92)	Ad hoc, incremental submaximal test on supine cycle divided into three 4-min constant stages increasing from 25 W to 75 W.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Sussman et al., (2019) ⁴⁹	23	2 time-points: 14-15 and 33-34 gw	30 (3)	YMCA protocol. Incremental test on semirecumbent to 60-80% HR _{Max} or RPE of 14 out 20.
Purdy et al., (2019) ⁵⁰	63	4 groups: 10-12, 20-27, 30-37	30.5 (4.5)	Ad hoc, incremental maximal test on recumbent cycle at 25 W at 50 rpm for 5 min. Then, to increase 25 W/min at same speed to volitional fatigue.
Bilodeau et al., (2019) ⁵¹	58	3 time-points: 16.5 (1.0), 35.6 (0.9); 39.8 (1.1) gw	30 (3.7)	Modified Bruce ramp protocol.
Matenchuk et al., (2019) ⁵²	47	4 groups: nonpregnant; 1 st trimester, 2 nd trimester, 3 rd trimester		Ad hoc, incremental maximal test at 25 W at 50 rpm for 5 min. Then, to increase 25 W/min to volitional fatigue.
Correa et al., (2020) ⁵³	48	2 time-points: 18; 36 gw.		Ad hoc, incremental ramp submaximal test at 4 W for 4 min. Then, to increase 20 W/min until symptom limitation or HR _{Max} (220-age).
Bijl et al., (2020) ⁵⁴	40	11 (1)		Ad hoc, incremental submaximal on an upright cycle ergometer for 3-min at 40rpm. Then, to increase at 60-70 rpm at 25 W followed by a rise of 5 Watt in every 12-s to 70% HR _{max} (Tanaka formula).

Treadmill protocol

Sibley et al., (1981) ⁵⁵	13	2 time-points: 21.9 (2.3); 33.9 (2.3)	24.3 (1.4)	Balke protocol to 140 bpm.
Veille, (1985) ⁴	17	35 (2)	31 (1)	Ad hoc, incremental submaximal walking test to 70% HR _{Max} (no equation to calculate HR _{max} shown).
Lewis et al., (1988) ⁵⁶	28	2 time-points: 22 wg and 30 wg	27.8 (3.3)	Modified Balke protocol.
Artal et al., (1989) ¹¹	37	30.3 (1.9)	25.9 (2.5)	Modified Balke protocol.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Clapp, Little & Capeless, (1993) ¹⁸	120	16-39	NR	1) Ad hoc, steady-state test at 40% ± 3% VO ₂ max for 30 min. 2) Idem at 60% ± 3% VO ₂ max.
Winn et al., (1994) ⁵⁷	12	26-36	32 (4)	Modified Bruce Protocol to 75% HR Max (220-age).
Marquez-Sterling et al., (2000) ⁵⁸	15	19.1 (2.15)	29.5 (3.1)	Ad hoc incremental test at 4 km/h and 0% grade for 2-min. Then, increasing 6 km/h and 2.5% every 2-min to 150 bpm.
Santos et al., (2005) ⁵⁹	72	17.9 (3.6)	27.3 (4.65)	Ad hoc, incremental ramp test from 2.4 km/h and 0% grade to AT.
Yeo et al., (2005) ⁶⁰	9	19 (5)	30 (3)	2 Cornell Protocol (85%MHR; Karvonen formula) with 2 systems (VO ₂ 000 and CPX/D).
Mottola et al., (2006) ⁶¹	156	16-22	30.8 (3.7)	Modified Balke protocol with this equation VO ₂ peak (predicted) = (0.055*peak HR) + (0.381* incline) + (5.541* speed (mph)) + (-0.090*BMI) -6.846 : incremental walking test at 3 mph for 5 min, 0% grade. Then, increase 2% every 2 min. Max inclination permitted 12% grade. Then, increasing speed 0.2 mph every 2-min to volitional fatigue.
Davenport, et al., (2008) ⁶²	106	16-20	20-39	Modified Balke protocol. Idem Mottola et al., 2006.
Oliveria et al., (2012) ⁶³	187	3 time-points: 13, 20, 28.	24.7 (5.5)	Modified Balke protocol. Idem Mottola et al. (2006).
Ruchat et al., (2012) ⁶⁴	44	2 time-points: 16-20 and 34-36	30.8 (4.2)	Modified Balke protocol. Idem Mottola et al. (2006).
Szymanski, (2012) ⁶⁵	45	30.4 (1)	33.36	Modified Balke protocol. Idem Mottola et al. (2006).
Salvesen et al., (2012) ⁶⁶	6	25.5	32	Ad hoc, incremental maximal test at 6% grade increasing speed in periods of 1km/h every 5-min to volitional fatigue.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Mottola et al., (2013) ⁶⁷	40	35.7 (0.4)	33.5 (0.7)	Ad hoc, steady-state test for 40-min, preceded by a 5-min warm-up increasing speed and inclination to 95% VT.
Bisson et al., (2013) ⁶⁸	65	16	29.9 (4.5)	Modified Balke protocol.
LeMoyné et al., (2014) ⁶⁹	67	1st trimester, 2nd trimester, 3rd trimester	29.6 (5.5) 30.1 (3.1) 32.3 (3.7)	Ebbeling single-stage submaximal treadmill walking test.
Bisson et al., (2014) ⁷⁰	61	16 (0.6)	30.0 (4.5)	Modified Balke protocol.
Marshall et al., (2015) ⁷¹	51	3 time-points: 20, 32	29.2(5.3)	Ad hoc, incremental submaximal test at 0% grade and 3.21 km/h for 5-min. Then, two 5-min stages with speed and grades self-administered to moderate (brisk walk) and vigorous (jog/run) respectively.
Santos et al., (2016) ⁷²	28	30.51 (3.3)	26 (6.9)	Modified Balke protocol.
Hesse et al., (2018) ⁷³	25	22.1 (1.4)	30 (3.6)	Bruce protocol until volitional fatigue.
Baena-García et al., (2020) ⁷⁴	127	16	32.9 (4.6)	Modified Bruce protocol until 85% HR _{Max}
Dobson et al., (2020) ⁷⁵	22	3 time-points: Early- (13–18 gw), mid- (24–28 gw) and late-pregnancy (34–37 gw).	31.4 (3.7)	Submaximal incremental Walking Exercise Test (SWET) during 21-min on a treadmill. From 3.2 km/hr at 4 min at 2% grade, to increase 2% every 3 min over seven stages.
<i>On track</i>				
Bung et al., (1991) ¹⁶	1	3 time-points(24, 28, 37)	25	Ad hoc, maximal test. 3 sprints of 200 m and one of 100 m on track

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Da Silva et al., (2010) ⁷⁶	74	37	21.5	6-minute walk test.
Ramírez-Vélez et al., (2011) ⁷⁷	64	2 time-points: 18.6 (3.4) and 16 weeks later.	19.5 (2.3)	6-minute walk test.
Hjorth et al., (2012) ⁷⁸	304	25.0 (7.3)	23.0	Ad hoc, steady-state walking test for 250 m on ground level at their normal walking pace.
Price et al., (2012) ⁷⁹	62	5 time-points: 12–14, 18–20, 24–26 and 30–32	29.05	Ad hoc test walking or running as fast as possible within comfort zone at a steady pace. Power = (weight x distance) / time.
Radzikowska et al., (2017) ⁸⁰	45	3-7	24-36	6-minute walk test.
Oviedo-Caro et al., (2018) ⁸¹	134	20	32.5 (4.2)	6-minute walk test.
Dennis et al., (2019) ⁸²	300	37 (1.3)	31 (4.2)	6-minute walk test.
Amola et al., (2019) ⁸³	34	3rd trimestre	25.1 (7.5)	6-minute walk test.
Birnbaumer et al., (2020) ⁸⁴	39	26 (7)	26 (3.4)	Ad hoc, incremental walking test on a 400 m track. Walking speed was paced by audio every 10 m and started at 3 km/h. Then, to increase 0.5 km/h every 50 m to participants were unable to walk the given pacer speed.

Step Protocol

Dibblee & Graham (1983) ⁸⁵	16	3 time-points: (the last month of each trimester)	23-31	Canadian Home Fitness Test.
Williams, Reilly et al. (1988) ⁸⁶	16 (10 pregnant)	First, second and third trimester.	25.6 (3.6)	Ad hoc, incremental test at 115, 135, and 155 bpm for 5 min.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

	and 6 non- pregnant)			
Melzer et al., (2010) ⁸⁷	44	38.27	31 (5.6)	Ad hoc, incremental test at 15-32.5 body lifts per minute (rate of change: 2.5 body lifts/ min ²). Mechanical power was calculated as: 9.81 m/s ² x step height (m) x lift frequency (number of body weight lifts/ min) and expressed in J/min/kg
<i>Muscular Fitness</i>				
Baker & Johnson (1994) ⁸⁸	200	NR	28-32	Hand Grip Sphygmomanometer Test: Pressing an inflated cuff of for 30-sec to MVCF over 3-min period.
Rogers & Tomilson (1998) ⁸⁹	20	NR	5 times: 12, 18, 24, 30, 36	Hand Grip Sphygmomanometer Test at 30% of MVCF for 2-min.
Feiner et al. (2000) ⁹⁰	34	22-36	22-35	Isometric Hand-Grip Test with dominant hand for 3 min at one-third of MVCF.
Gutke et al., (2008) ⁹¹	301	12-18	29	1) Maximal voluntary isometric hip extension test with a fixed sensor holding a sling around the thigh and pulling for 5 sec during 3 reps with 5-10-sec of rest. 2) Isometric back flexors endurance: Maintaining an abdominal crunch for a maximum of 120 sec.
Thorell et al., (2010) ⁴³	520	1 time-points: 10.9	29.0 (4.4)	Sit-up test. Supine position with the knees at a 90° angle and the feet flat on the floor. 3 sets per 5 repetitions, without a rest or to stop when they were unable to perform of 15 repetitions of sit-ups.
O'Connor et al., (2011) ⁹²	32	21-25	18-38	Ad hoc 5 tests: 1) Seated leg press; 2) Leg curls; 3) Leg extension; (4) Lat pull; (5) Back extension.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Hjorth et al., (2012) ⁷⁸	304	25.0 (7.3)	23.0	Hand-Grip maximal strength test twice on dominant and non-dominant side alternatively.
Price et al. (2012) ⁷⁹	62	5 time-points: 12–14, 18–20, 24–26 and 30–32	29.1	Ad hoc test. Lifting a 7-kg medicine ball from the floor to waist height as many times possible for 1 min.
Bisson et al. (2013) ⁶⁸	65	16	29.9 (4.5)	Hand-Grip maximal strength test twice on dominant and non-dominant side alternatively. Adjusting the handle of dynamometer.
Atay et al., (2015) ⁹³	37	2 time-points: 20 and 32	29.6 (5.9)	Hand-Grip maximal strength test in a sitting position.
Petrov et al., (2015) ⁹⁴	92	2 time-points: 13 and 35	30.7 (3.5)	Hand-Grip isometric peak strength.
Wickboldt (2015) ⁹⁵	43	32 (4)	37-42	Hand-Grip maximal strength test during the uterine contraction.
Kalliokoski et al. (2016) ⁹⁶	51	NR	28.3(6.4)	1) Hand-Grip maximal strength test for 10 sec 3-times in each hand. 2) Ad hoc upper leg performance test through 3 movements: a) To rise once after a squat b) to stand on one leg for 30 sec 3) Trendelenburg's test. It was evaluated able or unable.
Ngaka et al. (2016) ⁹⁷	50	>37	28.8 (5.7)	Hand-Grip maximal strength test in a supine position.
Rodriguez-Díaz et al., (2017) ⁹⁸	105	24-30	32.2 (4.7)	Hand-Grip maximal strength test for each hand.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Zelazniewicz, (2018) ⁹⁹	95	3 time-points (once in each trimester)	29.6 (3.4)	Hand-Grip maximal strength test twice on dominant and non-dominant side alternatively.
Takeda et al., (2019) ¹⁰⁰	21	22 and 23.25 gw.	32 (3.3)	1) Toe grip dynamometer 2) Hand-held dynamometer fixed to the legs of the chair with a belt not stretchable to assess quadriceps strength .
Baena-García et al., (2020) ⁷⁴	156	16	32.9 (4.6)	1) Hand-grip maximal strength twice on dominant and non-dominant side alternatively with 30 sec rest between them. 2) 30-sec Chair Stand Test
Yenisehir et al., (2020) ¹⁰¹	167	Second and third trimester.	28.4 (4.6)	5 Times Sit to Stand test, 5 repetitions of sit-to-stand maneuver as fast as possible with fold arms across the chest.
<i>Flexibility</i>				
Gilleard et al. (2002) ¹⁰²	21	4 time-points: 18 or less, 24, 32, 38	21-40	3 tests measured with Expert Vision™ Motion Analysis System: 1) Seated and standing forward flexion 2) Seated and standing side-to-side flexion 3) Seated axial rotation
Marnach et al. (2003) ¹⁰³	46	3 time-points: 8-12, 16-22, 34-36.	28.8 (0.8)	Wrist flexion-extension and medial-lateral deviation using goniometer
Garshasbi et al. (2005) ¹⁰⁴	212	17-22	26.4 (4.7)	Side bending test: Both sides.
rice et al., (2012) ⁷⁹	62	5 time-points: 12–14, 18–20, 24–26 and 30–32.	29.1	Sit-and-reach test.
Lindgren et al. (2014) ¹⁰⁵	200	3 time-points: 11, 24 and 36.	28.4 (5.9)	Ad hoc machine to test passive abduction of the left fourth finger.
Atay et al., (2015) ⁹³	37	2 time-points: 20 and 32,	29.6 (5.9)	Back scratch test.
Rodriguez-Díaz et al., (2017) ⁹⁸	105	24-30	32.2 (4.7)	Isquiosural flexibility test by goniometer.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Cherni et al., (2019) ¹⁰⁶	17	3 occasions: first, second and third trimester	36 (2)	4 tests measured with optoelectrical system: 1) Extensometer of the metacarpophalangeal joint of the index. 2) Figertrip to floor test: from 20cm platform, to reach the floor with knees extended; 3) Sit-and-reach test adapted on delivery bed 4) Beighton score
Baena-García et al., (2020) ⁷⁴	156	16	32.9 (4.6)	Back Scratch

Balance***Stabilometry – On force platform or pressures platform***

Butler et al., (2006) ¹⁰⁷	12	3 time-points: 11-14, 19-22, 36-39	32.9 (5.5)	Standing with eyes open and eyes-closed for 30 sec each. 3 trials. 1 piece. Force Platform.
Ribas et al., (2007) ¹⁰⁸	60	3 time-points: 1) Up to 12 week 2) 13-24 3) Upwards of 25 weeks	23.3 (4.8)	Standing with bipedal support and eyes open for 5 sec. 2 pieces at 40 Hz.
Nagai et al., (2009) ¹⁰⁹	43	30.3 (0.8)	33 (0.65)	Standing with feet parallel, gazing a black 12-cm circle fixed at a 1.5 m distance with eyes open and eyes closed for 1 min each. 1 piece.
Oliveira et al., (2009) ¹¹⁰	20	3 time-points: 15.1 (1.8); 24.0 (2.4); 34.5 (2.5)	28.7 (6.2)	Standing with 4 protocols at 50Hz and 2-min rest periods between them: 1) Eyes open with feet comfortably apart; 2) Eyes closed with feet comfortably apart; 3) Eyes open with feet together; 4) Eyes closed with feet together. 1 piece.
Karadag-Saygi et al., (2010) ¹¹¹	35	33 (3)	29.8 (4.5)	Standing for 60 sec.
Yu et al., (2013) ¹¹²	21	NR	30.2 (3.05)	Standing with heels on a line at 1.0 m from visual target with visual tasks and inspection tasks.
Ersal et al., (2014) ¹¹³	69	2 time-points: 20.9 (1.2) and 35.8 (1.5)	28.3 (5.0)	Standing with feet hip-width apart and staring straight ahead on Equitest platform.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Opala-Berdzik et al., (2014) ¹¹⁴	31	36.2 (1.2)	28.2 (3.6)	Standing with arms at both sides and in a comfortable stance on a stable force platform with eyes open and eyes closed for 2 trials of 30-sec and 1-min rest between them.
Opala-Berdzik et al., (2015) ¹¹⁵	45	2 time-points: 13.1 (2.5) and 36.2 (1.2)	28.2 (3.6)	Idem Opala-Berdzik et al., (2014)
Ozturk, (2016) ¹¹⁶	68	31.5 (4.73)	30.3 (3.6)	Standing and arms extended in 6 different positions for 32-sec: 1) facing forward eyes open and eyes closed; 2) Eyes closed head rotated at 45° to the right 3) Idem 45° to the left; 4) Eyes closed, head tilted at 30° backward and 5) Idem 30° forward; 6) Standing on an unstable cushion, facing forward eyes open and eyes closed. 4 pieces.
Shibayama et al., (2016) ¹¹⁷	161	28-33	33.3 (4.7)	Standing and feet together for 30 sec on force platform. 1 piece.
Takeda et al., (2018) ¹¹⁸	100	2 nd and 3 rd trimester	20-30	Standing with the medial malleoli 100 mm apart for 10-sec. Then, moving forward, backward, right and left for 10-sec each. 2 pieces.
Moreira et al., (2017) ¹¹⁹	30	1 st and 3 rd trimester	26.8 (5.1)	Standing with each foot positioned on each triaxial force plate (feet apart by ~20 cm) and arms along the body with eyes open focusing on a target located ~2 m in front and eyes closed for 3 trails of 60-sec each and 2-min rest. 2 pieces.
Opala-Berdzik et al., (2018) ¹²⁰	70	10.8 (1.6)	28.6 (4.4)	Standing with arms at both sides and in a comfortable stance on a stable force platform, with eyes open looking straight ahead at a wall 3m away for 2 trials of 30-sec and 1-min rest between them.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Catena et al., (2019) ¹²¹	17	9 time-points: 16-20 gw, 36-40 gw and 1 time per month up to 7 months postpartum	28.9 (4.0)	2 trials: 1) quiet static in anatomical position for 10 s on a force plate; 2) Idem 1 on a back-board spanning two force plates.
Fontana et al., (2020) ¹²²	24	23 (3)	30 (6)	Standing barefoot two-legged stance with arms at both sides with eyes open at 2 m from a cross placed on a wall at eye level during 3 x 30s trials with 30 s rest intervals. The mean was retained on force platform.
Valerio et al., (2020) ¹²³	40	30.8 (3.9)	28 (2.5)	Standing barefoot with freestanding supports inside the platform and arms by their sides. And staring at a mark on the opposite wall. 3 trials with the eyes open and three trials with eyes closed, with 30 s rest intervals.
Takeda et al., (2019) ¹⁰⁰	21	22 and 23.25 gw.	32 (3.3)	Standing barefoot on 2 stabilometers. 3 trials: 1) 10-sec standing position; 2) 10-sec moving in the anterior position; 3) 10-sec moving in the posterior position.

Others

Atay et al., (2015) ⁹³	37	2 time-points: 20 gw and 32 gw	29.6 (5.9)	One-legged stand test.
-----------------------------------	----	--------------------------------	------------	------------------------

Dynamic Balance***On platforms***

Davies et al., (2002) ¹²⁴	150	Day of labour	30.2 (5.8)	Balance Master Platform Tests: 1) Sit to Stand; 2) Walk Test, 3) Step and Quick Turn, 4) Step Up and Over.
--------------------------------------	-----	---------------	------------	------------------------------------------------------------------------------------------------------------

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Karadag-Saygi et al., (2010) ¹¹¹	35	33 (3)	29.75 (4.5)	Walking barefoot 4 m.
McCrorry et al, (2010) ¹²⁵	81	2 time-points: 20.9 (1.2) and 35.8 (1.5)	28 (5.7)	The Motor Control Test protocol with translational perturbations. Equitest posture platform.
Branco et al., (2013) ¹²⁶	22	27 (1.3)	32.5 (2.6)	Walking barefoot for 10 m between 2 points in a straight line at a natural and comfortable speed for 3 min.
Cakmak et al., (2014) ¹²⁷	41	6-12	26.5 (4.7)	Standing with knee flexed, arms placed across the chest and glare fixed ahead with open eyes on a movable platform provides up to 20° of surface tilt in a 360° range of motion for 3 trails of 20 sec each.
Inanir et al., (2014) ¹²⁸	110	3 groups: 1 st trimester, 2 nd trimester and 3 rd trimester.	24.7 (5.2)	Idem to Cakmak et al., (2014)

3-D Camera motion capture system

Wu et al., (2004) ¹²⁹	25	27	33.1	Walking on a treadmill at different velocities (incrementing 0.11 m/s, from 0.17 up to 1.72 m/s; for 3 min at each level).
Forczeck et al., (2012) ¹³⁰	13	NR	29.2 (3.5)	Walking barefoot at a self-selected speed across the room during 15 gait cycles.10
Takeda et al., (2012) ¹³¹	16	24.85 (1.95)	35 (1.4)	Stand-to-sit motion assessing the time taken to sit down; the leg joint moment; the antero-posterior and vertical floor reaction forces; and the range of motion of the lower limbs and trunk.
Gottschall et al., (2013) ¹³²	13	2 time-points: 20 and 32	31.3 (4.5)	Walking along 25 m on a custom-built portable apparatus composed of a 2.4 m ramp inclined at 15° continuous with a 4.8 m plateau.
McCrorry et al., (2014) ¹³³	69	28.35 (1.35)	28.0 (5.7)	Walking along the 8-m runway.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Krkeljas, (2018) ¹³⁴	35	3 time-points: 9-12 gw; 20-22 gw and 28-32 gw.	27 (6.1)	Walking on a straight line, at a self-selected pace along the 15-m walkway.
Catena et al., (2019) ¹³⁵	15	5 time-points: 16-20; 20-24, 24-28; 28-32; 32-36 gw.	29.3 (3.7)	60-second trial of semi-continuous stand-to-sit motion. 54 reflective markers were adhered to body land.
Catena et al., (2019) ¹³⁶	15	7 time-points: 12-16; 16-20; 20-24, 24-28; 28-32; 32-36; 36-40 gw	28.1 (4.3)	Walking on a treadmill for 60 seconds at a self-selected comfortable speed.
Forczek et al., (2019) ¹³⁷	30	3 time-points: 12, 25, 36 gw.	30.3 (3.4)	Walking barefoot at a self-selected speed during 12m intervals. 10 gait cycles.
Forczek et al., (2019) ¹³⁸	14	2 time-points: pre-pregnancy; 1 st trimester	20-40	Walking barefoot across room at a self-selected during 50 m with 1 min rest intervals. 10 gait cycles.
Catena et al., (2020) ¹³⁹	23	5 time-points: 18, 22, 26, 30, 34 gw.		Walking on a treadmill for 60 seconds at a self-selected comfortable speed.
Gimunova et al., (2020) ¹⁴⁰	41	4 time-points: 14, 28, 37 gw	30.5 (4.1)	Walking barefoot along a 6-meter walkway at a self-selected.
McCrorry et al., (2020) ¹⁴¹	95	2 time-points: 2 nd and 3 rd trimester	28.4 (5.5)	Walking along the 8m laboratory runway until walking speed stabilized.
Rothwell et al., (2020) ¹⁴²	17	2 time-points: 16-20; 36-40 gw	22-37	Walking on a treadmill for 60 seconds at a self-selected comfortable speed.
Forczek et al., (2019) ¹⁴³	36	3 time-points: 12; 25; 36 gw	30.3 (3.4)	Walking across the room 50 m with 1-min rest interval. 10 gait cycles.
Others				
Sawa et al., (2015) ¹⁴⁴	27	2 groups: early pregnancy (≤ 27 gw) or late pregnancy (≥ 27 gw)	30.9 (4.2)	Walking at self-pace speed along a 15-m smooth, horizontal corridor. It was recorded with 2 wireless motion-recording-sensor units and one piezo-resistive triaxial accelerometer.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

Błaszczyk et al., (2016) ¹⁴⁵	28	1 st trimester and 3 rd trimester	28.2 (3.4)	Walking along 10-m long walkway (back and forth 10 times) at self-space speed. It was recorded by custom made; self-adhesive copper foil electrodes attached to the soles of their shoes.
Speed				
Evensen et al., (2015) ¹⁴⁶	17		28.7 (7.4)	31.1(2.3) Ten-metres Timed walk Test (10mTWT)
Evensen et al., (2016) ¹⁴⁷	18		28.9 (7.3)	31.4 (2.7) 10mTWT
Multicomponent				
Evensen et al., (2015) ¹⁴⁶	17		28.7 (7.4)	31.1(2.3) Timed Up and Go Test (TUG)
Evensen et al., (2016) ¹⁴⁷	18		28.9 (7.3)	31.4 (2.7) TUG
Christensen et al., (2019) ¹⁴⁸	74	23	31.2 (3.7)	TUG

Ad hoc: test designed specifically for that study; NR: Not reported; PFS: Physical Fitness Score; kpm: kilopoundimeter; min: minutes; sec: seconds; HR_{Max}: Heart Rate Maximum; VO₂ max: oxygen consumption maximum; RPE: rate of perceived exertion; AT: anaerobic threshold; ICC: intraclass correlation coefficient; MVCF: maximal voluntary contraction force; HGS: hand-grip strength; m: meters; mm: millimeters; FP: force platform; PP: pressure platform.

1. Pomerance JJ, Gluck L, Lynch VA. Physical fitness in pregnancy: its effect on pregnancy outcome. *Am J Obstet Gynecol.* 1974;119(7):867-876. Accessed November 14, 2018. <https://www.sciencedirect.com/science/article/pii/0002937874900015?via%3Dihub>
2. Erkkola R. The influence of physical training during pregnancy on physical work capacity and circulatory parameters. *Scand J Clin Lab Invest.* 1976;36(8):747-754. doi:10.3109/00365517609081933
3. Morton MJ, Paul MS, Campos GR, Hart M V., Metcalfe J. Exercise dynamics in late gestation: Effects of physical training. *Am J Obstet Gynecol.* 1985;152(1):91-97. doi:10.1016/S0002-9378(85)80187-3
4. Veille J-CC, Hohimer ARR, Burry K, Speroff L. The effect of exercise on uterine activity in the last eight weeks of pregnancy. *Am J Obstet Gynecol.*

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

- 1985;151(6):727-730. doi:10.1016/0002-9378(85)90505-8
5. Jovanovic L, Kessler A, Peterson CM. Human maternal and fetal response to graded exercise. *J Appl Physiol.* 1985;58(5):1719-1722. doi:10.1152/jappl.1985.58.5.1719
 6. Wong SC, McKenzie DC. Cardiorespiratory fitness during pregnancy and its effect on outcome. *Int J Sports Med.* 1987;8(2):79-83. doi:10.1055/s-2008-1025645
 7. Kulpa PJ, White BM, Visscher R. Aerobic exercise in pregnancy. *Am J Obstet Gynecol.* 1987;156(6):1395-1403. doi:10.1016/0002-9378(87)90006-8
 8. Carpenter MW, Sady SP, Hoegsberg B, et al. Fetal heart rate response to maternal exertion. *JAMA J Am Med Assoc.* 1988;259(20):3006-3009. doi:10.1001/jama.259.20.3006
 9. Moore DH, Jarrett JC 2nd, Bendick PJ. Exercise-induced changes in uterine artery blood flow, as measured by Doppler ultrasound, in pregnant subjects. *Am J Perinatol.* 1988;5(2):94-97. doi:10.1055/s-2007-999663
 10. Sady SP, Carpenter MW, Sady MA, et al. Prediction of VO₂max during cycle exercise in pregnant women. *J Appl Physiol.* 1988;65(2):657-661. doi:10.1152/jappl.1988.65.2.657
 11. Artal R, Masaki DI, Khodiguian N, Romem Y, Rutherford SE, Wiswell RA. Exercise prescription in pregnancy: weight-bearing versus non-weight-bearing exercise. *Am J Obstet Gynecol.* 1989;161(6 Pt 1):1464-1469. doi:10.1016/0002-9378(89)90905-8
 12. Hume RFFJ, Bowie JDD, McCoy C, et al. Fetal umbilical artery Doppler response to graded maternal aerobic exercise and subsequent maternal mean arterial blood pressure: predictive value for pregnancy-induced hypertension. *Am J Obstet Gynecol.* 1990;163(3):826-829. doi:10.1016/0002-9378(90)91077-p
 13. Sady MA, Haydon BB, Sady SP, Carpenter MW, Thompson PD, Coustan DR. Cardiovascular response to maximal cycle exercise during pregnancy and at two and seven months post partum. *Am J Obstet Gynecol.* 1990;162(5):1181-1185. doi:10.1016/0002-9378(90)90012-v
 14. Field SK, Bell SG, Cenaiko DF, Whitelaw WA. Relationship between inspiratory effort and breathlessness in pregnancy. *J Appl Physiol.* 1991;71(5):1897-1902. doi:10.1152/jappl.1991.71.5.1897
 15. Rafla NM, Beazely JM. The effect of maternal exercise on fetal umbilical artery waveforms. *Eur J Obstet Gynecol Reprod Biol.* 1991;40(2):119-122. doi:10.1016/0028-2243(91)90102-Q
 16. Bung P, Huch R, Huch A. *Maternal and Fetal Heart Rate Patterns: A Pregnant Athlete during Training and Laboratory Exercise Tests; a Case Report.* Vol 39.; 1991. Accessed November 27, 2018. [https://www.ejog.org/article/0028-2243\(91\)90143-9/pdf](https://www.ejog.org/article/0028-2243(91)90143-9/pdf)

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

17. Young JC, Treadway JL. The effect of prior exercise on oral glucose tolerance in late gestational women. *Eur J Appl Physiol Occup Physiol*. 1992;64(5):130-133. doi:10.1136/bmj.39546.498796.34
18. Clapp JF, Little KD, Capeless EL. Fetal heart rate response to sustained recreational exercise. *Am J Obstet Gynecol*. 1993;168(1):198-206. doi:10.1016/S0002-9378(12)90914-X
19. Lotgering FK, Struijk PC, Doorn MBVAN, Spinnewijn WEM, Wallenburg HCS. Anaerobic threshold and respiratory compensation in pregnant women. *J Appl Physiol*. 1995;85(6):1772-1777. <http://www.ncbi.nlm.nih.gov/pubmed/7649911>
20. Artal R, Fortunato V, Welton A, et al. A comparison of cardiopulmonary adaptations to exercise in pregnancy at sea level and altitude. *Am J Obstet Gynecol*. 1995;172(4):1170-1180. doi:10.1016/0002-9378(95)91475-7
21. O'Neill ME. Maternal rectal temperature and fetal heart rate responses to upright cycling in late pregnancy. *Br J Sports Med*. 1996;30(1):32-35. doi:10.1136/bjbm.30.1.32
22. Soultanakis HN, Artal R, Wiswell RA. Prolonged exercise in pregnancy: glucose homeostasis, ventilatory and cardiovascular responses. *Semin Perinatol*. 1996;20(4):315-327. doi:10.1016/s0146-0005(96)80024-3
23. Manders MA. AM, Sonder GJ. JB, Mulder EJ. JH, Visser GH. HA. The effects of maternal exercise on fetal heart rate and movement patterns. *EARLY Hum Dev*. 1997;48(3):237-247. doi:10.1016/S0378-3782(96)01858-0
24. Kemp JG, Greer FA, Wolfe LA. Acid-base regulation after maximal exercise testing in late gestation. *J Appl Physiol*. 1997;83(2):644-651.
25. Mcgrath MJ, Wolfe LA, Preston RJ, Burggraf GW, Mcgrath MJ. Effects of pregnancy and chronic exercise on maternal cardiac structure and function. *Artic Can J Physiol Pharmacol*. Published online 1999. doi:10.1139/cjpp-77-11-909
26. Brenner IKM, Wolfe LA, Monga M, McGrath MJ. Physical conditioning effects on fetal heart rate responses to graded maternal exercise. *Med Sci Sports Exerc*. 1999;31(6):792-799. doi:10.1097/00005768-199906000-00006
27. MacPhail A, Davies GA., Victory R, Wolfe LA. Maximal exercise testing in late gestation: fetal responses. *Obstet Gynecol*. 2000;96(4):565-570. doi:10.1016/S0029-7844(00)00940-6
28. Heenan AP, Wolfe LA, Davies GA. Maximal exercise testing in late gestation: maternal responses. *Obstet Gynecol*. 2001;97(1):127-134. doi:10.1016/S0029-7844(00)01089-9
29. Kennelly MM, Geary M, McCaffrey N, McLoughlin P, Staines A, McKenna P. Exercise-related changes in umbilical and uterine artery waveforms as assessed by Doppler ultrasound scans. *Am J Obstet Gynecol*. 2002;187(3):661-666. doi:10.1067/MOB.2002.125741

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

30. Heenan AP, Wolfe LA. Plasma osmolality and the strong ion difference predict respiratory adaptations in pregnant and nonpregnant women. *Can J Physiol Pharmacol.* 2003;81(9):839-847. doi:10.1139/y03-072
31. Wolfe LA, Heenan AP, Bonen A. Aerobic conditioning effects on substrate responses during graded cycling in pregnancy. *Can J Physiol Pharmacol.* 2003;81(7):696-703. doi:10.1139/y03-059
32. Lindqvist PG, Marsal K, Merlo J, Pirhonen JP. Thermal response to submaximal exercise before, during and after pregnancy: a longitudinal study. *J Matern neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet.* 2003;13(3):152-156. doi:10.1080/jmf.13.3.152.156
33. Lynch A-MM, McDonald S, Magann EF, et al. Effectiveness and safety of a structured swimming program in previously sedentary women during pregnancy. *J Matern Neonatal Med.* 2003;14(3):163-169. doi:10.1080/jmf.14.3.163.169
34. Heenan AP, Wolfe LA, Davies GAL, McGrath MJ. Effects of human pregnancy on fluid regulation responses to short-term exercise. *J Appl Physiol.* 2003;95(6):2321-2327. doi:10.1152/jappphysiol.00984.2002
35. Pirhonen JP, Lindqvist PG, Marsal K. A longitudinal study of maternal oxygen saturation during short-term submaximal exercise. *Clin Physiol Funct Imaging.* 2003;23(1):37-41. doi:10.1046/j.1475-097X.2003.00467.x
36. Kardel KR. Effects of intense training during and after pregnancy in top-level athletes. *Scand J Med Sci Sport.* 2005;15(2):79-86. doi:10.1111/j.1600-0838.2004.00426.x
37. McAuley SE, Jensen D, McGrath MJ, Wolfe LA. Effects of human pregnancy and aerobic conditioning on alveolar gas exchange during exercise. *Can J Physiol Pharmacol.* 2005;83(7):625-633. doi:10.1139/y05-054
38. Weissgerber TL, Wolfe LA, Hopkins WG, Davies GALL. Serial respiratory adaptations and an alternate hypothesis of respiratory control in human pregnancy. *Respir Physiol Neurobiol.* 2006;153(1):39-53. doi:10.1016/j.resp.2005.09.004
39. Jensen D, Webb KA, Wolfe LA, O'Donnell DE. Effects of human pregnancy and advancing gestation on respiratory discomfort during exercise. *Respir Physiol Neurobiol.* 2007;156(1):85-93. doi:10.1016/J.RESP.2006.08.004
40. Jensen D, Webb KA, Davies GAL, O'Donnell DE. Mechanical ventilatory constraints during incremental cycle exercise in human pregnancy: implications for respiratory sensation. *J Physiol.* 2008;586(19):4735-4750. doi:10.1113/jphysiol.2008.158154
41. Kardel KR, Johansen BBBB, Voldner N, Iversen PO, Henriksen T. Association between aerobic fitness in late pregnancy and duration of labor in nulliparous women. *ACTA Obstet Gynecol Scand.* 2009;88(8):948-952. doi:10.1080/00016340903093583
42. Ong MJ, Guelfi KJ, Hunter T, Wallman KE, Fournier PA, Newnham JP. Supervised home-based exercise may attenuate the decline of glucose

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

- tolerance in obese pregnant women. *Diabetes Metab.* 2009;35(5):418-421. doi:10.1016/J.DIABET.2009.04.008
43. Thorell E, Svärdsudd K, Andersson K, Kristiansson P. Moderate impact of full-term pregnancy on estimated peak oxygen uptake, physical activity and perceived health. *Acta Obstet Gynecol Scand.* 2010;89(9):1140-1148. doi:10.3109/00016349.2010.487894
44. Vega SR, Kleinert J, Sulprizio M, Hollmann W, Bloch W, Strueder HK. Responses of serum neurotrophic factors to exercise in pregnant and postpartum women. *Psychoneuroendocrinology.* 2011;36(2):220-227. doi:10.1016/j.psyneuen.2010.07.012
45. Thorell E, Goldsmith L, Weiss G, Kristiansson P. Physical fitness, serum relaxin and duration of gestation. *BMC Pregnancy Childbirth.* 2015;15(1):1-7. doi:10.1186/s12884-015-0607-z
46. Kim J-H, Roberge RJ, Powell JB. Effect of external airflow resistive load on postural and exercise-associated cardiovascular and pulmonary responses in pregnancy: a case control study. *BMC Pregnancy Childbirth.* 2015;15(1):45. doi:10.1186/s12884-015-0474-7
47. Nakagaki A, Inami T, Minoura T, Baba R, Iwase S, Sato M. Differences in autonomic neural activity during exercise between the second and third trimesters of pregnancy. *J Obstet Gynaecol Res.* 2016;42(8):951-959. doi:10.1111/jog.12990
48. Jędrzejko M, Nowosielski K, Poręba R, Ulman-Włodarz I, Bobiński R. Physical efficiency and activity energy expenditure in term pregnancy females measured during cardiopulmonary exercise tests with a supine cycle ergometer. *J Matern Neonatal Med.* 2016;29(23):3800-3805. doi:10.3109/14767058.2016.1147550
49. Sussman D, Saini BS, Schneiderman JE, et al. Uterine artery and umbilical vein blood flow are unaffected by moderate habitual physical activity during pregnancy. *Prenat Diagn.* 2019;39(11):976-985. doi:10.1002/pd.5517
50. Purdy GM, James MA, Wakefield PK, et al. Maternal cardioautonomic responses during and following exercise throughout pregnancy. *Appl Physiol Nutr Metab.* 2019;44(3):263-270. doi:10.1139/apnm-2018-0397
51. Bilodeau J-F, Bisson M, Larose J, et al. Physical fitness is associated with prostaglandin F-2 alpha isomers during pregnancy. *PROSTAGLANDINS Leukot Essent Fat ACIDS.* 2019;145:7-14. doi:10.1016/j.plefa.2019.05.001
52. Matenchuk BA, James M, Skow RJ, et al. Longitudinal study of cerebral blood flow regulation during exercise in pregnancy. *J Cereb BLOOD FLOW Metab.* 2020;40(11):2278-2288. doi:10.1177/0271678X19889089
53. Correa M da S, Catai AM, Milan-Mattos JC, Porta A, Driusso P. Is pelvic floor muscle training able to alter the response of cardiovascular autonomic modulation and provide a possible cardiovascular benefit to pregnant women? *Neurourol Urodyn.* 2020;39(8):2272-2283. doi:10.1002/nau.24481
54. Bijl RC, Cornette MJM, van Der Ham K, et al. The physiological effect of early pregnancy on a woman's response to a submaximal cardiopulmonary exercise test. *Physiol Rep.* 2020;8(21). doi:10.14814/phy2.14624

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

55. Sibley L, Ruhling R, Ph D, Christensen C, Bolen T. Swimming and Physical fitness During pregnancy. 1981;26(6):3-12.
56. Lewis RD, Yates CY, Driskell JA. Riboflavin and thiamin status and birth outcome as a function of maternal aerobic exercise. *Am J Clin Nutr.* 1988;48(1):110-116. doi:10.1093/ajcn/48.1.110
57. WINN HN, HESS O, GOLDSTEIN I, WACKERS F, HOBBS JC. FETAL RESPONSES TO MATERNAL EXERCISE - EFFECT ON FETAL BREATHING AND BODY MOVEMENT. *Am J Perinatol.* 1994;11(4):263-266. doi:10.1055/s-2007-994588
58. Marquez-Sterling S, Perry AC, Kaplan TA, Halberstein RA, Signorile JF. Physical and psychological changes with vigorous exercise in sedentary primigravidae. *Med Sci Sports Exerc.* 2000;32(1):58-62. doi:10.1097/00005768-200001000-00010
59. Santos IA, Stein R, Fuchs SC, et al. Aerobic exercise and submaximal functional capacity in overweight pregnant women - A randomized trial. *Obstet Gynecol.* 2005;106(2):243-249. doi:10.1097/01.AOG.0000171113.36624.86
60. Yeo S, Ronis DL, Antonakos CL, Roberts K, Hayashi R. Need for population specific validation of a portable metabolic testing system: a case of sedentary pregnant women. *J Nurs Meas.* 2005;13(3):207-218. doi:10.1891/jnum.13.3.207
61. Mottola MF, Davenport MH, Brun CR, Inglis SD, Charlesworth S, Sopper MM. $\dot{V}O_2$ peak prediction and exercise prescription for pregnant women. *Med Sci Sports Exerc.* 2006;38(8):1389-1395. doi:10.1249/01.mss.0000228940.09411.9c
62. Davenport MH, Charlesworth S, Vanderspank D, Sopper MM, Mottola MF. Development and validation of exercise target heart rate zones for overweight and obese pregnant women. *Appl Physiol Nutr Metab.* 2008;33(5):984-989. doi:10.1139/H08-086
63. de Oliveria Melo AS, Silva JLPJLP, Tavares JS, Barros VO, Leite DFBB, Amorim MMRR. Effect of a Physical Exercise Program During Pregnancy on Uteroplacental and Fetal Blood Flow and Fetal Growth A Randomized Controlled Trial. *Obstet Gynecol.* 2012;120(2, 1):302-310. doi:10.1097/AOG.0b013e31825de592
64. Ruchat S-M, Davenport M, Giroux I, et al. Walking Program of Low or Vigorous Intensity During Pregnancy Confers an Aerobic Benefit. *Int J Sports Med.* 2012;33(08):661-666. doi:10.1055/s-0032-1304635
65. Szymanski LM, Satin AJ. Strenuous exercise during pregnancy: is there a limit? *Am J Obstet Gynecol.* 2012;207(3):179.e1-179.e6. doi:10.1016/j.ajog.2012.07.021
66. Salvesen K a. Å, Hem E, Sundgot-Borgen J. Fetal wellbeing may be compromised during strenuous exercise among pregnant elite athletes. *Br J Sports Med.* 2012;46(4):279-283. doi:10.1136/bjsem.2010.080259
67. Mottola MF, Inglis S, Brun CR, Hammond J-A. Physiological and metabolic responses of late pregnant women to 40 min of steady-state exercise followed by an oral glucose tolerance perturbation. *J Appl Physiol.* 2013;115(5):597-604. doi:10.1152/jappphysiol.00487.2013

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

68. Bisson M, Alméras N, Plaisance J, et al. Maternal fitness at the onset of the second trimester of pregnancy: Correlates and relationship with infant birth weight. *Pediatr Obes.* 2013;8(6):464-474. doi:10.1111/j.2047-6310.2012.00129.x
69. LeMoyné EL, Curnier D, Ellemberg D. Pregnancy and cognition: Deficits in inhibition are unrelated to changes in fitness. *J Clin Exp Neuropsychol.* 2014;36(2):178-185. doi:10.1080/13803395.2013.875520
70. Bisson M, Rhéaume C, Bujold E, Tremblay A, Marc I. Modulation of blood pressure response to exercise by physical activity and relationship with resting blood pressure during pregnancy. *J Hypertens.* 2014;32(7):1450-1457. doi:10.1097/HJH.0000000000000185
71. Marshall MR, Pivarnik JM. Perceived Exertion of Physical Activity During Pregnancy. *J Phys Act Heal.* 2015;12(7):1039-1043. doi:10.1123/jpah.2013-0458
72. Perales M, Santos-Lozano A, Ruiz JR, Lucia A, Barakat R. Benefits of aerobic or resistance training during pregnancy on maternal health and perinatal outcomes: A systematic review. *EARLY Hum Dev.* 2016;94:43-48. doi:10.1016/j.earlhumdev.2016.01.004
73. Hesse CM, Tinius RA, Pitts BC, et al. Assessment of endpoint criteria and perceived barriers during maximal cardiorespiratory fitness testing among pregnant women. *J Sports Med Phys Fitness.* 2018;58(12):1844-1851. doi:10.23736/S0022-4707.17.07750-7
74. Baena-García L, Coll-Risco I, Ocón-Hernández O, et al. Association of objectively measured physical fitness during pregnancy with maternal and neonatal outcomes. The GESTAFIT Project. *PLoS One.* 2020;15(2):e0229079. doi:10.1371/journal.pone.0229079
75. Dobson KL, da Silva DF, Dervis S, Mohammad S, Nagpal TS, Adamo KB. Physical activity and gestational weight gain predict physiological and perceptual responses to exercise during pregnancy. *Birth Defects Res.* Published online September 23, 2020:bdr2.1808. doi:10.1002/bdr2.1808
76. Da Silva EG, De Godoy I, De Oliveira Antunes LC, Da Silva EG, Peraçoli JC. Respiratory parameters and exercise functional capacity in preeclampsia. *Hypertens Pregnancy.* 2010;29(3):301-309. doi:10.3109/10641950902779271
77. Ramírez-Vélez R, Aguilar de Plata AC, Escudero MM, et al. Influence of regular aerobic exercise on endothelium-dependent vasodilation and cardiorespiratory fitness in pregnant women. *J Obstet Gynaecol Res.* 2011;37(11):1601-1608. doi:10.1111/j.1447-0756.2011.01582.x
78. Hjorth MF, Kloster S, Girma T, et al. Level and intensity of objectively assessed physical activity among pregnant women from urban Ethiopia. *BMC Pregnancy Childbirth.* 2012;12(1):154. doi:10.1186/1471-2393-12-154
79. Price BB, Amini SB, Kappeler K. Exercise in Pregnancy: Effect on Fitness and Obstetric Outcomes-A Randomized Trial. *Med Sci Sport Exerc.* 2012;44(12):2263-2269. doi:10.1249/MSS.0b013e318267ad67
80. Radzikowska E, Wiatr E, Franczuk M, Bestry I, Roszkowski-Sliz K. Lung Function in Pregnancy in Langerhans Cell Histiocytosis. In: Pokorski, M, ed. *PULMONARY DISORDERS AND THERAPY.* Vol 1023. Advances in Experimental Medicine and Biology. ; 2018:73-83.

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

doi:10.1007/5584_2017_72

81. Oviedo-Caro MA, Bueno-Antequera J, Munguía-Izquierdo D. Explanatory factors and levels of health-related quality of life among healthy pregnant women at midpregnancy: A cross-sectional study of The PregnActive Project. *J Adv Nurs*. 2018;74(12):2766-2776. doi:10.1111/jan.13787
82. Dennis AT, Salman M, Paxton E, et al. Resting Hemodynamics and Response to Exercise Using the 6-Minute Walk Test in Late Pregnancy: An International Prospective Multicentre Study. *Anesth Analg*. 2019;129(2):450-457. doi:10.1213/ANE.0000000000003818
83. Amola M, Pawara S, Kalra S. Effect of Inspiratory Muscle Training and Diaphragmatic Breathing Exercises on Dyspnea, Pulmonary Functions, Fatigue and Functional Capacity in Pregnancy during Third Trimester. *J Clin DIAGNOSTIC Res*. 2019;13(8):YC1-YC4. doi:10.7860/JCDR/2019/41732.13037
84. Birnbaumer P, Dietz P, Watson ED, et al. Absolute Accelerometer-Based Intensity Prescription Compared to Physiological Variables in Pregnant and Nonpregnant Women. *Int J Environ Res Public Health*. 2020;17(16). doi:10.3390/ijerph17165651
85. Dibblee L, Graham TEE. A longitudinal study of changes in aerobic fitness, body composition, and energy intake in primigravid patients. *Am J Obstet Gynecol*. 1983;147(8):908-914. doi:10.1016/0002-9378(83)90244-2
86. WILLIAMS A, Reilly T, CAMPBELL I, Sutherst J. Investigation of changes in responses to exercise and in mood during pregnancy. *Ergonomics*. 1988;31(11):1539-1549. doi:10.1080/00140138808966803
87. Melzer K, Schutz Y, Soehnchen N, et al. Effects of recommended levels of physical activity on pregnancy outcomes. *Am J Obstet Gynecol*. 2010;202(3):266.e1-6. doi:10.1016/j.ajog.2009.10.876
88. Baker PN, Johnson IR. The use of the hand-grip test for predicting pregnancy-induced hypertension. *Eur J Obstet Gynecol Reprod Biol*. 1994;56(3):169-172. doi:10.1016/0028-2243(94)90165-1
89. Rogers MS, Tomlinson B. Change in cardiovascular indices with position and isometric exercise throughout pregnancy: Assessment by impedance cardiography and oscillometric sphygmomanometry. *Hypertens Pregnancy*. 1998;17(2):191-202. doi:10.3109/10641959809006075
90. Feiner B, Weksler R, Ohel G, Degani S. The influence of maternal exercise on placental blood flow measured by Simultaneous Multigate Spectral Doppler Imaging (SM-SDI). *Ultrasound Obstet Gynecol*. 2000;15(6):498-501. doi:10.1046/j.1469-0705.2000.00146.x
91. Gutke A, Östgaard HC, Öberg B. Association between muscle function and low back pain in relation to pregnancy. *J Rehabil Med*. 2008;40(4):304-311. doi:10.2340/16501977-0170
92. O'Connor PJ, Poudevigne MS, Cress ME, Motl RW, Clapp JF. Safety and efficacy of supervised strength training adopted in pregnancy. *J Phys Act Health*. 2011;8(3):309-320. Accessed November 12, 2018. <http://www.ncbi.nlm.nih.gov/pubmed/21487130>

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

93. Atay E, Başalan İZ F. Investigation of the effect of changes in muscle strength in gestational age upon fear of falling and quality of life. *Turkish J Med Sci.* 2015;45(4):977-983. doi:10.3906/sag-1404-9
94. Petrov Fieril K, Glantz A, Fagevik Olsen M. The efficacy of moderate-to-vigorous resistance exercise during pregnancy: A randomized controlled trial. *Acta Obstet Gynecol Scand.* 2015;94(1):35-42. doi:10.1111/aogs.12525
95. Wickboldt N, Savoldelli G, Rehberg-Klug B. Continuous assessment of labour pain using handgrip force. *Pain Res Manag.* 2015;20(3):159-163. doi:10.1155/2015/281976
96. Kalliokoski P, Rodhe N, Bergqvist Y, Löfvander M. Long-term adherence and effects on grip strength and upper leg performance of prescribed supplemental vitamin D in pregnant and recently pregnant women of Somali and Swedish birth with 25-hydroxyvitamin D deficiency: a before-and-after treatment study. *BMC Pregnancy Childbirth.* 2016;16(1):353. doi:10.1186/s12884-016-1117-3
97. Ngaka TCC, Coetzee JFF, Dyer RAA. The Influence of Body Mass Index on Sensorimotor Block and Vasopressor Requirement During Spinal Anesthesia for Elective Cesarean Delivery. *Anesth Analg.* 2016;123(6):1527-1534. doi:10.1213/ANE.0000000000001568
98. Rodriguez-Diaz L, Ruiz-Frutos C, Maria Vazquez-Lara J, Ramirez-Rodrigo J, Villaverde-Gutierrez C, Torres-Luque G. Effectiveness of a physical activity programme based on the Pilates method in pregnancy and labour. *Enferm Clin.* 2017;27(5):271-277. doi:10.1016/j.enfcl.2017.05.007
99. Żelaźniewicz A, Pawłowski B. Maternal hand grip strength in pregnancy, newborn sex and birth weight. *Early Hum Dev.* 2018;119:51-55. doi:10.1016/j.earlhumdev.2018.03.004
100. Takeda K, Yoshikata H, Imura M. Do Squat Exercises With Weight Shift During Pregnancy Improve Postural Control? *Int J WOMENS Heal Reprod Sci.* 2019;7(1):10-16. doi:10.15296/ijwhr.2019.02
101. Yenişehir S, Çıtak Karakaya İ, Sivashoğlu AA, Özen Oruk D, Karakaya MG. Reliability and validity of Five Times Sit to Stand Test in pregnancy-related pelvic girdle pain. *Musculoskelet Sci Pract.* 2020;48:102157. doi:10.1016/j.msksp.2020.102157
102. Gilleard W, Crosbie J, Smith R. Effect of pregnancy on trunk range of motion when sitting and standing. *Acta Obstet Gynecol Scand.* 2002;81(11):1011-1020. doi:10.1034/j.1600-0412.2002.811104.x
103. Marnach ML, Ramin KD, Ramsey PS, Song S-W, Stensland JJ, An K-N. Characterization of the relationship between joint laxity and maternal hormones in pregnancy. *Obstet Gynecol.* 2003;101(2):331-335. doi:10.1016/s0029-7844(02)02447-x
104. Garshasbi A, Faghil Zadeh S. The effect of exercise on the intensity of low back pain in pregnant women. *Int J Gynecol Obstet.* 2005;88(3):271-275. doi:10.1016/j.ijgo.2004.12.001
105. Lindgren A, Kristiansson P. Finger joint laxity, number of previous pregnancies and pregnancy induced back pain in a cohort study. *BMC Pregnancy*

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

- Childbirth*. 2014;14(1):61. doi:10.1186/1471-2393-14-61
106. Cherni Y, Desseauve D, Decatoire A, et al. Evaluation of ligament laxity during pregnancy. *J Gynecol Obstet Hum Reprod*. 2019;48(5):351-357. doi:10.1016/j.jogoh.2019.02.009
107. Butler EE, Colon I, Druzin ML, Rose J. Postural equilibrium during pregnancy: Decreased stability with an increased reliance on visual cues. *Am J Obstet Gynecol*. 2006;195(4):1104-1108. doi:10.1016/j.ajog.2006.06.015
108. Ribas I S, Guirro ECO. Analysis of plantar pressure and postural balance during different phases of pregnancy. *BRAZILIAN J Phys Ther*. 2007;11(5):391-396.
109. Nagai M, Isida M, Saitoh J, Hirata Y, Natori H, Wada M. Characteristics of the control of standing posture during pregnancy. *Neurosci Lett*. 2009;462(2):130-134. doi:10.1016/j.neulet.2009.06.091
110. Oliveira LF, Vieira TMM, Macedo AR, Simpson DM, Nadal J. Postural sway changes during pregnancy: A descriptive study using stabilometry. *Eur J Obstet Gynecol Reprod Biol*. 2009;147(1):25-28. doi:10.1016/J.EJOGRB.2009.06.027
111. Karadag-Saygi E, Unlu-Ozkan F, Basgul A. Plantar Pressure and Foot Pain in the Last Trimester of Pregnancy. *FOOT ANKLE Int*. 2010;31(2):153-157. doi:10.3113/FAI.2010.0153
112. Yu Y, Chung HC, Hemingway L, Stoffregen TA. Standing body sway in women with and without morning sickness in pregnancy. *Gait Posture*. 2013;37(1):103-107. doi:10.1016/j.gaitpost.2012.06.021
113. Ersal, T., McCrory, J. L., & Sienko KH, Ersal T, McCrory JL, Sienko KH. Theoretical and experimental indicators of falls during pregnancy as assessed by postural perturbations. *Gait Posture*. 2014;39(1):218-223. doi:10.1016/j.gaitpost.2013.07.011
114. Opala-Berdzik A, Bacik B, Markiewicz A, et al. Comparison of static postural stability in exercising and non-exercising women during the perinatal period. *Med Sci Monit*. 2014;20:1865-1870. doi:10.12659/MSM.890846
115. Opala-Berdzik A, Błaszczuk JW, Bacik B, et al. Static Postural Stability in Women during and after Pregnancy: A Prospective Longitudinal Study. McCrory JL, ed. *PLoS One*. 2015;10(6):e0124207. doi:10.1371/journal.pone.0124207
116. Öztürk G, Geler Külcü D, Aydoğ E, Kaspar Ç, Uğurel B. Effects of lower back pain on postural equilibrium and fall risk during the third trimester of pregnancy. *J Matern Neonatal Med*. 2016;29(8):1358-1362. doi:10.3109/14767058.2015.1049148
117. Shibayama Y, Kuwata T, Yamaguchi J, et al. Changes in standing body sway of pregnant women after long-term bed rest. *J Obstet Gynaecol (Lahore)*. 2016;36(4):479-482. doi:10.3109/01443615.2015.1086983

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

118. Takeda K, Yoshikata H, Imura M. Changes in Posture Control of Women That Fall During Pregnancy. Published online 2018. doi:10.15296/ijwhr.2018.43
119. Moreira LS, Elias LA, Gomide AB, Vieira MF, Do Amaral WN. A longitudinal assessment of myoelectric activity, postural sway, and low-back pain during pregnancy. *Acta Bioeng Biomech.* 2017;Vol. 19(nr 3). doi:10.5277/ABB-00753-2016-02
120. Opala-Berdzik A, Błaszczuk JW, Świder D, Cieslińska-Świder J. Trunk forward flexion mobility in reference to postural sway in women after delivery: A prospective longitudinal comparison between early pregnancy and 2- and 6-month postpartum follow-ups. *Clin Biomech.* 2018;56:70-74. doi:10.1016/j.clinbiomech.2018.05.009
121. Catena RD, Campbell N, Wolcott WC, Rothwell SA. Anthropometry, standing posture, and body center of mass changes up to 28 weeks postpartum in Caucasians in the United States. *Gait Posture.* 2019;70:196-202. doi:10.1016/j.gaitpost.2019.03.009
122. Fontana Carvalho AP, Dufresne SS, de Oliveira M, et al. Effects of lumbar stabilization and muscular stretching on pain, disabilities, postural control and muscle activation in pregnant woman with low back pain. *Eur J Phys Rehabil Med.* 2020;56(3):297-306. doi:10.23736/S1973-9087.20.06086-4
123. Valerio PM, Goncalves VE, Zordao CC, Rezende MS, Dantas Moises EC, de Oliveira Guirro EC. Influence of type 1 diabetes on the postural control of women in the third gestational trimester. *Clin Biomech.* 2020;77. doi:10.1016/j.clinbiomech.2020.105062
124. Davies J, Fernando R, McLeod A, Verma S, Found P. Postural stability following ambulatory regional analgesia for labor. *Anesthesiology.* 2002;97(6):1576-1581. doi:10.1097/00000542-200212000-00033
125. McCrory JLL, Chambers AJJ, Daftary A, Redfern MSS. Dynamic postural stability during advancing pregnancy. *J Biomech.* 2010;43(12):2434-2439. doi:10.1016/j.jbiomech.2009.09.058
126. Branco M, Santos-Rocha R, Aguiar L, Vieira F, Veloso A. Kinematic analysis of gait in the second and third trimesters of pregnancy. *J Pregnancy.* 2013;2013:718095. doi:10.1155/2013/718095
127. Cakmak B, Inanir A, Nacar MC, Filiz B. The Effect of Maternity Support Belts on Postural Balance in Pregnancy. *PM&R.* 2014;6(7):624-628. doi:10.1016/j.pmrj.2013.12.012
128. Inanir A, Cakmak B, Hisim Y, Demirturk F. Evaluation of postural equilibrium and fall risk during pregnancy. *Gait Posture.* 2014;39(4):1122-1125. doi:10.1016/j.gaitpost.2014.01.013
129. Wu X, Yeoh HT. Intrinsic factors associated with pregnancy falls. *Workplace Health Saf.* 2014;62(10):403-408. doi:10.3928/21650799-20140902-04
130. Forczek W, Staszkiwicz R. Changes of kinematic gait parameters due to pregnancy. *Acta Bioeng Biomech.* 2012;14(4):113-119. doi:10.5277/abb120413

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

131. Takeda K. A Kinesiological Analysis of the Stand-to-Sit during the Third Trimester. *J Phys Ther Sci.* 2012;24(7):621-624. doi:10.1589/jpts.24.621
132. Gottschall JS, Sheehan RC, Downs DS. Pregnant women exaggerate cautious gait patterns during the transition between level and hill surfaces. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol.* 2013;23(5):1237-1242. doi:10.1016/j.jelekin.2013.04.011
133. McCrory JL, Chambers AJ, Daftary A, Redfern MS. The pregnant “waddle”: An evaluation of torso kinematics in pregnancy. *J Biomech.* 2014;47(12):2964-2968. doi:10.1016/j.jbiomech.2014.07.009
134. Krkeljas Z. Changes in gait and posture as factors of dynamic stability during walking in pregnancy. *Hum Mov Sci.* 2018;58:315-320. doi:10.1016/j.humov.2017.12.011
135. Catena RD, Bailey JP, Campbell N, Music HE. Stand-to-sit kinematic changes during pregnancy correspond with reduced sagittal plane hip motion. *Clin Biomech.* 2019;67:107-114. doi:10.1016/j.clinbiomech.2019.05.014
136. Catena RD, Campbell N, Werner AL, Iverson KM. Anthropometric Changes During Pregnancy Provide Little Explanation of Dynamic Balance Changes. *J Appl Biomech.* 2019;35(3):232-239. doi:10.1123/jab.2018-0345
137. Forczek W, Ivanenko Y, Curyło M, et al. Progressive changes in walking kinematics throughout pregnancy-A follow up study. *Gait Posture.* 2019;68:518-524. doi:10.1016/j.gaitpost.2019.01.004
138. Forczek W, Masłoń A, Frączek B, Curyło M, Salamaga M, Suder A. Does the first trimester of pregnancy induce alterations in the walking pattern? *PLoS One.* 2019;14(1):e0209766. doi:10.1371/journal.pone.0209766
139. Catena RD, Bailey JP, Campbell N, Stewart BC, Marion SJ. Correlations between joint kinematics and dynamic balance control during gait in pregnancy. *Gait Posture.* 2020;80:106-112. doi:10.1016/j.gaitpost.2020.05.025
140. Gimunova M, Zvonar M, Sebera M, Turcinek P, Kolarova K. Special footwear designed for pregnant women and its effect on kinematic gait parameters during pregnancy and postpartum period. *PLoS One.* 2020;15(5). doi:10.1371/journal.pone.0232901
141. McCrory JL, Chambers AJ, Daftary A, Redfern MS. Torso kinematics during gait and trunk anthropometry in pregnant fallers and non-fallers. *Gait Posture.* 2020;76:204-209. doi:10.1016/j.gaitpost.2019.11.012
142. Rothwell SA, Eckland CB, Campbell N, Connolly CP, Catena RD. An analysis of postpartum walking balance and the correlations to anthropometry. *Gait Posture.* 2020;76:270-276. doi:10.1016/j.gaitpost.2019.12.017
143. Forczek W, Ivanenko Y, Salamaga M, et al. Pelvic movements during walking throughout gestation - the relationship between morphology and kinematic parameters. *Clin Biomech.* 2020;71:146-151. doi:10.1016/j.clinbiomech.2019.11.001

Assessing physical fitness during pregnancy: validity and reliability of fitness tests, and relationship with health-related outcomes. A systematic review.

144. Sawa R, Doi T, Asai T, Watanabe K, Taniguchi T, Ono R. Differences in trunk control between early and late pregnancy during gait. *Gait Posture*. 2015;42(4):455-459. doi:10.1016/j.gaitpost.2015.07.058
145. Opala-Berdzik A, Bacik B, Markiewicz A, et al. Comparison of Static Postural Stability in Exercising and Non-Exercising Women During the Perinatal Period. *Med Sci Monit*. 2014;20:1865-1870. doi:10.12659/MSM.890846
146. Evensen NM, Kvale A, Braekken IH, Kvåle A, Braekken IH. Reliability of the Timed Up and Go test and Ten-Metre Timed Walk Test in Pregnant Women with Pelvic Girdle Pain. *Physiother Res Int*. 2015;20(3):158-165. doi:10.1002/pri.1609
147. Evensen NM, Kvåle A, Brækken IH. Convergent validity of the Timed Up and Go Test and Ten-metre Timed Walk Test in pregnant women with pelvic girdle pain. *Man Ther*. 2016;21:94-99. Accessed November 10, 2018. <https://www.sciencedirect.com/science/article/pii/S1356689X15001319>
148. Christensen L, Vollestad NK, Veierod MB, Stuge B, Cabri J, Robinson HS. The Timed Up & Go test in pregnant women with pelvic girdle pain compared to asymptomatic pregnant and non-pregnant women. *Musculoskelet Sci Pract*. 2019;43:110-116. doi:10.1016/j.msksp.2019.03.006