

新しいサイトカイン・IL8とMCAFの基礎ならびに前臨床実験

著者	松島 綱治
著者別表示	Matsushima Kouji
雑誌名	平成4(1992)年度 科学研究費補助金 国際学術研究 研究成果報告書概要
巻	1991 1992
ページ	3p.
発行年	1994-03-23
URL	http://doi.org/10.24517/00067401



1992 Fiscal Year Final Research Report Summary

Basic and preclinical experiments of IL 8 and MCAF

Research Project

Project/Area Number

03044066

Research Category

Grant-in-Aid for international Scientific Research

Allocation Type

Single-year Grants

Section

Joint Research

Research Institution

Cancer Research Institute, Kanazawa University

Principal Investigator

MATSUSHIMA Kouji Professor, Cancer Research Institute, Kanazawa University, がん研究所, 教授 (50222427)

Co-Investigator(Kenkyū-buntansha)

CHRISTIAN G. デンマークオーフス大学, 医学部, 臨床研究員

ETTORE APPEL 米国国立がん研究所, 主任研究員

JOOST J. OPP 米国国立がん研究所, 研究室主任

MUKAIDA Naofumi Associate Prof., Cancer Research Institute, Kanazawa University, がん研究所, 助教授 (30182067)

KASHARA Tadashi Associate Prof., Jichi Medical School, 医学部, 助教授 (60049096)

MURAKAMI Seishi Associate Prof., Cancer Research Institute, Kanazawa University, がん研究所, 助教授 (90019878)

OPPENHEIM Joost J. Laboratory Chief, National Cancer Institute, USA

LARSEN Christian Clinical Staff Fellow, Aarhus University School of Medicine, DK

APPELLA Ettore Section Chief, National Cancer Institute, USA

Project Period (FY)

1991 - 1992

Keywords

Leukocyte Chemotactic factors / cytokine / interleukin 8 / MCAF / structural analysis / receptors / gene expression / pathophysiological roles

Research Abstract

BASIC AND PRECLINICAL EXPERIMENTS OF IL 8 AND MCAF

Novel chemotactic cytokines, IL 8 and MCAF were identified, purified, and molecularly cloned by us at National Cancer Institute, USA in 1987-1989, belong a family of emerging chemotactic cytokine family, CHEMOKINE. In this international collaborative research, the following projects were performed.

- 1) Structural analysis of IL 8/MCAF and determining their active site(s)-----Human IL 8 was expressed in E coli and purified to homogeneity in a large scale. The structure of IL 8 was analyzed by both NMR and X-ray crystallography. The proposed model shows that IL 8 exists as dimer through hydrogen bonding and two symmetry-related anti-parallel α -helices, 24Å long and separated by 14Å, lie on top of a six-stranded anti-parallel β -sheet platform. The active site of IL 8 was presumed to be the region surrounding His at 33 as well as N-terminal region based on the mutation studies and structural analyses.
- 2) Structure of the receptors for IL 8 and MCAF-----The MW of the receptors for IL 8 on human neutrophils was estimated to be 60,000 with Kd of 1nM. The MW of the receptors for MCAF on human monocytes was estimated to be 40,000 with Kd of 25nM. Although the cDNA for IL 8 was cloned by other groups, we have cloned several novel cDNA which belong to the family of chemotactic cytokine receptors.
- 3) Molecular analysis of the regulation of the production of IL 8-----After cDNA cloning of human IL 8, we revealed the induction of the production of IL 8 by various types of cells after stimulating with IL 1, TNF, endotoxin, exotoxin, viral proteins, heavy metals, and superoxide generating substances. We also showed the suppression of the production of IL 8 by glucocorticoids, vitamin D3, lipoxygenase inhibitors, and FK506. We have determined the enhancer of the human IL 8 gene to be consisted of AP-1+NFKB. IL 8 production suppressive agents seem to inhibit the activation of nuclear factors which bind to these regions.
- 4) Establishment of pathophysiological roles of IL 8/MCAF and examining possible clinical application of IL 8/MCAF-----We have revealed the production of IL 8/MCAF in various human inflammatory diseases including ulcerative colitis, psoriasis, glomerulonephritis, urinary tract infection, and arthritis. Essential involvement of IL 8 in recruiting neutrophils in acute inflammation models in rabbits, such as dermatitis, arthritis, and lung reperfusion was established using monoclonal neutralizing antibody against rabbit IL 8. IL 8/MCAF cDNA transfection into tumor cells showed significant anti-tumor effect of these cytokines in mice. MCAF is further shown to prime murine macrophages to be tumor cytotoxic and also have anti-infectious activity in mice. Possible hematopoietic activity of these cytokines was not proved either in vitro or in vivo.▲ Less

Research Products (15 results)

All Other

All Publications (15 results)

- [Publications] Mahe, Y. et al.: "Hepatitis B virus X protein transactivates human interleukin 8 gene" *J. Biol. Chem.* 266. 13759-13763 (1991) ▼
- [Publications] Smyth, M. J., et al.: "Interleukin 8 gene expression and production in human peripheral blood lymphocyte subsets." *J. Immunol.* 146. 3815-3823 (1991) ▼
- [Publications] Ko, Y., et al.: "A sensitive enzyme-linked immunosorbent assay for human interleukin 8." *J. Immunol. Methods.* 149. 227-235 (1992) ▼
- [Publications] YASUMOTO, K., et al.: "Tumor necrosis factor alpha and interferon gamma synergistically induce interleukin 8 production in a human gastric cancer cell line" *J. Biol. Chem.* 267. 22506-22511 (1992) ▼
- [Publications] Thompson, H. L., MATSUSHIMA, K.: "Human polymorphonuclear leukocytes stimulated by tumor necrosis factor-alpha show increased adherence to extracellular matrix proteins" *Clin. Exp. Immunol.* 90. 280-285 (1992) ▼
- [Publications] Ko, Y., et al.: "Elevated IL-8 level in the urine of patients with urinary tract infections." *Infect. Immun.* (1993) ▼
- [Publications] Oppenheim, J. J., Zachariae, C. O. C., Mukaida, N., Matsushima, K.: "Properties of the novel proinflammatory supergene"intercrine"cytokine family." *Annu. Rev. Immunol.*, 31 (1991) ▼
- [Publications] MATSUSHIMA, K., BALDWIN, E. T., MUJAJIDA, N.: "Chemical Immunol. In INTERLEUKINS: MOLECULAR BIOLOGY AND IMMUNOLOGY Kishimoto, T. Ed. Novel leukocyte recruitment and activating cytokines." Karger Publ., Basel, SW, 29 (1992) ▼
- [Publications] Mahe, Y. et al.: "Hepatitis B virus X protein transactivates human interleukin 8 gene" *J. Biol. Chem.* 266. 13759-13763 (1991) ▼
- [Publications] Smyth, M. J., et al.: "Interleukin 8 gene expression and production in human peripheral blood lymphocyte subset" *J. Immunol.* 146. 3815-3823 (1991) ▼
- [Publications] Ko, Y., et al.: "A sensitive enzyme-linked immunosorbent assay for human interleukin 8" *J. Immunol. Methods.* 149. 227-235 (1992) ▼
- [Publications] Yasumoto, K., et al.: "Tumor necrosis factor alpha and interferon gamma synergistically induce interleukin 8 production in a human gastric cancer cell line" *J. Biol. Chem.* 267. 22506-22511 (1992) ▼
- [Publications] Ko, Y., et al.: "Elevated IL 8 level in the urine of patients with urinary tract infections" *Infect. Immun.* ▼
- [Publications] Oppenheim, J. J. et al.: *Annu. Rev. Immunol.* Properties of the novel proinflammatory supergene"intercrine"cytokine family, 617-648 (1991) ▼

[Publications] Matsushima, K., et al.: Kishimoto, T.Edt. "Novel leukocyte recruitment and activating cytokines". Chemical Immunology, In INTERLEUKINS : MOLECULAR BIOLOGY AND IMMUNOLOGY, 236-265 (1992)



URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-03044066/030440661992kenkyu_seika_hokoku

Published: 1994-03-23