CLUSTERING OF LOCAL GROUP DISTANCES: PUBLICATION BIAS OR CORRELATED MEASUREMENTS? VII. A DISTANCE FRAMEWORK OUT TO 100 MPC

RICHARD DE GRIJS^{1,2,3} AND GIUSEPPE BONO^{4,5}

¹Department of Physics & Astronomy, Macquarie University, Balaclava Road, Sydney, NSW 2109, Australia

²Research Centre for Astronomy, Astrophysics & Astrophotonics, Macquarie University, Balaclava Road, Sydney, NSW 2109, Australia

³International Space Science Institute–Beijing, 1 Nanertiao, Zhongguancun, Hai Dian District, Beijing 100190, China

 4 Dipartimento di Fisica, Università di Roma Tor Vergata, via Della Ricerca Scientifica 1, 00133, Roma, Italy

⁵INAF, Rome Astronomical Observatory, via Frascati 33, 00078 Monte Porzio Catone, Italy

ABSTRACT

We consider the body of published distance moduli to the Fornax and Coma galaxy clusters, with specific emphasis on the period since 1990. We have carefully homogenized our final catalogs of distance moduli onto the distance scale established in the previous papers in this series. We assessed systematic differences associated with the use of specific tracers, and consequently discarded results based on application of the Tully–Fisher relation and of globular cluster and planetary nebula luminosity functions. We recommend 'best' weighted relative distance moduli for the Fornax and Coma clusters with respect to the Virgo cluster benchmark of $\Delta(m-M)_0^{\text{Fornax-Virgo}} = 0.18 \pm 0.28$ mag and $\Delta(m-M)_0^{\text{Coma-Virgo}} = 3.75 \pm 0.23$ mag. The set of weighted mean distance moduli (distances) we derived as most representative of the clusters' distances is,

$$(m - M)_0^{\text{Formax}} = 31.41 \pm 0.15 \text{ mag} (D = 19.1^{+1.4}_{-1.2} \text{ Mpc}) \text{ and}$$

= 31.21 ± 0.28 mag (D = 17.5^{+2.4}_{-2.2} \text{ Mpc});
(m - M)_0^{\text{Coma}} = 34.99 \pm 0.38 \text{ mag} (D = 99.5^{+19.0}_{-15.9} \text{ Mpc}) \text{ and}
= 34.78 ± 0.27 mag (D = 90.4^{+11.9}_{-10.6} \text{ Mpc}),

where the first distance modulus for each cluster is the result of our analysis of the direct, absolute distance moduli, while the second modulus is based on distance moduli relative to the Virgo cluster. While the absolute and relative distance moduli for both clusters are mutually consistent within the uncertainties, the relative distance moduli yield shorter absolute distances by $\sim 1\sigma$. It is unclear what may have caused this small difference for both clusters; lingering uncertainties in the underlying absolute distance scale appear to have given rise to a systematic uncertainty on the order of 0.20 mag.

Keywords: Astronomical reference materials — Astronomy databases — Distance measure — Galaxy distances — Coma Cluster

1. A ROBUST FRAMEWORK OF EXTRAGALACTIC DISTANCES

Over the course of the past decade, we have established a robust distance framework to galaxies in the Local Group and beyond, based on a set of mutually and internally consistent distance moduli that were validated on a robust statistical basis. In de Grijs et al. (2014, henceforth Paper I), we explored the presence of 'publication bias' in the body of published distance determinations for the Large Magellanic Cloud (LMC), as suggested by Schaefer (2008). While we did not find any evidence of authors have jumped on this proverbial bandwagon, we put Freedman et al. (2001)'s canonical LMC distance modulus of $(m - M)_0^{\text{LMC}} = 18.50 \pm 0.10$ mag on a well-established statistical footing, recommending $(m - M)_0^{\text{LMC}} = 18.49 \pm 0.09 \text{ mag}$ (see also Crandall & Ratra 2015).

This was followed by a series of papers aimed at both exploring the reality and/or the presence of publication bias among published distance measurements and establishing a robust local distance framework. In order of increasing distance, we applied the same analysis as developed in Paper I to the Galactic Center (de Grijs & Bono 2016, Paper IV) and the Galactic rotation constants (de Grijs & Bono 2017, Paper V), the Small Magellanic Cloud (de Grijs & Bono 2015, Paper III; see also Crandall & Ratra 2015), the M31 group (de Grijs & Bono 2014, Paper II), and the Virgo cluster (de Grijs & Bono 2020, Paper VI). In addition, we strongly recommend the independent, geometric distance measurement to the maser host galaxy NGC 4258 published by Herrnstein et al. (1999) as additional, intermediatedistance stepping stone. Our full, internally consistent distance framework thus far established is summarized in Table 3 of Paper VI.

In this paper, we expand our previous analyses by focusing on two additional, rich benchmark galaxy clusters. These include the Fornax cluster as southernhemisphere benchmark counterpart to the Virgo cluster in the northern hemisphere, as well as the Coma cluster. This takes our internally consistent 'local' distance framework out to distances of order 100 Mpc. At those distances, a significant fraction (although still a large minority) of articles citing distance estimates refer to redshifts rather than linear scales. This will therefore conclude our efforts to establish a benchmark set of statistically validated distance estimates in the local Universe.

We have organized this paper as follows. In Section 2 we briefly summarize our data-mining approach and describe the resulting catalogs containing Fornax and Coma distances. We examine trends in distance determinations for both clusters in Section 4. Then, in Section 3, we analyze the systematic differences, if any, among tracer populations for both clusters. This eventually results in a set of recommended benchmark distances, which we summarize in Section 5. These should be combined with the distance moduli summarized in Paper VI (see that paper's Table 3) to gain a full overview of our extragalactic distance framework.

2. OUR DATABASE

Similarly to the previous papers in this series, we mined the NASA/Astrophysics Data System (ADS) for newly derived or recalibrated/updated distance measures to the Fornax and Coma clusters. We used as search terms 'Fornax Cluster' and 'Coma Cluster.' We adopted the same criteria as in our previous papers for inclusion of any new values in our final database (for a detailed description, see, e.g., Paper I). We included both measurements to the galaxy clusters as a whole, as well as to individual galaxies in the cluster cores (indicated separately in our final catalogs). For the Fornax cluster, we hence included distance measures to NGC 1316, NGC 1326A, NGC 1365, and NGC 1399 (as well as NGC 1404). The relevant galaxies in the Coma cluster for which distance moduli are included in our final database are NGC 4874, NGC 4881, NGC 4889, NGC 4921, NGC 4923, and IC 4051.

For the 'modern' period, from 1990 onward, we carefully perused all articles resulting from our NASA/ADS queries; prior to 1990, we followed the reference trail. As of 2019 December 19, when we completed our data mining, the numbers of hits in the NASA/ADS for the Fornax and Coma clusters were 1849 and 5357, respectively. The resulting numbers of absolute/relative distance measures included in our final catalogs are 140/62 and 95/56 for the Fornax and Coma clusters, respectively. For the Fornax cluster, we retrieved distance moduli relative to that of the Virgo cluster; for the Coma cluster, we retrieved relative distance measures with respect to the Virgo (53), Leo I (2 direct, 10 indirect; see Section 4.4), and Fornax (1) clusters.

Our final database, sorted by year and by tracer for both galaxy clusters separately, is available online through http://astro-expat.info/Data/pubbias.html,¹ For graphical depictions of the clusters' distance moduli as a function of year of publication, see Figures 1 and 2. We will discuss the trends and any evidence of publication bias versus correlated measures in Section 3.

3. TRENDS IN FORNAX AND COMA CLUSTER DISTANCE DETERMINATIONS

Figures 1 and 2 show both the overall distribution of distance moduli and the distance measures for selected individual tracers for the Fornax and Coma clusters, respectively. The bottom panels in both figures show the sets of published relative distance moduli of our target clusters with respect to the Virgo cluster.

Since we have focused our detailed data mining on the 'modern' period from 1990 onward, we will concentrate on those measures in our analysis of the distances implied by the individual tracers. However, careful assessment of both figures shows that hidden trends in the distance moduli may be present for Fornax distance measures based on application of the Tully–Fisher relation (TFR), surface brightness fluctuations (SBF), and the planetary nebula luminosity function (PNLF). Hence, for the Fornax distances based on the TFR and SBF, we analyzed the period from 2000 onward, while for PNLFbased measures we used the timeframe since 1995. For the Coma cluster, we restricted our analysis of its TFRbased distances to the period from 2000 given the large scatter of the individual data points prior to 2000, which may imply the presence of unaccounted-for biases and measurement errors. Note that all data points included in Figures 1 and 2 reflect the assumptions made by their original authors. In Section 4 we will first homogenize the data points used for further analysis before drawing our conclusions.

¹ A permanent link to this page can be found at http://web.archive.org/web/20200331174040/http://astro-expat.info/Data/pubbias.html; members of the community are encouraged to send us updates or missing information.




Figure 1. Published distance moduli (original values and original error bars, where available) to the Fornax cluster and its central galaxies, NGC 1316, NGC 1326A, NGC 1365, and NGC 1399 (as well as NGC 1404). The horizontal dotted lines in panels (a)–(e) represent $(m - M)_0 = 31.4$ mag and are meant to guide the eye. GCLF: Globular cluster luminosity function; TFR: Tully–Fisher relation; TRGB: Tip of the red-giant branch; SBF: Surface brightness fluctuations; SNe Ia: Type Ia supernovae. 'Averages' in panel (a) include weighted and unweighted means of different methods of distance determination, as well as mean values of the distance moduli to samples of central Fornax cluster galaxies, as published by the original authors (see Section 10 in our externally linked data table); 'Uncategorized' distance moduli include any measurements that are not already included in the other panels, mostly because of the scarcity of data points for a particular measurement approach (see Section 11 of the same table). Panel (f) shows the set of published relative Fornax–Virgo Cluster distance moduli (any tracer), where positive values reflect a greater distance to the Fornax cluster compared with Virgo.

Although we retain the Fornax distance modulus suggested by Drinkwater et al. (2001) in our online catalog (http://astro-expat.info/Data/pubbias.html), we did not include this measurement in our analysis. It is based on the mean distance modulus resulting from three Cepheid-based measurements available at the time of publication, to NGC 1365, NGC1326A, NGC 1425. These authors suggest that neither NGC 1326A nor NGC 1425 may be representative of the cluster core (but note that our final catalog retains the Cepheid distance to NGC 1326A, since it does not appear out of place with respect to the overall body of measurements), thus only leaving the Cepheid distance to NGC 1365 (Madore et al. 1999). The latter is already included in our Fornax catalog, thus rendering the Drinkwater et al. (2001) result superfluous.

Similarly, we retain two distances based on globular cluster luminosity functions (GCLFs) in our final catalog of Fornax cluster distance measures, which we however do not use in our analysis. This relates to the distance moduli of Whitmore (1997) and Dirsch et al. (2003). The former reference does not provide sufficient information for us to include its distance determination in our analysis; the latter distance determination is, in essence, based on an assessment of the measured peak of the GCLF as being 'consistent' with the SBF-based distance measurements of Tonry et al. (2001) and Liu et al. (2002). As such, it is not a firm determination.

4. STATISTICALLY VALIDATED DISTANCES

4.1. Systematic differences affecting individual tracers

In the previous section, we identified a number of individual distance tracers for both galaxy clusters for which we had collected a sufficient number of measurements to analyze their distance distributions separately. We show their distance measures separately in panels (b)–(e) in both Figures 1 and 2. Table 1 includes the weighted means for each tracer and for the relevant period of interest, as well as the relevant 1σ uncertainties.

However, before compiling Table 1, we ensured that all distance moduli contributing to the weighted means were carefully homogenized onto our overall distance framework as determined in this series of papers. Tables 2 and 3 include the numerical basis of this homogenization, showing both the zero-point calibrations used by the original authors and our adjustments of their distance moduli to match our distance framework thus far established (see footnote a to Table 2 for a quick overview). The homogenized distance moduli (as well as the original values) are displayed as a function of publication date and for each of the main tracers separately in Figure 3.

From a sociological perspective, we note that distance measures using a specific tracer are often dominated by articles published by the same group and their junior team members. This is not unexpected, of course, since this practice reflects the central expertise of the different groups of authors. Nevertheless, we considered the effects of including series of results from the same group on the overall value of the resulting distance modulus.

4.2. Fornax

First, we considered the post-1990 Cepheid-based distance moduli for the Fornax cluster. Comparing author lists, combined with a careful perusal of the papers in question, it is clear that among our set of 13 Cepheid distances to the Fornax cluster, the only truly independent measurement was provided by Riess et al. (2016), i.e., $(m - M)_0 = 31.21 \pm 0.06$ mag (after adjustment of its zero-point calibration). Their value falls within the mutual 1σ uncertainties of the Cepheid-based distance included in Table 1; one should, of course, keep in mind that all other measurements contributing to that distance are correlated and not independent.

Among the eight post-2000 TFR-based distances to the Fornax cluster, five were published by the same group. The weighted average of those five determinations is $(m - M)_0 = 31.22 \pm 0.25$ mag. Of the remaining three values, two fall comfortably within the 1σ uncertainties following homogenization: $(m - M)_0 =$ 31.21 ± 0.13 mag (Bernardi et al. 2002) and $(m - M)_0 =$ 31.20 ± 0.33 mag (Allen & Shanks 2004). The third value, $(m - M)_0 = 31.45 \pm 0.14$ mag (Masters et al. 2006), is larger although still consistent with the bulk of our values.

Nevertheless, inspection of Table 1 shows that the TFR-based weighted mean distance to the Fornax cluster is significantly shorter than the equivalent distance estimates based on most other tracers (with the notable exception of GCLF-based distance moduli; see below). A combination of effects may have given rise to this difference. First, Allen & Shanks (2004) concluded that for their sample of 18 galaxies at distances of $(m-M)_0 > 29.5$ mag hosting Cepheid variables and which were observed with the Hubble Space Telescope, the TFR-based distance moduli yield distances that are shorter by $\Delta(m - M)_0 = -0.44 \pm 0.09$ mag (see also Shanks 1997). They suggested that at least some of this effect may be owing to unaccounted-for metallicity differences (e.g. Kennicutt et al. 1998) and sample incompleteness, leading to a significant scale error in TFRbased distances. We note, however, that addition of the offset in distance modulus suggested by Shanks (1997) and Allen & Shanks (2004) would lead to overestimated TFR distances compared with Fornax distance moduli based on other tracers. This situation is exacerbated by the often convoluted calibration approaches, often involving at least some iterative means to tie Cepheid, Type Ia supernovae (SNe Ia), and TFR distances to the same scale.

In addition, TFR distances tend to differ depending on the operating wavelength. Tully & Pierce (2000) found for the Ursa Major galaxy cluster that although the overall agreement among the distance moduli resulting from analysis of different passbands is good, their *I*band analysis yielded shorter distance moduli than the weighted mean by 0.02 mag, while in the *B* band their moduli were overestimated by 0.04 mag. In the *R* and K' bands, their estimates were 0.03 mag larger and 0.05 mag smaller, respectively, than the mean. Finally, a de-




Figure 2. As Figure 1, but for the Coma cluster and its central galaxies, NGC 4874, NGC 4881, NGC 4889, NGC 4921, NGC 4923, and IC 4051. Fund. Plane: Fundamental Plane. Panel (f) shows the relative Coma–Virgo cluster distance moduli (as in Figure 1f).

gree of publication bias could have crept into our sample of TFR-based Fornax cluster distance measures, given that some authors confidently state that their derived TFR-based distance moduli comfortably agree with previously published measures, but without comparing the underlying calibrations applied (e.g. Bureau et al. 1996).

Our SBF-based distance measures to Fornax represent the largest subsample. Although they do exhibit some spread about the weighted mean, the distribution's standard deviation is small, 0.14 mag, and therefore not indicative of statistical anomalies. Nevertheless, 21 of the 28 post-2000 values considered here were published by the same team (and are, hence, likely correlated). However, the remaing seven values (Ferrarese et al. 2000; Liu et al. 2002; Jerjen 2003; Dunn & Jerjen 2006) are all fully consistent with the overall weighted mean and its 1σ uncertainty. Both data sets are statistically indistinguishable. We note that our three TRGB-based Fornax distances were all published by the same team, but we included them in our analysis because they provide an independent stellar population tracer. Both the SBF technique and the TRGB-based distances rely on

Fornax				Coma				
Tracer	Period	N	$(m - M)_0$	Tracer	Period	N	$(m - M)_0$	
			(mag)				(mag)	
$Cepheids^*$	1990 - 2019	13	31.38 ± 0.14	Hubble law ^{a}	1990 - 2019	8	35.02 ± 0.06	
TFR	2000 - 2019	8	31.25 ± 0.24	TFR^*	2000 - 2019	33	34.72 ± 0.18	
SBF	2000 - 2019	28	31.44 ± 0.19	SBF^*	1990 - 2019	7	34.98 ± 0.37	
TRGB	1990 - 2019	3	31.41 ± 0.09	GCLF^*	1990 - 2019	15	34.90 ± 0.17	
GCLF^*	1990 - 2019	19	31.22 ± 0.21					
PNLF	1995 - 2019	5	31.36 ± 0.09					

 Table 1. Mean, post-1990 published distance measures to the centers of the Fornax and Coma Clusters as a function of tracer population.

^a Adopting $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$; a reduction (increase) to $H_0 = 67.3$ (72) km s⁻¹ Mpc⁻¹ (spanning the range implied by the current 'Hubble tension') would result in an increase (decrease) in the Coma distance modulus by 0.09 (0.06) mag. The uncertainty represents the Gaussian width (σ) of the distribution, given that none of the published Coma distances based on application of the Hubble law included estimates of the associated uncertainties.

* Since a number of published distance measures did not include associated uncertainties, the central values are based on the full set of published measurements, while the uncertainties only include those values that were published with their associated uncertainties. This affects the following numbers of measurements: (1) Fornax – Cepheids: 1; GCLF: 3; (2) Coma – GCLF: 3; SBF: 1; TFR: 8.

red-giant stars; it is therefore comforting to see that the distance moduli resulting from independent application of these techniques are indeed very close to one another.

The GCLF-based data set comprises 19 distance moduli. The author lists of the contributing papers are more diverse than for the previously discussed tracers. The largest single group of collaborators contributed to nine of the measures included in our catalog. Overall, assuming a Gaussian distribution of distance measures, the 1σ spread is 0.19 mag, similar to the uncertainty on the mean. This indicates that the intrinsic spread among the contributing values is more significant than the equivalent spreads resulting from the other tracers used here. Some of the most significantly deviating values result from calibrations using non-standard calibrators, specifically the poorly defined B- or I-band GCLFs pertaining to the Milky Way or M31 (e.g. Kohle et al. 1996; Ferrarese et al. 2000; Gómez et al. 2001). In addition, calibration of the (V-band) GCLF in the Milky Way relies on accurate distance determinations to the contributing globular clusters (and, in fact, a reliable distance determination to the Galactic Center; see Paper IV). Moreover, one must make assumptions regarding the shape (width) of the GCLF, which may differ among different galaxy types (e.g. Jordán et al. 2006).

Finally, as we discussed in the context of the distance to the Galactic Center (Paper IV), distances based on GCLFs tend to be systematically smaller than most other distance measurements (for the Fornax cluster, see also Villegas et al. 2010). This could be caused by incomplete corrections for internal or foreground extinction, or because of incomplete samples of objects, in the sense that our observational data may be biased toward objects in the foreground of the host galaxy. In view of these lingering uncertainties, we will refrain from further consideration of the GCLF-based distances.

The five articles yielding PNLF-based distances to Fornax were published by four different groups, yet following homegenization the weighted mean is welldefined with a small uncertainty (0.09 mag). Nevertheless, we are reluctant to place too much emphasis on this result, given that – like the GCLF – planetary nebulae samples are often dominated by objects located predominantly in the foreground of their host galaxies, thus resulting in underestimated distances (for a discussion, see Paper VI).

In summary, we argue that the most reliable Fornax distance moduli among the values in our database are those resulting from analyses of Cepheid distances, SBF, and the TRGB. Their weighted mean results in

$$(m-M)_0^{\text{Fornax}} = 31.41 \pm 0.15 \text{ mag}$$

or $D = 19.1_{-1.2}^{+1.4} \text{ Mpc.}$

For completeness, we also considered the post-1990 distance moduli that were not included in our analysis, including those based on SNe Ia, novae, appli-

7

Date (yyyy/mm)	$(m - M)_0^{\text{orig}}$ (mag)	Orig. calibration	Correction ^{a} (m (mag)	(mag)	Target	Notes	Reference
			Cepheids				
1996/12		$(m - M)_0^{ m LMC} = 18.50 \text{ mag}$	-0.01	31.31	NGC 1365		Madore et al. (1996)
1998/09		$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	31.34	NGC 1365		Madore et al. (1998)
1999/04		$(m - M)_0^{\rm LMC} = 18.50 \text{ mag}$	-0.01	31.34	NGC 1365		Madore et al. (1999)
1999/04	31.31 ± 0.20	$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	31.30	NGC 1365		Silbermann et al. (1999
1999/04	31.26 ± 0.10	$(m - M)_0^{LMC} = 18.50 \text{ mag}$	-0.01	31.25	NGC 1365		Silbermann et al. (1999
1999/11	31.36 ± 0.17	$(m - M)_0^{\rm LMC} = 18.50 \text{ mag}$	-0.01	31.35	NGC 1326A		Prosser et al. (1999)
2000/02		$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	31.59	Fornax		Ferrarese et al. (2000)
2000/06	31.43 ± 0.07	$(m - M)_0^{\rm LMC} = 18.50 \text{ mag}$	-0.01	31.42	NGC 1365		Ferrarese et al. (2000)
2001/05	31.32 ± 0.17	$(m - M)_0^{\rm LMC} = 18.50 \text{ mag}$	-0.01	31.31	Fornax		Freedman et al. (2001)
2001/05		$(m - M)_{\rm L}^{\rm DMC} = 18.50 \text{ mag}$	-0.01	31.38	Fornax	Corr. for metallicity	Freedman et al. (2001)
2003/01		$M_I^{\text{TRGB}} = -4.05 \text{ mag}$	0.00	31.62	Fornax		Jerjen (2003)
2016/07		$(m - M)_0^{N4258} = 29.387 \text{ mag}$	-0.10	31.21	NGC 1365		Riess et al. (2016)
2018/01	31.26 ± 0.05	$(m - M)_0^{\text{LMC}} = 18.49 \text{ mag}$	0.00	31.26	NGC 1365		Jang et al. (2018)
		(SBF				
1991/03	30.85 ± 0.04	$(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	30.88	Fornax		Tonry (1991a)
1991/05		$(m - M)_0^{M31} = 24.43 \text{ mag}$ $(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	30.88	Fornax		Tonry (1991b)
1991/08		$(m - M)_0^{M31} = 24.43 \text{ mag}$ $(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	31.38	Fornax		
1991/08 1993/12		$(m - M)_0 = 24.43 \text{ mag}$ $(m - M)_0^{M31} = 24.43 \text{ mag}$		31.38	NGC 1316		Bothun et al. (1991) Ciardullo et al. (1993)
1993/12 1993/12		$(m - M)_0^{M31} = 24.43 \text{ mag}$ $(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03		NGC 1310 NGC 1399		
	30.99 ± 0.10	$(m - M)_0^{M31} = 24.43 \text{ mag}$ $(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	31.02			Ciardullo et al. (1993)
1996/05	31.23 ± 0.06	$(m - M)_{0}^{00} = 24.43 \text{ mag}$	+0.03	31.26	Fornax		Tonry (1997)
1998/09	31.22 ± 0.06	$(m - M)_0^{\text{Virgo}} = 31.00 \text{ mag}$ $(m - M)_0^{\text{Virgo}} = 31.00 \text{ mag}$	+0.03	31.25	Fornax	Ι	Jensen et al. (1998)
1998/09		$(m - M)_0^{+1.95} = 31.00 \text{ mag}$	+0.03	31.35	Fornax	K'	Jensen et al. (1998)
2000/02	31.41 ± 0.06		+0.01	31.42	Fornax	Ι	Tonry et al. (2000)
2000/02	31.50 ± 0.16	в	+0.01	31.51	NGC 1399		Tonry et al. (2000)
2000/02		+0.05 mag w.r.t. Tonry et al. (2001)	-0.04	31.55	Fornax	Ι	Ferrarese et al. (2000)
2000/02		+0.05 mag w.r.t. Tonry et al. (2001)	-0.04	31.47	Fornax	K'	Ferrarese et al. (2000)
2001/01	31.40 ± 0.11	$(m - M)_0^{\text{Virgo}} = 31.03 \text{ mag}$	0.00	31.40	Fornax	Ι	Blakeslee et al. (2001)
2001/10	31.21 ± 0.23	based on Ferrarese et al. (2000)	-0.04	31.17	NGC 1316		Ajhar et al. (2001)
2002/01	31.4 ± 0.2	Tonry et al. (2000) calibration	+0.01	31.41	NGC 1399		Liu et al. (2002)
2003/01		$M_I^{\text{TRGB}} = -4.05 \text{ mag}$	0.00	31.54	Fornax	Dwarf galaxies	Jerjen (2003)
2003/01	31.50 ± 0.04	$M_I^{\text{TRGB}} = -4.05 \text{ mag}$	0.00	31.50	Fornax	Early-type galaxies	Jerjen (2003)
2003/02		$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	31.49	NGC 1316		Jensen et al. (2003)
2006/09	31.51 ± 0.04	$M_I^{\text{TRGB}} = -4.05 \text{ mag}$	0.00	31.51	Fornax		Dunn & Jerjen (2006)
2006/09	31.52 ± 0.04	$M_I^{\text{TRGB}} = -4.05 \text{ mag}$	0.00	31.52	NGC 1399		Dunn & Jerjen (2006)
2007/10	31.78 ± 0.09	-0.21 mag w.r.t. Ferrarese et al. (2000)	-0.25	31.53	NGC 1316		Cantiello et al. (2007)
2007/10	31.62 ± 0.09	-0.21 mag w.r.t. Ferrarese et al. (2000)	-0.25	31.37	NGC 1316		Cantiello et al. (2007)
2007/10	31.62 ± 0.09	-0.21 mag w.r.t. Ferrarese et al. (2000)	-0.25	31.37	NGC 1316		Cantiello et al. (2007)
2007/10	31.53 ± 0.13	-0.21 mag w.r.t. Ferrarese et al. (2000)	-0.25	31.28	NGC 1316		Cantiello et al. (2007)
2007/10	31.59 ± 0.08	-0.21 mag w.r.t. Ferrarese et al. (2000)	-0.25	31.34	NGC 1316		Cantiello et al. (2007)
2007/10		-0.21 mag w.r.t. Ferrarese et al. (2000)		31.05	Fornax		Cantiello et al. (2007)
2009/03		$(m - M)_0^{\text{Virgo}} = 31.09 \text{ mag}$	-0.06	31.45	Fornax	z	Blakeslee et al. (2009)
2009/03	31.606 ± 0.065	$(m - M)_0^{\text{Virgo}} = 31.09 \text{ mag}$	-0.06	31.55	NGC 1316	z	Blakeslee et al. (2009)
2009/03	31.596 ± 0.091	$(m - M)_0^{\text{Virgo}} = 31.09 \text{ mag}$	-0.06	31.54	NGC 1399		Blakeslee et al. (2009)
2003/03		$(m - M)_0$ = 51.05 mag +0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.57	NGC 1399		Blakeslee et al. (2000) Blakeslee et al. (2010)
2010/11 2010/11		+0.06 mag w.r.t. Tonry et al. (2001) +0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.59	NGC 1399 NGC 1316		Blakeslee et al. (2010) Blakeslee et al. (2010)
						1.01410	
2013/04		+0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.54	NGC 1316		Cantiello et al. (2013)
2013/04		+0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.55	NGC 1316		Cantiello et al. (2013)
2013/04		+0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.61	NGC 1316		Cantiello et al. (2013)
2013/04	31.3 ± 0.2	+0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.25	NGC 1316		Cantiello et al. (2013)
2013/04	31.4 ± 0.2	+0.06 mag w.r.t. Tonry et al. (2001)	-0.05	31.35	NGC 1316	z	Cantiello et al. (2013)

 Table 2. Corrections to published Fornax cluster distance moduli.

Table 2. (Continued)

Date	$(m - M)_0^{\mathrm{orig}}$	Orig. calibration	$Correction^a$	$(m - M)_0^{\text{scaled}}$	Target	Notes	Reference
(yyyy/mm)	(mag)		(mag)	(mag)			
			GCLF				
1990/02		$R_0 = 8.0 \text{ kpc}$	+0.08	31.08	Fornax		Geisler & Forte (1990)
1990/02		$R_0 = 8.0 \text{ kpc}$	+0.08	30.93	NGC 1399		Geisler & Forte (1990)
1991/02		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.36 \mathrm{~mag}$	-0.14	31.16	NGC 1399		Bridges et al. (1991)
1991/02		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.36 \mathrm{~mag}$	-0.14	31.31	Fornax		Bridges et al. (1991)
1991/02		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.36 \mathrm{mag}$	-0.14	31.16	Fornax		Bridges et al. (1991)
1996/05	31.20 ± 0.13	$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.40 \mathrm{~mag}$	+0.10	31.30	Fornax	V	Kohle et al. (1996)
1996/05		$M_{V, TO}^{MW} = -7.40 \text{ mag}$	+0.10	30.99	Fornax	Ι	Kohle et al. (1996)
1998/12	31.02 ± 0.2	$M_{V,\text{TO}}^{\text{MW}} = -7.40 \text{ mag}$	+0.10	31.12	NGC 1399		Ostrov et al. (1998)
1999/01	30.91 ± 0.19	$(m-M)_0^{M31} = 24.43 \text{ mag}$	+0.03	30.94	NGC 1399		Grillmair et al. (1999)
1999/03	31.30 ± 0.13	$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.61 \mathrm{mag}$	-0.11	31.19	Fornax		Richtler et al. (2000)
1999/03		$M_{V,TO}^{MW} = -7.61 \text{ mag}$	-0.11	31.21	NGC 1316		Richtler et al. (2000)
1999/03		$M_{VTO}^{\rm NW} = -7.4 \text{ mag}$	+0.10	31.10	Fornax		Kissler-Patig et al. (1997)
1999/07		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.4 \mathrm{mag}$	+0.10	31.40	Fornax		Hilker et al. (1999)
2000/02		V,10 $M_{V,TO}^{MW} = -7.60 \text{ mag}$	-0.10	31.28	Fornax	V	Ferrarese et al. (2000)
2000/02		$M_{V,TO}^{MW} = -7.60 \text{ mag}$	-0.10	31.66		B	Ferrarese et al. (2000)
2000/02		$M_{V,\rm TO}^{\rm MW} = -7.60 \text{ mag}$	-0.10	31.48	NGC 1316		Gómez et al. (2001)
2001/06		$M_{V,\text{TO}} = -7.60 \text{ mag}$ $M_{V,\text{TO}}^{\text{MW}} = -7.60 \text{ mag}$		31.48	NGC 1316 NGC 1316		Gómez et al. (2001) Gómez et al. (2001)
			-0.10				· · ·
2001/06		$M_{V,TO}^{MW} = -7.60 \text{ mag}$	-0.10	31.09	NGC 1316	1	Gómez et al. (2001)
2005/01	31.4	$M_{V,\rm TO}^{\rm MW} = -7.50 \rm mag$	0.00	31.4	NGC 1399		Forte et al. (2005)
		M31	PNLF				
1990/09		$(m - M)_0^{M31} = 24.27 \text{ mag}$ $(m - M)_0^{M31} = 24.27 \text{ mag}$	+0.19	31.07	Fornax		Ciardullo et al. (1990)
1991/00 1993/10		$(m - M)_0^{M31} = 24.27 \text{ mag}$ $(m - M)_0^{M31} = 24.27 \text{ mag}$	+0.19	31.06 31.33	Fornax		cited by de Vaucouleurs (199
1993/10 1993/10		$(m - M)_0^{-1} = 24.27 \text{ mag}$ $(m - M)_0^{M31} = 24.27 \text{ mag}$	+0.19 +0.19	31.33	Fornax NGC 1316		McMillan et al. (1993) McMillan et al. (1993)
1993/10 1993/10		$(m - M)_0 = 24.27 \text{ mag}$ $(m - M)_0^{M31} = 24.27 \text{ mag}$	+0.19 +0.19	31.32	NGC 1310 NGC 1399		McMillan et al. (1993)
1993/10	31.17 ± 0.00 31.19 ± 0.07	$(m - M)_0^{M31} = 24.32 \text{ mag}$ $(m - M)_0^{M31} = 24.32 \text{ mag}$	+0.19 +0.14	31.33	NGC 1355 NGC 1316		Ciardullo et al. (1993)
1993/12		$(m - M)_0^{M31} = 24.32 \text{ mag}$ $(m - M)_0^{M31} = 24.32 \text{ mag}$	+0.14 +0.14	31.36	NGC 1310		Ciardullo et al. (1993)
1996/05		$(m - M)_0^{M31} = 24.24 \text{ mag}$	+0.19	31.43	Fornax		Jacoby (1997)
1999/03		$(m - M)_0^{M31} = 24.44 \text{ mag}$	+0.02	31.35	Fornax		Richtler et al. (2000)
1999/09		+0.06 mag w.r.t. McMillan et al. (1993)	+0.13	31.33	Fornax		Lindblad (1999)
2000/02		$M^* = -4.58 \text{ mag}$	-0.10	31.10	Fornax		Ferrarese et al. (2000)
2007/03		$M^* = -4.47 \text{ mag}$	+0.01	31.27	NGC 1316		Feldmeier et al. (2007)
			TFR				. ,
1996/05	30.94 ± 0.33	$(m - M)_0^{\text{Virgo}} = 31.00 \text{ mag}$	+0.03	30.97	Fornax		Bureau et al. (1996)
1999/03	31.43 ± 0.12	$(m - M)_0^{\text{Virgo}} = 31.00 \text{ mag}$ $(m - M)_0^{\text{Virgo}} = 31.58 \text{ mag}$ $(m - M)_0^{\text{Virgo}} = 31.39 \text{ mag}$	-0.55	30.88	Fornax		Richtler et al. (2000)
1999/09	31.37 ± 0.35	$(m-M)_0^{ m Virgo} = 31.39 m mag$	-0.36	31.01	Fornax		Lindblad (1999)
2000/04	31.25 ± 0.50	$(m - M)_{\rm LMC}^{\rm UMC} = 18.50 \text{ mag}$	-0.01	31.24	Fornax		Tully & Pierce (2000)
2002/06	31.22 ± 0.13	$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	31.21	Fornax		Bernardi et al. (2002)
2004/01	31.57 ± 0.33	$(m-M)_0^{\text{Virgo}} = 31.40 \text{ mag}$	-0.37	31.20	Fornax		Allen & Shanks (2004)
2006/12	31.46 ± 0.14	$(m - M)_0^{\rm LMC} = 18.50 \text{ mag}$	-0.01	31.45	Fornax		Masters et al. (2006)
2009/08	31.38 ± 0.06	$(m - M)_0^{\rm LMC} = 18.50 \rm mag$	-0.01	31.37	NGC 1365		Tully et al. (2009)
2012/04		$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	31.18	Fornax		Tully & Courtois (2012)
2013/03		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	31.21	Fornax	$3.6~\mu{ m m}$	Sorce et al. (2013)
2014/10	31.16 ± 0.13	$(m - M)_0^{\rm LMC} = 18.48 \text{ mag}$	+0.01	31.17	Fornax		Sorce et al. (2014)

^a The calibration of our distance framework is based on distance moduli to the LMC, M31, NGC 4258, and the Virgo cluster of $(m - M)_0^{\text{LMC}} = 18.49 \text{ mag}$ (Paper I), $(m - M)_0^{\text{M31}} = 24.46 \text{ mag}$ (Paper II), $(m - M)_0^{\text{Virgo}} = 31.06 \text{ mag}$ (Paper VI), and $(m - M)_0^{\text{N4258}} = 29.29 \text{ mag}$ (Herrnstein et al. 1999). In addition, we have adopted $M_I^{\text{TRGB}} = -4.05 \text{ mag}$ (TRGB magnitude in the *I* band; Bellazzini et al. 2004), $M^* = -4.67 \text{ mag}$ (PNLF cut-off magnitude at 5007 Å; Ciardullo et al. 1989), $M_{V,\text{TO}}^{\text{MW}} = -7.50 \text{ mag}$ (GCLF turnover magnitude in the Milky Way; Harris 1996), and a Galactic Center distance $R_0 = 8.3 \text{ kpc}$ (paper IV).

^(paper 17) ^b This calibration corresponds to $(m - M)_0^{\text{Virgo}} = 31.03 \text{ mag}$ (based on group membership) and $(m - M)_0^{\text{M31}} = 24.44 \text{ mag}$. We adopted a mean adjustment of +0.01 mag to reconcile these calibration choices with our distance framework.

	(, orig	Onia	Compari: (Miscaled	Tran (Net	D-f-
Date	$(m - M)_0^{\text{orig}}$	Orig. calibration	Correction (m	. 0	Target	Notes	Reference
(yyyy/mm)	(mag)		(mag) SBF	(mag)			
1997/07	25.04 ± 0.21	$(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	35.07	NGC 4881		Thomsen et al. (1997)
1997/07	35.04 ± 0.31	$(m - M)_0^{M31} = 24.43 \text{ mag}$ $(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03 +0.03	35.13	NGC 4881 NGC 4881		Brocato et al. (1998)
1999/00	33.1 ± 0.3	$(m - M)_0^{\text{Virgo}} = 31.06 \text{ mag}$	-0.03	34.61	NGC 4881 NGC 4889		Jensen et al. (1999)
2000/02		$(m - m)_0$ = 51.00 mag +0.05 mag w.r.t. Tonry et al. (1997)	-0.02	35.08	Coma		Ferrarese et al. (2000)
2000/02		+0.05 mag w.r.t. Tonry et al. (1997) +0.05 mag w.r.t. Tonry et al. (1997)	-0.02	35.08	NGC 4881		Freedman et al. (2001)
2001/03		+0.05 mag w.r.t. Tonry et al. (1997) +0.05 mag w.r.t. Tonry et al. (1997)	-0.02	34.97	Coma	K	Liu & Graham (2001)
2001/03	34.55 ± 0.21 35.	$(m - M)_0^{\text{Virgo}} = 31.09 \text{ mag}$	-0.06	34.94	Coma	K	Blakeslee (2013)
2010/02	55.	$(m - m)_0 = 51.05 \text{ mag}$	GCLF	04.04	Coma		Blakesiee (2013)
1995/12	35.16 ± 0.23	$(m - M)_0^{M31} = 24.6 \text{ mag}$	-0.14	35.02	NGC 4881	ower limit	Baum et al. (1995)
1996/12	35.15 ± 0.06	$(m - M)_0^{M31} = 24.6 \text{ mag}$	-0.14	35.01	IC 4051		Baum et al. (1996)
1996/12		$(m-M)_0^{M31} = 24.6 \text{ mag}$	-0.14	34.93	IC 4051		Baum et al. (1996)
1997/05		$(m-M)_0^{M31} = 24.6 \text{ mag}$	-0.14	35.01	IC 4051		Baum et al. (1997)
1997/05		$(m-M)_0^{M31} = 24.6 \text{ mag}$	-0.14	34.93	IC 4051		Baum et al. (1997)
1997/05		$(m-M)_0^{M31} = 24.6 \text{ mag}$	-0.14	34.97	IC 4051		Baum et al. (1997)
1998/07		$(m-M)_0^{M31} = 24.6 \text{ mag}$	-0.14	34.99	IC 4051		Baum (1998)
1998/07		$(m - M)_0^{M31} = 24.6 \text{ mag}$	-0.14	35.24	IC 4051		Baum (1998)
1999/00		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.62 \mathrm{mag}$	-0.12	34.22	\mathbf{Coma}		Tammann & Sandage (1999)
2000/04		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.26 \mathrm{mag}$	+0.14	35.19	Coma		Kavelaars et al. (2000)
2011/00	34.9	$M_{\rm WTO}^{\rm NW} = -7.5 \text{ mag}$	0.00	34.9	NGC 4921		Tikhonov & Galazutdinova (2011)
2011/00	35.0	$M_{V,TO}^{MW} = -7.5 \text{ mag}$	0.00	35.0	NGC 4923		Tikhonov & Galazutdinova (2011)
2011/00	34.93 ± 0.11	$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.5 \mathrm{mag}$	0.00	34.93	Coma		Tikhonov & Galazutdinova (2011)
2016/03		$M_{V,\mathrm{TO}}^{\mathrm{MW}} = -7.66 \mathrm{mag}$	-0.16	34.64	\mathbf{Coma}		Lee & Jang (2016)
2016/03	34.65	$M_{V,TO}^{MW} = -7.66 \text{ mag}$	-0.16	34.49	Coma		Lee & Jang (2016)
		V,10 -	TFR				
1991/07	34.5 ± 0.4	$(m - M)_0^{M31} = 24.2 \text{ mag}$	+0.26	34.76	Coma	В	Fukugita et al. (1991)
1991/07		$(m - M)_0^{M31} = 24.2 \text{ mag}$	+0.26	34.63	Coma	В	Fukugita et al. (1991)
1991/07	33.9	$(m-M)_0^{M31} = 24.2 \text{ mag}$	+0.26	34.16	Coma	В	Fukugita et al. (1991)
1991/07	34.5	$(m - M)_0^{M31} = 24.2 \text{ mag}$	+0.26	34.76	Coma	В	Fukugita et al. (1991)
1993/07		$(m-M)_0^{M31} = 24.37 \text{ mag}$	+0.09	34.41	Coma	В	Rood & Williams (1993)
1993/07	34.56 ± 0.16	$(m-M)_0^{M31} = 24.37 \text{ mag}$	+0.09	34.65	Coma	Н	Rood & Williams (1993)
1994/06	34.86 ± 0.14	$(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	34.89	Coma		Bernstein et al. (1994)
1994/06		$(m-M)_0^{M31} = 24.43 \text{ mag}$	+0.03	34.57	Coma		Bernstein et al. (1994)
1994/06		$(m - M)_0^{M31} = 24.43 \text{ mag}$	+0.03	33.93	\mathbf{Coma}		Bernstein et al. (1994)
1998/00		$(m - M)_0^{\rm LMC} = 18.50 \text{ mag}$	-0.01	34.66	\mathbf{Coma}	BRI	Tully (1998)
2000/04		$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	34.67	\mathbf{Coma}	Ι	Tully & Pierce (2000)
2001/05	34.66	$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	34.65	Coma	Ι	Freedman et al. (2001)
2001/07	34.94 ± 0.13	$(m - M)_0^{M31} = 24.44 \text{ mag}$	+0.02	34.96	Coma	IR	Watanabe et al. (2001)
2004/05	34.58 ± 0.30	$(m - M)_0^{M31} = 24.48 \text{ mag}$	-0.02	34.56	\mathbf{Coma}		Russell (2004)
2004/05	34.76 ± 0.33	$(m - M)_0^{M31} = 24.48 \text{ mag}$	-0.02	34.74	\mathbf{Coma}		Russell (2004)
2007/10	34.80 ± 0.11	$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	34.79	\mathbf{Coma}		Springob et al. (2007)
2009/08	34.65	$(m - M)_0^{\rm LMC} = 18.39 \rm mag$	+0.11	34.76	NGC 4881	K	Russell (2009)
2009/08	34.59	$(m - M)_0^{\rm LMC} = 18.39 {\rm mag}$	+0.11	34.70	NGC 4881	K	Russell (2009)
2009/08	34.70	$(m - M)_0^{\rm LMC} = 18.39 {\rm mag}$	+0.11	34.81	\mathbf{Coma}	K	Russell (2009)
2009/08	34.77	$(m - M)_0^{\rm LMC} = 18.39 \rm mag$	+0.11	34.88	\mathbf{Coma}	K	Russell (2009)
2011/06	34.40 ± 0.08	$M_I^{\text{TRGB}} = -4.05 \text{ mag}$	0.00	34.40	\mathbf{Coma}		Hislop et al. (2011)
2012/04	34.77 ± 0.10	$(m - M)_0^{ m LMC} = 18.50 \text{ mag}$	-0.01	34.76	\mathbf{Coma}	Ι	Tully & Courtois (2012)
2012/04	34.83 ± 0.06	$(m - M)_0^{\text{LMC}} = 18.50 \text{ mag}$	-0.01	34.82	\mathbf{Coma}		Courtois & Tully (2012)
2012/10	34.90 ± 0.13	$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.91	\mathbf{Coma}	$3.6~\mu{ m m}$	Sorce et al. (2012)
2013/03		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.91	\mathbf{Coma}	mid-IR	Sorce et al. (2013)
2013/10		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	35.11	NGC 4889	Ι	Tully et al. (2013)
2013/10		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.83	NGC 4874	Ι	Tully et al. (2013)
2014/09		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.92	\mathbf{Coma}	W1	Neill et al. (2014)
2014/09		$(m - M)_0^{ m LMC} = 18.48 \text{ mag}$	+0.01	34.95	\mathbf{Coma}	W2	Neill et al. (2014)
2014/09		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.87	\mathbf{Coma}	W1	Neill et al. (2014)
2014/09		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.88	\mathbf{Coma}	W2	Neill et al. (2014)
2014/09		$(m - M)_0^{\text{LMC}} = 18.48 \text{ mag}$	+0.01	34.78	\mathbf{Coma}	Ι	Neill et al. (2014)
2014/10		$(m - M)_0^{LMC} = 18.48 \text{ mag}$	+0.01	34.79	Coma		Sorce et al. (2014)

 Table 3. Corrections to published Coma cluster distance moduli.

Table 3. (Continued)

Date	$(m - M)_0^{\text{orig}}$	Orig. calibration	$Correction^a$	$(m - M)_0^{\text{scaled}}$	Target Notes	s Reference
(yyyy/mm)	(mag)		(mag)	(mag)		
			Hubble l	aw		
1998/06	34.21	$H_0 = 100 \rm km s^{-1} \; Mpc^{-1}$	+0.77	34.98	\mathbf{Coma}	Kashikawa et al. (1998)
1999/12	34.2	$H_0 = 100 \rm km s^{-1} \; Mpc^{-1}$	+0.77	34.97	\mathbf{Coma}	Terlevich et al. (1999)
2002/03	35.13	$H_0 = 70 \rm km s^{-1} Mpc^{-1}$	0.00	35.13	\mathbf{Coma}	Brighenti & Mathews (2002)
2003/08	35.03	$H_0 = 70 \rm km s^{-1} Mpc^{-1}$	0.00	35.03	\mathbf{Coma}	Lokas & Mamon (2003)
2008/06	34.98	$H_0 = 71 {\rm km s^{-1} Mpc^{-1}}$	+0.03	35.01	\mathbf{Coma}	Carter et al. (2008)
2009/02	34.97 ± 0.13	$H_0 = 72 \rm km s^{-1} Mpc^{-1}$	+0.06	35.03	\mathbf{Coma}	Harris et al. (2009)
2015/01	34.96	$H_0 = 70 \rm km s^{-1} Mpc^{-1}$	0.00	34.96	\mathbf{Coma}	van Dokkum et al. (2015)
2015/01	35.06	$H_0 = 70 \rm km s^{-1} Mpc^{-1}$	0.00	35.06	\mathbf{Coma}	van Dokkum et al. (2015)




Figure 3. Original and homogenized distance moduli published since 1990 to (left; a–d) the Fornax and (right; e–h) the Coma clusters for specific tracers. The horizontal dotted lines, meant to guide the eye, are drawn at $(m - M)_0 = 31.4$ and 34.8 mag for the Fornax and Coma clusters, respectively. Open circles represent original, published values; solid bullets have been corrected to a common distance framework (see text). Error bars are included where they were provided by the original authors.

cation of the Hubble law, and other, less commonly used tracers (see the online table at http://astroexpat.info/Data/pubbias.html). With few exceptions, the vast majority of these post-1990 measures were comfortably consistent with the weighted mean distance modulus derived above.

4.3. Coma

We will now biefly review the published Coma cluster distance moduli along the same lines as we just did for the Fornax cluster. Table 1 includes four different distance measures. As we argued above for the Fornax cluster, use of GCLF-based distances is fraught with lingering uncertainties, and so we will not consider those measurements here. Of the remaining three tracers, use of the Hubble law requires a somewhat different analysis. The Coma cluster distance coincides with the distance where the 'smooth' Hubble flow starts, i.e., where redshifts of field galaxies become reasonably reliable proxies of their distances. Riess et al. (2009) suggested a minimum redshift of z = 0.023 ($D \sim 100$ Mpc) for the smooth Hubble flow. At D = 100 Mpc, the Hubble-flow

velocity is around 7000 km s⁻¹ and peculiar velocities will typically amount to a 5% contribution.

The main uncertainty in this context relates to the value of the Hubble parameter; radial velocities to individual galaxies and even to entire galaxy clusters can be determined to high accuracy and precision. The mean distance modulus to the Coma cluster based on its recession velocity has been homogenized to a Hubble parameter of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, adopted as compromise value given the prevailing $1-2\sigma$ tension remaining in this field (e.g. Riess et al. 2019, and references therein). A reduction (increase) to $H_0 = 67.3$ (72) km s^{-1} Mpc⁻¹ would result in an increase (decrease) in the Coma distance modulus by 0.09 (0.06) mag. The 'uncertainty' associated with this method included in the table reflects the Gaussian σ of the distribution (since onle one of the measurements included in our final catalog quoted uncertainties). Using the single uncertainty estimate published in this context (Harris et al. 2009), a more realistic uncertainty would require addition in quadrature of this 0.13 mag uncertainty, resulting in a total error of order 0.14 mag.

As for the Fornax cluster, the TFR-based distances to the Coma cluster are systematically shorter than our other distances. We will therefore not include TFRbased distance measures in our analysis. This thus leaves the SBF-based Coma cluster distances. Four of the seven values relate to the SBF distance to NGC 4881, with the remaining three referring to the Coma cluster as a whole. None of the groups contributing to its weighted mean dominate the set of values, so we have no reason to suspect correlated measurements.

In summary, if we adopt the Hubble law- and SBFbased distance moduli (with our updated uncertainty for the Hubble distances), we find a Coma cluster distance modulus of

$$(m-M)_0^{\text{Coma}} = 34.99 \pm 0.38 \text{ mag}$$

or $D = 99.5^{+19.0}_{-15.9} \text{ Mpc.}$

For completeness, we again considered the post-1990 distance moduli that were not included in our analysis, including those based on SNe Ia, Fundamental Plane scaling, and other, less commonly used tracers (see the online table at http://astroexpat.info/Data/pubbias.html). All of the latter post-1990 measures were comfortably consistent with the weighted mean distance modulus derived above.

4.4. Relative distance moduli

At distances equivalent to or beyond that of the Virgo cluster, it has become relatively common to quote distance measures relative to the Virgo cluster. The main advantage of using relative rather than absolute distances is that the number of assumptions one has to make is significantly reduced, provided that the physical basis on which the distances are compared is similar. For the Fornax cluster, our database includes 43 relative Fornax–Virgo cluster distance moduli published since 1990. A straight weighted mean yields $\Delta(m-M)_0^{\rm Fornax-Virgo} = 0.18 \pm 0.28$ mag. We used all 43 values to determine the central value of this relative distance modulus; since five measurements do not have any uncertainties associated with them, we used the remaining 38 values to determine the uncertainty pertaining to our weighted mean. Both clusters thus appear to be located at very similar distance, with Fornax placed marginally more distant.

To avoid any effects associated with small-number statistics, we subdivided our 43 relative distance moduli into three groups of similar tracers, yielding post-1990 weighted-mean relative distance moduli of $\Delta(m - M)_0^{\text{Fornax-Virgo}} = 0.23 \pm 0.18$ mag for the 11 SBF-based measurements, $\Delta(m - M)_0^{\text{Fornax-Virgo}} = 0.10 \pm 0.12$ mag for the 10 values based on luminosity functions (GCLF, PNLF), and $\Delta(m - M)_0^{\text{Fornax-Virgo}} = 0.15 \pm 0.15$ mag for the 11 relative distance measures based on kinematic tracers (TFR, the $D_n - \sigma$ relation, and Fundamental Plane scaling). Combined with our recommended Virgo cluster distance modulus of $(m - M)_0^{\text{Virgo}} = 31.03 \pm 0.14$ mag (Paper VI), these relative measurements convert to absolute Fornax cluster distance moduli of

- $(m M)_0^{\text{Fornax}} = 31.21 \pm 0.28 \text{ mag} \text{ (all values)};$
- $(m-M)_0^{\text{Fornax,SBF}} = 31.26 \pm 0.23 \text{ mag} \text{ (SBF-based measures);}$
- $(m M)_0^{\text{Fornax,LFs}} = 31.13 \pm 0.12 \text{ mag}$ (luminosity functions); and
- $(m M)_0^{\text{Fornax,kin}} = 31.18 \pm 0.15 \text{ mag}$ (kinematic tracers);

The weighted mean of the latter three values is

$$(m-M)_0^{\text{Fornax,stat}} = 31.18 \pm 0.17 \text{ mag}$$

or $D = 17.2_{-1.3}^{+1.4} \text{ Mpc.}$

As for the Coma cluster, relative distance moduli have been determined with respect to the Virgo and Fornax clusters and the Leo I group. The 23 post-1990 relative Coma–Virgo distance moduli (of which 22 have associated uncertainties) yield a straight weighted mean of $\Delta(m-M)_0^{\text{Coma-Virgo}} = 3.75 \pm 0.23$ mag. The subset of six kinematics-based relative distance measures (TFR, Fundamental Plane) result in $\Delta(m-M)_0^{\text{Coma-Virgo}} =$ 3.63 ± 0.22 mag. (None of the other possible subsets of Coma–Virgo relative distance determinations reach the threshold where a statistical analysis becomes meaningful.) Combining the weighted mean of the full set of measurements with our preferred Virgo cluster distance modulus yields

$$(m-M)_0^{\text{Coma}} = 34.78 \pm 0.27 \text{ mag}$$

or $D = 90.4^{+11.9}_{-10.6} \text{ Mpc},$

while the kinematics-based distances yield

$$(m-M)_0^{\text{Coma}} = 34.66 \pm 0.26 \text{ mag}$$

or $D = 85.5^{+10.9}_{-9.6} \text{ Mpc.}$

Second, a significant number of relative distance moduli with respect to the Leo I group were published by Gregg (1997), in the form of Coma distance moduli based on a variety of independent calibrations. His unweighted mean distance modulus to the Coma cluster, $(m - M)_0 = 34.90 \pm 0.13$ mag, is based on a mean distance ratio of $D_{\text{Coma}}/D_{\text{Leo I}} = 8.84\pm0.23$, corresponding to $\Delta(m - M)_0^{\text{Coma-Leo I}} = 4.73 \pm 0.06$ mag. However, on closer inspection, the data set underlying these values raises a number of concerns. If one calculates the individual Coma/Leo I distance ratios using the values included in his Table 2, the central value of the resulting ratio is $D_{\text{Coma}}/D_{\text{Leo I}} = 8.84$ for every single, presumably independent calibration method.

In the preamble to his Section 3, Gregg (1997) states that he derived the Leo I distances included in his Table 2 on the basis of four different calibration methods. In addition, he states that the Coma cluster distances included are based on the zero-point calibration of the distance–velocity dispersion relation for Coma. If the Leo I calibration methods applied were indeed independent (which we have no reason to doubt), this very tightly defined central value is statistically highly unlikely. Therefore, we decided to discard Gregg (1997)'s measurements, since we cannot ascertain their integrity. This has unintended consequences, however, because a number of subsequently published Coma distance moduli were also based on this result (Cassisi & Salaris 1998; Salaris & Cassisi 1998), and so we were forced to discard them.

Thomsen et al. (1997) provided the only independent relative distance modulus between the Coma cluster and the Leo I group, $\Delta (m-M)_0^{\text{Coma-Leo I}} = 4.89 \pm 0.30$ mag. Although this measurement is consistent, within the 1σ uncertainties, with the value promoted by Gregg (1997), we will nevertheless refrain from further analysis of the Coma-Leo I distance differential.

Finally, van den Bergh (1994) cited a distance ratio of $D_{\rm Coma}/D_{\rm Fornax} = 5.25 \pm 0.38$, although without providing provenance. This corresponds to a relative distance modulus between the Coma and Fornax clusters of $\Delta (m-M)_0^{\rm Coma-Fornax} = 3.60 \pm 0.15$ mag. Combining this with the Fornax cluster distance moduli obtained above, we obtain

$$(m - M)_0^{\text{Coma}} = 34.78 \pm 0.23 \text{ mag},$$

= $34.81 \pm 0.32 \text{ mag}, \text{ and}$
= $35.01 \pm 0.21 \text{ mag}$

for a subset of relative Coma–Virgo distance moduli, the full set of Coma–Virgo measures, and our best direct estimate for the Virgo cluster distance, respectively.

5. A DISTANCE FRAMEWORK OUT TO 100 MPC

In this paper, we have considered the body of published distance moduli to the Fornax and Coma clusters, with specific emphasis on the period since 1990. We carefully homogenized our final catalogs of distance moduli onto the distance scale established in Papers I through VI. We assessed systematic differences associated with the use of specific tracers, and consequently discarded results based on application of the TFR and of luminosity functions.

We recommend 'best' weighted relative distance moduli for the Fornax and Coma clusters with respect to the Virgo cluster benchmark of $\Delta (m-M)_0^{\rm Fornax-Virgo} =$ 0.18 ± 0.28 mag and $\Delta (m-M)_0^{\rm Coma-Virgo} = 3.75 \pm 0.23$ mag. On balance, the set of weighted mean distance moduli we derived as most representative of the clusters' distances is as follows:

$$(m - M)_0^{\text{Fornax}} = 31.41 \pm 0.15 \text{ mag and}$$

= $31.21 \pm 0.28 \text{ mag};$
 $(m - M)_0^{\text{Coma}} = 34.99 \pm 0.38 \text{ mag and}$
= $34.78 \pm 0.27 \text{ mag}.$

For each cluster, this first distance modulus is the result of our analysis of the direct, absolute distance moduli published since 1990, while the second modulus is based on the relative measures published during the same period.

Interestingly, while the absolute and relative distance moduli for both clusters are mutually consistent within the uncertainties, the relative distance moduli yield absolute distances that are shorter by ~ 0.20 mag, or about 1σ . It is unclear what may have caused this small difference for both clusters; investigation of the cause is beyond the scope of the present paper since it requires careful examination of the individual distances comprising the tracers commonly used in this field. It is unlikely that line-of-sight depth effects are to blame (e.g. Jerjen 2003; Dunn & Jerjen 2006; Blakeslee et al. 2009), given that most individual galaxies and galaxy samples in both of our clusters comprise the same or similar objects. We speculate that lingering uncertainties in the underlying absolute distance scale appear to have given rise to a systematic uncertainty of order 0.20 mag.

This concludes our series of papers aimed at establishing a robust and internally consistent, statistically validated distance framework out to distances of order 100 Mpc. Our recommended distances to the Fornax and Coma clusters quoted above should be read in tandem with the distance moduli we derived and recommend for the Galactic Center (Paper IV), the Magellanic Clouds (Papers I and III), the M31 group (Paper II), NGC 4258 (Herrnstein et al. 1999), and the Virgo cluster (Paper VI): see Table 3 in Paper VI for the full set of recommended distance moduli.

ACKNOWLEDGEMENTS

This research has made extensive use of NASA's Astrophysics Data System Abstract Service.

REFERENCES

- Ajhar, E. A., Tonry, J. L., Blakeslee, J. P., et al. 2001, ApJ, 559, 584
- Allen, P. D., & Shanks, T. 2004, MNRAS, 347, 1011
- Baum, W. A. 1998, AJ, 116, 31
- Baum, W. A., Hammergren, M., Groth, E. J., et al. 1995, AJ, 110, 2537
- Baum, W. A., Hammergren, M., Thomsen, B., et al. 1996, Am. Astron. Soc. Mtg Abstr. 189, 12.04
- Baum, W. A., Hammergren, M., Thomsen, B., et al. 1997, AJ, 113, 1483
- Bellazzini, M., Ferraro, F. R., Sollima, A., et al. 2004, A&A, 424, 199
- Bernardi, M., Alonso, M. V., da Costa, L. N., et al. 2002, AJ, 123, 2990
- Bernstein, G. M., Guhathakurta, P., Raychaudhury, S., et al. 1994, AJ, 107, 1962
- Blakeslee, J. P. 2013, in Advancing the Physics of Cosmic Distances, ed. R. de Grijs, IAU Symp. 289, 304
- Blakeslee, J. P., Cantiello, M., Mei, S., et al. 2010, ApJ, 724, 657
- Blakeslee, J. P., Jordán, A., Mei, S., et al. 2009, ApJ, 694, 556
- Blakeslee, J. P., Vazdekis, A., & Ajhar, E. A. 2001, MNRAS, 320, 193
- Bothun, G. D., Impey, C. D., & Malin, D. F. 1991, ApJ, 376, 404
- Bridges, T. J., Hanes, D. A., & Harris, W. E. 1991, AJ, 101, 469
- Brighenti, F., & Mathews, W. G. 2002, ApJ, 567, 130
- Brocato, E., Capaccioli, M., & Condelli, M. 1998,
- Mem. Soc. Astron. Italiana, 69, 155
- Bureau, M., Mould, J. R., & Staveley-Smith, L. 1996, ApJ, 463, 60
- Cantiello, M., Blakeslee, J., Raimondo, G., et al. 2007, ApJ, 668, 130
- Cantiello, M., Grado, A., Blakeslee, J. P., et al. 2013, A&A, 552, A106
- Carter, D., Goudfrooij, P., Mobasher, B., et al. 2008, ApJS, 176, 424
- Cassisi, S., & Salaris, M. 1998, Mem. Soc. Astron. Italiana, 69, 127
- Ciardullo, R., Jacoby, G. H., & Engle, K. A. 1990, BAAS22, 1312
- Ciardullo, R., Jacoby, G. H., Ford, H. C., et al. 1989, ApJ, 339, 53
- Ciardullo, R., Jacoby, G. H., & Tonry, J. L. 1993, ApJ, 419, 479
- Courtois, H. M., & Tully, R. B. 2012, ApJ, 749, 174
- Crandall, S., & Ratra, B. 2015, ApJ, 815, 87
- de Grijs, R., & Bono, G. 2014, AJ, 148, 17 (Paper II)
- de Grijs, R., & Bono, G. 2015, AJ, 149, 179 (Paper III)
- de Grijs, R., & Bono, G. 2016, ApJS, 227, 5 (Paper IV)
- de Grijs, R., & Bono, G. 2017, ApJS, 232, 22 (Paper V)
- de Grijs, R., & Bono, G. 2020, ApJS, 246, 3 (Paper VI)
- de Grijs, R., Wicker, J. E., & Bono, G. 2014, AJ, 147, 122 (Paper I)
- de Vaucouleurs, G. 1993, ApJ, 415, 10
- Dirsch, B., Richtler, T., Geisler, D., et al. 2003, AJ, 125, 1908

- Drinkwater, M. J., Gregg, M. D., & Colless, M. 2001, ApJL, 548, L139
- Dunn, L. P., & Jerjen, H. 2006, AJ, 132, 1384
- Feldmeier, J. J., Jacoby, G. H., & Phillips, M. M. 2007, ApJ, 657, 76
- Ferrarese, L., Mould, J. R., Kennicutt, R. C., et al. 2000, ApJ, 529, 745
- Forte, J. C., Faifer, F., & Geisler, D. 2005, MNRAS, 357, 56
- Freedman, W. L., Madore, B. F., Gibson, B. K., et al. 2001, ApJ, 553, 47
- Fukugita, M., Okamura, S., Tarusawa, K., et al. 1991, ApJ, 376, 8
- Geisler, D., & Forte, J. C. 1990, ApJL, 350, L5
- Gómez, M., Richtler, T., Infante, L., et al. 2001, A&A, 371, 875
 Gregg, M. D. 1997, NewA, 1, 363
- Grillmair, C. J., Forbes, D. A., Brodie, J. P., et al. 1999, AJ, 117, 167
- Harris, W. E. 1996, AJ, 112, 1487
- Harris, W. E., Kavelaars, J. J., Hanes, D. A., et al. 2009, AJ, 137, 3314
- Herrnstein, J. R., Moran, J. M., Greenhill, L. J., et al. 1999, Nature, 400, 539
- Hilker, M., Infante, L., & Richtler, T. 1999, A&AS, 138, 55
- Hislop, L., Mould, J., Schmidt, B., et al. 2011, ApJ, 733, 75
- Jacoby, G. H. 1997, in The Extragalactic Distance Scale, eds. M. Livio, M. Donahue, & N. Panagia, Cambridge, UK: Cambridge Univ. Press, p. 197
- Jang, I. S., Hatt, D., Beaton, R. L., et al. 2018, ApJ, 852, 60
- Jensen, J. B., Tonry, J. L., Barris, B. J., et al. 2003, ApJ, 583, 712
- Jensen, J. B., Tonry, J. L., & Luppino, G. A. 1998, ApJ, 505, 111
- Jensen, J. B., Tonry, J. L., & Luppino, G. A. 1999, ApJ, 510, 71
- Jerjen, H. 2003, A&A, 398, 63
- Jordán, A., McLaughlin, D. E., Côté, P., et al. 2006, ApJL, 651, L25
- Kashikawa, N., Sekiguchi, M., Doi, M., et al. 1998, ApJ, 500, 750
- Kavelaars, J. J., Harris, W. E., Hanes, D. A., et al. 2000, ApJ, 533, 125
- Kennicutt, R. C., Stetson, P. B., Saha, A., et al. 1998, ApJ, 498, 181
- Kissler-Patig, M., Kohle, S., Hilker, M., et al. 1997, A&A, 319, 470
- Kohle, S., Kissler-Patig, M., Hilker, M., et al. 1996, A&A, 309, L39
- Lee, M. G., & Jang, I. S. 2016, ApJ, 819, 77
- Lindblad, P. O. 1999, A&A Rv, 9, 221
- Liu, M. C., & Graham, J. R. 2001, ApJL, 557, L31
- Liu, M. C., Graham, J. R., & Charlot, S. 2002, ApJ, 564, 216
- Lokas, E. L., & Mamon, G. A. 2003, MNRAS, 343, 401
- Madore, B. F., Freedman, W. L., Kennicutt, R. C., et al. 1996, Am. Astron. Soc. Mtg Abstr. 189, 108.04
- Madore, B. F., Freedman, W. L., Silbermann, N., et al. 1998, Nature, 395, 47

15

- Madore, B. F., Freedman, W. L., Silbermann, N., et al. 1999, ApJ, 515, 29
- Masters, K. L., Springob, C. M., Haynes, M. P., et al. 2006, ApJ, 653, 861
- McMillan, R., Ciardullo, R., & Jacoby, G. H. 1993, ApJ, 416, 62
- Neill, J. D., Seibert, M., Tully, R. B., et al. 2014, ApJ, 792, 129
- Ostrov, P. G., Forte, J. C., & Geisler, D. 1998, AJ, 116, 2854
- Prosser, C. F., Kennicutt, R. C., Bresolin, F., et al. 1999, ApJ, 525, 80
- Richtler, T., Drenkhahn, G., Gómez, M., et al. 2000, in From Extrasolar Planets to Cosmology: The VLT Opening Symposium, eds J. Bergeron & A. Renzini, p. 259
- Riess, A. G., Casertano, S., Yuan, W., et al. 2019, ApJ, 876, 85
- Riess, A. G., Macri, L., Casertano, S., et al. 2009, ApJ, 699, 539
- Riess, A. G., Macri, L. M., Hoffmann, S. L., et al. 2016, ApJ,
- 826, 56
- Rood, H. J., & Williams, B. A. 1993, MNRAS, 263, 211
- Russell, D. G. 2004, ApJ, 607, 241
- Russell, D. G. 2009, J. Astrophys. Astron., 30, 93
- Salaris, M., & Cassisi, S. 1998, MNRAS, 298, 166
- Schaefer, B. E. 2008, AJ, 135, 112
- Shanks, T. 1997, MNRAS, 290, L77
- Silbermann, N. A., Harding, P., Ferrarese, L., et al. 1999, ApJ, 515, 1
- Sorce, J. G., Courtois, H. M., Tully, R. B., et al. 2013, ApJ, 765, 94
- Sorce, J. G., Tully, R. B., & Courtois, H. M. 2012, ApJL, 758, L12
- Sorce, J. G., Tully, R. B., Courtois, H. M., et al. 2014, MNRAS, 444, 527
- Springob, C. M., Masters, K. L., Haynes, M. P., et al. 2007, ApJS, 172, 599

- Tammann, G. A., & Sandage, A. 1999, in Harmonizing Cosmic Distance Scales in a Post-hipparcos Era, eds D. Egret & A. Heck, ASP Conf. Ser. 167, 204
- Terlevich, A. I., Kuntschner, H., Bower, R. G., et al. 1999, MNRAS, 310, 445
- Thomsen, B., Baum, W. A., Hammergren, M., et al. 1997, ApJL, 483, L37
- Tikhonov, N. A., & Galazut
dinova, O. A. 2011, Astron. Lett., $37,\,766$
- Tonry, J. L. 1991a, BAAS, 23, 956
- Tonry, J. L. 1991b, ApJL, 373, L1
- Tonry, J. L. 1997, in The Extragalactic Distance Scale, eds. M. Livio, M. Donahue, & N. Panagia, Cambridge, UK: Cambridge Univ. Press, p. 297
- Tonry, J. L., Blakeslee, J. P., Ajhar, E. A., et al. 1997, ApJ, 475, 399
- Tonry, J. L., Blakeslee, J. P., Ajhar, E. A., et al. 2000, ApJ, 530, 625
- Tonry, J. L., Dressler, A., Blakeslee, J. P., et al. 2001, ApJ, 546, 681
- Tully, R. B. 1998, Mem. Soc. Astron. Italiana, 69, 237
- Tully, R. B., & Courtois, H. M. 2012, ApJ, 749, 78
- Tully, R. B., & Pierce, M. J. 2000, ApJ, 533, 744
- Tully, R. B., Courtois, H. M., Dolphin, A. E., et al. 2013, AJ, 146, 86
- Tully, R. B., Rizzi, L., Shaya, E. J., et al. 2009, AJ, 138, 323
- van den Bergh, S. 1994, PASP, 106, 1113
- van Dokkum, P. G., Abraham, R., Merritt, A., et al. 2015, ApJL, 798, L45
- Villegas, D., Jordán, A., Peng, E. W., et al. 2010, ApJ, 717, 603
- Watanabe, M., Yasuda, N., Itoh, N., et al. 2001, ApJ, 555, 215
- Whitmore, B. C. 1997, in The Extragalactic Distance Scale, eds. M. Livio, M. Donahue, & N. Panagia, Cambridge, UK: Cambridge Univ. Press, p. 254