

ODD LENGTH: ODD DIAGRAMS AND DESCENT CLASSES

FRANCESCO BRENTI AND ANGELA CARNEVALE

ABSTRACT. We define and study odd analogues of classical geometric and combinatorial objects associated to permutations, namely odd Schubert varieties, odd diagrams, and odd inversion sets. We show that there is a bijection between odd inversion sets of permutations and acyclic orientations of the Turán graph, that the dimension of the odd Schubert variety associated to a permutation is the odd length of the permutation, and give several necessary conditions for a subset of $[n] \times [n]$ to be the odd diagram of a permutation. We also study the sign-twisted generating function of the odd length over descent classes of the symmetric groups.

1. INTRODUCTION

Motivated by questions in enumerative geometry a new statistic on the symmetric groups was introduced and studied in [7]. This statistic combines combinatorial and parity conditions and is now known as the odd inversion number or odd length (see, e.g., [2]). It was conjectured in [7, Conjecture C] that the sign-twisted generating function of this new statistic on any quotient of any symmetric group is given by an explicit product formula. This conjecture was proved in [2]. An odd length statistic has also been defined and studied on the hyperoctahedral groups by Stasinski and Voll in [12] and [13], on the even hyperoctahedral groups by the authors in [3], and on all Weyl groups in [4] and [14].

Our purpose in this paper is to carry out a further study of this statistic on the symmetric groups from the combinatorial, enumerative and geometric point of view. More precisely, we show that, given any permutation σ , there is a complex projective variety $X_o(\sigma)$ (which we call an *odd Schubert variety*) whose dimension is the odd length of σ . We also define and study the odd analogues of two other familiar combinatorial objects associated to a permutation, namely diagrams and inversion sets. We show that there is a simple transformation connecting odd inversion sets and odd diagrams, we characterize the subsets of $[n] \times [n]$ that are odd inversion sets of permutations, and we give several necessary conditions for a subset of $[n] \times [n]$ to be the odd diagram of a permutation. Also, we study the sign-twisted generating function of the odd length over descent classes of the symmetric groups. In particular, we give sufficient conditions for the generating function to be zero, and compute it explicitly for the alternating permutations and for a family of descent classes which includes all quotients.

The organization of the paper is as follows. In the next section we recall definitions and results that we use in the sequel. In § 3 we introduce and study odd Schubert varieties, odd inversion sets, and odd diagrams. More precisely, we show that the odd length of a permutation σ is the dimension of its associated odd Schubert variety, and that this variety depends on a subset of the diagram of σ (which we call the *odd diagram* of σ). Furthermore, that there is a simple transformation relating the odd diagram with a subset of the inversion set (which we call the *odd inversion set*) of σ , that there is a bijection between odd inversion sets of permutations and acyclic orientations of the Turán graph, and give several necessary conditions for a subset of $[n] \times [n]$ to be the odd diagram of a permutation. In § 4 we study the effect that some operations, that can be performed on a descent class, have on

the corresponding sign-twisted generating function of the odd length. In § 5 we give sufficient conditions on a descent class for its sign-twisted generating function to be zero and we compute it explicitly for the descent classes of the alternating permutations, and for a general family of descent classes which includes all quotients. Finally, in § 6, we present some conjectures and open problems arising from the present work.

2. PRELIMINARIES

In this section we collect some notation and basic facts about symmetric groups that we use in the sequel. Besides the combinatorial aspects we recall the geometric facts and definitions about the Schubert variety associated with a permutation.

For $X \subseteq \mathbb{N}$ we let X_0 denote $X \cup \{0\}$. For $m, n \in \mathbb{Z}$, $m \leq n$, we let $[m, n]$ denote the set $\{m, m+1, \dots, n-1, n\}$ and for $n \in \mathbb{N}$ we let $[n] = [1, n]$. If $\{a_1, \dots, a_n\} \subseteq \mathbb{Z}$ we write $\{a_1, \dots, a_n\}_<$ to mean that $a_1 < \dots < a_n$. Given $J \subseteq [n-1]$ there are unique integers $a_1 < \dots < a_s$ and $b_1 < \dots < b_s$ such that $J = [a_1, b_1] \cup \dots \cup [a_s, b_s]$ and $b_i + 1 < a_{i+1}$ for $i = 1, \dots, s-1$. We call the intervals $[a_1, b_1], \dots, [a_s, b_s]$ the *connected components of J* .

For $n \in \mathbb{N}$ we let $[n]_q := (1 - q^n)/(1 - q)$ (so $[0]_q = 0$), and $[n]_q! := \prod_{i=1}^n [i]_q$ (so $[0]_q! = 1$). For $n_1, \dots, n_k \in \mathbb{N}$ such that $\sum_{i=1}^k n_i = n$ we let

$$\left[\begin{array}{c} n \\ n_1, \dots, n_k \end{array} \right]_q := \frac{[n]_q!}{[n_1]_q! \cdots [n_k]_q!}.$$

We refer to [1] for notation, terminology and basic facts about Coxeter groups. The symmetric group S_n is the group of permutations of $[n]$. We let $S = \{s_1, \dots, s_{n-1}\}$ denote the set of standard generators of S_n , where s_i denotes the i -th transposition $(i, i+1)$. It is well known that S_n is a Coxeter group with respect to this set of generators and that for $\sigma \in S_n$ the Coxeter length $\ell(\sigma)$ and the descent set $\text{Des}(\sigma)$ have combinatorial interpretations

$$\ell(\sigma) = |\{(i, j) \in [n]^2 : i < j, \sigma(i) > \sigma(j)\}|$$

and

$$(2.1) \quad \text{Des}(\sigma) = |\{i \in [n-1] : \sigma(i) > \sigma(i+1)\}|,$$

respectively. In the sequel we often identify the generating set S with the set $[n-1]$. For $\sigma \in S_n$ the *diagram* of σ is

$$D(\sigma) := \{(i, j) \in [n]^2 : j < \sigma(i), \sigma^{-1}(j) > i\},$$

and the *inversion set* of σ is

$$\text{Inv}(\sigma) := \{(i, j) \in [n]^2 : i < j, \sigma(i) > \sigma(j)\}.$$

Note that $|D(\sigma)| = |\text{Inv}(\sigma)| = \ell(\sigma)$, and that $(i, \sigma(j)) \in D(\sigma)$ if and only if $(i, j) \in \text{Inv}(\sigma)$, for all $(i, j) \in [n]^2$.

Let $n \in \mathbb{N}$ and $\{e_1, \dots, e_n\}$ be the canonical basis of \mathbb{C}^n . A *flag* in \mathbb{C}^n is a sequence (U_1, \dots, U_n) of subspaces of \mathbb{C}^n such that $U_1 \subset \dots \subset U_n$ and $\dim(U_i) = i$ for all $i = 1, \dots, n$. The set F_n of all flags in \mathbb{C}^n is called the *flag manifold* of \mathbb{C}^n . For $\sigma \in S_n$ we now recall the definition of the the *Schubert cell* of σ , which we denote by $C(\sigma)$. Namely, $(U_1, \dots, U_n) \in C(\sigma)$ if and only if there are $(a_{i,j})_{(i,j) \in D(\sigma)} \in \mathbb{C}^{D(\sigma)}$ such that

$$U_k = \langle \{e_{\sigma(i)} + \sum_{\{j:(i,j) \in D(\sigma)\}} a_{i,j} e_j\}_{1 \leq i \leq k} \rangle$$

for $k \in [n]$. It is well known (see, e.g., [9, (A.4)]) and not hard to see, that the map $(a_{i,j})_{(i,j) \in D(\sigma)} \mapsto (U_1, \dots, U_n)$ is injective. In particular, $C(\sigma)$ is isomorphic to an affine space of dimension $\ell(\sigma)$.

Recall (see, e.g., [5, p. 209]) that for $i \in [n]$ the *Plücker embedding* π_i associates to any i -dimensional subspace U of \mathbb{C}^n a point in the projective space $\mathbb{P}(\Lambda^i(\mathbb{C}^n)) = \mathbb{P}^{\binom{n}{i}-1}$. More precisely, the image of U under π_i is the $\binom{n}{i}$ -tuple $(U_I)_{\{I \subseteq [n]: |I|=i\}}$ where, for $I \subseteq [n]$, $|I| = i$, U_I is the minor whose columns are indexed by the elements in I of a matrix which has as rows a basis of U .

One may thus associate to any flag (U_1, \dots, U_n) of \mathbb{C}^n a point in the Cartesian product $\mathbb{P}(\Lambda^1(\mathbb{C}^n)) \times \mathbb{P}(\Lambda^2(\mathbb{C}^n)) \times \dots \times \mathbb{P}(\Lambda^{n-1}(\mathbb{C}^n))$. In turn, to any point in this product the *Segre embedding* (see, e.g., [6, Chap. I, Ex. 2.14] for the definition and information about the Segre embedding) associates a point in the projective space $\mathbb{P}(E)$ where $E := \mathbb{C}^n \otimes \Lambda^2(\mathbb{C}^n) \otimes \dots \otimes \Lambda^{n-1}(\mathbb{C}^n)$. The image of F_n under this composite embedding (which we denote by π) is a complex projective algebraic variety.

The *Schubert variety* $X(\sigma)$ is the closure of $\pi(C(\sigma))$ in $\pi(F_n)$. It is well known that $X(\sigma)$ is a complex projective variety of dimension $\ell(\sigma)$.

One of our results concerns generating functions on descent classes of the symmetric groups, which we now define. For $I, J \subseteq S$, $J \subseteq S \setminus I$ we let

$$(2.2) \quad \mathcal{D}_J^I(S_n) := \{\sigma \in S_n \mid J \subseteq \text{Des}(\sigma) \subseteq S \setminus I\},$$

$$(2.3) \quad S_n^I := \mathcal{D}_\emptyset^I(S_n).$$

Similarly, for subsets $X \subseteq S_n$, we denote $\mathcal{D}_J^I(X) := X \cap \mathcal{D}_J^I(S_n)$.

To state the main result of [2], which is a special case of one of our main results, we need the following definitions. Let $n \in \mathbb{N}$. Set:

$$C_{n,+} := \{\sigma \in S_n : i + \sigma(i) \equiv 0 \pmod{2}, i = 1, \dots, n\},$$

$$C_{n,-} := \{\sigma \in S_n : i + \sigma(i) \equiv 1 \pmod{2}, i = 1, \dots, n\},$$

$$C_n := C_{n,+} \cup C_{n,-}.$$

Note that

$$C_n = \{\sigma \in S_n : i \equiv j \pmod{2} \Rightarrow \sigma(i) \equiv \sigma(j) \pmod{2}, \text{ for all } i, j \in [n]\}.$$

Elements in $C_{n,+}$ are called *even* chessboard elements, those in $C_{n,-}$ *odd* chessboard elements. Informally, a chessboard element is a permutation whose matrix fits either on all black or on all white squares of a chessboard. Note that $|C_{n,+}| = \lfloor \frac{n}{2} \rfloor! \lceil \frac{n}{2} \rceil!$ (see also [10, A010551]). For $n = 2m + 1$, clearly $C_{n,-} = \emptyset$ and therefore $C_n = C_{n,+}$.

Note that the set of chessboard elements C_n is a subgroup of S_n and that the set of even chessboard elements $C_{n,+}$ is a subgroup of C_n isomorphic to $S_{\lfloor \frac{n}{2} \rfloor} \times S_{\lceil \frac{n}{2} \rceil}$. Relatives of these groups feature in our proof of Proposition 3.5.

The odd length is defined as follows (see also [7] and [2]).

Definition 2.1. Let $n \in \mathbb{N}$ and $\sigma \in S_n$. The *odd length* of σ is

$$(2.4) \quad L(\sigma) := |\{(i, j) \in [n]^2 : i < j, \sigma(i) > \sigma(j), i \not\equiv j \pmod{2}\}|.$$

Informally, the statistic L counts inversions between values in positions with opposite parity. In the next proposition we collect some properties satisfied by L .

Proposition 2.2. Let $n \in \mathbb{N}$ and let w_0 be the unique longest element of S_n . Then

$$(i) \quad L(e) = 0,$$

- (ii) $L(s_i) = 1$, for $i = 1, \dots, n-1$,
- (iii) $L(\sigma w_0) = L(w_0 \sigma) = L(w_0) - L(\sigma)$ for all $\sigma \in S_n$,
- (iv) w_0 is the unique element on which L attains its maximum, and $L(w_0) = \lfloor \frac{n}{2} \rfloor \lfloor \frac{n}{2} \rfloor$.

Proof. The only non-trivial point is the last one. It follows from (iii) and the fact that the identity is the unique element on which L is zero. The last statement comes from the fact that, by definition, $L(w_0) = \sum_{i=1}^{n-1} \lfloor \frac{i}{2} \rfloor$. \square

The following result, conjectured in [7] and proved in [2], gives explicit product formulas for the sign-twisted distribution of the odd length on all parabolic quotients of the symmetric groups.

Theorem 2.3. *Let $n \in \mathbb{N}$, $I \subseteq [n-1]$, and I_1, \dots, I_s be the connected components of I . Then*

$$(2.5) \quad \sum_{\sigma \in S_n^I} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} \left[\begin{matrix} m \\ \lfloor \frac{|I_1|+1}{2} \rfloor, \dots, \lfloor \frac{|I_s|+1}{2} \rfloor \end{matrix} \right]_{x^2} \prod_{k=b+1}^{\lfloor \frac{n-1}{2} \rfloor} (1 - x^{2k}), \\ \text{if } n \equiv 1 \pmod{2}, \text{ or if } n = 2b, \\ (1 + x^m) \left[\begin{matrix} m \\ \lfloor \frac{|I_1|+1}{2} \rfloor, \dots, \lfloor \frac{|I_s|+1}{2} \rfloor \end{matrix} \right]_{x^2} \prod_{k=b+1}^{\lfloor \frac{n-1}{2} \rfloor} (1 - x^{2k}), \\ \text{otherwise,} \end{cases}$$

where $b := \sum_{k=1}^s \lfloor \frac{|I_k|+1}{2} \rfloor$.

More precisely, the following result is what is proved in [2].

Theorem 2.4. *Let $n \in \mathbb{N}$, $I \subseteq [n-1]$, and I_1, \dots, I_s be the connected components of I . Then*

$$(2.6) \quad \sum_{\sigma \in \mathcal{D}_\emptyset^I(C_{n,+})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \left[\begin{matrix} b \\ \lfloor \frac{|I_1|+1}{2} \rfloor, \dots, \lfloor \frac{|I_s|+1}{2} \rfloor \end{matrix} \right]_{x^2} \prod_{k=b+1}^{\lfloor \frac{n-1}{2} \rfloor} (1 - x^{2k}),$$

and

$$(2.7) \quad \sum_{\sigma \in \mathcal{D}_\emptyset^I(C_{n,-})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} 0, & \text{if } n \equiv 1 \pmod{2}, \text{ or} \\ -x^m \sum_{\sigma \in \mathcal{D}_\emptyset^I(C_{n,+})} (-1)^{\ell(\sigma)} x^{L(\sigma)}, & \text{if } n \equiv 0 \pmod{2} \text{ and } b = m, \\ \text{otherwise,} \end{cases}$$

where $b := \sum_{k=1}^s \lfloor \frac{|I_k|+1}{2} \rfloor$, and $m := \lfloor \frac{n}{2} \rfloor$.

3. ODD DIAGRAMS, ODD INVERSION SETS, AND ODD SCHUBERT VARIETIES

In this section we define and study odd analogues of familiar combinatorial and geometric objects which are associated to permutations. More precisely, we define and study odd diagrams, odd Schubert varieties, and odd inversion sets. In particular, we give a geometric interpretation of the odd length function $L : S_n \rightarrow \mathbb{N}_0$ as the dimension of the corresponding odd Schubert variety.

Let $n \in \mathbb{N}$ and $\sigma \in S_n$. We define the *odd diagram* of σ to be

$$D_o(\sigma) := \{(i, j) \in D(\sigma) : \sigma^{-1}(j) \not\equiv i \pmod{2}\}.$$

Clearly $D_o(\sigma) \subset D(\sigma)$ for all $\sigma \in S_n$. Also, note that $|D_o(\sigma)| = L(\sigma)$.




FIGURE 1. The diagram (a) and odd Schubert cell (b) of $\sigma = [4, 1, 3, 2] \in S_4$. Elements of the diagram are indicated by a \bullet , while the permutation is represented by \circ .

Let $(U_1, \dots, U_n) \in C(\sigma)$ and $(a_{i,j})_{(i,j) \in D(\sigma)} \in \mathbb{C}^{D(\sigma)}$ be the corresponding set of complex numbers (see § 2). We define the *odd Schubert cell* of σ to be

$$C_o(\sigma) := \{(U_1, \dots, U_n) \in C(\sigma) : a_{i,j} = 0 \text{ if } (i,j) \in D(\sigma) \setminus D_o(\sigma)\}.$$

So if $(U_1, \dots, U_n) \in C_o(\sigma)$ then there are $(a_{i,j})_{(i,j) \in D_o(\sigma)} \in \mathbb{C}^{D_o(\sigma)}$ such that

$$U_k = \langle \{e_{\sigma(i)} + \sum_{\{j:(i,j) \in D_o(\sigma)\}} a_{i,j} e_j\}_{1 \leq i \leq k} \rangle$$

for all $k \in [n]$, and all flags of this form are in $C_o(\sigma)$. In particular, $C_o(\sigma)$ is isomorphic to an affine space of dimension $L(\sigma)$. We then define the *odd Schubert variety* $X_o(\sigma)$ associated with σ to be the closure of $\pi(C_o(\sigma))$ in $\pi(F_n)$, where π is the embedding defined in § 2. The next result then follows from standard facts (see, e.g., [6, Chap. I, Ex. 2.17]).

Proposition 3.1. *Let $\sigma \in S_n$. Then $X_o(\sigma)$ is a complex projective variety of dimension $L(\sigma)$.*

We illustrate our definitions with an example.

Example 3.2. Let $\sigma := [4, 1, 3, 2]$. Then $C_o(\sigma)$ consists of all flags $(U_1, U_2, U_3, U_4) \in F_4$ for which there are complex numbers $a, b, c \in \mathbb{C}$ such that $U_1 = \langle \{e_4 + ae_2 + be_1\} \rangle$, $U_2 = \langle \{e_4 + ae_2 + be_1, e_1\} \rangle$, and $U_3 = \langle \{e_4 + ae_2 + be_1, e_1, e_3 + ce_2\} \rangle$. The Plücker coordinates of these subspaces are, respectively, $(b, a, 0, 1)$, $(-a, 0, -1, 0, 0, 0)$, and $(0, 1, c, -a)$. The Segre embedding of this triple of points is, after removing 0's,

$$(3.1) \quad (-ab, -abc, ab^2, a^2b, -b, -bc, ab, -a^2, -a^2c, a^3, -a, -ac, a^2, -a, -ac, a^2, -1, -c, a).$$

Therefore, the odd Schubert cell $C_o([4, 1, 3, 2])$ may be identified with all the points in $\mathbb{P}(\mathbb{C}^{18})$ of the form (3.1) where $a, b, c \in \mathbb{C}$.

It is easy to characterize the permutations for which the diagram and the odd diagram coincide.

Proposition 3.3. *Let $\sigma \in S_n$. We have $D(\sigma) = D_o(\sigma)$ if and only if $\sigma(k-2) < \sigma(k)$ for all $3 \leq k \leq n$.*

The following is an immediate consequence.

Corollary 3.4. *Let $\sigma \in S_n$. The odd Schubert variety $X_o(\sigma)$ coincides with the Schubert variety $X(\sigma)$ if and only if $\sigma(k-2) < \sigma(k)$ for all $k \in [3, n]$.*

For $n \in \mathbb{N}$, there are $\binom{n}{\lfloor \frac{n}{2} \rfloor}$ permutations of degree n satisfying the last property, and thus for which L and ℓ coincide; see also the sequence [10, A001405]. We write

$$G_n := \{\sigma \in S_n : \sigma(k-2) < \sigma(k) \text{ for all } 3 \leq k \leq n\} = \{\sigma \in S_n : \ell(\sigma) = L(\sigma)\}$$

for the set of permutations for which all inversions are odd inversions. In particular, the previous proposition implies that for permutations in G_n the odd diagram “faithfully” encodes the permutation itself. It would be interesting to characterize, for every n , the largest subset of S_n on which the map associating to a permutation its odd diagram is injective. In § 6 we put forward a conjecture in this direction. Now, inspired by Proposition 3.3, we prove a product formula for the distribution of the difference of length and odd length, namely of the *even inversions* einv over the symmetric groups.

Proposition 3.5. *Let $n \in \mathbb{N}$. Then*

$$\sum_{\sigma \in S_n} x^{\ell(\sigma) - L(\sigma)} = \sum_{\sigma \in S_n} x^{\text{einv}(\sigma)} = \binom{n}{\lfloor \frac{n}{2} \rfloor} \prod_{i=1}^n \left(\frac{1 - x^{\lfloor \frac{i}{2} \rfloor}}{1 - x} \right).$$

Proof. Consider the subgroups

$$S_o := \langle (i, i+2) : i \equiv 1 \pmod{2} \rangle \simeq S_{\lfloor \frac{n}{2} \rfloor}$$

and

$$S_e := \langle (i, i+2) : i \equiv 0 \pmod{2} \rangle \simeq S_{\lfloor \frac{n}{2} \rfloor}.$$

Let π_o and π_e denote the natural projections from S_n onto S_o and S_e , respectively. It is easy to see that $\text{einv}(\sigma) = \text{inv}(\pi_o(\sigma)) + \text{inv}(\pi_e(\sigma))$. Therefore,

$$(3.2) \quad \sum_{\sigma \in S_o \times S_e} x^{\text{einv}(\sigma)} = \left(\sum_{\sigma \in S_{\lfloor \frac{n}{2} \rfloor}} x^{\text{inv}(\sigma)} \right) \left(\sum_{\sigma \in S_{\lfloor \frac{n}{2} \rfloor}} x^{\text{inv}(\sigma)} \right) = \prod_{i=1}^n \left(\frac{1 - x^{\lfloor \frac{i}{2} \rfloor}}{1 - x} \right).$$

The proposition follows, as

$$S_n = \bigcup_{\tau \in G_n} \tau(S_o \times S_e)$$

and the identity in (3.2) also holds on all of the $\binom{n}{\lfloor \frac{n}{2} \rfloor}$ cosets. \square

The following is a straightforward corollary of this formula.

Corollary 3.6. *The polynomial $\sum_{\sigma \in S_n} x^{\text{einv}(\sigma)}$ is symmetric and unimodal for all $n \in \mathbb{N}$.*

Remark 3.7. The proof of Proposition 3.5 shows that similar results hold for the polynomials giving the distribution of inversions between positions which are congruent modulo any positive integer. More precisely, for $k, n \in \mathbb{N}$ and $\sigma \in S_n$, let $\text{inv}_{k,0}(\sigma)$ denote the number of inversions between positions congruent modulo k in σ ,

$$(3.3) \quad \text{inv}_{k,0}(\sigma) = |\{(i, j) \in \text{Inv}(\sigma) : j - i \equiv 0 \pmod{k}\}|.$$

If $n = mk + r$ for some $m \in \mathbb{N}_0$ and $0 \leq r < k$ then

$$\begin{aligned} \sum_{\sigma \in S_n} x^{\text{inv}_{k,0}(\sigma)} &= \binom{n}{\underbrace{m, \dots, m}_{k-r}, \underbrace{m+1, \dots, m+1}_r} \left(\sum_{\sigma \in S_m} x^{\text{inv}(\sigma)} \right)^{k-r} \left(\sum_{\sigma \in S_{m+1}} x^{\text{inv}(\sigma)} \right)^r \\ &= \binom{n}{\underbrace{m, \dots, m}_{k-r}, \underbrace{m+1, \dots, m+1}_r} \prod_{i=1}^n \left[\left[\frac{i}{k} \right] \right]_x. \end{aligned}$$

It is clear that the odd Schubert variety attached to a permutation $\sigma \in S_n$ does not really depend on the permutation, but only on its odd diagram. While it is well known that diagrams are in bijection with permutations, this is not the case for odd diagrams. Indeed, already in S_3 there are two permutations with the same odd diagram: $D_o([2, 1, 3]) = \{(1, 1)\} = D_o([3, 1, 2])$. It is therefore a natural and interesting problem to characterize odd diagrams of permutations. This is probably not an easy task since no characterization of (ordinary) diagrams seems to be known. Also, odd diagrams are probably not equivalent to any known combinatorial objects. In fact, if we let o_n be the number of different odd diagrams of permutations in S_n then the first values of the sequence $\{o_n\}_{n \in \mathbb{N}}$ are 1, 2, 5, 17, 70, 351, 2041, 13732, and this sequence is new in the OEIS ([10, A335926]).

While we are unable to characterize odd diagrams, we can characterize a closely related set. For $\sigma \in S_n$ let

$$\text{Inv}_o(\sigma) := \{(i, j) \in \text{Inv}(\sigma) : j \not\equiv i \pmod{2}\}.$$

We call $\text{Inv}_o(\sigma)$ the *odd inversion set* of σ . Note that $|\text{Inv}_o(\sigma)| = L(\sigma)$ and that $\text{Inv}_o(\sigma) \subset OS_n$ where $OS_n := \{(i, j) \in [n]^2 : i < j, j \not\equiv i \pmod{2}\}$ is the *odd staircase* of size n .

The following result follows easily from our definitions.

Lemma 3.8. *Let $\sigma \in S_n$ and $(i, j) \in [n]^2$. Then $(i, \sigma(j)) \in D_o(\sigma)$ if and only if $(i, j) \in \text{Inv}_o(\sigma)$.*

We now characterize the odd inversion sets of permutations. Recall that the *Turán graph* (see, e.g., [8]) is the complete bipartite graph $T_n := ([n], E_n)$ where, if $i, j \in [n]$, then $\{i, j\} \in E_n$ if and only if $i \not\equiv j \pmod{2}$.

Given $I \subseteq OS_n$ we define an orientation A_I of T_n as follows. Let $\{i, j\}_< \in E_n$. Then we let $i \rightarrow j$ in A_I if and only if $(i, j) \in I$. So, for example, for $I = \{(1, 4), (2, 3), (2, 5), (3, 4)\}$ we get the orientation of T_5 in Figure 2. We then have the following simple characterization of odd inversion sets of permutations in terms of orientations of T_n .

Proposition 3.9. *Let $I \subseteq OS_n$. Then there is a permutation $\sigma \in S_n$ such that $I = \text{Inv}_o(\sigma)$ if and only if A_I is acyclic.*

Proof. Suppose first that $I = \text{Inv}_o(\sigma)$ for some $\sigma \in S_n$. Then we have that, for all $\{i, j\}_< \in E_n$, $i \rightarrow j$ in A_I if and only if $\sigma(i) > \sigma(j)$, so A_I is acyclic.

Conversely, suppose that A_I is acyclic. It is then easy to see, by induction on the number of vertices, that given any acyclic orientation of a graph $D = (V, E)$ there is a bijection $f : V \rightarrow [|V|]$ such that if $\{x, y\} \in E$ then $x \rightarrow y$ if and only if $f(x) > f(y)$. Indeed, as the orientation is acyclic there is either a source or a sink $v \in V$. Say v is a source. Now define $f(v) := |V|$, remove v and all edges incident to it from D and argue by induction. In particular, there is $\sigma \in S_n$ such that $i \rightarrow j$ if and only if $\sigma(i) > \sigma(j)$ for all $\{i, j\}_< \in E_n$. So $I = \text{Inv}_o(\sigma)$. \square

We illustrate the preceding result with an example.

Example 3.10. Given $\sigma = [3, 5, 4, 1, 2] \in S_5$ we have $\text{Inv}_o(\sigma) = \{(1, 4), (2, 3), (2, 5), (3, 4)\}$, which defines the acyclic orientation in Figure 2. Conversely, given $I = \{(1, 4), (2, 3), (2, 5), (3, 4)\}$, following the steps of the induction and maintaining notation from the above proof we get: $f(2) = 5$, $f(1) = 4$, $f(3) = 3$, $f(5) = 2$ and $f(4) = 1$, which defines the permutation $\tau = [4, 5, 3, 1, 2]$ with $\text{Inv}_o(\tau) = I$. As expected, this is not the only permutation of S_5 with this odd inversion set. There are 6 permutations with odd inversion set equal to I : $[2, 5, 3, 1, 4]$, $[2, 5, 4, 1, 3]$, $[3, 5, 2, 1, 4]$, $[4, 5, 2, 1, 3]$, σ and τ .




FIGURE 2. Acyclic orientation of T_5 defined by $I = \{(1, 4), (2, 3), (2, 5), (3, 4)\}$

We conclude this section by giving several necessary conditions on a subset of $[n]^2$ to be the odd diagram of a permutation. It is easy to see that to characterize subsets of $[n]^2$ which are odd diagrams of a permutation it is enough to consider those that have at least one element in the first row or column.

Proposition 3.11. *Let $S \subseteq [2, n]^2$. Then there exists $\sigma \in S_n$ such that $D_o(\sigma) = S$ if and only if there exists $\tau \in S_{n-1}$ such that $D_o(\tau) = \{(x-1, y-1) : (x, y) \in S\}$.*

Proof. Suppose there is $\sigma \in S_n$ such that $S = D_o(\sigma)$. Then $\sigma^{-1}(1) = 1$ (otherwise $(\sigma^{-1}(1) - 1, 1) \in D_o(\sigma)$). So $\tau = [\sigma(2) - 1, \dots, \sigma(n) - 1] \in S_{n-1}$ is the desired permutation. Conversely, if $\tau \in S_{n-1}$ is such that $D_o(\tau) = \{(x-1, y-1) : (x, y) \in S\}$ then $\sigma = [1, \tau(1) + 1, \dots, \tau(n-1) + 1]$ has odd diagram equal to S . \square

Note that the previous proof implies that if an odd diagram does not have any elements in the first column, then it has no elements in the first row.

The following result gives some necessary enumerative conditions for a subset of $[n]^2$ to be the odd diagram of a permutation in S_n .

Proposition 3.12. *Let $\sigma \in S_n$ and $S := D_o(\sigma)$. Then:*

- (i) *if $(i, j) \in S$ then $|\{k \in [j-1] : (i, k) \notin S\}| \leq \min\{\lfloor \frac{n+i-2}{2} \rfloor, j-1\}$;*
- (ii) *if $(i, j) \in S$ then $|\{k \in [i-1] : (k, j) \notin S\}| \leq \min\{\lceil \frac{i+2j-3}{2} \rceil, i-1\}$;*
- (iii) *if $i \in [n]$ then $|\{j \in [n] : (i, j) \in S\}| \leq \lceil \frac{n-i}{2} \rceil$;*
- (iv) *if $j \in [n]$ then $|\{i \in [n] : (i, j) \in S\}| \leq \min\{\lceil \frac{n-1}{2} \rceil, n-j\}$.*

Proof. Suppose that $(i, j) \in S$. Let $k \in [j-1]$ be such that $(i, k) \notin S$. Then either $\sigma^{-1}(k) < i$ or $\sigma^{-1}(k) > i$ and $\sigma^{-1}(k) \equiv i \pmod{2}$. But there are at most $i-1$ possibilities in the first case, and at most $\lfloor \frac{n-i}{2} \rfloor$ in the second case. This proves (i).

Similarly, let $(i, j) \in S$, and $k \in [i-1]$ be such that $(k, j) \notin S$. Then either $\sigma(k) < j$ or $\sigma(k) > j$ and $\sigma(k) \equiv j \pmod{2}$, and there are at most $j-1$ possibilities in the first case and at most $\lceil \frac{i-1}{2} \rceil$ in the second one.

Finally, if $i, j \in [n]$ are such that $(i, j) \in S$ then $\sigma^{-1}(j) > i$ and $\sigma^{-1}(j) \not\equiv i \pmod{2}$, which proves (iii). The proof of (iv) is analogous and is omitted. \square

The next proposition collects a number of configurations which cannot occur in odd diagrams (see also Figures 3 and 4).

Proposition 3.13. *Let $\sigma \in S_n$ and $S := D_o(\sigma)$. Then:*

- (i) *if $(i, j), (k, l) \in S$ with $i \leq k, j \geq l$, and $i \equiv k \pmod{2}$ then $(i, l) \in S$;*




FIGURE 3. A forbidden configuration for odd diagrams

- (ii) if $(i, j), (k, l) \in S$ with $i < k$, $j \geq l$, and $i \not\equiv k \pmod{2}$ then $(i, l) \notin S$;
- (iii) if $\{i_1, \dots, i_{k+1}\}_{<} \subseteq [n]$ and $j \in [n]$ are such that $i_r \equiv i_{r+1} \pmod{2}$, $\{i_r\} \times [j, j + k - 1] \cap S \neq \emptyset$, and $(i_r, j + k) \notin S$ for $r \in [k]$, then $(i_{k+1}, j + k) \notin S$;
- (iv) if $\{j_1, \dots, j_{\lceil \frac{k+1}{2} \rceil}\}_{<} \subseteq [n]$, and $i_1, \dots, i_{\lceil \frac{k+1}{2} \rceil} \in [i, i + k - 1]$ are such that $(i_r, j_r) \in S$, $(i + k, j_r) \notin S$, and $i_r \equiv i + k \pmod{2}$ for $r = 1, \dots, \lceil \frac{k+1}{2} \rceil$, then $(i + k, j_{\lceil \frac{k+1}{2} \rceil}) \notin S$;
- (v) if j is the minimum index for which $([n] \times [j]) \cap S \neq \emptyset$, then if $i \in [n]$ is such that $(i, j) \in S$ and $(\{i + 1\} \times [j + 1, n]) \cap S \neq \emptyset$ then $(i + 2, j) \in S$;
- (vi) if $(i, j) \in [n - 2] \times [2, n - 1]$ are such that $(i, j), (i + 2, j - 1) \in S$ and $(i + 1, j), (i + 2, j) \notin S$ then $\{i + 1\} \times [j + 1, n] \cap S = \emptyset$.

Proof. We first prove (i). Since $(i, j), (k, l) \in S$ we have that $\sigma(i) > j$, $\sigma^{-1}(l) > k$, and $\sigma^{-1}(l) \not\equiv k \pmod{2}$. Hence $\sigma^{-1}(l) \not\equiv i \pmod{2}$ so $(i, l) \in D_o(\sigma)$.

The proof of (ii) is identical, except that now $\sigma^{-1}(l) \equiv i \pmod{2}$ so $(i, l) \notin D_o(\sigma)$.

We now prove (iii) (see Figure 3). Suppose, by contradiction, that $(i_{k+1}, j + k) \in D_o(\sigma)$. Then $\sigma^{-1}(j + k) > i_{k+1}$ and $\sigma^{-1}(j + k) \not\equiv i_{k+1} \pmod{2}$. Let $r \in [k]$. Since $(i_r, j + k) \notin S$, by what we have just observed we have that $\sigma(i_r) < j + k$. On the other hand, since $\{i_r\} \times [j, j + k - 1] \cap S \neq \emptyset$, $\sigma(i_r) > j$. So $\sigma(i_r) \in [j + 1, j + k - 1]$ for all $r \in [k]$, which is a contradiction.

To prove (iv) suppose, by contradiction, that $(i + k, j_{\lceil \frac{k+1}{2} \rceil}) \in S$. Then $\sigma(i + k) > j_{\lceil \frac{k+1}{2} \rceil}$. Let $r \in [\lceil \frac{k+1}{2} \rceil]$. Since $(i_r, j_r) \in S$, we have that $\sigma(i_r) > j_r$, $\sigma^{-1}(j_r) \not\equiv i_r \pmod{2}$, and $\sigma^{-1}(j_r) > i_r$. So $\sigma^{-1}(j_r) \not\equiv i + k \pmod{2}$. On the other hand, since $(i + k, j_r) \notin S$, $\sigma^{-1}(j_r) < i + k$. So $\sigma^{-1}(j_r) \in [i + 1, i + k - 1]$ and $\sigma^{-1}(j_r) \not\equiv i + k \pmod{2}$ for all $r \in [\lceil \frac{k+1}{2} \rceil]$, which is a contradiction. This proves (iv).

Parts (v) and (vi) are easy to check (see Figure 4). □

Note that for $j = l$ part (ii) of Proposition 3.13 implies that if $(i + 1, j) \in S$ then $(i, j) \notin S$.

In the following result, we collect a few more conditions satisfied by odd diagrams which say that some configurations can only appear in certain areas of the square grid.

Proposition 3.14. *Let $\sigma \in S_n$ and $S := D_o(\sigma)$. Then:*

- (i) if $i, j, k \in [n]$ are such that $(i + 2k - 1, j + k - 1) \in [n]^2$, $\{i\} \times [j, j + k - 1] \subseteq S$, and $[i + 1, i + 2k - 1] \times [j, j + k - 1] \subseteq [n]^2 \setminus S$, then $j \geq k$;
- (ii) if $i, j \in [n]$, $k \in \mathbb{N}$ are such that $(i + 2k, j + 2k) \in [n]^2$, and

$$(i + a, j + b) \in S \Leftrightarrow a \equiv b \equiv 0 \pmod{2}, \text{ and } a + b \leq 2k$$





FIGURE 4. Forbidden configurations for odd diagrams

FIGURE 5. This configuration can only appear if $i + j \geq 4$

for all $(a, b) \in [0, 2k]^2$, then $i + j \geq k + 2$.

Proof. We first prove (i). Since $\{i\} \times [j, j+k-1] \subseteq S$ we have that $\sigma^{-1}(j+r-1) > i$ and $\sigma^{-1}(j+r-1) \not\equiv i \pmod{2}$ for all $r \in [k]$. Therefore there is $r_0 \in [k]$ such that $\sigma^{-1}(j+r_0-1) \geq i+2k-1$. Hence, since $\{i+2t\} \times [j, j+k-1] \subseteq [n]^2 \setminus S$ for all $t \in [k-1]$, $\sigma(i+2t) \leq j+k-1$ for all $t \in [k-1]$, so $\sigma(i+2t) < j$ for all $t \in [k-1]$, and the result follows.

We now prove (ii) (see Figure 5). We show that

$$(3.4) \quad |\{a \in [k] : \sigma(i+2a-1) < j\} \cup \{b \in [k] : \sigma^{-1}(j+2b-1) < i\}| \geq k,$$

which implies our claim.

We proceed by induction on $k \geq 0$. Let $k = 1$. Assume, by contradiction, that (3.4) fails. Then $\sigma(i+1) \geq j$ and $\sigma^{-1}(j+1) \geq i$. Hence, since $(i+2, j) \in S$ and $(i, j+1) \notin S$, $\sigma(i+1) \geq j+2$. Similarly, since $(i, j+2) \in S$ and $(i, j+1) \notin S$, $\sigma^{-1}(j+1) \geq i+2$. But then either $(i, j+1) \in S$ (if $\sigma^{-1}(j+1) \not\equiv i \pmod{2}$) or $(i+1, j+1) \in S$ (if $\sigma^{-1}(j+1) \equiv i \pmod{2}$), which is a contradiction. So assume $k \geq 2$. Since $(i, j+2k) \in S$ we have that $\sigma(i) > j+2k$, $\sigma^{-1}(j+2k) > i$, and $\sigma^{-1}(j+2k) \not\equiv i \pmod{2}$.

If $\sigma^{-1}(j+2k) = i+1$ then $\sigma^{-1}(j+2b-1) < i$ for all $b \in [k]$ (else either $(i, j+2b-1) \in S$ or $(i+1, j+2b-1) \in S$ for some $b \in [k]$) and the claim holds.

Assume now that $\sigma^{-1}(j+2k) = i+2b+1$ for some $b \in [k-1]$ then $j+2k-1 = \sigma(i+2)$ (else $(i+2, j+2k) \in S$). Hence $\sigma(i+1) < j$ (for if $\sigma(i+1) > j+2k$ then $(i+1, j+2k-1) \in S$, while if $j \leq \sigma(i+1) < j+2k$ then necessarily $\sigma(i+1) = j+2a-1$ for some $a \in [k-1]$, which implies that $(i, j+2a-1) \in S$, which again contradicts our hypotheses). In an analogous way one concludes that $\sigma^{-1}(j+1) < i$ (for if $\sigma^{-1}(j+1) \geq i+2k$ then either $(i, j+1) \in S$ or $(i+2b+1, j+1) \in S$, while if $i \leq \sigma^{-1}(j+1) < i+2k$ then $\sigma^{-1}(j+1) \not\equiv i \pmod{2}$, so again $(i, j+1) \in S$). Now, by our induction hypothesis (applied to $i+2, j+2, k-2$) we conclude

that

$$|\{a \in [k-2] : \sigma(i+2a+1) < j+2\} \cup \{b \in [k-2] : \sigma^{-1}(j+2b+1) < i+2\}| \geq k-2.$$

But if $\sigma(i+2a+1) < j+2$ for some $a \in [k-2]$ then $\sigma(i+2a+1) < j$ (for if $\sigma(i+2a+1) = j+1$ then $(i, j+1) \in S$, while $\sigma(i+2a+1) \neq j$ since $(i+2k, j) \in S$). Also, since $\sigma(i+1) < j$, and $\sigma(i) > j+2k$, if $\sigma^{-1}(j+2b+1) < i+2$ for some $b \in [k-2]$, then $\sigma^{-1}(j+2b+1) < i$. Therefore

$$|\{a \in [k-2] : \sigma(i+2a+1) < j\} \cup \{b \in [k-2] : \sigma^{-1}(j+2b+1) < i\}| \geq k-2,$$

and this implies (3.4) since $\sigma(i+1) < j$ and $\sigma^{-1}(j+1) < i$.

Finally, assume that $\sigma^{-1}(j+2k) = i+2b+1$ for some $b \geq k$. Then as in the previous case we conclude that $j+2k-1 = \sigma(i+2)$ and $\sigma(i+1) < j$. Therefore $\sigma(i+4) = j+2k-3$ (for if $\sigma(i+4) > j+2k$ then $(i+4, j+2k) \in S$, while if $\sigma(i+4) = j+2k-2$ then $(i+2, j+2k-2) \notin S$). This implies that $\sigma(i+3) < j$ (for if $\sigma(i+3) > j+2k-3$ then $(i+3, j+2k-3) \in S$). Now, by our induction hypothesis (applied to $i+4, j, k-2$) we have that

$$|\{a \in [k-2] : \sigma(i+2a+3) < j\} \cup \{b \in [k-2] : \sigma^{-1}(j+2b-1) < i+4\}| \geq k-2.$$

But if $\sigma^{-1}(j+2b-1) < i+4$ for some $b \in [k-2]$ then $\sigma^{-1}(j+2b-1) < i$ so

$$|\{a \in [k-2] : \sigma(i+2a+3) < j\} \cup \{b \in [k-2] : \sigma^{-1}(j+2b-1) < i\}| \geq k-2$$

and this proves (3.4) since $\sigma(i+1) < j$ and $\sigma(i+3) < j$.

This concludes the induction step and hence the proof. \square

The conditions in Propositions 3.12, 3.13 and 3.14 are also sufficient for $S \subseteq [n]^2$, with $n \leq 4$, to be the odd diagram of a permutation. However, for $n \geq 5$ they fail to characterize these subsets. For instance, $\{(1, 1), (1, 2), (3, 2), (4, 4)\}$ is not the odd diagram of any permutation.

4. SHIFTING AND REVERSING

In this section we derive a number of results concerning operations that can be performed on the subsets defining a descent class, after which the sign-twisted generating function of the odd length remains the same or changes in a controlled way. We also give sufficient conditions on a descent class for the corresponding sign-twisted generating function to be zero, and we compute it explicitly for the descent class of the alternating permutations.

Recall that a permutation in the descent class $\mathcal{D}_J^I(S_n)$ is a permutation which is increasing in the positions corresponding to $I \cup (I+1)$ and decreasing in $J \cup (J+1)$.

The proofs of the following two results are similar to those of [2, Lemma 3.1 and Proposition 3.3]. However, for the reader's convenience, and for completeness, we provide proofs here. Our first lemma shows that the sign-twisted generating function of the odd length is zero on the non-chessboard elements of a descent class in which the ascents and the descents are disjoint.

Lemma 4.1. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_J^I(C_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

Proof. Let $\sigma \in \mathcal{D}_J^I(S_n) \setminus \mathcal{D}_J^I(C_n)$. Then there exists $i \in [n-1]$ such that $\sigma^{-1}(i) \equiv \sigma^{-1}(i+1) \pmod{2}$ (else either $\sigma^{-1}(i) \equiv i \pmod{2}$ for all $i \in [n]$ or $\sigma^{-1}(i) \equiv i+1 \pmod{2}$ for all $i \in [n]$ so $\sigma \in C_n$). Let i be minimal with this property and define $\sigma^* = s_i \sigma$. This is a well

defined involution on $\mathcal{D}_J^I(S_n) \setminus \mathcal{D}_J^I(C_n)$ since $|\sigma^{-1}(i) - \sigma^{-1}(i+1)| \geq 2$. But $L(\sigma^*) = L(\sigma)$ and $\ell(\sigma^*) = \ell(\sigma) \pm 1$, which implies the result. \square

The next result is the first of a series of invariance results for the sign-twisted generating function of the odd length over a descent class $\mathcal{D}_J^I(S_n)$. It shows that a connected component of odd cardinality of the ascents can be shifted or enlarged of one unit to the right without changing the generating function, as long as it remains a connected component.

Proposition 4.2. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$. Let $i \in \mathbb{N}$, $k \in \mathbb{N}_0$ be such that $[i, i+2k] \subseteq I$ is a connected component of $I \cup J$ and $i+2k+2 \notin I \cup J$.*

Then

$$(4.1) \quad \sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_J^{I \cup \tilde{I}}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_J^{\tilde{I}}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}$$

where $\tilde{I} := (I \setminus \{i\}) \cup \{i+2k+1\}$.

Proof. First note that, by our hypotheses, $(I \cup \tilde{I}) \cap J = \emptyset$. We have

$$(4.2) \quad \begin{aligned} \sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} &= \sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \\ \sigma(i) > \sigma(i+2k+2)}} (-1)^{\ell(\sigma)} x^{L(\sigma)} + \sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \sigma(i+2k+1) < \\ \sigma(i+2k+2)}} (-1)^{\ell(\sigma)} x^{L(\sigma)} \\ &+ \sum_{j=1}^{2k+1} \left(\sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \sigma(i+j-1) < \\ \sigma(i+2k+2) < \sigma(i+j)}} (-1)^{\ell(\sigma)} x^{L(\sigma)} \right). \end{aligned}$$

Let $r \in [k]$. Note that, by our hypotheses, $i-1 \notin J$ and $i+2k+1 \notin J$. Therefore the map $\sigma \mapsto \tilde{\sigma} := \sigma(i+2k+2, i+2r)$ is a bijection between $\{\sigma \in \mathcal{D}_J^I(S_n) : \sigma(i+2r) < \sigma(i+2k+2) < \sigma(i+2r+1)\}$ and $\{\sigma \in \mathcal{D}_J^I(S_n) : \sigma(i+2r-1) < \sigma(i+2k+2) < \sigma(i+2r)\}$. Furthermore, $\ell(\tilde{\sigma}) = \ell(\sigma) + 1$ and $L(\tilde{\sigma}) = L(\sigma)$ so

$$\sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \sigma(i+2r) < \\ \sigma(i+2k+2) < \sigma(i+2r+1)}} (-1)^{\ell(\sigma)} x^{L(\sigma)} = - \sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \sigma(i+2r-1) < \\ \sigma(i+2k+2) < \sigma(i+2r)}} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

Similarly, the bijection $\sigma \mapsto \sigma(i+2k+2, i)$ shows that

$$\sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \\ \sigma(i+2k+2) < \sigma(i)}} (-1)^{\ell(\sigma)} x^{L(\sigma)} = - \sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \sigma(i) < \\ \sigma(i+2k+2) < \sigma(i+1)}} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

Therefore, by (4.2),

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\substack{\sigma \in \mathcal{D}_J^I(S_n): \sigma(i+2k+1) < \\ \sigma(i+2k+2)}} (-1)^{\ell(\sigma)} x^{L(\sigma)}$$

and the first equality in (4.1) follows.

The proof of the second equality is similar, and is therefore omitted. \square

Note that the proof of the previous result actually yields that if $I, J \subseteq [n-1]$ are such that $I \cap J = \emptyset$, and if $i \in \mathbb{N}$ and $k \in \mathbb{N}_0$ are such that $[i, i+2k+1]$ is a connected component of $I \cup J$ and $i+2k+1 \in J$, $[i, i+2k] \subseteq I$, then $\tilde{I} \cap J \neq \emptyset$, hence

$$\sum_{\sigma \in \mathcal{D}_J^{\tilde{I}}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = 0.$$

This is a special case of a more general fact (see Proposition 5.1).

The following is the “left” version of Proposition 4.2. Informally, it shows that a connected component of odd cardinality of the ascents can be shifted or enlarged of one unit to the left without changing the sign-twisted generating function, as long as it remains a connected component.

Proposition 4.3. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$. Let $i \in \mathbb{N}$, $k \in \mathbb{N}_0$ be such that $[i+1, i+2k+1] \subseteq I$ is a connected component of $I \cup J$, and $i-1 \notin I \cup J$. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_J^{I \cup \bar{I}}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_J^{\bar{I}}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}$$

where $\bar{I} := (I \setminus \{i+2k+1\}) \cup \{i\}$.

Proof. Under our hypotheses we have that $(I \cup \bar{I}) \cap J = \emptyset$, $[i, i+2k]$ is a connected component of $\bar{I} \cup J$, $[i, i+2k] \subseteq \bar{I}$, and $i+2k+2 \notin \bar{I} \cup J$, so the result follows from Proposition 4.2. \square

We now show that a connected component of even cardinality of the descents can be “transformed” (or “reversed”) into a connected component of the ascents, by changing the generating function by a simple factor.

Lemma 4.4. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$, and $i, k \in \mathbb{N}$ be such that $K := [i, i+2k-1]$ is a connected component of $I \cup J$, $K \subseteq J$. Then*

$$(4.3) \quad \sum_{\sigma \in \mathcal{D}_J^I(C_{n,\pm})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = (-1)^k x^{k(k+1)} \sum_{\sigma \in \mathcal{D}_{J \setminus K}^{I \cup K}(C_{n,\pm})} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

In particular,

$$(4.4) \quad \sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = (-1)^k x^{k(k+1)} \sum_{\sigma \in \mathcal{D}_{J \setminus K}^{I \cup K}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

Proof. We have

$$\sum_{\sigma \in \mathcal{D}_J^I(C_{n,+})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\tau \in \mathcal{D}_{J \setminus K}^{I \cup K}(C_{n,+})} (-1)^{\ell(\bar{\tau})} x^{L(\bar{\tau})},$$

where $\bar{\tau} := [\tau(1), \dots, \tau(i-1), \tau(i+2k), \dots, \tau(i+1), \tau(i), \tau(i+2k+1), \dots, \tau(n)]$. But $\ell(\bar{\tau}) = \ell(\tau) + (2k+1)k$ and, by Proposition 2.2 $L(\bar{\tau}) = L(\tau) + k(k+1)$, thus

$$\sum_{\tau \in \mathcal{D}_{J \setminus K}^{I \cup K}(C_{n,+})} (-1)^{\ell(\bar{\tau})} x^{L(\bar{\tau})} = (-1)^k x^{k(k+1)} \sum_{\tau \in \mathcal{D}_{J \setminus K}^{I \cup K}(C_{n,+})} (-1)^{\ell(\tau)} x^{L(\tau)}$$

as desired. Similarly for $C_{n,-}$. \square

In a similar way, it is easy to determine the generating function on the descent class obtained by transforming all the descents into ascents, and conversely, as shown in the following result.

Proposition 4.5. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = (-1)^{\ell(w_0)} x^{L(w_0)} \sum_{\sigma \in \mathcal{D}_I^J(S_n)} (-1)^{\ell(\sigma)} \left(\frac{1}{x}\right)^{L(\sigma)}.$$

Proof. It is clear that the map $\sigma \mapsto w_0\sigma$ is a bijection from $\mathcal{D}_J^I(S_n)$ to $\mathcal{D}_I^J(S_n)$. Therefore, by Proposition 2.2 we have

$$\begin{aligned} \sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} &= \sum_{\tau \in \mathcal{D}_I^J(S_n)} (-1)^{\ell(w_0\tau)} x^{L(w_0\tau)} \\ &= (-1)^{\ell(w_0)} x^{L(w_0)} \sum_{\tau \in \mathcal{D}_I^J(S_n)} (-1)^{\ell(\tau)} \left(\frac{1}{x}\right)^{L(\tau)}. \end{aligned}$$

□

Remark 4.6. The bijection $\sigma \mapsto w_0\sigma$ in the proof of Proposition 4.5 restricts to a bijection between chessboard elements of the relevant descent classes. In particular, if n is even it is a bijection between $\mathcal{D}_J^I(C_{n,+})$ and $\mathcal{D}_I^J(C_{n,-})$.

The sign-twisted generating function is also invariant under left and right shifting of connected components of the descents, under certain hypotheses. The next two results are analogous to Proposition 4.2 and 4.3, respectively. The first shows that a connected component of odd cardinality of the descents can be shifted (or enlarged of one unit) to the right, as long as it remains a connected component.

Proposition 4.7. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$. Let $i \in \mathbb{N}$, $k \in \mathbb{N}_0$ be such that $[i, i+2k]$ is a connected component of $I \cup J$, $[i, i+2k] \subseteq J$, and $i+2k+2 \notin I \cup J$. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{J \cup \tilde{J}}^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{\tilde{J}}^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)},$$

where $\tilde{J} := (J \setminus \{i\}) \cup \{i+2k+1\}$.

Proof. By Proposition 4.2 we have

$$\sum_{\sigma \in \mathcal{D}_I^J(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_I^{J \cup \tilde{J}}(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{\tilde{J}}^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}$$

so the result follows from Proposition 4.5. □

In a similar way, using Proposition 4.3, we obtain the following invariance result under left shifting of a connected component of odd cardinality of the descents.

Proposition 4.8. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$, and $i \in \mathbb{N}$, $k \in \mathbb{N}_0$ be such that $[i+1, i+2k+1]$ is a connected component of $I \cup J$, $[i+1, i+2k+1] \subseteq J$, and $i-1 \notin I \cup J$. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{J \cup \tilde{J}}^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{\tilde{J}}^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)},$$

where $\tilde{J} := (J \setminus \{i+2k+1\}) \cup \{i\}$.

Computer calculations suggest that the operation of shifting can be performed under weaker hypotheses, namely even if the connected component to be shifted is not contained in I (as required in Proposition 4.2) and therefore not contained in J (as in Proposition 4.7). More precisely, we conjecture the following.

Conjecture 4.9. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$. Let $i \in \mathbb{N}$, $k \in \mathbb{N}_0$ be such that $i+2k+2 \notin I \cup J$ and $[i, i+2k]$ is a connected component of $I \cup J$, say $[i, i+2k] = A \cup B$, where $A \subseteq I$ and $B \subseteq J$. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{\tilde{J}}^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}$$

where $\tilde{I} := (I \setminus A) \cup (A+1)$ and $\tilde{J} := (J \setminus B) \cup (B+1)$.

5. DESCENT CLASSES

In this section we investigate the sign-twisted generating function of the odd length over descent classes. More precisely, we give sufficient conditions on a descent class for the generating function to be zero, and we compute it explicitly for the alternating permutations and for a general family of descent classes which includes all quotients.

Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$, and $i \in [n]$. We say that i is a *peak* of $\mathcal{D}_J^I(S_n)$ if $i \in (I+1) \setminus I$ or $i \in J \setminus (J+1)$. Similarly, i is a *valley* if $i \in I \setminus (I+1)$ or $i \in (J+1) \setminus J$.

Proposition 5.1. *Let $I, J \subseteq [n-1]$, $I \cap J = \emptyset$, and $i \in \mathbb{N}$, $k \in \mathbb{N}_0$ be such that $[i, i+2k+1]$ is a connected component of $I \cup J$ and $v \not\equiv p \pmod{2}$ for any $v, p \in [i, i+2k+2]$, v valley, p peak. Then*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = 0.$$

Proof. Let $\sigma \in \mathcal{D}_J^I(S_n)$. Let $\{a_1, \dots, a_{2k+3}\}_< := \{\sigma(i), \sigma(i+1), \dots, \sigma(i+2k+2)\}$. Let $v := \sigma^{-1}(a_1)$. Then v is a valley (for if $i < v < i+2k+2$ then $\sigma(v-1) > \sigma(v) < \sigma(v+1)$ so $v \in I \cap (J+1)$, while if $v = i$ then $\sigma(v) < \sigma(v+1)$ so $v \in I \setminus (I+1)$, and if $v = i+2k+2$ then $\sigma(v-1) > \sigma(v)$ so $v \in (J+1) \setminus J$). Similarly, $\sigma^{-1}(a_{2k+3})$ is a peak. Therefore, by our hypotheses, $\sigma^{-1}(a_1) \not\equiv \sigma^{-1}(a_{2k+3}) \pmod{2}$.

Let $j := \min\{r \in [2k+2] : \sigma^{-1}(a_r) \equiv \sigma^{-1}(a_{r+1}) \pmod{2}\}$ (note that j certainly exists for if $\sigma^{-1}(a_1) \not\equiv \sigma^{-1}(a_2) \not\equiv \dots \not\equiv \sigma^{-1}(a_{2k+3}) \pmod{2}$ then $\sigma^{-1}(a_1) \equiv \sigma^{-1}(a_{2k+3}) \pmod{2}$ which is a contradiction), and $\hat{\sigma} := (a_j, a_{j+1}) \sigma$. Then $\hat{\sigma} \in \mathcal{D}_J^I(S_n)$, $\ell(\hat{\sigma}) = \ell(\sigma) \pm 1$, $L(\hat{\sigma}) = L(\sigma)$ and the map $\sigma \mapsto \hat{\sigma}$ is an involution. The result follows. \square

Note that the converse of the previous result does not hold. For example, if $n = 8$, $I = \{1, 2, 4\}$, and $J = \{3, 5, 6\}$ then the sign-twisted generating function for $\mathcal{D}_J^I(S_8)$ is zero but $\mathcal{D}_J^I(S_8)$ has peaks $\{3, 5\}$ and valleys $\{1, 4, 7\}$. On the other hand, under the weaker hypothesis that there exist at least one peak and one valley with different parities the generating function is not, in general, zero. For example, if $n = 8$, $I = \{1, 2, 4\}$, and $K = \{3, 5, 6, 7\}$ then $\mathcal{D}_K^I(S_8)$ has peaks $\{3, 5\}$ and valleys $\{1, 4, 8\}$ but the corresponding generating function is $-x^6(1+x^2+x^4)$. It would be interesting to find necessary and sufficient conditions on I and J for the sign-twisted generating function on $\mathcal{D}_J^I(S_n)$ to be zero.

Proposition 5.1 implies that if $I \cup J$ has a “zig-zag” connected component K of even cardinality (i.e., if all even elements of K are in I and all odd ones are in J , or conversely) then the corresponding sign-twisted generating function is zero. Thus, this is in particular true for the alternating permutations of a symmetric group of odd degree. This makes it natural to investigate the corresponding generating function for all alternating permutations. For $n \in \mathbb{N}$ we let

$$E_n^- := \{\sigma \in S_n : \sigma(1) > \sigma(2) < \sigma(3) > \dots\},$$

and

$$E_n^+ := \{\sigma \in S_n : \sigma(1) < \sigma(2) > \sigma(3) < \dots\}.$$

We call the elements of E_n^- (resp. E_n^+) *alternating* (resp. *reverse alternating*) permutations (we refer the reader to, e.g., [11, §1.6] for further information about alternating permutations).

Proposition 5.2. *Let $n \in \mathbb{N}$. Then*

$$(5.1) \quad \sum_{\sigma \in E_n^-} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} 0, & \text{if } n \equiv 1 \pmod{2}, \\ (-x)^{\frac{n}{2}}, & \text{if } n \equiv 0 \pmod{2}, \end{cases}$$

and

$$(5.2) \quad \sum_{\sigma \in E_n^+} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} 0, & \text{if } n \equiv 1 \pmod{2}, \\ x^{\frac{n}{2}(\frac{n}{2}-1)}, & \text{if } n \equiv 0 \pmod{2}. \end{cases}$$

Proof. Note that $E_n^- = \mathcal{D}_J^I(S_n)$ where $I := \{i \in [n-1] : i \equiv 0 \pmod{2}\}$ and $J := \{i \in [n-1] : i \equiv 1 \pmod{2}\}$ so the first equation in (5.1) follows from Proposition 5.1. So assume that $n \equiv 0 \pmod{2}$, say $n = 2m$ for some $m \in \mathbb{N}$. By Lemma 4.1 we have

$$\sum_{\sigma \in E_n^-} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_J^I(C_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

We claim that $\mathcal{D}_J^I(C_{n,+}) = \emptyset$. Let $\sigma \in \mathcal{D}_J^I(C_{n,+})$. Let $i := \sigma^{-1}(1)$. Then $i \equiv 1 \pmod{2}$ so $i \in J$ and hence $\sigma(i) > \sigma(i+1)$ which is a contradiction. Let now $\sigma \in \mathcal{D}_J^I(C_{n,-})$. We claim that then

$$\sigma = [2, 1, 4, 3, 6, 5, \dots, 2m, 2m-1].$$

We prove this claim by induction on $m \in \mathbb{N}$. If $m = 1$ the claim is clear. Let $m \geq 2$. Let $a := \sigma^{-1}(2m-1)$. Then $a \equiv 0 \pmod{2}$ so $a = 2m$ (else $\sigma(a-1), \sigma(a+1) > \sigma(a) = 2m-1$) and hence $\sigma(2m-1) = 2m$. But $\sigma|_{[2m-2]} \in \mathcal{D}_{J \cap [n-3]}^{I \cap [n-3]}(C_{n-2,-})$ so the claim follows by induction. Since $\ell([2, 1, 4, 3, \dots, 2m, 2m-1]) = m = L([2, 1, 4, 3, \dots, 2m, 2m-1])$ the second equation in (5.1) follows.

Since the map $\sigma \mapsto w_0 \sigma$ is an involution between E_n^+ and E_n^- , the equations in (5.2) follow from those in (5.1) and Proposition 2.2. \square

We now consider a general family of descent classes which includes all quotients. Let $I, J \subseteq [n-1]$. We say that I and J are *unmixed* if

$$(5.3) \quad I \cap J = (I+1) \cap J = I \cap (J+1) = \emptyset.$$

Let $I, J \subseteq [n-1]$ be unmixed. Let I_1, \dots, I_s be the connected components of I and J_1, \dots, J_t be those of J . We say that (I, J) is *compressed* if $|I_1| \equiv \dots \equiv |I_s| \equiv |J_1| \equiv \dots \equiv |J_t| \equiv 1 \pmod{2}$ and $|[n-1] \setminus (I \cup J)| = s + t - 1$. For instance, $(\{1, 7, 8, 9\}, \{3, 4, 5, 11, 12, 13\})$ is compressed for $n = 14$ while $(\{1, 3\}, \{7, 8, 9, 11, 12, 13\})$ is not. Note that if $I, J \subseteq [n-1]$ are unmixed and (I, J) is compressed then $n-1 = |I| + |J| + s + t - 1 \equiv 1 \pmod{2}$ so n is even.

Let now $n = 2m \in \mathbb{N}$ and let I, J be unmixed with connected components I_1, \dots, I_s , and J_1, \dots, J_t , respectively. Then $I_1, \dots, I_s, J_1, \dots, J_t$ are the connected components of $I \cup J$. Therefore $\sum_{j=1}^s \binom{|I_j|+1}{2} + \sum_{k=1}^t \binom{|J_k|+1}{2} \leq m$, with equality holding if and only if (I, J) is compressed.

We can now state one of the main results of this section.

Theorem 5.3. *Let $I, J \subseteq [n-1]$ be unmixed. Let I_1, \dots, I_s be the connected components of I and J_1, \dots, J_t be the connected components of J . Then we have*

$$(5.4) \quad \sum_{\sigma \in \mathcal{D}_J^I(C_{n,+})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = (-x)^d x^{\alpha(J)} \begin{bmatrix} b+d \\ \mathbf{b}, \mathbf{d} \end{bmatrix}_{x^2} \prod_{k=b+d+1}^m (1-x^{2k}),$$

if n is odd, while

$$(5.5) \quad \sum_{\sigma \in \mathcal{D}_J^I(C_{n,+})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} (-x)^d x^{\alpha(J)} \begin{bmatrix} [b]_{x^2} \\ [m]_{x^2} \end{bmatrix}_{x^2} \begin{bmatrix} b+d \\ \mathbf{b}, \mathbf{d} \end{bmatrix}_{x^2}, & \text{if } m = b+d, \\ (-x)^d x^{\alpha(J)} \begin{bmatrix} b+d \\ \mathbf{b}, \mathbf{d} \end{bmatrix}_{x^2} \prod_{k=b+d+1}^{m-1} (1-x^{2k}), & \text{otherwise,} \end{cases}$$

and

(5.6)

$$\sum_{\sigma \in \mathcal{D}_J^I(C_n, -)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} (-1)^d x^{b+\alpha(J)} \frac{[d]_{x^2}}{[m]_{x^2}} \left[\begin{matrix} b+d \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2}, & \text{if } m = b+d, \\ -(-x)^d x^{m+\alpha(J)} \left[\begin{matrix} b+d \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2} \prod_{k=b+d+1}^{m-1} (1-x^{2k}), & \text{otherwise,} \end{cases}$$

if n is even, where $m := \lfloor \frac{n}{2} \rfloor$, $b_j := \lfloor \frac{|J_j|+1}{2} \rfloor$, for $j = 1, \dots, s$, $d_k := \lfloor \frac{|J_k|+1}{2} \rfloor$, for $k = 1, \dots, t$, $b := \sum_{i=1}^s b_i$, $d := \sum_{k=1}^t d_k$, $\mathbf{b} := b_1, \dots, b_s$, $\mathbf{d} := d_1, \dots, d_t$, and $\alpha(J) := \sum_{k=1}^t d_k^2$.

Proof. We let, for convenience, $\bar{b}_j := b_j + 1$, $\bar{d}_k := d_k + 1$, for $j \in [s]$ and $k \in [t]$, $\hat{\alpha}(J) := \alpha(J) + d$, and $\check{\alpha}(J) := \alpha(J) - d$.

Before delving into the proof we think it useful to sketch the idea of it. If J has at least one connected component of even size then by Lemma 4.4 this can be changed to a connected component of I and we can proceed by induction. If I has a connected component of even size (and all connected components of J have odd size) then by Propositions 4.2 and 4.3 we can remove one of the endpoints from this connected component and then shift some of the other connected components of I and J so that the resulting ‘‘empty spot’’ sits next to a connected component of J , to which it can then be ‘‘added’’ by Propositions 4.7 or 4.8. The resulting descent class now has a connected component of the descents of even size so can be computed by induction as in the previous case. If all the connected components of $I \cup J$ are of odd size but (I, J) is not compressed then there is either an ‘‘empty spot’’ to the right of the rightmost connected component of $I \cup J$, or to the left of the leftmost, or there are two consecutive connected components of $I \cup J$ separated by at least two empty spots. By shifting the connected components of $I \cup J$ we can ‘‘move’’ this extra empty spot so that it sits next to a connected component of J , to which it can then be ‘‘added’’, and we can conclude as in the previous case. If (I, J) is compressed then n is even and must appear immediately to the right of a connected component of the ascents which allows us to ‘‘delete’’ n and compute the generating function as a sum of generating functions of unmixed descent classes of S_{n-1} .

We proceed by induction on $t \in \mathbb{N}_0$, the number of connected components of the descents. Let $t = 0$ (i.e., $J = \emptyset$). Then (I, \emptyset) is compressed if and only if $b = \frac{n}{2}$ so Theorem 5.4 reduces to Theorem 2.4 in this case. Let now $t \geq 1$.

Assume first that there exists $i \in [t]$ such that $|J_i| \equiv 0 \pmod{2}$. Then by Lemma 4.4 and our induction hypothesis we have

$$\begin{aligned} \sum_{\sigma \in \mathcal{D}_J^I(C_n, +)} (-1)^{\ell(\sigma)} x^{L(\sigma)} &= (-1)^{\frac{|J_i|}{2}} x^{\frac{|J_i|(|J_i|+2)}{4}} \sum_{\sigma \in \mathcal{D}_{J \setminus J_i}^{I \cup J_i}(C_n, +)} (-1)^{\ell(\sigma)} x^{L(\sigma)} \\ &= (-1)^{d_i} x^{d_i \bar{d}_i} (-1)^{d-d_i} x^{\hat{\alpha}(J)-d_i \bar{d}_i} \left[\begin{matrix} b+d \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2} \prod_{k=b+d+1}^{\lfloor \frac{n-1}{2} \rfloor} (1-x^{2k}), \end{aligned}$$

so (5.4) and the second formula in (5.5) follow in this case.

Under the same hypothesis, for the odd chessboard elements we have

$$\begin{aligned} \sum_{\sigma \in \mathcal{D}_J^I(C_{2m}, -)} (-1)^{\ell(\sigma)} x^{L(\sigma)} &= (-1)^{\frac{|J_i|}{2}} x^{\frac{|J_i|(|J_i|+2)}{4}} \sum_{\sigma \in \mathcal{D}_{J \setminus J_i}^{I \cup J_i}(C_{2m}, -)} (-1)^{\ell(\sigma)} x^{L(\sigma)} \\ &= -(-1)^{d_i} x^{d_i \bar{d}_i} (-1)^{d-d_i} x^{m+\hat{\alpha}(J)-d_i \bar{d}_i} \left[\begin{matrix} b+d \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2} \prod_{k=b+d+1}^{m-1} (1-x^{2k}), \end{aligned}$$

yielding the second formula in (5.6).

We may therefore assume that $|J_1| \equiv |J_2| \equiv \cdots \equiv |J_t| \equiv 1 \pmod{2}$.

Assume now that there exists $r \in [s]$ such that $|I_r| \equiv 0 \pmod{2}$. Then by repeated application of Proposition 4.7 and 4.8, we have

$$\sum_{\sigma \in \mathcal{D}_J^I(C_n, \pm)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{\tilde{J}}^{\tilde{I}}(C_n, \pm)} (-1)^{\ell(\sigma)} x^{L(\sigma)},$$

where \tilde{I} has connected components $\tilde{I}_1 \cup \cdots \cup \tilde{I}_s$, where $|\tilde{I}_r| = |I_r| - 1$ and $|\tilde{I}_k| = |I_k|$, for $k \in [s] \setminus \{r\}$ and \tilde{J} has connected components $\tilde{J}_1 \cup \cdots \cup \tilde{J}_t$, where $|\tilde{J}_1| = |J_1| + 1$ and $|\tilde{J}_k| = |J_k|$, for $k \in [2, t]$, and the connected components of $\tilde{I} \cup \tilde{J}$ are $\tilde{I}_1, \dots, \tilde{I}_s, \tilde{J}_1, \dots, \tilde{J}_t$. Since \tilde{J} has a connected component of even cardinality, reasoning as in the previous case, and observing that $\lfloor \frac{|\tilde{J}_1|+1}{2} \rfloor = \lfloor \frac{|J_1|+1}{2} \rfloor = d_1$ and $\lfloor \frac{|\tilde{I}_r|+1}{2} \rfloor = \lfloor \frac{|I_r|+1}{2} \rfloor = b_r$, we conclude again by induction.

We may therefore assume that $|I_1| \equiv \cdots \equiv |I_s| \equiv |J_1| \equiv \cdots \equiv |J_t| \equiv 1 \pmod{2}$.

Suppose first that $|[n-1] \setminus (I \cup J)| > s + t - 1$. Therefore either $1 \notin I \cup J$ or $n-1 \notin I \cup J$ or there exists $i \in [n-1]$ such that $i, i+1 \notin I \cup J$. In any of these cases we can apply Propositions 4.7 and 4.8 to get

$$\sum_{\sigma \in \mathcal{D}_J^I(C_n, \pm)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{\sigma \in \mathcal{D}_{\bar{J}}^{\bar{I}}(C_n, \pm)} (-1)^{\ell(\sigma)} x^{L(\sigma)},$$

where \bar{I} has connected components $\bar{I}_1, \dots, \bar{I}_s$ such that $|\bar{I}_j| = |I_j|$ for $j \in [s]$ and \bar{J} has connected components $\bar{J}_1, \dots, \bar{J}_t$ such that $|\bar{J}_1| = |J_1| + 1$ and $|\bar{J}_l| = |J_l|$ for $l \in [2, t]$. Then, again, \bar{J} has a connected component of even size so, reasoning as above (5.4), and the second equations in (5.5) and (5.6) follow by induction, since $\lfloor \frac{|\bar{J}_1|+1}{2} \rfloor = \lfloor \frac{|J_1|+1}{2} \rfloor = d_1$.

We may therefore assume that $|I_1| \equiv \cdots \equiv |I_s| \equiv |J_1| \equiv \cdots \equiv |J_t| \equiv 1 \pmod{2}$ and $|[n-1] \setminus (I \cup J)| = s + t - 1$, i.e., that (I, J) is compressed. Then $n \equiv 0 \pmod{2}$, say $n = 2m$, and $m = b + d$, and both the leftmost and the rightmost elements of any connected component of $J \cup J$ are odd.

For $i \in [s]$ let $a_i := \max I_i + 1$ and for $i \in [t]$ let $c_i := \min J_i$. Then $a_1 \equiv \cdots \equiv a_s \equiv 0 \pmod{2}$ and $c_1 \equiv \cdots \equiv c_t \equiv 1 \pmod{2}$. Therefore, if $\sigma \in \mathcal{D}_J^I(C_{2m, +})$, then $\sigma^{-1}(2m) \in \{a_1, \dots, a_s\}$. Hence

$$\sum_{\sigma \in \mathcal{D}_J^I(C_{2m, +})} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \sum_{j=1}^s \sum_{\substack{\sigma \in \mathcal{D}_J^I(C_{2m, +}): \\ \sigma^{-1}(2m) = a_j}} (-1)^{\ell(\sigma)} x^{L(\sigma)}.$$

Fix $j \in [s]$. Let $k := \max\{i \in [t] : c_i < a_j\}$ (where $k := 0$ if $\{i \in [t] : c_i < a_j\} = \emptyset$). So J_1, \dots, J_k are to the left of a_j , while J_{k+1}, \dots, J_t are to the right. Let $\bar{\tau}$ be obtained from τ by removing the maximum (which is in position a_j) and reversing the elements in each of the blocks of ascents and descents that are to the right of a_j , so reversing the elements in positions $[\min I_i, a_i]$ for each $i = j+1, \dots, s$, and those in positions $[c_i, \max J_i + 1]$ for each $i = k+1, \dots, t$. Then the map $\tau \mapsto \bar{\tau}$ is a bijection between $\{\sigma \in \mathcal{D}_J^I(C_{2m, +}) : \sigma^{-1}(2m) = a_j\}$ and $\mathcal{D}_{\varphi_j(J)}^{\varphi_j(I)}(C_{2m-1})$, where $\varphi_j(I) := I_1 \cup \cdots \cup I_{j-1} \cup (I_j \setminus \{a_j - 1\}) \cup (J_{k+1} - 1) \cup \cdots \cup (J_t - 1)$ and $\varphi_j(J) := J_1 \cup \cdots \cup J_k \cup (I_{j+1} - 1) \cup \cdots \cup (I_s - 1)$.

Furthermore, we have $\ell(\bar{\tau}) = \ell(\tau) + A$ and $L(\bar{\tau}) = L(\tau) + B$, where, by Proposition 2.2

$$\begin{aligned} A &= \sum_{r=j+1}^s \binom{|I_r|+1}{2} - \sum_{h=k+1}^t \binom{|J_h|+1}{2} - (2m - a_j) = \sum_{r=j+1}^s b_r(2b_r - 1) - \sum_{h=k+1}^t d_h(2d_h - 1) - (2m - a_j) \\ &= \sum_{r=j+1}^s b_r(2b_r - 3) - \sum_{h=k+1}^t d_h(2d_h + 1), \\ B &= \sum_{r=j+1}^s \left(\frac{|I_r|+1}{2}\right)^2 - \sum_{h=k+1}^t \left(\frac{|J_h|+1}{2}\right)^2 - \frac{2m - a_j}{2} = \sum_{r=j+1}^s b_r(b_r - 1) - \sum_{h=k+1}^t d_h(d_h + 1), \end{aligned}$$

since $2m - a_j = 2\left(\sum_{r=j+1}^s b_r + \sum_{h=k+1}^t d_h\right)$. Therefore, by our induction hypothesis (5.4),

$$\begin{aligned} \sum_{\substack{\tau \in \mathcal{D}_J^I(C_{2m,+}) \\ \sigma^{-1}(2m) = a_j}} (-1)^{\ell(\tau)} x^{L(\tau)} &= (-1)^A x^{-B} \sum_{\bar{\tau} \in \mathcal{D}_{\varphi_j(J)}^{\varphi_j(I)}(C_{2m-1})} (-1)^{\ell(\bar{\tau})} x^{L(\bar{\tau})} \\ (5.7) \qquad \qquad \qquad &= (-1)^d x^{\hat{\alpha}(\varphi_j(J)) - B} \left[\begin{matrix} m-1 \\ b_1, \dots, b_{j-1}, b_j - 1, b_{j+1}, \dots, b_s, \mathbf{d} \end{matrix} \right]_{x^2}. \end{aligned}$$

But $\hat{\alpha}(\varphi_j(J)) = \sum_{r=1}^k d_r(d_r + 1) + \sum_{r=j+1}^s b_r(b_r + 1)$, so $\hat{\alpha}(\varphi_j(J)) - B = \hat{\alpha}(J) + 2\sum_{r=j+1}^s b_r$.

Thus, the sum in (5.7) becomes

$$(-1)^d x^{\hat{\alpha}(J)} x^{2\sum_{r=j+1}^s b_r} \left[\begin{matrix} m-1 \\ b_1, \dots, b_{j-1}, b_j - 1, b_{j+1}, \dots, b_s, \mathbf{d} \end{matrix} \right]_{x^2}.$$

Therefore

$$\begin{aligned} \sum_{\sigma \in \mathcal{D}_J^I(C_{2m,+})} (-1)^{\ell(\sigma)} x^{L(\sigma)} &= (-1)^d x^{\hat{\alpha}(J)} \sum_{j=1}^s x^{2\sum_{r=j+1}^s b_r} \left[\begin{matrix} m-1 \\ b_1, \dots, b_{j-1}, b_j - 1, b_{j+1}, \dots, b_s, \mathbf{d} \end{matrix} \right]_{x^2} \\ &= (-1)^d x^{\hat{\alpha}(J)} \frac{[b]_{x^2}}{[m]_{x^2}} \left[\begin{matrix} m \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2} \end{aligned}$$

as desired.

Under the same hypothesis, for the sum over odd chessboard elements we have, by Proposition 4.5 and Remark 4.6

$$\begin{aligned} \sum_{\sigma \in \mathcal{D}_J^I(C_{2m,-})} (-1)^{\ell(\sigma)} x^{L(\sigma)} &= (-1)^{\ell(w_0)} x^{L(w_0)} \sum_{\tau \in \mathcal{D}_I^J(C_{2m,+})} (-1)^{\ell(\tau)} x^{-L(\tau)} \\ &= (-1)^{\binom{2m}{2}} x^{m^2} \sum_{\tau \in \mathcal{D}_I^J(C_{2m,+})} (-1)^{\ell(\tau)} x^{-L(\tau)} \\ &= (-1)^m x^{m^2} (-1)^b x^{-\sum_{j=1}^s b_j \bar{b}_j} \frac{[d]_{x^{-2}}}{[m]_{x^{-2}}} \left[\begin{matrix} m \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^{-2}} \\ &= (-1)^d x^{m + \hat{\alpha}(J)} \frac{[d]_{x^2}}{[m]_{x^2}} \left[\begin{matrix} m \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2} \end{aligned}$$

and the result follows. This concludes the proof of the first equations in (5.5) and (5.6) and hence of the result. \square

By Lemma 4.1 the preceding result implies the following one, which computes the sign-twisted generating function of the odd length over any unmixed descent class.

Theorem 5.4. *Let $I, J \subseteq [n-1]$ be unmixed. Then, keeping the same notation as in Theorem 5.3*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = \begin{cases} (-1)^d x^{\alpha(J)} \frac{x^d [b]_{x^2} + x^b [d]_{x^2}}{[b+d]_{x^2}} \left[\begin{matrix} b+d \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2}, & \text{if } n = 2(b+d), \\ (-x)^d x^{\alpha(J)} \left[\begin{matrix} b+d \\ \mathbf{b}, \mathbf{d} \end{matrix} \right]_{x^2} \prod_{k=2b+2d+2}^n (1 + (-1)^{k-1} x^{\lfloor \frac{k}{2} \rfloor}), & \text{otherwise.} \end{cases}$$

6. OPEN PROBLEMS

In this section we collect some conjectures and open problems arising from this work.

For $\sigma \in S_n$, we let $\text{cl}_n(\sigma) = \{\tau \in S_n : D_o(\tau) = D_o(\sigma)\}$ denote the equivalence class of permutations in S_n with the same odd diagram as σ . Clearly, the problem of characterizing the odd diagrams is closely related to that of identifying these equivalence classes.

Recall that a permutation $\sigma \in S_n$ is said to *contain the pattern* $\alpha = \alpha_1 \cdots \alpha_k$ if there exist $1 \leq i_1 < \cdots < i_k \leq n$ such that $\sigma(i_1), \dots, \sigma(i_k)$ are in the same relative order as $\alpha_1, \dots, \alpha_k$. A permutation $\sigma \in S_n$ is said to *avoid* the pattern α if it does not contain the pattern α . We denote with $\text{Av}_n(\alpha) = \{\sigma \in S_n : \sigma \text{ avoids } \alpha\}$ the set of permutations of degree n avoiding α . We conjecture that odd diagrams faithfully encode permutations avoiding some patterns of length 3. More precisely, we conjecture the following.

Conjecture 6.1. *Let $\alpha \in \{213, 312\}$. The map $D_o: \text{Av}_n(\alpha) \rightarrow [n]^2$, $\sigma \mapsto D_o(\sigma)$ is injective. More precisely, for a permutation $\sigma \in S_n$, the class $\text{cl}_n(\sigma)$ contains at most one permutation avoiding the pattern 213 and at most one avoiding 312. If they exist, they are respectively the longest and the shortest element of $\text{cl}_n(\sigma)$.*

We have verified Conjecture 6.1 for $n \leq 7$.

In light of Proposition 3.5 and Remark 3.7, it is natural to investigate the polynomials giving the (non-twisted) distribution of the odd inversions. For $n \in \mathbb{N}$ we denote this polynomial by $L_n(x) := \sum_{\sigma \in S_n} x^{L(\sigma)}$. Properties (iii) and (iv) in Proposition 2.2 imply that $L_n(x)$ is monic and symmetric for all $n \in \mathbb{N}$. For small values of n we have:

$$\begin{aligned} L_3(x) &= 1 + 4x + x^2 \\ L_4(x) &= 1 + 8x + 6x^2 + 8x^3 + x^4 \\ L_5(x) &= 1 + 12x + 23x^2 + 48x^3 + 23x^4 + 12x^5 + x^6 \\ L_6(x) &= 1 + 16x + 59x^2 + 137x^3 + 147x^4 + 147x^5 + 137x^6 + 59x^7 + 16x^8 + x^9 \end{aligned}$$

With the exception of $n = 4$, for $n \leq 11$ the polynomials $L_n(x)$ are unimodal. We therefore conjecture the following.

Conjecture 6.2. *Let $n \geq 5$. Then the polynomial $L_n(x)$ is unimodal.*

The first rows of the associated triangle are recorded in [10, A289511].

Generalizing further from Remark 3.7, we let, for $k, n \in \mathbb{N}$, $h \in \mathbb{Z}/k\mathbb{Z}$ and $\sigma \in S_n$,

$$(6.1) \quad \text{inv}_{k,h}(\sigma) = |\{(i, j) \in [n]^2 : i < j, \sigma(i) > \sigma(j), j - i \equiv h \pmod{k}\}|.$$

Note that $L(\sigma) = \text{inv}_{2,1}(\sigma)$. Also, note that for $k \geq n - 1$, the polynomials of the distributions of the statistic $\text{inv}_{k,1}$ over S_n coincide with the Eulerian polynomials:

$$\sum_{\sigma \in S_n} x^{\text{inv}_{k,1}(\sigma)} = \sum_{\sigma \in S_n} x^{\text{des}(\sigma)},$$

where $\text{des}(\sigma) = |\text{Des}(\sigma)|$ denotes the descent number of the permutation σ . Inspired by this fact and Conjecture 6.2, we put forward the following general conjecture.

Conjecture 6.3. *The polynomials*

$$\sum_{\sigma \in S_n} x^{\text{inv}_{k,1}(\sigma)}$$

are unimodal for all $n \in \mathbb{N}$, and all $k \geq 3$.

We have verified that Conjecture 6.3 holds for $n \leq 9$, and all relevant k .

We have seen in Proposition 5.1 some sufficient conditions for the sign-twisted generating function of the odd length to be zero on a descent class. This, together with the comments following Proposition 5.1, suggests the following natural problem.

Problem 6.4. *Let $I, J \subseteq [n]$, $I \cap J = \emptyset$. Give necessary and sufficient conditions on I and J such that*

$$\sum_{\sigma \in \mathcal{D}_J^I(S_n)} (-1)^{\ell(\sigma)} x^{L(\sigma)} = 0.$$

Acknowledgments. The authors would like to thank the referee for useful comments. The first author would like to thank A. Postnikov, A. Rapagnetta, R. Stanley, M. Wachs, and L. Williams for interesting and useful conversations. FB was partially supported by the MIUR Excellence Department Project CUP E83C18000100006. The second named author was partially supported by the German-Israeli Foundation for Scientific Research and Development through grant no. 1246. AC would also like to thank the Erwin Schrödinger International Institute for Mathematics and Physics (Vienna), where part of this research was carried out.

REFERENCES

- [1] A. Björner, F. Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, **231**, Springer-Verlag, New York, 2005.
- [2] F. Brenti, A. Carnevale, *Proof of a conjecture of Klopsch-Voll on Weyl groups of type A*, Trans. Amer. Math. Soc. **369** (2017), 7531–7547.
- [3] F. Brenti, A. Carnevale, *Odd length for even hyperoctahedral groups and signed generating functions*, Discrete Math., **340** (2017), 2822–2833.
- [4] F. Brenti, A. Carnevale, *Odd length in Weyl groups*, Algebraic Comb., **2** (2019), no. 6, 1125–1147.
- [5] P. Griffiths, J. Harris, *Principles of algebraic geometry*, Pure and Applied Mathematics, Wiley-Interscience, New York, 1978.
- [6] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, **52**, Springer-Verlag, New York-Heidelberg, 1977.
- [7] B. Klopsch, C. Voll, *Igusa-type functions associated to finite formed spaces and their functional equations*, Trans. Amer. Math. Soc., **361** (2009), no. 8, 4405–4436.
- [8] T. Kövari, V. T. Sós, P. Turán, *On a problem of K. Zarankiewicz*, Colloquium Math., **3**, (1954), 50–57.
- [9] I.G. Macdonald, *Notes on Schubert polynomials*, Publications du LaCIM, Montreal, 1991.
- [10] N. J. A. Sloane, editor, *The On-Line Encyclopedia of Integer Sequences*, published electronically at <https://oeis.org>, 2019.
- [11] R. P. Stanley, *Enumerative Combinatorics*, vol.2, Cambridge Studies in Advanced Mathematics, no.62, Cambridge Univ. Press, Cambridge, 1999.
- [12] A. Stasinski, C. Voll, *A new statistic on the hyperoctahedral groups*, Electronic J. Combin., **20** (2013), no. 3, Paper 50, 23 pp.

- [13] A. Stasinski, C. Voll, *Representation zeta functions of nilpotent groups and generating functions for Weyl groups of type B*, Amer. J. Math., **136** (2) (2014), 501–550.
- [14] J. Stembridge, *Sign-twisted Poincaré series and odd inversions in Weyl groups*, Algebraic Comb., **2** (2019), no. 4, 621–644.

FRANCESCO BRENTI, DIPARTIMENTO DI MATEMATICA UNIVERSITÀ DI ROMA “TOR VERGATA” VIA DELLA RICERCA SCIENTIFICA, 1 00133 ROMA, ITALY

Email address: `brenti@mat.uniroma2.it`

ANGELA CARNEVALE, SCHOOL OF MATHEMATICS, APPLIED MATHEMATICS AND STATISTICS, NATIONAL UNIVERSITY GALWAY, GALWAY, IRELAND

Email address: `angela.carnevale@nuigalway.ie`