



Article

# Validation of StepTest4all for Assessing Cardiovascular Capacity in Young Adults

José A. Bragada <sup>1,2,\*</sup>, Raul F. Bartolomeu <sup>1,2,3</sup>, Pedro M. Rodrigues <sup>1</sup>, Pedro M. Magalhães <sup>1</sup>, João P. Bragada <sup>4</sup> and Jorge E. Morais <sup>1,2</sup>

<sup>1</sup> Department of Sport Sciences, Instituto Politécnico de Bragança, 5300-253 Bragança, Portugal

<sup>2</sup> Research Centre in Sports, Health and Human Development (CIDESD), 5001-801 Vila Real, Portugal

<sup>3</sup> Department of Sport Sciences, Polytechnic Institute of Guarda, 6300-559 Guarda, Portugal

<sup>4</sup> North East Local Health Unit—Health Care Unit of Santa Maria, 5301-852 Bragança, Portugal

\* Correspondence: jbragada@ipb.pt

**Abstract:** Background: Cardiovascular capacity, expressed as maximal oxygen uptake ( $VO_{2max}$ ), is a strong predictor of health and fitness and is considered a key measure of physiological function in the healthy adult population. The purpose of this study was to validate a specific step test (StepTest4all) as an adequate procedure to estimate cardiovascular capacity in young adults. Methods: The sample was composed of 56 participants, including 19 women (aged  $21.05 \pm 2.39$  years, body mass =  $57.50 \pm 6.64$  kg, height =  $1.62 \pm 0.05$  m, body mass index =  $22.00 \pm 2.92$  kg/m<sup>2</sup>) and 37 men (aged  $22.05 \pm 3.14$  years, body mass =  $72.50 \pm 7.73$  kg, height =  $1.76 \pm 0.07$  m, body mass index =  $23.34 \pm 2.17$  kg/m<sup>2</sup>). Participants were included in one of the following groups: (i) the group used to predict the  $VO_{2max}$ , and (ii) the group used to validate the prediction model. All participants performed the StepTest4all protocol. The step height and the intensity of the effort was determined individually. Heart rate and oxygen uptake were measured continuously during rest, effort, and recovery phases. The validation process included the following three stages: (i) mean data comparison, (ii) simple linear regression, and (iii) Bland–Altman analysis. Results: The linear regression retained, as significant predictors of the  $VO_{2max}$ , sex ( $p < 0.001$ ) and heart rate recovery for one minute ( $p = 0.003$ ). The prediction equation revealed a high relationship between measurements ( $R^2 = 63.0\%$ ,  $SEE = 5.58$ ). The validation procedure revealed non-significant differences ( $p > 0.05$ ) between the measured and estimated maximal oxygen uptake, high relationship ( $R^2 = 63.3\%$ ), and high agreement with Bland–Altman plots. Thus,  $VO_{2max}$  can be estimated with the formula:  $VO_{2max} = 22 + 0.3 \cdot (HRR_{1min}) + 12 \cdot (sex)$ , where  $HRR_{1min}$  is the magnitude of the HR decrease (bpm) in one minute immediately after the step was stopped, and sex: men = 1, women = 0. Conclusions: The StepTest4all is an adequate procedure to estimate cardiovascular capacity, expressed as  $VO_{2max}$ , in young adults. In addition, it is possible to determine the qualitative level of cardiovascular capacity from the heart rate recovery for one minute, more specifically, poor:  $<20$ , moderate: 20 to 34, good: 35 to 49, and excellent:  $\geq 50$ . This procedure has the benefit of being simple to apply and can be used by everyone, even at home, without specialist supervision.

**Keywords:** heart rate recovery; exercise and health; StepTest4all; cardiovascular capacity; cardiovascular classification; validation;  $VO_{2max}$



**Citation:** Bragada, J.A.; Bartolomeu, R.F.; Rodrigues, P.M.; Magalhães, P.M.; Bragada, J.P.; Morais, J.E. Validation of StepTest4all for Assessing Cardiovascular Capacity in Young Adults. *Int. J. Environ. Res. Public Health* **2022**, *19*, 11274. <https://doi.org/10.3390/ijerph191811274>

Academic Editor: Paul B. Tchounwou

Received: 29 July 2022

Accepted: 3 September 2022

Published: 8 September 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Cardiovascular capacity, expressed as  $VO_{2max}$ , is a strong predictor of fitness and has been considered a key physiological measure in the healthy adult population [1]. Testing and maintaining good cardiorespiratory fitness could help to prevent declines in health status within the general population [2].

Exercise effort testing is a procedure that can be used to estimate cardiovascular fitness, diagnose cardiovascular diseases, and predict mortality associated with cardiovascular

problems [3,4]. Studies have demonstrated that exercise testing data have powerful prognostic properties, particularly when focused on functional capacity and exercise heart rate dynamics, such as heart rate recovery (HRR) [5,6].

The heart rate (HR) increases with exercise intensity to respond to greater metabolic demand of muscles and other tissues. This increase in HR is tightly regulated by the action of central and peripheral mechanisms that project afferent inputs to medullary centers in the brain. These afferent inputs result in an appropriate efferent response of the autonomic nervous system branches, i.e., a decrease in parasympathetic and an increase in sympathetic activity [7]. After the exercise ends, there is a progressive reduction in metabolic activity and, consequently, HR also decreases. A slow HRR after exercise is a strong indicator of cardiac autonomic dysfunction [7,8]. A delayed decrease in HR during the first minute after graded exercise is a powerful predictor of overall mortality independent of workload, the presence or absence of myocardial perfusion defects, and changes in HR during exercise [1,7,9]. Better levels of aerobic fitness act beneficially on the autonomic control of post-exercise HR, preserving the vagal reentry velocity in healthy middle-aged adults [10]. Thus, there is scientific evidence to accept the association between HRR and prognosis of cardiovascular disease, which supports the recommendation of recording HRR for risk assessment in clinical practice, as a routine [11,12].

Among the variety of tests described for this purpose, the step tests are the most accessible to the general population and have been used for many years [13–15]. Step tests can be considered of major importance, as they allow the assessment of cardiovascular capacity as well as identification of the dynamics of HR during effort and in the recovery period [16]. The HR decrease during recovery, and in the period immediately after the end of the exercise can provide a simple, effective, and ecologically valid method of submaximal assessment of cardiorespiratory fitness that can be implemented in a variety of situations [14]. Heart rate recovery can be defined as the reduction in heart rate immediately after the cessation of exercise, usually for one minute [8,16,17].

The use of step tests may be appropriate for the following reasons: (i) they can be used to monitor training status over time [18], (ii) they can estimate the cardiovascular capacity from the HR recovery [19–21], and (iii) they allow an appreciation of the functionality of the sympathetic and parasympathetic nervous systems [22]. Thus, its use makes sense both for the assessment of physical capacity, and in clinical practice. Additionally, step tests have the advantage of requiring minimal and portable equipment and marginal space compared to tests utilizing treadmills, shuttle walks, or cycle ergometers. However, most of the existing step tests have one or more of the following limitations: (i) pre-established durations; (ii) frequently eliciting efforts above 85% of maximal oxygen uptake ( $VO_{2max}$ ), which are not recommended for sedentary or elderly people [23]; and (iii) a fixed step height, which is often unsuited for people with short stature, excess weight, or poor physical capacity.

Due to all these limitations, we developed a new step test procedure, StepTest4all, that solves those constraints and can also be adjusted to individual capacity. In StepTest4all, the step height, duration, and level of difficulty can be adjusted to each person regardless of their age, weight, height, or level of physical fitness. Thus, the purpose of this study was to validate the StepTest4all as an adequate procedure to estimate cardiovascular capacity in young adults.

## 2. Material and Methods

### 2.1. Participants and Samples

The sample was composed of 56 participants, including 19 women (aged  $21.05 \pm 2.39$  years, body mass =  $57.50 \pm 6.64$  kg, height =  $1.62 \pm 0.05$  m, body mass index =  $22.00 \pm 2.92$  kg/m<sup>2</sup>) and 37 men (aged  $22.05 \pm 3.14$  years, body mass =  $72.50 \pm 7.73$  kg, height =  $1.76 \pm 0.07$  m, body mass index =  $23.34 \pm 2.17$  kg/m<sup>2</sup>). The sample recruitment excluded people who had any physical limitation that made it impossible for them to go up and down the step, or any other medical contraindication to performing moderate physical effort. Participants signed an informed consent form. All procedures were in accordance with the Declaration

of Helsinki regarding human research, and the Polytechnic Ethics Board approved the research design.

Participants were included in one of the following groups: (i) the group for  $VO_{2max}$  prediction equation (equation group—EG), and (ii) the group for validation of the prediction model (Validation group—VG). The participants' characteristics are presented in Table 1.

**Table 1.** Participants' characteristics. Equation group—group for  $VO_{2max}$  prediction equation and Validation group—group for validation of prediction model.

	Equation Group			Validation Group		
		Mean $\pm$ 1SD		Mean $\pm$ 1SD		
	Women (n = 10)	Men (n = 18)	Total EG	Women (n = 9)	Men (n = 19)	Total VG
Age [years]	21.60 $\pm$ 1.84	21.17 $\pm$ 1.98	21.32 $\pm$ 1.90	20.44 $\pm$ 2.88	22.89 $\pm$ 3.80	22.11 $\pm$ 3.67
Body mass [kg]	61.45 $\pm$ 5.06	72.06 $\pm$ 8.83	68.27 $\pm$ 9.19	53.11 $\pm$ 5.42	72.92 $\pm$ 6.75	66.55 $\pm$ 13.31
Height [m]	1.63 $\pm$ 0.04	1.77 $\pm$ 0.07	1.72 $\pm$ 0.09	1.60 $\pm$ 0.04	1.75 $\pm$ 0.07	1.71 $\pm$ 0.10
BMI [kg/m <sup>2</sup> ]	23.13 $\pm$ 3.02	22.93 $\pm$ 2.22	23.00 $\pm$ 2.48	20.75 $\pm$ 2.38	23.72 $\pm$ 2.11	22.77 $\pm$ 2.58
HRR <sub>1min</sub> [bpm]	35.50 $\pm$ 10.23	40.22 $\pm$ 10.99	38.54 $\pm$ 10.79	34.89 $\pm$ 10.71	38.42 $\pm$ 10.42	37.29 $\pm$ 10.45
$VO_{2max}$ [mL/kg/min]	34.41 $\pm$ 6.39	47.75 $\pm$ 6.71	42.99 $\pm$ 9.18	33.51 $\pm$ 5.64	46.15 $\pm$ 6.23	42.09 $\pm$ 8.45
HRR <sub>rest</sub> [bpm]	81.90 $\pm$ 12.00	69.11 $\pm$ 10,94	73.68 $\pm$ 12.75	80.76 $\pm$ 10.60	64.37 $\pm$ 10.88	69.64 $\pm$ 13.16
$VO_{2,rest}$ [mL/kg/min]	3.60 $\pm$ 0.86	3.61 $\pm$ 0.54	3.61 $\pm$ 0.66	3,54 $\pm$ 0.44	3.66 $\pm$ 0.64	3.62 $\pm$ 0.58

## 2.2. Data Collection

Anthropometric variables were measured using a digital stadiometer (Seca 242, Hamburg, Germany) and an electronic scale (Seca 884, Hamburg, Germany).  $VO_2$  and HR were measured using a stationary breath-by-breath electronic metabolic device (Cortex, Model MetaLyzer 3B, Leipzig, Germany). The device includes a heart rate transmitter (Polar Electro Oy, Kempele, Finland). The apparatus was calibrated with standard gases before each test. According to the manufacturer's manual, the standard error is 0.1% for oxygen and carbon dioxide sensors.

The  $VO_2$  and HR were measured continuously for each participant while they performed the following activities in sequence: rest, StepTest4all protocol, and recovery. The HR and  $VO_2$  values obtained as follows, were considered for further analysis: resting values—average of the last minute of resting period, values obtained during StepTest4all—the average values obtained in the last 5 s of each intensity level, recovery phase—the average values obtained in the last 5 s of the first minute of recovery. Resting  $VO_2$  and resting HR were collected continuously during 10 min sitting on a chair in a silent and dimly lit room. Participants were not allowed to sleep. The values obtained in the last minute were used for data analysis. During the recovery phase, although the HR values were recorded after the first two minutes, we only considered the value after the first minute. We chose one minute recovery time because it is the usually chosen time and has a higher reproducibility [24].

## 2.3. StepTest4all Protocol

For the StepTest4all protocol, each participant performed a continuous progressive test that consisted of going up and down on a stable step. After the step-up phase, the opposite leg also stepped up to the platform, so the participant stood vertically, supported by both legs, before beginning the step-down phase. The step-down phase began with the

same leg as the previous phase and ended when the participant was at the beginning point, also standing in a vertical position, supported by both legs.

The step height was calculated for each participant individually, based on characteristics of their cardiovascular capacity. The variables chosen were sex, age, physical fitness, height, body mass index (BMI), and smoking status. For each variable, a numerical value was assigned as follows: (i) sex (women = 0.5; men = 1), (ii) age (senior = 0, adult = 0.5, young = 1), (iii) physical fitness (sedentary = 0, active = 0.5, very active = 1, athlete = 1.5), (iv) body mass index (BMI < 25 = 0.5, BMI < 30 = 0, BMI ≥ 30 = −0.5), and (v) smoking status (smoker = 0, non-smoker = 0.5). From these data, the step height was calculated using the formula: step height (cm) = 4 × sum of those variables + 15, and it could range from 15 to 40 cm.

This formula was the result of multiple tests on people with different characteristics and of different physical ability. The height of the step, although important, may vary slightly, because the control of the load progression intensity until the desired value will be done mainly by increasing the ascent and descent pace.

A height of 40 to 45 cm has been used in other protocols, for example in the Harvard step test (see: <https://www.brianmac.co.uk/havard.html> (accessed on 28 July 2022)) [25]. In the present study, the step height of 40 cm combined with a high pace, led to an intensity of 80% of the estimated HR<sub>max</sub> in 5–10 min. This occurs even in subjects with good physical ability and tall stature.

The test started at a rate of 15 cycles per minute (0.25 Hz), controlled by a metronome. In each cycle, the participant went up and down the step, so the cycle ended at the same time the second leg reached the floor. Every minute, the cadence was raised by 2.5 cycles per minute. The maximum expected duration of the test was 10 min.

The lower limit of 15 cycles per minute is a very slow ascent and descent pace that can be performed by anyone and serves as a warm-up. The upper limit of 37.5 cycles per minute is a pace that can only be performed by subjects of high physical capacity.

The test ended when one of the following criteria was met: (i) the HR reached 80% of the maximal heart rate (HR<sub>max</sub>), (ii) when asked by the participant upon feeling uncomfortable with the effort, or (iii) when the participant was unable to perform the exercise at the correct cadence. Immediately after the end of the test, the participants were to stand for two minutes. Although a standing position was mandatory, the participants were encouraged to be at ease, in order to recover as much as possible, and were not permitted to talk, grab, or hold onto anything.

The combination of the step's height with the rhythm's increments, associated with a limitation in intensity to finish the test (80% of HR<sub>max</sub>), made it possible to achieve the test objectives in a reasonable amount of time (5 to 10 min) on all kinds of people. In the recovery phase the HR was collected continuously, also with the Garmin Fenix 6, and its HR belt (Garmin International, Inc., Olathe, KS, USA).

The HR<sub>max</sub> and VO<sub>2max</sub> determination were made as follows: HR<sub>max</sub> was estimated according to the formula: HR<sub>max</sub> = 208 − 0.7 × age [26].

The VO<sub>2max</sub> was estimated from the individual equation of the regression line associating HR–VO<sub>2</sub>, obtained from the rest data and during 3 or more steps of StepTest4all, more precisely, by calculating the value of VO<sub>2</sub> corresponding to HR<sub>max</sub> [27]. We assumed this value as VO<sub>2max</sub> measured. Individual linear regressions (R<sup>2</sup>) ranged between 0.97 and 0.99 (almost perfect relationship). This is a usual and suitable procedure for evaluating VO<sub>2max</sub> in subjects for whom a maximal test to exhaustion may have some inconvenience.

The use of submaximal tests to estimate VO<sub>2max</sub> from the association of HR–VO<sub>2</sub> is recurrent and has been shown to be adequate [28,29]. Evans and co-workers [30], in a systematic review, found that non-significant differences were reported between the measured and predicted VO<sub>2max</sub> in 28 equations. The variable most used in the predictive equations was the HR (N = 19). Additionally, Bennett and co-workers [31] validated a submaximal treadmill-based protocol. The authors noted that VO<sub>2max</sub> was better estimated when calculated from the projection of HR<sub>max</sub> [31].

With the participants of the present study, it would not be appropriate to perform maximal testing to exhaustion. They were categorized either as sedentary or active people, that is, they either did not practice any physical activity or did one or two sessions of recreative exercise, such as walking or cycling.

#### 2.4. Statistical Analysis

Initially, Kolmogorov-Smirnov and Levene tests assessed normality and homoscedasticity, respectively. Descriptive data means and one standard deviation (1SD) were calculated. For the prediction of  $VO_{2max}$  a stepwise regression (backward elimination) was computed with inclusion of all the variables in the study (sex, body mass, height, BMI, and HRR). The final model retained only significant predictors ( $p < 0.05$ ).

The validation process included three stages: (i) mean data comparison between the measured and estimated  $VO_{2max}$ , (ii) simple linear regression, and (iii) Bland–Altman analysis. For the mean data comparison, the paired samples  $t$ -test ( $p < 0.05$ ) between the measured and estimated  $VO_{2max}$  was computed. The mean difference, 95% confidence intervals, and Cohen’s  $d$  as effect size index were used.

Cohen’s  $d$  was deemed as: (i) trivial, if  $0 \leq d < 0.20$ ; (ii) small, if  $0.20 \leq d < 0.60$ ; (iii) moderate, if  $0.60 \leq d < 1.20$ ; (iv) large, if  $1.20 \leq d < 2.00$ ; (v) very large, if  $2.00 \leq d < 4.00$ ; (vi) nearly perfect, if  $d \geq 4.00$  [32]. Simple linear regression between the  $VO_{2max}$  measured and estimated was computed. The trendline, determination coefficient ( $R^2$ ), standard error of estimation (SEE), 95% of confidence (95CI) and prediction (95PI) intervals were calculated. The qualitative interpretation of the relationship was defined as: (i) very weak, if  $R^2 < 0.04$ ; (ii) weak, if  $0.04 \leq R^2 < 0.16$ ; (iii) moderate, if  $0.16 \leq R^2 < 0.49$ ; (iv) high, if  $0.49 \leq R^2 < 0.8$ ; and (v) very high, if  $0.81 \leq R^2 < 1.0$  [33]. Bland–Altman analysis included the plots of the difference and average of the  $VO_{2max}$  measured and estimated [34]. For qualitative assessment it was considered that at least 80% of the plots were within the  $\pm 1.96$  standard deviation of the difference (95CI).

### 3. Results

The multiple linear regression retained, as significant, these  $VO_{2max}$  predictors: sex ( $p < 0.001$ ) and  $HRR_{1min}$  ( $p = 0.003$ ). The age, body mass, height, and BMI were not significant, in this model.

The prediction equation ( $R^2 = 63.0\%$ ,  $SEE = 5.58$ ) revealed a close relationship between measurements and can be expressed as:

$$VO_{2max} = 22.246 + 0.343 \cdot (HRR_{1min}) + 11.722 \cdot (sex) \tag{1}$$


where  $VO_{2max}$  corresponds to the maximum oxygen uptake (mL/kg/min),  $HRR_{1min}$  corresponds to heart rate recovery (beats per minute) for one minute immediately after finishing the step test, sex corresponds to the value of zero for women and 1 for men. Table 2 presents the comparison between measured and estimated  $VO_{2max}$ . The results showed non-significant differences with a trivial effect size.

**Table 2.**  $t$ -test paired samples comparison between the  $VO_{2max}$  measured and estimated in validation group. Effect size index (Cohen’s  $d$ ) is also presented.

$VO_{2max}$ Measured [mL/kg/min]	$VO_{2max}$ Estimated [mL/kg/min]	Mean difference (95CI)	$t$ -test ( $p$ value)	$d$ [descriptor]
Mean $\pm$ 1SD	Mean $\pm$ 1SD			
42.09 $\pm$ 8.45	42.99 $\pm$ 7.10	−0.899 (−2.889 to 1.089)	−0.928 (0.362)	0.12 [trivial]

$VO_{2max}$ —maximal oxygen uptake.

Figure 1 presents the simple linear regression (panel A) between both variables. This revealed a close relationship ( $R^2 = 63.3\%$ ,  $p < 0.001$ ,  $SEE = 5.22$ ). Bland–Altman plots (panel B) also fulfilled the agreement criteria, where more than 80% of the plots were within 95CI. In this case, all plots were within the 95CI.



**Figure 1.** Linear regression between  $VO_{2max}$  measured and estimated by equation 1 (panel (A)), and Bland–Altman plots (panel (B)). All plots were within 95CI. This data refers to the validation group.

#### 4. Discussion

Our study confirmed the StepTest4all as a valid procedure in estimating  $VO_{2max}$  in the young adult population. Consequently, it was found that the magnitude of HR decrease immediately after exercise can be used to assess cardiovascular capacity. This procedure may be suitable for monitoring the level of cardiovascular capacity from an individual point of view, especially over time. This can be done by performing the StepTest4all periodically. However, some caution is needed when comparing  $VO_{2max}$  between different people. The same  $VO_{2max}$  value can mean different levels of physical capacity for a young person, an adult, a male, or a female. Thus, individual values must be checked against benchmark tables available in the literature [35] to see if they match the recommendations.

This test does not measure the performance in going up and down the step but reflects the cardiac recovery capacity after effort. The rise and drop of HR are controlled mainly by the sympathetic and parasympathetic nervous systems. Evaluating the magnitude of the decrease after exertion can also provide an assessment of nervous system function [36]. It is known that too small of a decrease in HR during the minutes following exertion cessation is associated with an increased likelihood of cardiovascular problems [37] and is even associated with premature death [7]. Conversely, a more rapid HR reduction after exercise is associated with a greater cardiovascular capacity [38]. However, there is still no consensus on the threshold value associated with a high risk of cardiovascular disease, the minimum value associated with an acceptable cardiovascular capacity, or nervous system dysfunction. For instance, Adabag et al. [39] in an article review, refer to values of 12–13 bpm of HRR in 1 min as threshold values. However, given the wide diversity of tests used and level of demand, some care is needed in defining the cut-off values (between normal and abnormal). As for the optimal recovery values, we know that healthy athletes can recover 60 or more bpm in one minute. Thus, the quality of recovery can be benchmarked with values between 12 and 60 bpm. Values close to 12 bpm may warn of a higher risk of cardiovascular disease or parasympathetic nervous system dysfunction; progressively higher values reflect very good cardiovascular capacity and good autonomic nervous system function [39,40]. The average HRR values found in our participants was

39 ± 10 bpm. This value is well above the minimum values aforementioned. Thus, they seem to agree with their age group and active lifestyle.

To provide a qualitative classification of cardiovascular capacity in this population group (young adults), we subdivided the range of HRR variation (12 to 60 bpm of recovery over one minute) usually found in these people and defined four categories, as can be observed in Table 3. This table shows a cardiovascular capacity classification and the reference values for  $VO_{2max}$  of participants between 20 and 29 years of age, obtained in our study, and the values of similar categories proposed by McArdle et al. [41]. Although the values are not completely coincident for men, in the case of women they are very close. This fact may mean that the values of  $VO_{2max}$  estimated by equation 1 are not discrepant in relation to other estimates.

In this regard, our experience with the StepTest4all allowed us to see that those values lower than 20 were usually associated with sedentary lifestyles and the presence of other risk factors (such as smoking and obesity) and values higher than 50 were usually present in people with high daily physical activity and a healthy lifestyle.

**Table 3.** Cardiovascular capacity (CVC) classification based on  $HRR_{1min}$  and the respective HRR cut-off values. It also presents the  $VO_{2max}$  values predicted by equation 1 (from our study) and the values proposed by McArdle et al. [41] for similar categories. These values are also used by a company of worldwide repute in the evaluation of body composition (Tanita: <https://tanita.eu/blog/could-improving-your-vo2-max-be-the-secret-of-success> (accessed on 28 July 2022)) [42]. These reference values are for the age group 20 to 29 years.

CVC_Classification	HRR <sub>1min</sub>	Men		Women	
		VO <sub>2max</sub>	McArdle et al., 2003 [42]	VO <sub>2max</sub>	McArdle et al., 2003 [42]
Poor	<20	<40	<36.5	<28	<29
Moderate	20–34	42–44.2	36.5–42.4	28–32.2	29–32
Good	35–49	44.3–49	42.5–46.4	32.3–36.9	33–36
Excellent	≥50	≥49	≥46.5	≥37	≥37

Because of its characteristics, the StepTest4all can be applied to people with different levels of physical ability and different somatic characteristics. To the best of our knowledge, this is the first step test that calculates the step height from several variables determining cardiovascular capacity. This calculation allows a primary adjustment, so that the test does not become too difficult/short or too easy/long. The finer adjustment is made by the precise control of the pace and its increment during the whole test. In this way, it is possible to reach the intensity of effort associated with 80% of  $HR_{max}$  (upper limit) for all in an adequate period (4 to 10 min).

In our study, age was an excluded variable from the  $VO_{2max}$  prediction model. This occurrence may be because our sample was made up of young adults. Probably, in samples with a larger range, age will also be part of the equation. The use of step tests, in addition to the aforementioned advantages, is appropriate because it is simple to apply, requires little space or means to carry out and can even be performed at home by anyone (see: <https://www.facebook.com/StepTest4all> (accessed on 28 July 2022)) [43]. The present study has some limitations: (i) it was carried out with a sample restricted to young adults, (ii) the sample was relatively small, and (iii) the amount and intensity of physical activity was not rigorously accounted for and did not enter the multiple linear regression as a possible factor. However, regarding this aspect, we can say that most participants in the sample had a similar weekly physical activity. Thus, further studies should be conducted to overcome these limitations. It should also be mentioned that the  $VO_{2max}$  and  $HR_{max}$  values of the participants were obtained by estimation. Notwithstanding, it was an estimation based on the evolution of the real individual values of  $VO_2$  and HR, both at rest and at

different intensity levels. As mentioned earlier, this is a standard procedure generally used in non-athletic participants or in special population groups that should not be submitted to maximal test to exhaustion.

## 5. Conclusions

The StepTest4all was shown to be an adequate procedure to estimate cardiovascular capacity, expressed by  $VO_{2max}$ , in young adults. The validation procedure indicated a high-level of agreement between the  $VO_2$  measured and estimated. Additionally, it is possible to determine the qualitative level of cardiovascular capacity from the  $HRR_{1min}$  more specifically, poor: <20 bpm, moderate: 20 to 34 bpm, good: 35 to 49 bpm, and excellent:  $\geq 50$  bpm. This procedure has the benefit of being simple to apply and can be used by everyone, even at home, without specialist supervision.

**Author Contributions:** Conceptualization, J.A.B.; methodology, J.A.B.; investigation, J.A.B., R.F.B., P.M.R., P.M.M., J.P.B. and J.E.M.; resources, R.F.B., P.M.R., P.M.M., J.P.B. and J.E.M.; data curation, J.A.B.; writing—original draft preparation, J.A.B., R.F.B., P.M.R., P.M.M., J.P.B. and J.E.M. The paper and data have not been published before, are not under consideration for publication elsewhere, and have been approved by all co-authors. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Project “GreenHealth-Digital Strategies in biological assets to improve well-being and promote green health” (Norte-01-0145-FEDER-000042), supported by the Regional Operational Program of Northern Portugal (NORTE 2020), under the PORTUGAL 2020 Partnership Agreement”. This research was supported by Portuguese Foundation for Science and Technology, I. P. (project UIDB/DTP/04045/2020)”.

**Institutional Review Board Statement:** All procedures were in accordance with the Declaration of Helsinki regarding human research, and the Polytechnic Ethics Board approved the research design (Nº 43/2021).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** Data is available under request to the contact author.

**Acknowledgments:** The authors express acknowledgement of the cooperation of all subjects who voluntarily participated in the collection procedures.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Cole, C.R.; Blackstone, E.H.; Pashkow, F.J.; Snader, C.E.; Lauer, M.S. Heart-rate recovery immediately after exercise as a predictor of mortality. *N. Engl. J. Med.* **1999**, *341*, 1351–1357. [[CrossRef](#)]
2. Kodama, S.; Saito, K.; Tanaka, S. Cardiorespiratory Fitness as a Quantitative Predictor of All-Cause Mortality and Cardiovascular Events in Healthy Men and Women: A Meta-Analysis. *JAMA* **2009**, *301*, 2024–2035. [[CrossRef](#)] [[PubMed](#)]
3. Jae, S.Y.; Kurl, S.; Laukkanen, J.A.; Zaccardi, F.; Choi, Y.H.; Fernhall, B.; Carnethon, M.; Franklin, B.A. Exercise Heart Rate Reserve and Recovery as Predictors of Incident Type 2 Diabetes. *Am. J. Med.* **2016**, *129*, 536.e7–536.e12. [[CrossRef](#)] [[PubMed](#)]
4. Yu, T.Y.; Jee, J.H.; Bae, J.C.; Hong, W.J.; Jin, S.M.; Kim, J.H.; Lee, M.K. Delayed heart rate recovery after exercise as a risk factor of incident type 2 diabetes mellitus after adjusting for glycometabolic parameters in men. *Int. J. Cardiol.* **2016**, *221*, 17–22. [[CrossRef](#)] [[PubMed](#)]
5. Messinger-Rapport, B.; Pothier Snader, C.E.; Blackstone, E.H.; Yu, D.; Lauer, M.S. Value of exercise capacity and heart rate recovery in older people. *J. Am. Geriatr. Soc.* **2003**, *51*, 63–68. [[CrossRef](#)] [[PubMed](#)]
6. Kurl, S.; Jae, S.Y.; Voutilainen, A.; Hagnäs, M.; Laukkanen, J.A. Exercise heart rate reserve and recovery as risk factors for sudden cardiac death. *Prog. Cardiovasc. Dis.* **2021**, *68*, 7–11. [[CrossRef](#)] [[PubMed](#)]
7. Peçanha, T.; Bartels, R.; Brito, L.C.; Paula-Ribeiro, M.; Oliveira, R.S.; Goldberger, J.J. Methods of assessment of the post-exercise cardiac autonomic recovery: A methodological review. *Int. J. Cardiol.* **2017**, *227*, 795–802. [[CrossRef](#)]
8. Adali, M.K.; Varol, E.; Aksoy, F.; Icli, A.; Ersoy, I.H.; Ozaydin, M.; Erdogan, D.; Dogan, A. Impaired heart rate recovery in patients with endemic fluorosis. *Biol. Trace Elem. Res.* **2013**, *152*, 310–315. [[CrossRef](#)]
9. Sydó, N.; Sydó, T.; Gonzalez Carta, K.A.; Hussain, N.; Farooq, S.; Murphy, J.G.; Merkely, B.; Lopez-Jimenez, F.; Allison, T.G. Prognostic Performance of Heart Rate Recovery on an Exercise Test in a Primary Prevention Population. *J. Am. Heart Assoc.* **2018**, *7*, e008143. [[CrossRef](#)]



10. Trevizani, G.A.; Benchimol-Barbosa, P.R.; Nadal, J. Effects of age and aerobic fitness on heart rate recovery in adult men. *Arq. Bras. Cardiol.* **2012**, *99*, 802–810. [[CrossRef](#)]
11. Qiu, S.; Cai, X.; Sun, Z.; Li, L.; Zuegel, M.; Steinacker, J.M.; Schumann, U. Heart Rate Recovery and Risk of Cardiovascular Events and All-Cause Mortality: A Meta-Analysis of Prospective Cohort Studies. *J. Am. Heart Assoc.* **2017**, *6*, e005505. [[CrossRef](#)]
12. Jin, Q.; Li, X.; Zhang, Y.; Zhao, Z.; Zhao, Q.; Yan, L.; Duan, A.; Luo, Q.; Liu, Z. Heart Rate Recovery at 1 Min after Exercise Is a Marker of Disease Severity and Prognosis in Chronic Thromboembolic Pulmonary Hypertension. *Respiration* **2022**, *101*, 455–464. [[CrossRef](#)]
13. Petrella, R.; Koval, J.; Cunningham, D.; Paterson, D. A self-paced step test to predict aerobic fitness in older adults in the primary care clinic. *J Am Geriatr Soc* 2001; 49:632e8. Bennett, H.; Parfitt, G.; Davison, K.; Eston, R. Validity of Submaximal Step Tests to Estimate Maximal Oxygen Uptake in Healthy Adults. *Sports Med.* **2016**, *46*, 737–750. [[CrossRef](#)]
14. Hughes, A.D.; Chaturvedi, N. Estimation of Maximal Oxygen Consumption and Heart Rate Recovery Using the Tecumseh Sub-Maximal Step Test and their Relationship to Cardiovascular Risk Factors. *Artery Res.* **2017**, *18*, 29–35. [[CrossRef](#)]
15. Pepera, G.; Panagiota, Z. Comparison of heart rate response and heart rate recovery after step test among smoker and non-smoker athletes. *Afr. Health Sci.* **2021**, *21*, 105–111. [[CrossRef](#)] [[PubMed](#)]
16. Morita, A.A.; Silva, L.K.O.; Bisca, G.W.; Oliveira, J.M.; Hernandes, N.A.; Pitta, F.; Furlanetto, K.C. Heart Rate Recovery, Physical Activity Level, and Functional Status in Subjects with COPD. *Respir. Care.* **2018**, *63*, 1002–1008. [[CrossRef](#)]
17. Haraldsdottir, K.; Brickson, S.; Sanfilippo, J.; Dunn, W.; Watson, A. In-season changes in heart rate recovery are inversely related to time to exhaustion but not aerobic capacity in rowers. *Scand. J. Med. Sci. Sports.* **2018**, *28*, 418–424. [[CrossRef](#)] [[PubMed](#)]
18. Daanen, H.A.; Lamberts, R.P.; Kallen, V.L.; Jin, A.; Van Meeteren, N.L. A systematic review on heart-rate recovery to monitor changes in training status in athletes. *Int. J. Sports Physiol. Perform.* **2012**, *7*, 251–260. [[CrossRef](#)]
19. Rodríguez-Fernández, A.; Sanchez-Sanchez, J.; Ramirez-Campillo, R.; Nakamura, F.Y.; Rodríguez-Marroyo, J.A.; Villa-Vicente, J.G. Relationship between Repeated Sprint Ability, Aerobic Capacity, Intermittent Endurance, and Heart Rate Recovery in Youth Soccer Players. *J. Strength Cond. Res.* **2019**, *33*, 3406–3413. [[CrossRef](#)]
20. Harry, K.; Booyesen, M.J. Faster Heart Rate Recovery Correlates with High-Intensity Match Activity in Female Field Hockey Players-Training Implications. *J. Strength Cond. Res.* **2020**, *34*, 1150–1157. [[CrossRef](#)]
21. Suzic Latic, J.; Dekleva, M.; Soldatovic, I.; Leischik, R.; Suzic, S.; Radovanovic, D.; Djuric, B.; Nestic, D.; Latic, M.; Mazic, S. Heart rate recovery in elite athletes: The impact of age and exercise capacity. *Clin. Physiol. Funct. Imaging* **2017**, *37*, 117–123. [[CrossRef](#)] [[PubMed](#)]
22. Armstrong, R.; Wheen, P.; Brandon, L.; Finucane, C.; Kenny, R.A.; Maree, A. Impaired autonomic function after incomplete revascularisation. *Open Heart* **2021**, *8*, e001835. [[CrossRef](#)] [[PubMed](#)]
23. Hansen, D.; Jacobs, N.; Bex, S.; D’Haene, G.; Dendale, P.; Claes, N. Are fixed-rate step tests medically safe for assessing physical fitness? *Eur. J. Appl. Physiol.* **2011**, *111*, 2593–2599. [[CrossRef](#)]
24. Fecchio, R.Y.; Brito, L.; Leicht, A.S.; Forjaz, C.L.M.; Peçanha, T. Reproducibility of post-exercise heart rate recovery indices: A systematic review. *Auton Neurosci.* **2019**, *221*, 102582. [[CrossRef](#)]
25. Mackenzie, B. Harvard Step Test [WWW]. 2007. Available online: <https://www.brianmac.co.uk/havard.htm> (accessed on 29 August 2022).
26. Tanaka, H.; Monahan, K.D.; Seals, D.R. Age-predicted maximal heart rate revisited. *J. Am. Coll. Cardiol.* **2001**, *37*, 153–156. [[CrossRef](#)]
27. Heyward, V.H. *Advanced Fitness Assessment and Exercise Prescription*, 3rd ed.; Library of Congress: Washington, DC, USA, 1997.
28. Jamnick, N.A.; By, S.; Pettitt, C.D.; Pettitt, R.W. Comparison of the YMCA and a Custom Submaximal Exercise Test for Determining VO<sub>2</sub>max. *Med. Sci. Sports Exerc.* **2016**, *48*, 254–259. [[CrossRef](#)]
29. Andersen, R.E.; Wadden, T.A. Validation of a cycle ergometry equation for predicting steady-rate VO<sub>2</sub> in obese women. *Med. Sci. Sports Exerc.* **1995**, *27*, 1457–1460. [[CrossRef](#)]
30. Evans, H.J.; Ferrar, K.E.; Smith, A.E.; Parfitt, G.; Eston, R.G. A systematic review of methods to predict maximal oxygen uptake from submaximal, open circuit spirometry in healthy adults. *J. Sci. Med. Sport.* **2015**, *18*, 183–188. [[CrossRef](#)]
31. Bennett, H.; Davison, K.; Parfitt, G.; Eston, R. Validity of a perceptually-regulated step test protocol for assessing cardiorespiratory fitness in healthy adults. *Eur. J. Appl. Physiol.* **2016**, *116*, 2337–2344. [[CrossRef](#)]
32. Hopkins, W.G. A scale of magnitude for effect statistics. In *A New View of Statistics*; Will, G., Ed.; Hopkins: Melbourne, Australia, 2002; p. 50.
33. Barbosa, T.M.; Ramos, R.; Silva, A.J.; Marinho, D.A. Assessment of passive drag in swimming by numerical simulation and analytical procedure. *J. Sports Sci.* **2018**, *36*, 492–498. [[CrossRef](#)]
34. Bland, J.M.; Altman, D. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* **1986**, *327*, 307–310. [[CrossRef](#)]
35. ACSM. *Guidelines for Exercise Testing and Prescription*, 10th ed.; Library of Congress: Washington, DC, USA, 2018.
36. Okutucu, S.; Karakulak, U.N.; Aytimir, K.; Oto, A. Heart rate recovery: A practical clinical indicator of abnormal cardiac autonomic function. *Expert Rev. Cardiovasc. Ther.* **2011**, *9*, 1417–1430. [[CrossRef](#)] [[PubMed](#)]
37. Latchman, P.L.; Yang, Q.; Kong, L.; Zhang, H.; Sebagisha, J.; De Meersman, R.E. Heart Rate Recovery, Central Systolic Pressure, and Augmentation Index in Young Healthy Individuals. *Vasc. Health Risk Manag.* **2022**, *18*, 17–25. [[CrossRef](#)]

38. Jaafar, Z.; Lim, Y.Z. A comparison of low and high dose of moderate intensity aerobic exercise on heart rate recovery of the sedentary adults: A pragmatic randomised controlled trial. *J. Sports Med. Phys. Fitness.* **2022**, *27*, 35620954. [[CrossRef](#)]
39. Adabag, S.; Pierpont, G.L. Exercise heart rate recovery: Analysis of methods and call for standards. *Heart* **2013**, *99*, 1711–1712. [[CrossRef](#)]
40. Gupta, M.; Bansal, V.; Chhabra, S.K. Abnormal heart rate recovery and chronotropic incompetence on exercise in chronic obstructive pulmonary disease. *Chron. Respir. Dis.* **2013**, *10*, 117–126. [[CrossRef](#)]
41. McArdle, W.D. *Fisiologia do Exercício: Energia, Nutrição e Desempenho Humano*; Guanabara Koogan: Rio de Janeiro, Brazil, 2003.
42. TANITA. Available online: <https://tanita.eu/blog/could-improving-your-vo2-max-be-the-secret-of-success> (accessed on 28 July 2022).
43. Bragada et al. Available online: <https://www.facebook.com/StepTest4all> (accessed on 28 July 2022).