

CATEGORICAL MILNOR SQUARES AND K-THEORY OF ALGEBRAIC STACKS

TOM BACHMANN, ADEEL A. KHAN, CHARANYA RAVI, AND VLADIMIR SOSNILO

ABSTRACT. We introduce a notion of Milnor square of stable ∞ -categories and prove a criterion under which algebraic K-theory sends such a square to a cartesian square of spectra. We apply this to prove Milnor excision and proper excision theorems in the K-theory of algebraic stacks with affine diagonal and nice stabilizers. This yields a generalization of Weibel's conjecture on the vanishing of negative K-groups for this class of stacks.

CONTENTS

Introduction	2
1. Milnor squares of stable ∞ -categories	8
1.1. The base change property	8
1.2. Precartesian squares	9
1.3. Milnor squares	10
1.4. Pro-Milnor squares	11
2. Quasi-coherent sheaves on algebraic stacks	12
2.1. Nisnevich, cdh, and Milnor squares of stacks	12
2.2. Proto-excision statements	13
2.3. Formal completions of squares	15
2.4. Formal Milnor and finite excision	17
3. Categorical Milnor excision	19
3.1. Adjustments of squares	19
3.2. The \odot -construction	20
3.3. Characterization of Milnor squares	23
3.4. Categorical Milnor excision	25
3.5. Categorical pro-Milnor excision	26
4. Localizing invariants of algebraic stacks	31
4.1. Étale excision	32
4.2. Formal Milnor and finite excision	32
4.3. Derived blow-ups	34
4.4. Formal proper excision	38
5. Negative K-theory	39
5.1. Nil-invariance of negative K-groups	39
5.2. Killing by blow-ups	40
5.3. Weibel's conjecture (I)	41
5.4. Weibel's conjecture (II)	42
Appendix A. Algebraic stacks	44
A.1. ANS stacks	44
A.2. The resolution property	46
A.3. Compact generation	48
A.4. Dimension of algebraic stacks	49

Appendix B. Formal stacks	51
B.1. Formal completion	51
B.2. Quasi-coherent sheaves	53
Appendix C. Weak pro-Milnor squares	53
C.1. Connected invariants	53
C.2. Weak pro-Milnor squares	54
C.3. Weak pro-excision	56
References	57

INTRODUCTION

A *Milnor square* of rings, following [Mil71, Sect. 2], is a cartesian square

$$\begin{array}{ccc}
 A & \twoheadrightarrow & A/I \\
 \downarrow & & \downarrow \\
 B & \twoheadrightarrow & B/J
 \end{array} \tag{0.0.a}$$

where $I \subseteq A$ and $J \subseteq B$ are two-sided ideals. The starting point of this paper is the following result of Land and Tamme, building on work of Morrow, Geisser and Hesselholt, and Suslin (see Corollaries 2.10 and 2.33 in [LT19], as well as [Tam18], [Mor18], [GH06] and [Sus95]):

Theorem 0.0.1.

- (i) *If the square (0.0.a) is Tor-independent, i.e., the group $\mathrm{Tor}_i^A(A/I, B)$ vanishes for all $i > 0$, then the square*

$$\begin{array}{ccc}
 \mathrm{K}(A) & \longrightarrow & \mathrm{K}(A/I) \\
 \downarrow & & \downarrow \\
 \mathrm{K}(B) & \longrightarrow & \mathrm{K}(B/J)
 \end{array}$$

is cartesian.

- (ii) *If the pro-system $\{\mathrm{Tor}_i^A(A/I^n, B)\}_{n>0}$ vanishes for all $i > 0$, then the square of pro-spectra*

$$\begin{array}{ccc}
 \{\mathrm{K}(A)\} & \longrightarrow & \{\mathrm{K}(A/I^n)\}_{n>0} \\
 \downarrow & & \downarrow \\
 \{\mathrm{K}(B)\} & \longrightarrow & \{\mathrm{K}(B/J^n)\}_{n>0}
 \end{array}$$

is cartesian.

Remark 0.0.2. A sufficient condition for the pro-Tor-independence in (ii) is pro-Tor-unity of I (see e.g. [LT19, Lem. 2.14]). As observed by Morrow, the latter condition is equivalent to the vanishing of $\{\mathrm{Tor}_i^A(A/I^n, A/I^n)\}_{n>0}$ for all $i > 0$ (see [Mor18, Thm. 0.2]). Moreover, this is automatic in the case of noetherian commutative rings (see [Mor18, Thm. 0.3]).

Our first goal in this paper is to prove a categorical version of Theorem 0.0.1.

Definition 0.0.3.

- (i) Let Δ be a commutative square of presentable stable ∞ -categories and colimit-preserving functors of the form

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array} \quad (0.0.b)$$

- (a) We say Δ is a *precartesian* if the canonical functor $(p^*, f^*) : \mathcal{A} \rightarrow \mathcal{A}' \times_{\mathcal{B}'} \mathcal{B}$ is fully faithful.
- (b) We say Δ is a *Milnor square* if it is precartesian, and each of the functors f^* , g^* , p^* and q^* is compact and generates its codomain under colimits.
- (c) We say Δ *satisfies base change* if it is vertically right-adjointable; that is, the base change transformation $f^*p_* \rightarrow q_*g^*$ is invertible (where p_* and q_* are the right adjoints of p^* and q^* , respectively).
- (ii) Let Δ be a commutative square of pro-systems in the ∞ -category of presentable stable ∞ -categories and compact colimit-preserving functors. We say Δ is a *pro-Milnor square*, resp. *satisfies pro-base change*, if it can be represented by a cofiltered system $\{\Delta_n\}_n$ where each Δ_n is a Milnor square, resp. satisfies base change.

Given a presentable stable ∞ -category \mathcal{A} , we write simply $K(\mathcal{A})$ for the nonconnective algebraic K-theory spectrum of the full subcategory \mathcal{A}^ω of compact objects. Our first main result is as follows (see Theorems 3.4.3 and 3.5.11):

Theorem A (Categorical Milnor excision).

- (i) Suppose Δ is a Milnor square of compactly generated stable ∞ -categories. If Δ satisfies base change, then the induced square

$$\begin{array}{ccc} K(\mathcal{A}) & \xrightarrow{f^*} & K(\mathcal{B}) \\ \downarrow p^* & & \downarrow q^* \\ K(\mathcal{A}') & \xrightarrow{g^*} & K(\mathcal{B}') \end{array}$$

is cartesian.

- (ii) Suppose Δ is a pro-Milnor square satisfying the projective generation and boundedness hypotheses of 3.5.11. If Δ satisfies pro-base change, then the induced square of pro-spectra $K(\Delta)$ is cartesian.

Example 0.0.4. Let $\mathbf{D}(A)$ denote the derived ∞ -category of left A -modules over a ring A . For any Milnor square as in (0.0.a), the induced square

$$\begin{array}{ccc} \mathbf{D}(A) & \longrightarrow & \mathbf{D}(A/I) \\ \downarrow & & \downarrow \\ \mathbf{D}(B) & \longrightarrow & \mathbf{D}(B/J) \end{array}$$

is a Milnor square of stable ∞ -categories. It satisfies base change precisely when $\mathrm{Tor}_i^A(A/I, B) = 0$ for all $i > 0$, and the induced square of pro- ∞ -categories (formed by taking the ideal I^n for $n > 0$) satisfies pro-base change precisely when $\{\mathrm{Tor}_i^A(A/I^n, B)\}_{n>0} = 0$ for all $i > 0$. Thus Theorem A may be regarded as a generalization of Theorem 0.0.1.

Remark 0.0.5. Land and Tamme prove Theorem 0.0.1 as a consequence of a more general result [LT19, Thm. A] which applies to cartesian squares of \mathcal{E}_1 -ring spectra. Theorem A also recovers this result when applied to stable ∞ -categories of module spectra.

Remark 0.0.6. In Theorem A, nonconnective algebraic K-theory can be replaced by any localizing invariant of stable ∞ -categories (which is not required to preserve filtered colimits).

Theorem A allows us to prove excision statements in the K-theory of algebraic stacks. For a (quasi-compact quasi-separated) algebraic stack \mathcal{X} , we let $\mathbf{D}(\mathcal{X})$ denote the derived ∞ -category of quasi-coherent sheaves on \mathcal{X} , $\mathrm{Perf}(\mathcal{X}) \subseteq \mathbf{D}(\mathcal{X})$ the full subcategory of perfect complexes, and $\mathbf{K}(\mathcal{X})$ the nonconnective algebraic K-theory spectrum of $\mathrm{Perf}(\mathcal{X})$. If \mathcal{Z} is a closed substack, we regard the formal completion $\mathcal{X}_{\mathcal{Z}}^{\wedge}$ as an ind-algebraic stack so that there is naturally associated to it a ‘‘continuous K-theory’’ pro-spectrum $\widehat{\mathbf{K}}(\mathcal{X}_{\mathcal{Z}}^{\wedge})$. For example, for I an ideal in a noetherian commutative ring A , the continuous K-theory of the formal completion of $\mathrm{Spec}(A)$ along the vanishing locus of I is the pro-spectrum $\{\mathbf{K}(A/I^n)\}_n$. We then have:

Theorem B (Milnor excision). *Let Δ be a commutative square*

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

of noetherian ANS stacks (algebraic stacks with affine diagonal and nice stabilizers). Assume that Δ is a Milnor square: that is, it is cartesian and cocartesian, f is an affine morphism, and i is a closed immersion.

(i) *If Δ is Tor-independent, then the square*

$$\begin{array}{ccc} \mathbf{K}(\mathcal{X}) & \longrightarrow & \mathbf{K}(\mathcal{Z}) \\ \downarrow & & \downarrow \\ \mathbf{K}(\mathcal{X}') & \longrightarrow & \mathbf{K}(\mathcal{Z}') \end{array}$$

is cartesian.

(ii) *The induced square of pro-spectra*

$$\begin{array}{ccc} \{\mathbf{K}(\mathcal{X})\} & \longrightarrow & \widehat{\mathbf{K}}(\mathcal{X}_{\mathcal{Z}}^{\wedge}) \\ \downarrow & & \downarrow \\ \{\mathbf{K}(\mathcal{X}')\} & \longrightarrow & \widehat{\mathbf{K}}(\mathcal{X}'_{\mathcal{Z}'}^{\wedge}) \end{array}$$

is cartesian.

Theorem C (Proper excision). *Consider a cartesian square of ANS noetherian algebraic stacks*

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

where i is a closed immersion and f is a proper representable morphism which is an isomorphism away from \mathcal{Z} . Then the induced square of pro-spectra

$$\begin{array}{ccc} \{K(\mathcal{X})\} & \longrightarrow & \widehat{K}(\mathcal{X}_{\mathcal{Z}}^{\wedge}) \\ \downarrow & & \downarrow \\ \{K(\mathcal{X}')\} & \longrightarrow & \widehat{K}(\mathcal{X}'_{\mathcal{Z}'}^{\wedge}) \end{array}$$

is cartesian.

Remark 0.0.7. Theorem C can be viewed as a generalization of [KST17, Thm. A], with the minor caveat that a scheme is ANS if and only if it has affine diagonal. However, the statement for arbitrary noetherian schemes can be deduced from the case of schemes with affine diagonal using Zariski descent.

Remark 0.0.8. Theorems B and C also hold for arbitrary localizing invariants of stable ∞ -categories. This is an improvement on documented results even in the case of schemes. In particular, we find that Theorem 0.0.1(ii) generalizes to arbitrary localizing invariants.

Remark 0.0.9. The condition on nice stabilizers in Theorem B, and in the finite case of Theorem C, can be relaxed. In fact, Theorem B(i) holds for all perfect stacks in the sense of Definition A.3.1. In Theorem B(ii) and the finite case of Theorem C, it suffices to assume that \mathcal{X} admits a scallop decomposition $(\mathcal{U}_i, \mathcal{V}_i, u_i)_i$ as in [KR21, Def. 2.7] where \mathcal{V}_i are quotients of affines by actions of embeddable group schemes that are *linearly reductive* (but not necessarily nice; compare Theorem A.1.8, Proposition A.1.9). The stronger condition of niceness of all stabilizers is important in our proof of the general case of Theorem C because it ensures that linear reductivity of the stabilizers is preserved under blow-ups.

Finally, we apply Theorem C to generalize to stacks the proof in [KST17] of Weibel’s conjecture on negative K-theory (see [Wei80, 2.9]).

Theorem D (Weibel’s conjecture). *Let \mathcal{X} be a noetherian ANS stack of covering dimension d .*

- (i) *The negative K-groups $K_{-n}(\mathcal{X})$ vanish for all $n > d$.*
- (ii) *For every vector bundle $\pi : \mathcal{E} \rightarrow \mathcal{X}$, the map*

$$\pi^* : K_{-n}(\mathcal{X}) \rightarrow K_{-n}(\mathcal{E})$$

is bijective for all $n \geq d$.

Remark 0.0.10. For Deligne–Mumford stacks, the covering dimension coincides with the Krull dimension and with the usual dimension as in [Stacks, Tag 0AFL]. In general, the latter can be negative and so is not suitable for the purposes of Theorem D. See Subsect. A.4 for the definitions.

Example 0.0.11 (Equivariant K-theory). Let k be a field and G a group scheme over k acting on a noetherian finite-dimensional k -scheme X with affine diagonal. If G is finite of order prime to the characteristic of k , or if G is a torus, then the quotient stack $\mathcal{X} = [X/G]$ is noetherian and ANS (see Subsect. A.1 for more examples) and $K(\mathcal{X})$ is the algebraic K-theory $K^G(X)$ of G -equivariant perfect complexes on X as in [KR18b] (see also [Tho87]). Thus Theorem D implies that $K_{-n}^G(X)$ vanishes for all $n > \dim(X)$.

The following example was pointed out to us by B. Antieau.

Example 0.0.12 (Twisted K-theory). Let \mathcal{X} be a noetherian ANS stack of covering dimension d and let $\mathcal{Y} \rightarrow \mathcal{X}$ be a \mathbf{G}_m -gerbe over \mathcal{X} . Then \mathcal{Y} is also of covering dimension d , since $\mathcal{Y} \rightarrow \mathcal{X}$ is smooth and \mathcal{Y} is étale-locally on \mathcal{X} a trivial \mathbf{G}_m -gerbe. Therefore by Theorem D, $K_{-n}(\mathcal{Y})$ vanishes for all $n > d$. For any Azumaya algebra \mathcal{A} over a d -dimensional noetherian scheme X , we may apply this to the \mathbf{G}_m -gerbe \mathcal{Y} over X associated to the Brauer class representing the Azumaya algebra \mathcal{A} . Then $\text{Perf}(\mathcal{Y})$ is equivalent to the stable ∞ -category of \mathcal{A} -twisted perfect complexes on X , and in particular we find that the \mathcal{A} -twisted K-groups $K_{-n}^{\mathcal{A}}(X)$ of X vanish for all $n > d$. This recovers the main result of [Sta20] for Azumaya algebras.

Remark 0.0.13. To our knowledge, Theorem D is the first result in the literature about negative K-groups of singular stacks, or even negative equivariant K-groups of singular schemes with group

action. Recall that in the nonsingular case, all the groups $K_{-n}(\mathcal{X})$ vanish for $n > 0$. See [Ker18] for some discussion about the relationship between negative K-groups and singularities in the setting of schemes.

Outline. We define categorical Milnor squares, and their pro-versions, in Sect. 1.

In Sect. 2 we give many examples of categorical Milnor and pro-Milnor squares of derived ∞ -categories of (derived) algebraic stacks. The appearance of pro- ∞ -categories comes from the key observation that failure of the base change property can often be rectified by passing to formal completions. Let $\widehat{\mathbf{D}}(\mathcal{X})$ denote the canonical pro- ∞ -categorical refinement of the derived ∞ -category $\mathbf{D}(\mathcal{X})$ of a formal stack (or ind-algebraic stack) \mathcal{X} . Then we prove, for instance, that if

$$\begin{array}{ccc} z' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow \\ z & \longrightarrow & \mathcal{X} \end{array}$$

is either a Milnor square or finite cdh square of noetherian algebraic stacks, then the induced square of pro- ∞ -categories

$$\begin{array}{ccc} \{\mathbf{D}(\mathcal{X})\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}_z^\wedge) \\ \downarrow & & \downarrow \\ \{\mathbf{D}(\mathcal{X}')\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}'_{z'}^\wedge) \end{array}$$

is not only a pro-Milnor square but also satisfies pro-base change (see Corollaries 2.4.2 and 2.4.3). These results can be viewed as pro-refinements of some results of Halpern-Leistner and Preygel [HLP14].

Sect. 3 deals with the proof of Theorem A. Our main tool is a categorical version of the \odot -construction introduced in [LT19]. For every categorical Milnor square Δ of the form (0.0.b), we construct a new square Δ_0

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q_0^* \\ \mathcal{A}' & \xrightarrow{g_0^*} & \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}, \end{array}$$

which is isomorphic to Δ precisely when the latter satisfies base change (Theorem 3.3.1). This is the generalization of [LT19, Main Theorem] in this setting. The proof of the pro-variant is somewhat involved, as it requires a user-friendly criterion for a cofiltered system of compact colimit-preserving functors of presentable stable ∞ -categories to induce a pro-equivalence (see Corollary 3.5.10). This question is surprisingly subtle and forces us to impose strong projective generation hypotheses on our categories, which roughly puts us in the situation of “weighted” ∞ -categories (see Remark 3.5.7). The reader willing to accept Theorem A as a black box can safely skip this section.

Theorems B and C are proven in Sect. 4. This involves three main steps:

- Applying Theorem A to the categorical pro-Milnor squares constructed in Corollaries 2.4.2 and 2.4.3, we get Theorem B as well as Theorem C in the case of finite morphisms. See Corollaries 4.2.2 and 4.2.3.
- Next we prove a formally completed version of [Kha20, Thm. A], which yields formal excision for blow-ups in quasi-smooth derived centres (Proposition 4.3.2). Combining this with finite excision yields the case of blow-ups (in arbitrary centres).

- For the general case of Theorem C, we use a generalization of the arguments of [KST17] to reduce to the case of a blow-up. The key ingredient, as in [HK19] and [Kha20, 5.3.4], is Rydh’s stacky extension of the flatification theorem of Raynaud and Gruson. See Theorem 4.4.1.

The proof of Theorem D is accomplished in Sect. 5. As in [KST17], the proof is a relatively straightforward consequence of proper excision (Theorem C) and a “killing lemma” that allows one to kill negative K-classes by blowing up (see [KS17, Prop. 5], [HK19, Prop. 7.3] for stacks, and Proposition 5.2.1 for our version with supports). To use the latter, we also need a nil-invariance result for low enough K-groups (Corollary 5.1.4).

Appendices A and B develop some preliminary material on (derived) algebraic and formal stacks. In appendix C, we study a “weak” version of categorical pro-Milnor squares, where pro- ∞ -categories are not regarded up to isomorphism but rather only up to weak pro-equivalence. This can be used to drop the boundedness condition imposed in Theorem A, and hence also the boundedness conditions on the structure sheaves of derived stacks in some statements in Sects. 2 and 4.

Related work. Theorem A can be contrasted with a recent result of [Tam18, Thm. 18] (cf. [Hoy18, Cor. 13]), which provides a different criterion for a square of compactly generated stable ∞ -categories as in (0.0.b) to induce a cartesian square in algebraic K-theory: namely, it suffices that the square be cartesian and the functor p_* fully faithful. Our base change condition can be regarded as a generalization of this, since for a Milnor square with p_* fully faithful, the base change property is automatic (see Lemma 1.3.4). However, Theorem A is not strictly more general than Tamme’s criterion because the definition of Milnor square also requires that the functors f^* and g^* generate under colimits (or equivalently, that their right adjoints f_* and g_* are conservative).

The idea of passing to formal completions to prove excision statements for derived ∞ -categories of quasi-coherent sheaves is present in the work of Halpern-Leistner and Preygel (compare Corollary 2.4.3 with [HLP14, Lem. 3.3.4], for example). To our knowledge, the more refined invariant $\widehat{\mathbf{D}}(\mathcal{X}_2^\wedge)$ (see Subsect. B.2) has not been considered in the literature before.

The analogue of Theorem C in homotopy invariant K-theory was obtained recently by Hoyois and Krishna [HK19], under slightly weaker hypotheses. Another proof in homotopy invariant K-theory (that applies more generally to truncating invariants in the sense of [LT19]) was given in [Kha20, Thm. 5.6, Rmk. 5.11(iii)], modulo a derived invariance property that was later established independently in [ES20] and [KR21, Cor. F]. Our proof combines the arguments of [KST17] and [Kha20].

Hoyois and Krishna also proved a variant of Theorem D in homotopy invariant K-theory (see [HK19, Thm. 1.1]). Compared to their result, our Theorem D applies to a larger class of stacks at the cost of a possibly less sharp bound (using covering dimension instead of blow-up dimension).

Notation and conventions. We freely use the language of ∞ -categories and derived algebraic geometry. We generally follow the notation of [HTT, HA], [GR17] and [HLP14]. Some exceptions are as follows:

- In an ∞ -category \mathcal{C} , we write $\mathrm{Maps}_{\mathcal{C}}(C, D)$ for the mapping space between any two objects C and D .
- We write Spc and Spt for the ∞ -categories of spaces and spectra, respectively. The ∞ -category of presentable ∞ -categories and colimit-preserving functors (see [HTT, Defn. 5.5.3.1])

will be denoted Pres . The (non-full) subcategory of Pres where the morphisms are *compact* colimit-preserving functors will be denoted Pres_c . Recall that a colimit-preserving functor is compact if its right adjoint preserves filtered colimits (see e.g. [SAG, Defn. C.3.4.2]).

- A *derived commutative ring* is an object of the nonabelian derived ∞ -category of ordinary commutative rings. This ∞ -category can be realized as the localization of the category of simplicial commutative rings. See [SAG, 25.1] for details. The reader may also choose to read the term “derived commutative ring” as “connective \mathcal{E}_∞ -ring” as in [HA, Chap. 7].
- We write LMod_R for the stable ∞ -category of left modules over an \mathcal{E}_1 -ring spectrum R . If R is a derived commutative ring, we write Mod_R for the stable ∞ -category of modules over the underlying \mathcal{E}_∞ -ring spectrum.
- Given a derived commutative ring R and a collection of elements $f_1, \dots, f_n \in \pi_0(R)$, we write $R//\langle f_1, \dots, f_n \rangle$ for the derived commutative ring of functions on the derived zero locus of f_1, \dots, f_n . That is, $R//\langle f_1, \dots, f_n \rangle$ is the derived tensor product of R and \mathbf{Z} over $\mathbf{Z}[T_1, \dots, T_n]$ as in [KR18a, 2.3.1].
- A derived algebraic stack is a derived 1-Artin stack as in [GR17, Chap. 2, 4.1]. All derived algebraic stacks are assumed quasi-compact and quasi-separated. A derived algebraic stack \mathcal{X} is *bounded* if it admits a smooth surjection $\text{Spec}(A) \rightarrow \mathcal{X}$ where A is a bounded derived commutative ring ($\pi_i(A) = 0$ for $i \gg 0$).
- We write $\mathbf{D}(\mathcal{X})$ for the derived ∞ -category of quasi-coherent sheaves on a derived algebraic stack \mathcal{X} , defined as in [GR17, Chap. 3, 1.1.4]. When \mathcal{X} is a classical algebraic stack, this agrees with the derived ∞ -category of $\mathcal{O}_{\mathcal{X}}$ -modules with quasi-coherent cohomology (see [HR17, Prop. 1.3]).

Acknowledgments. We would like to thank the organizers of the event “Algebraic Groups and Motives” in St. Petersburg in September 2019, where the preliminary versions of these results were first obtained. We thank David Rydh for helpful discussions about [AHR19] and [AHHLR], and for extensive comments on a draft. We thank Charles Weibel for his comments on a draft. The third author thanks Benjamin Antieau for discussions related to the ‘many-objects version’ of the Land–Tamme excision theorem. The fourth author thanks Elden Elmanto for discussing the relations between various excision results.

The second author acknowledges partial support from SFB 1085 Higher Invariants, Universität Regensburg, the Simons Collaboration on Homological Mirror Symmetry, and MOST 110-2115-M-001-016-MY3. The third author also acknowledges support from SFB 1085 Higher Invariants. The fourth author’s work on sections 1, 3, 4, 5, and Appendix C was completed under the support of Russian Science Foundation grant 20-41-04401.

1. MILNOR SQUARES OF STABLE ∞ -CATEGORIES

1.1. The base change property.

Notation 1.1.1. Recall that any colimit-preserving functor between presentable ∞ -categories admits a right adjoint by the adjoint functor theorem. We will usually use a symbol of the form f^* to denote such a functor, and f_* for its right adjoint. We also write $\eta_f : \text{id} \rightarrow f_*f^*$ and $\varepsilon_f : f^*f_* \rightarrow \text{id}$ for the unit and co-unit transformations. This is just a notational device: we have not assigned any meaning to the symbol “ f ” itself.

Construction 1.1.2 (Exchange transformation). Let Δ be a commutative square

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array}$$

of presentable ∞ -categories and colimit-preserving functors. The *exchange transformation* associated to Δ is the canonical natural transformation

$$\mathrm{Ex}_\Delta : f^* p_* \rightarrow q_* g^*$$

of functors $\mathcal{A}' \rightarrow \mathcal{B}$ defined as the composite

$$f^* p_* \xrightarrow{\eta_q} q_* q^* f^* p_* \simeq q_* g^* p^* p_* \xrightarrow{\varepsilon_p} q_* g^*.$$

Definition 1.1.3. When Ex_Δ is invertible, we say that the square Δ *satisfies base change*¹.

Example 1.1.4. Suppose given a commutative square

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ A' & \longrightarrow & B' \end{array}$$

of rings or more generally of \mathcal{E}_1 -ring spectra. This induces a square of stable ∞ -categories of module spectra

$$\begin{array}{ccc} \mathrm{LMod}_A & \longrightarrow & \mathrm{LMod}_B \\ \downarrow & & \downarrow \\ \mathrm{LMod}_{A'} & \longrightarrow & \mathrm{LMod}_{B'}, \end{array}$$

where the functors are each given by (derived) extension of scalars. Then the exchange transformation evaluated on any object $M' \in \mathrm{LMod}_{A'}$ is the canonical B -module morphism

$$B \otimes_A M' \rightarrow B' \otimes_{A'} M'.$$

Since $\mathrm{LMod}_{A'}$ is generated under colimits by $A' \in \mathrm{LMod}_{A'}$, we see that this square satisfies base change if and only if the canonical morphism

$$B \otimes_A A' \rightarrow B'$$

is invertible.

1.2. Precartesian squares.

Definition 1.2.1. We say that a commutative square in Pres

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array}$$

is *precartesian* if the canonical functor

$$(p^*, f^*) : \mathcal{A} \rightarrow \mathcal{A}' \times_{\mathcal{B}'} \mathcal{B}$$

¹It is also common to say that the square satisfies the *Beck–Chevalley condition*, or that it is vertically *right-adjointable*.

is fully faithful. This is equivalent to the condition that the square of natural transformations

$$\begin{array}{ccc} \mathrm{id} & \xrightarrow{\eta_f} & f_* f^* \\ \downarrow \eta_p & & \downarrow \eta_q \\ p_* p^* & \xrightarrow{\eta_g} & f_* q_* q^* f^* \end{array} \quad (1.2.a)$$

is cartesian.

Example 1.2.2. In the situation of Example 1.1.4, the square is precartesian if and only if the original square of \mathcal{E}_1 -rings is cartesian.

Warning 1.2.3. A cartesian square of ordinary rings

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ A' & \longrightarrow & B' \end{array}$$

is not necessarily cartesian as a square of \mathcal{E}_1 -rings, as the fibred product $A' \times_{B'} B$ may acquire negative homotopy groups when taken in the ∞ -category of \mathcal{E}_1 -rings. A sufficient condition is that $A' \rightarrow B'$ is surjective. For example, Milnor squares of rings give rise to cartesian squares of \mathcal{E}_1 -rings and hence to precartesian squares of presentable stable ∞ -categories.

Example 1.2.4. If

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f} & \mathcal{B} \\ \downarrow p & & \downarrow q \\ \mathcal{A}' & \xrightarrow{g} & \mathcal{B}' \end{array}$$

is a cartesian square of small stable ∞ -categories and exact functors, then the induced square

$$\begin{array}{ccc} \mathrm{Ind}(\mathcal{A}) & \xrightarrow{f^*} & \mathrm{Ind}(\mathcal{B}) \\ \downarrow p^* & & \downarrow q^* \\ \mathrm{Ind}(\mathcal{A}') & \xrightarrow{g^*} & \mathrm{Ind}(\mathcal{B}') \end{array}$$

is precartesian. (Indeed, the fully faithful functor

$$\mathcal{A} \xrightarrow{\sim} \mathcal{A}' \times_{\mathcal{B}'} \mathcal{B} \hookrightarrow \mathrm{Ind}(\mathcal{A}') \times_{\mathrm{Ind}(\mathcal{B}')} \mathrm{Ind}(\mathcal{B})$$

factors through the full subcategory of compact objects, since filtered colimits commute with finite limits of spaces.) However, it is cartesian only under additional hypotheses: for example, when g_* is fully faithful [EHIK20, Lem. 4].

1.3. Milnor squares.

Lemma 1.3.1. *Let $f^* : \mathcal{A} \rightarrow \mathcal{B}$ be a colimit-preserving functor between presentable stable ∞ -categories. Then the following conditions are equivalent:*

- (i) *If $\mathcal{B}_0 \subseteq \mathcal{B}$ is a cocomplete stable subcategory containing the essential image of $f^* : \mathcal{A} \rightarrow \mathcal{B}$, then $\mathcal{B}_0 = \mathcal{B}$.*
- (ii) *The right orthogonal to the essential image of $f^* : \mathcal{A} \rightarrow \mathcal{B}$ is the zero subcategory.*
- (iii) *The right adjoint $f_* : \mathcal{B} \rightarrow \mathcal{A}$ is conservative.*

Proof. See e.g. [MNN17, Lemma 7.6] for the equivalence of the first two conditions. The second and third are equivalent by adjunction. \square

Definition 1.3.2. If $f^* : \mathcal{A} \rightarrow \mathcal{B}$ satisfies the equivalent conditions of Lemma 1.3.1, then we say f^* generates \mathcal{B} under colimits.

Definition 1.3.3. Let Δ be a commutative square in Pres of the form

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array}$$

We say Δ is a *Milnor square* if it is precartesian and each of the functors f^* , g^* , p^* and q^* is compact and generates its codomain under colimits.

Lemma 1.3.4. Suppose given a Milnor square Δ of presentable stable ∞ -categories as above. If the functor p^* is a localization, i.e., its right adjoint p_* is fully faithful, then Δ satisfies base change.

Proof. The exchange transformation is by definition the composite

$$f^* p_* \xrightarrow{\eta_q} q_* q^* f^* p_* \simeq q_* g^* p^* p_* \xrightarrow{\varepsilon_p} q_* g^*.$$

The counit ε_p is invertible by assumption, so it will suffice to show that the first arrow induced by the unit η_q is invertible. Since Δ is precartesian, we have by precomposition of (1.2.a) with p_* the cartesian square

$$\begin{array}{ccc} p_* & \xrightarrow{\eta_f} & f_* f^* p_* \\ \downarrow \eta_p & & \downarrow \eta_q \\ p_* p^* p_* & \xrightarrow{\eta_g} & f_* q_* q^* f^* p_* \end{array}$$

Since f_* is conservative (Lemma 1.3.1), it will suffice to show that the right-hand arrow is invertible. The left-hand arrow in the square is invertible by the assumption that ε_p is invertible and the adjunction identities. By stability, the claim follows. \square

1.4. Pro-Milnor squares. We now define a pro-version of Definition 1.3.3.

For a presentable ∞ -category \mathcal{C} , we let $\text{Pro}(\mathcal{C})$ denote the ∞ -category of pro-objects in \mathcal{C} as in [SAG, A.8.1]. Any pro-object $X \in \text{Pro}(\mathcal{C})$ can be represented (non-uniquely) by a cofiltered system $\{X_i\}_{i \in \mathcal{J}}$, i.e., a diagram $\mathcal{J} \rightarrow \mathcal{C}$ from a cofiltered ∞ -category \mathcal{J} . If $X \in \text{Pro}(\mathcal{C})$ and $Y \in \text{Pro}(\mathcal{C})$ are represented by cofiltered systems $\{X_i\}_i$ and $\{Y_j\}_j$, respectively, then the mapping space can be computed by the formula

$$\text{Maps}(X, Y) \simeq \lim_{\leftarrow j} \lim_{\rightarrow i} \text{Maps}_{\mathcal{C}}(X_i, Y_j),$$

see [SAG, Rem. A.8.1.5]. From this one sees that the functor $\mathcal{C} \rightarrow \text{Pro}(\mathcal{C})$ sending an object $C \in \mathcal{C}$ to the constant pro-system $\{C\}$ is fully faithful. It also implies that for any cofiltered ∞ -category \mathcal{J} , the functor of “passage to pro-objects”

$$\text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \text{Pro}(\mathcal{C})$$

commutes with finite limits and colimits, since filtered colimits of spaces commute with finite limits.

Definition 1.4.1. A commutative square Δ in $\text{Pro}(\text{Pres}_c)$ is *pro-precartesian*, *pro-Milnor*, or *satisfies pro-base change* if it can be represented by a cofiltered system $\{\Delta_n\}_n$ of commutative squares

$$\begin{array}{ccc} \mathcal{A}_n & \xrightarrow{f_n^*} & \mathcal{B}_n \\ \downarrow p_n^* & & \downarrow q_n^* \\ \mathcal{A}'_n & \xrightarrow{g_n^*} & \mathcal{B}'_n \end{array}$$

such that every Δ_n has the respective property.

2. QUASI-COHERENT SHEAVES ON ALGEBRAIC STACKS

In this section, we give many examples of Milnor and pro-Milnor squares coming from squares of (derived) schemes and stacks.

2.1. Nisnevich, cdh, and Milnor squares of stacks.

Definition 2.1.1. A *Nisnevich square* of derived algebraic stacks is a cartesian square

$$\begin{array}{ccc} \mathcal{U}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{U} & \xrightarrow{j} & \mathcal{X}, \end{array} \quad (2.1.a)$$

where j is a quasi-compact open immersion and f is a representable étale morphism of finite presentation, inducing an isomorphism $f^{-1}(\mathcal{Z}) \rightarrow \mathcal{Z}$ for some closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ complementary to j . An *affine Nisnevich square* is a Nisnevich square where f is affine.

Definition 2.1.2. A *Milnor square* of algebraic stacks is a commutative square

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array} \quad (2.1.b)$$

which is cartesian and cocartesian, where f is an affine morphism and i is a closed immersion with quasi-compact open complement.

Definition 2.1.3. A *proper cdh square* (or *abstract blow-up square*) of derived algebraic stacks is a commutative square

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array} \quad (2.1.c)$$

satisfying the following properties:

- (i) The square is cartesian on classical truncations, i.e. $\mathcal{Z}' \rightarrow \mathcal{Z} \times_{\mathcal{X}} \mathcal{X}'$ induces an isomorphism $\mathcal{Z}'_{\text{cl}} \simeq (\mathcal{Z} \times_{\mathcal{X}} \mathcal{X}')_{\text{cl}}$.
- (ii) The morphism f is representable and proper, and i is a closed immersion with quasi-compact open complement.
- (iii) The induced map $f_{\mathcal{U}} : \mathcal{U}' \rightarrow \mathcal{U}$ is invertible, where \mathcal{U} (resp. \mathcal{U}') is the open complement of \mathcal{Z} in \mathcal{X} (resp. of \mathcal{Z}' in \mathcal{X}').

A *finite cdh square* is a proper cdh square as above where the morphism f is finite. The class of *cdh squares* is the union of Nisnevich squares and proper cdh squares.

Example 2.1.4. Given a diagram of algebraic spaces or stacks

$$\mathcal{X}_0 \xleftarrow{i_0} \mathcal{X}_{01} \xrightarrow{f} \mathcal{X}_1,$$

where i_0 is a closed immersion with quasi-compact open complement and f is a finite morphism, the operation of forming the pushout $\mathcal{X} = \mathcal{X}_0 \sqcup_{\mathcal{X}_{01}} \mathcal{X}_1$ is often called “pinching”. The square

$$\begin{array}{ccc} \mathcal{X}_{01} & \xrightarrow{i_0} & \mathcal{X}_0 \\ \downarrow f & & \downarrow f' \\ \mathcal{X}_1 & \longrightarrow & \mathcal{X} \end{array}$$

is a Milnor square by [Ryd11, Thm. A.4]. See also [Rao74, Prop. 1] for the case of (separated noetherian) algebraic spaces. Note that, by [Ryd11, Thm. A.4], the square is also almost a finite cdh square, except that f' is only integral but not necessarily of finite type. If all stacks involved are of finite type over a noetherian base, then f' is of finite type and hence finite.

Example 2.1.5. In Example 2.1.4, if the morphism f is only affine, then the square is called a *Ferrand pushout* after [Fer03]. See [Art70, Thm. 6.1] and [AHHLR] for the theory of Ferrand pushouts in the setting of algebraic spaces and stacks, respectively.

2.2. Proto-excision statements. In this subsection we begin studying the behaviour of the functor $\mathcal{X} \mapsto \mathbf{D}(\mathcal{X})$ with respect to various classes of squares of stacks.

Remark 2.2.1. Let \mathcal{X} and \mathcal{Y} be derived algebraic stacks. For any representable morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$, the functor $f^* : \mathbf{D}(\mathcal{Y}) \rightarrow \mathbf{D}(\mathcal{X})$ is compact (see e.g. [HLP14, Prop. A.1.5]). If f is affine (or quasi-affine), then f^* generates under colimits.

Example 2.2.2 (Base change). Suppose given a commutative square of derived algebraic stacks

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{g} & \mathcal{Y}' \\ \downarrow q & & \downarrow p \\ \mathcal{X} & \xrightarrow{f} & \mathcal{Y} \end{array}$$

where p is representable. If the square is homotopy cartesian, then the induced square

$$\begin{array}{ccc} \mathbf{D}(\mathcal{Y}) & \xrightarrow{f^*} & \mathbf{D}(\mathcal{X}) \\ \downarrow p^* & & \downarrow q^* \\ \mathbf{D}(\mathcal{Y}') & \xrightarrow{g^*} & \mathbf{D}(\mathcal{X}') \end{array}$$

satisfies base change by [SAG, Cor. 3.4.2.2]. In fact, this remains valid for non-algebraic stacks (i.e., for arbitrary derived prestacks, see [HLP14, Prop. A.1.5]). Note moreover that this condition is both sufficient and necessary for base change.

Affine Nisnevich squares give rise to categorical Milnor squares. More generally, we have:

Theorem 2.2.3 (Excision). *Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a representable morphism of derived algebraic stacks. Suppose there exists a closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ with quasi-compact open complement such that f is an isomorphism infinitely near \mathcal{Z} , i.e., the induced morphism $\mathcal{X}'_{f^{-1}(\mathcal{Z})} \rightarrow \mathcal{X}_{\mathcal{Z}}$ is*

invertible. Then the induced square

$$\begin{array}{ccc} \mathbf{D}(\mathcal{X}) & \xrightarrow{j^*} & \mathbf{D}(\mathcal{X} \setminus \mathcal{Z}) \\ \downarrow f^* & & \downarrow \\ \mathbf{D}(\mathcal{X}') & \longrightarrow & \mathbf{D}(\mathcal{X}' \setminus f^{-1}(\mathcal{Z})) \end{array}$$

is a cartesian square in Pres_c satisfying base change. If f is affine, then it is moreover a Milnor square.

Proof. Since open immersions are representable, the base change formula holds by Example 2.2.2 applied to the homotopy cartesian square

$$\begin{array}{ccc} \mathcal{X}' \setminus f^{-1}(\mathcal{Z}) & \xrightarrow{j'} & \mathcal{X}' \\ \downarrow f' & & \downarrow f \\ \mathcal{X} \setminus \mathcal{Z} & \xrightarrow{j} & \mathcal{X}. \end{array}$$

By Remark 2.2.1 it remains to show the cartesianness. For this, consider the adjunction

$$\mathbf{D}(\mathcal{X}) \rightleftarrows \mathbf{D}(\mathcal{X} \setminus \mathcal{Z}) \times_{\mathbf{D}(\mathcal{X}' \setminus f^{-1}(\mathcal{Z}))} \mathbf{D}(\mathcal{X}').$$

It will suffice to show that the unit and counit are invertible. We will make use of the canonical exact triangle of endofunctors of $\mathbf{D}(\mathcal{X})$

$$\hat{i}_\# \hat{i}^* \rightarrow \text{id} \rightarrow j_* j^* \tag{2.2.a}$$

where $\hat{i} : \mathcal{X}_Z^\wedge \hookrightarrow \mathcal{X}$ is the inclusion and $\hat{i}_\#$ is left adjoint to \hat{i}^* ; see [GR14, 7.1] or [HLP14, Thm. 2.2.3].

The unit evaluated on $\mathcal{F} \in \mathbf{D}(\mathcal{X})$ is the canonical morphism

$$\mathcal{F} \rightarrow f_* j^*(\mathcal{F}) \times_{g_* g^*(\mathcal{F})} j_* j^*(\mathcal{F}),$$

where $g : \mathcal{X}' \setminus f^{-1}(\mathcal{Z}) \rightarrow \mathcal{X}$. Using (2.2.a), it will suffice to show that it is invertible after applying either j^* or \hat{i}^* . By the base change formula for j_* and f_* (Example 2.2.2), the map becomes the identity of $j^*(\mathcal{F})$ in the former case and the unit $\hat{i}^*(\mathcal{F}) \rightarrow \hat{f}_* \hat{f}^* \hat{i}^*(\mathcal{F})$ in the latter, where $\hat{f} : \mathcal{X}'_{f^{-1}(\mathcal{Z})} \rightarrow \mathcal{X}_Z^\wedge$ is the base change (which is invertible by assumption).

For the counit, let $\mathcal{F}_U \in \mathbf{D}(\mathcal{X} \setminus \mathcal{Z})$, $\mathcal{F}_{X'} \in \mathbf{D}(\mathcal{X}')$, and $\mathcal{F}_{U'} \in \mathbf{D}(\mathcal{X}' \setminus f^{-1}(\mathcal{Z}))$ such that there are isomorphisms $f'^*(\mathcal{F}_U) \simeq \mathcal{F}_{U'} \simeq j'^*(\mathcal{F}_{X'})$. It will suffice to show that the canonical morphisms

$$\begin{aligned} j^*(j_* \mathcal{F}_U \times_{g_* \mathcal{F}_{U'}} f_* \mathcal{F}_{X'}) &\rightarrow \mathcal{F}_U \\ f^*(j_* \mathcal{F}_U \times_{g_* \mathcal{F}_{U'}} f_* \mathcal{F}_{X'}) &\rightarrow \mathcal{F}_{X'} \end{aligned}$$

are invertible. The first is straightforward using base change, and the second can be checked using (2.2.a) again (applied to \mathcal{X}' and \mathcal{Z}' this time instead of \mathcal{X} and \mathcal{Z}). \square

Specializing to étale neighbourhoods, we get:

Corollary 2.2.4 (Étale excision). *Suppose given a Nisnevich square of derived algebraic stacks of the form (2.1.a). Then the induced square*

$$\begin{array}{ccc} \mathbf{D}(\mathcal{X}) & \xrightarrow{j^*} & \mathbf{D}(\mathcal{U}) \\ \downarrow f^* & & \downarrow \\ \mathbf{D}(\mathcal{X}') & \longrightarrow & \mathbf{D}(\mathcal{U}') \end{array}$$

is cartesian and satisfies base change. In particular, if (2.1.a) is an affine Nisnevich square, then the above is a Milnor square satisfying base change.

In contrast with the case of Nisnevich squares, Milnor squares of stacks are not generally homotopy cartesian. Therefore the induced square of stable ∞ -categories does not usually satisfy base change. However, it will at least be precartesian in the following class of examples:

Theorem 2.2.5. *Suppose given a Ferrand pushout square of derived algebraic stacks*

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X}. \end{array} \tag{2.2.b}$$

That is, (2.2.b) is cocartesian, i' is a closed immersion, and g is affine. Then the induced square in Pres_c

$$\begin{array}{ccc} \mathbf{D}(\mathcal{X}) & \xrightarrow{i^*} & \mathbf{D}(\mathcal{Z}) \\ \downarrow f^* & & \downarrow g^* \\ \mathbf{D}(\mathcal{X}') & \xrightarrow{i'^*} & \mathbf{D}(\mathcal{Z}') \end{array}$$

is a Milnor square. It satisfies base change if and only if (2.2.b) is homotopy cartesian.

Proof. By Remark 2.2.1 it is enough to show precartesianness. By fpqc descent we immediately reduce to the case where \mathcal{X} is affine, hence so are \mathcal{X}' , \mathcal{Z} and \mathcal{Z}' . Consider the corresponding cartesian square of derived commutative rings

$$\begin{array}{ccc} \mathcal{O}_{\mathcal{X}} & \longrightarrow & \mathcal{O}_{\mathcal{Z}} \\ \downarrow & & \downarrow \\ \mathcal{O}_{\mathcal{X}'} & \longrightarrow & \mathcal{O}_{\mathcal{Z}'}. \end{array}$$

Since i is a closed immersion, the homomorphism of derived commutative rings $\mathcal{O}_{\mathcal{X}} \rightarrow \mathcal{O}_{\mathcal{Z}}$ is surjective on π_0 , and thus the square is also cartesian on underlying (nonconnective) \mathcal{E}_1 -rings. Hence the claim follows from [SAG, Thm. 16.2.0.2]. The last part follows from Example 2.2.2. \square

2.3. Formal completions of squares.

Construction 2.3.1. Suppose given a commutative square of derived algebraic stacks

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

where i and i' are closed immersions with quasi-compact open complements. Formally completing the horizontal arrows (see Subject. B.1) gives rise to a commutative square

$$\begin{array}{ccc} \mathcal{X}'_{\mathcal{Z}'} \hookrightarrow & \mathcal{X}' & \\ \downarrow f^\wedge & & \downarrow f \\ \mathcal{X}'_{\mathcal{Z}} \hookrightarrow & \mathcal{X} & \end{array} \quad (2.3.a)$$

If the original square is cartesian on underlying classical stacks as in Definition 2.1.3(i), then the formally completed square is homotopy cartesian (Remark B.1.2). In particular, this holds for cdh squares and Milnor squares.

We now show that the square (2.3.a) is also *cocartesian* when the original square is a finite cdh square or a Milnor square (under mild finiteness assumptions).

Lemma 2.3.2. *Suppose given a finite cdh square of bounded noetherian derived algebraic stacks of the form (2.1.c). Then the formally completed square (2.3.a) is cocartesian.*

Proof. Since \mathcal{X} is algebraic, it will suffice to show that the square is cocartesian after smooth base change to any affine derived scheme. Since the latter operation preserves cocartesian squares, we may assume \mathcal{X} is affine. Let $\mathcal{X} = \mathrm{Spec}(A)$, $\mathcal{X}' = \mathrm{Spec}(A')$, and choose $f_1, \dots, f_m \in \pi_0(A)$ such that $\mathcal{Z}_{\mathrm{cl}} = \mathrm{Spec}(\pi_0(A)/(f_1, \dots, f_m))$. Let $f'_1, \dots, f'_m \in \pi_0(A')$ denote their respective images. By [HA, 7.2.4.31], A' is almost of finite presentation over A . By Example B.1.4 it will suffice to show that the commutative squares

$$\begin{array}{ccc} A & \longrightarrow & A//\langle f_1^n, \dots, f_m^n \rangle \\ \downarrow & & \downarrow \\ A' & \longrightarrow & A'//\langle f_1'^n, \dots, f_m'^n \rangle \end{array}$$

induce a cartesian square of pro-derived commutative rings as n varies. For this it is enough that the underlying square of pro-spectra is cartesian, or equivalently that the induced morphism $F \rightarrow F'$ on horizontal homotopy fibres induces isomorphisms of pro-abelian groups $\pi_k(F) \rightarrow \pi_k(F')$ for all $k \geq 0$ (since the derived commutative rings in the square are all bounded above). By the five lemma it suffices to show that the right- and left-hand vertical arrows are invertible in the commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Coker}(\phi_{k+1}) & \longrightarrow & \pi_k(F) & \longrightarrow & \mathrm{Ker}(\phi_k) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathrm{Coker}(\phi'_{k+1}) & \longrightarrow & \pi_k(F') & \longrightarrow & \mathrm{Ker}(\phi'_k) \longrightarrow 0 \end{array}$$

where ϕ_k and ϕ'_k denote the induced morphisms $\{\pi_k(A)\} \rightarrow \{\pi_k(A//\langle f_1^n, \dots, f_m^n \rangle)\}_n$ and $\{\pi_k(A')\} \rightarrow \{\pi_k(A'//\langle f_1'^n, \dots, f_m'^n \rangle)\}_n$, respectively. By [SAG, Lem. 8.4.4.5] or [KST17, Lem. 4.10] the canonical morphisms

$$\begin{aligned} \{\pi_k(A//\langle f_1^n, \dots, f_m^n \rangle)\}_n &\rightarrow \{\pi_k(A)/\langle f_1^n, \dots, f_m^n \rangle\}_n, \\ \{\pi_k(A'//\langle f_1'^n, \dots, f_m'^n \rangle)\}_n &\rightarrow \{\pi_k(A')/ \langle f_1'^n, \dots, f_m'^n \rangle\}_n \end{aligned}$$

are invertible for all $k \geq 0$. In particular, the morphisms ϕ_k and ϕ'_k are all surjective, with kernels $\{(f_1^n, \dots, f_m^n) \cdot \pi_k(A)\}_n$ and $\{(f_1'^n, \dots, f_m'^n) \cdot \pi_k(A')\}_n$, respectively. Since A' is noetherian and the morphism $f : \mathrm{Spec}(A') \rightarrow \mathrm{Spec}(A)$ is finite, the homotopy groups $\pi_k(A')$ are finitely generated

over $\pi_0(A')$ and hence over $\pi_0(A)$. Now it is straightforward to check, using the assumption that $f : \mathrm{Spec}(A') \rightarrow \mathrm{Spec}(A)$ induces an isomorphism away from Z , that the canonical morphisms

$$\{(f_1^n, \dots, f_m^n) \cdot \pi_k(A)\}_n \rightarrow \{(f_1'^n, \dots, f_m'^n) \cdot \pi_k(A')\}_n,$$

are invertible. Indeed for each i , the morphism $\pi_k(A)[1/f_i] \rightarrow \pi_k(A')[1/f_i]$ is invertible by assumption. This implies in particular that there exists an m such that for all $m \geq n$, (f_1^n, \dots, f_m^n) annihilates the kernel (which is finitely generated by the noetherian assumption) and $(f_1^n, \dots, f_m^n) \cdot \pi_k(A) \rightarrow \{(f_1'^n, \dots, f_m'^n) \pi_k(A')\}$ is surjective (since $\pi_k(A')$ is finitely generated over $\pi_0(A)$). By the former claim and Artin-Rees lemma we can also ensure (by choosing a larger m if necessary) that the morphism is injective. The claim follows. \square

Corollary 2.3.3. *Let \mathcal{X} be a bounded noetherian derived algebraic stack. For any closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ with 0-truncated quasi-compact open complement $\mathcal{X} \setminus \mathcal{Z}$, the square*

$$\begin{array}{ccc} (\mathcal{X}_{\mathrm{cl}})_{\mathcal{Z}_{\mathrm{cl}}}^{\wedge} & \hookrightarrow & \mathcal{X}_{\mathrm{cl}} \\ \downarrow & & \downarrow \\ \mathcal{X}_{\mathcal{Z}}^{\wedge} & \hookrightarrow & \mathcal{X} \end{array}$$

is cocartesian, where the right-hand vertical arrow is the inclusion of the classical truncation.

Lemma 2.3.4. *Suppose given a Milnor square of noetherian algebraic stacks of the form (2.1.b). Then the formally completed square (2.3.a) is cocartesian.*

Proof. As in the proof of Lemma 2.3.2, we may assume that \mathcal{X} (and hence \mathcal{X}') is affine. Let $\mathcal{X} = \mathrm{Spec}(A)$, $\mathcal{X}' = \mathrm{Spec}(A')$, and $f_1, \dots, f_m \in A$ such that $\mathcal{Z} = \mathrm{Spec}(A/(f_1, \dots, f_m))$ and $\mathcal{Z}' = \mathrm{Spec}(A'/(f_1, \dots, f_m)A')$. By assumption, the square

$$\begin{array}{ccc} A & \longrightarrow & A/(f_1^n, \dots, f_m^n) \\ \downarrow & & \downarrow \\ A' & \longrightarrow & A'/(f_1^n, \dots, f_m^n) \end{array}$$

is cartesian for $n = 1$, i.e., $A \rightarrow A'$ sends the ideal (f_1, \dots, f_m) isomorphically onto $(f_1, \dots, f_m)A'$. Then the same holds for all $n > 0$ and hence by Example B.1.4 it follows that the square

$$\begin{array}{ccc} \mathrm{Spec}(A')^{\wedge} & \hookrightarrow & \mathrm{Spec}(A') \\ \downarrow & & \downarrow \\ \mathrm{Spec}(A)^{\wedge} & \hookrightarrow & \mathrm{Spec}(A) \end{array}$$

is cocartesian. \square

2.4. Formal Milnor and finite excision. In this subsection we prove the following result, which shows that if we pass to formal completions in Theorem 2.2.5, then we get a pro-Milnor square.

Theorem 2.4.1. *Suppose given a square of bounded noetherian derived algebraic stacks of the form*

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X}. \end{array}$$

Assume that i is a closed immersion, f is affine, the square is cartesian on classical truncations, and the formally completed square (2.3.a) is cocartesian. Then the induced commutative square in $\text{Pro}(\text{Pres}_c)$

$$\begin{array}{ccc} \{\mathbf{D}(\mathcal{X})\} & \xrightarrow{\hat{i}^*} & \widehat{\mathbf{D}}(\mathcal{X}^\wedge) \\ \downarrow f^* & & \downarrow (f^\wedge)^* \\ \{\mathbf{D}(\mathcal{X}')\} & \xrightarrow{\hat{i}'^*} & \widehat{\mathbf{D}}(\mathcal{X}'^\wedge). \end{array} \quad (2.4.a)$$

is a pro-Milnor square satisfying pro-base change.

Proof. By Construction 2.3.1 and the assumption, the square

$$\begin{array}{ccc} \mathcal{X}'_{\tilde{z}'} & \hookrightarrow & \mathcal{X}' \\ \downarrow f^\wedge & & \downarrow f \\ \mathcal{X}'_{\tilde{z}} & \hookrightarrow & \mathcal{X} \end{array}$$

is both cartesian and cocartesian. Thus by Remark B.1.3 it can be represented by either of the following filtered systems

$$\begin{array}{ccc} \tilde{\mathcal{Z}}' & \hookrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \tilde{\mathcal{Z}} & \hookrightarrow & \mathcal{X}, \end{array} \quad \begin{array}{ccc} \tilde{\mathcal{Z}}' & \hookrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \tilde{\mathcal{Z}} & \hookrightarrow & \tilde{\mathcal{Z}} \sqcup_{\tilde{\mathcal{Z}}} \mathcal{X}', \end{array}$$

indexed by $\tilde{\mathcal{Z}}$ as in Remark B.1.3, where $\tilde{\mathcal{Z}}' = \tilde{\mathcal{Z}} \times_{\mathcal{X}} \mathcal{X}'$. The left-hand squares are levelwise cartesian and the right-hand squares are levelwise cocartesian. Thus by Example 2.2.2 and Theorem 2.2.5 the induced square (2.4.a) can be represented alternatively by squares that satisfy levelwise base change or are levelwise precartesian. In particular, it is a pro-Milnor square. \square

Combining this with Lemmas 2.3.2 and 2.3.4 yields:

Corollary 2.4.2 (Formal Milnor excision). *Suppose given a Milnor square of noetherian algebraic stacks of the form (2.1.b). Then the induced commutative square in $\text{Pro}(\text{Pres}_c)$*

$$\begin{array}{ccc} \{\mathbf{D}(\mathcal{X})\} & \xrightarrow{\hat{i}^*} & \widehat{\mathbf{D}}(\mathcal{X}^\wedge) \\ \downarrow f^* & & \downarrow (f^\wedge)^* \\ \{\mathbf{D}(\mathcal{X}')\} & \xrightarrow{\hat{i}'^*} & \widehat{\mathbf{D}}(\mathcal{X}'^\wedge). \end{array}$$

is a pro-Milnor square satisfying pro-base change.

Corollary 2.4.3 (Formal finite excision). *Suppose given a finite cdh square of bounded noetherian derived algebraic stacks of the form (2.1.c). Then the induced commutative square in $\text{Pro}(\text{Pres}_c)$*

$$\begin{array}{ccc} \{\mathbf{D}(\mathcal{X})\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}'_{\tilde{z}}) \\ \downarrow & & \downarrow \\ \{\mathbf{D}(\mathcal{X}')\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}'_{\tilde{z}'}) \end{array}$$

is a pro-Milnor square satisfying pro-base change.

Corollary 2.4.4 (Formal nil-excision). *Let \mathcal{X} be a bounded noetherian derived algebraic stack with classical truncation \mathcal{X}_{cl} . Then for any closed immersion $\mathcal{Z} \hookrightarrow \mathcal{X}$ with 0-truncated quasi-compact open complement $\mathcal{X} \setminus \mathcal{Z}$, the induced commutative square in $\text{Pro}(\text{Pres}_{\mathbb{C}})$*

$$\begin{array}{ccc} \{\mathbf{D}(\mathcal{X})\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}_{\mathcal{Z}}^{\wedge}) \\ \downarrow & & \downarrow \\ \{\mathbf{D}(\mathcal{X}_{\text{cl}})\} & \longrightarrow & \widehat{\mathbf{D}}((\mathcal{X}_{\text{cl}})_{\mathcal{Z}_{\text{cl}}}^{\wedge}) \end{array}$$

is a pro-Milnor square satisfying pro-base change.

3. CATEGORICAL MILNOR EXCISION

In this section we prove categorical Milnor excision (Theorem A). We will first prove a characterization of Milnor squares in terms of the “ \ominus -construction” (see Theorem 3.3.1).

3.1. Adjustments of squares. Of particular interest for us will be the behaviour of Milnor squares as \mathcal{B}' is allowed to vary (especially among categories of non-geometric origin). For convenience, we refer to the operation of extending the square Δ along a colimit-preserving functor $a^* : \mathcal{B}' \rightarrow \mathcal{B}''$ as “adjustment”.

Definition 3.1.1. Let Δ be a commutative square in Pres of the form

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array} \quad (3.1.a)$$

Any choice of another presentable ∞ -category \mathcal{B}'' and a colimit-preserving functor $a^* : \mathcal{B}' \rightarrow \mathcal{B}''$ determines a new square Δ_a of the form

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q_a^* \\ \mathcal{A}' & \xrightarrow{g_a^*} & \mathcal{B}'' \end{array},$$

where $q_a^* = a^* \circ q^*$ and $g_a^* = a^* \circ g^*$. We call Δ_a the *adjustment of Δ by a^** .

Lemma 3.1.2. *Let Δ be a commutative square in Pres of the form (3.1.a). Let Δ_a be the adjustment of Δ by a colimit-preserving functor $a^* : \mathcal{B}' \rightarrow \mathcal{B}''$. Then the exchange transformations for the respective squares fit into a canonical commutative square*

$$\begin{array}{ccc} f^* p_* & \xrightarrow{\text{Ex}_{\Delta}} & q_* g^* \\ \parallel & & \downarrow \\ f^* p_* & \xrightarrow{\text{Ex}_{\Delta_a}} & q_{a,*} g_a^* \end{array}$$

where the vertical arrow is the morphism $q_* g^* \rightarrow q_{a,*} a^* g^* \simeq q_{a,*} g_a^*$ induced by the unit η_a .

Proof. The diagram in question can be subdivided as follows

$$\begin{array}{ccccccc}
 f^*p_* & \xrightarrow{\eta_q} & q_*q^*f^*p_* & \xrightarrow{\sim} & q_*g^*p^*p_* & \xrightarrow{\varepsilon_p} & q_*g^* \\
 \parallel & & \downarrow \eta_a & & \downarrow \eta_a & & \downarrow \eta_a \\
 f^*p_* & \xrightarrow{\eta_{aq}} & q_*a_*a^*q^*f^*p_* & \xrightarrow{\sim} & q_*a_*a^*g^*p^*p_* & \xrightarrow{\varepsilon_p} & q_*a_*a^*g^*
 \end{array}$$

where each square is tautologically commutative. \square

Corollary 3.1.3. *Let Δ be a commutative square of presentable stable ∞ -categories and colimit-preserving functors of the form (3.1.a). Suppose that Δ satisfies base change. For any presentable stable ∞ -category \mathcal{B}'' and colimit-preserving functor $a^* : \mathcal{B}' \rightarrow \mathcal{B}''$, consider the following conditions:*

- (i) *The functor $a^* : \mathcal{B}' \rightarrow \mathcal{B}''$ is fully faithful.*
- (ii) *The adjustment Δ_a satisfies base change.*

The implication (i) \implies (ii) always holds, and the converse holds if g^ and q^* generate \mathcal{B}' under colimits.*

Proof. Recall that the first condition is equivalent to invertibility of the unit $\eta_a : \text{id} \rightarrow a_*a^*$. Therefore the claim follows immediately from Lemma 3.1.2. \square

Lemma 3.1.4. *Let Δ be a commutative square in Pres of the form (3.1.a). If a colimit-preserving functor $a^* : \mathcal{B}' \rightarrow \mathcal{B}''$ induces monomorphisms on mapping spaces (e.g. it is fully faithful, or more generally if it is a monomorphism of ∞ -categories), the following conditions are equivalent:*

- (i) *The square Δ is precartesian.*
- (ii) *The adjustment Δ_a is precartesian.*

Proof. This follows from the following basic fact: given a diagram of spaces $X \rightarrow Z \leftarrow Y$ and a map $f : Z \rightarrow Z'$, the induced map

$$X \times_Z Y \rightarrow X \times_{Z'} Y$$

is invertible when $Z \rightarrow Z'$ is a monomorphism. \square

3.2. The \ominus -construction. In this subsection we introduce the main tool in our analysis of categorical Milnor squares (see Construction 3.2.5). This material closely follows [LT19], where it is developed in the special case of ∞ -categories of module spectra.

Fix a commutative square Δ in $\text{Pres}_{\mathcal{C}}$ of the form

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\
 \downarrow p^* & & \downarrow q^* \\
 \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}'
 \end{array}$$

We begin with the following preliminary construction.

Construction 3.2.1. Consider the ∞ -category \mathcal{C} of triples (X', Y, θ) , where $X' \in \mathcal{A}'$ and $Y \in \mathcal{B}$ are objects and $\theta : g^*(X') \rightarrow q^*(Y)$ is a morphism in \mathcal{B}' . A morphism $(X'_1, Y_1, \theta_1) \rightarrow (X'_2, Y_2, \theta_2)$

in \mathcal{C} is the data of a morphism $\alpha : X'_1 \rightarrow X'_2$ in \mathcal{A}' , a morphism $\beta : Y_1 \rightarrow Y_2$ in \mathcal{B} , and a commutative square

$$\begin{array}{ccc} g^*(X'_1) & \xrightarrow{g^*(\alpha)} & g^*(X'_2) \\ \downarrow \theta_1 & & \downarrow \theta_2 \\ q^*(Y_1) & \xrightarrow{q^*(\beta)} & q^*(Y_2) \end{array}$$

in \mathcal{B}' . More precisely, \mathcal{C} may be defined by the cartesian square

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \text{Fun}(\Delta^1, \mathcal{B}') \\ \downarrow & & \downarrow \\ \mathcal{A}' \times \mathcal{B} & \longrightarrow & \mathcal{B}' \times \mathcal{B}'. \end{array}$$

This construction is called the *lax pullback* or *oriented fibre product*. Note that the usual fibre product can be identified with the full subcategory of \mathcal{C} spanned by objects (X', Y, θ) such that θ is an isomorphism.

Remark 3.2.2. By construction, the ∞ -category \mathcal{C} is stable and presentable (see [Tam18, Lemma 8]). The full subcategory of \mathcal{C} spanned by objects of the form $(X', 0, 0)$ is canonically identified with \mathcal{A}' , and similarly \mathcal{B} is identified with the full subcategory spanned by objects of the form $(0, Y, 0)$. Every object $(X', Y, \theta) \in \mathcal{C}$ fits functorially into a cofibre sequence

$$(0, Y, 0) \rightarrow (X', Y, \theta) \rightarrow (X', 0, 0). \quad (3.2.a)$$

In fact, there is a semi-orthogonal decomposition $\mathcal{C} = \langle \mathcal{B}, \mathcal{A}' \rangle$, see [Tam18, Proposition 10].

Remark 3.2.3. Consider the functor $d_* : \mathcal{A} \rightarrow \mathcal{C}$ defined as the composite

$$(p^*, f^*) : \mathcal{A} \rightarrow \mathcal{A}' \times_{\mathcal{B}'} \mathcal{B} \subseteq \mathcal{C}.$$

whose essential image is spanned by objects of the form $(p^*(X), f^*(X), \theta_\Delta) \in \mathcal{C}$, where $X \in \mathcal{A}$ and $\theta_\Delta : g^*p^*(X) \simeq q^*f^*(X)$ is the isomorphism determined by the commutative square Δ . Note that d_* is fully faithful if and only if the square Δ is precartesian. Since d_* preserves colimits it admits a right adjoint² $d^! : \mathcal{C} \rightarrow \mathcal{A}$ whose value on any object $(X', Y, \theta) \in \mathcal{C}$ can be computed by the cartesian square

$$\begin{array}{ccc} d^!(X', Y, \theta) & \longrightarrow & f_*(Y) \\ \downarrow & & \downarrow \eta_q \\ & & f_*q_*q^*(Y) \\ & & \parallel \\ p_*(X') & \xrightarrow{\eta_g} & p_*g_*g^*(X') \xrightarrow{\theta} p_*g_*q^*(Y). \end{array}$$

Remark 3.2.4. For every object $X \in \mathcal{A}$, the cofibre sequence (3.2.a) may be evaluated at the object $d_*(X) = (p^*(X), f^*(X), \theta_\Delta) \in \mathcal{C}$ to get a cofibre sequence

$$(0, f^*(X), 0) \rightarrow d_*(X) \rightarrow (p^*(X), 0, 0)$$

with boundary map

$$\partial : (p^*(X), 0, 0) \rightarrow \Sigma(0, f^*(X), 0) \simeq (0, \Sigma f^*(X), 0).$$

²The departure from our usual notation (1.1.1) is because of the semi-orthogonal decomposition we will see below (Remark 3.2.7).

As the construction is functorial in X , ∂ defines a natural transformation of functors $\mathcal{A} \rightarrow \mathcal{C}$, up to which the square

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & \nearrow & \downarrow (0, \Sigma, 0) \\ \mathcal{A}' & \xrightarrow{(\text{id}, 0, 0)} & \mathcal{C}. \end{array} \quad (3.2.b)$$

commutes. Note that the adjustment of this square by the canonical colimit-preserving functor $c^* : \mathcal{C} \rightarrow \mathcal{B}'$, sending (X', Y, θ) to the fibre of θ , is canonically identified with our original square Δ .

The \odot -construction is obtained by forcing the square (3.2.b) to become commutative, which amounts to killing the essential image of $d_* : \mathcal{A} \rightarrow \mathcal{C}$.

Construction 3.2.5 (\odot -Construction). We let $\mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ \mathcal{B} denote the Verdier quotient of the presentable stable ∞ -category \mathcal{C} by the essential image of the functor $d_* : \mathcal{A} \rightarrow \mathcal{C}$. We write $a^* : \mathcal{C} \rightarrow \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ \mathcal{B} for the quotient functor. Thus we have a canonical commutative square Δ_0

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q_0^* \\ \mathcal{A}' & \xrightarrow{g_0^*} & \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}, \end{array} \quad (3.2.c)$$

where $g_0^* = a^*(\text{id}, 0, 0)$ and $q_0^* = \Sigma a^*(0, \text{id}, 0)$. Since the composite

$$\mathcal{A} \xrightarrow{d_*} \mathcal{C} \xrightarrow{c^*} \mathcal{B}'$$

is canonically null-homotopic, the colimit-preserving functor $c^* : \mathcal{C} \rightarrow \mathcal{B}'$ descends to a canonical colimit-preserving functor

$$b^* : \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B} \rightarrow \mathcal{B}'.$$

By construction, the adjustment of Δ_0 by b^* is canonically identified with Δ .

Remark 3.2.6. Upon application of the quotient functor $a^* : \mathcal{C} \rightarrow \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ \mathcal{B} , the cofibre sequences of Remark 3.2.2 induce cofibre sequences of the form

$$\Omega q_0^*(Y) \rightarrow a^*(X', Y, \theta) \rightarrow g_0^*(X')$$

for every object $(X', Y, \theta) \in \mathcal{C}$.

Remark 3.2.7. By construction, we have a semi-orthogonal decomposition $\mathcal{C} = \langle \mathcal{A}, \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ $\mathcal{B} \rangle$. In particular there is a cofibre sequence

$$d_* d^! \xrightarrow{\varepsilon_d} \text{id} \xrightarrow{\eta_a} a_* a^*$$

of natural transformations $\mathcal{C} \rightarrow \mathcal{C}$.

Lemma 3.2.8.

- (i) If $p^* : \mathcal{A} \rightarrow \mathcal{A}'$ generates its codomain under colimits, then so does $q_0^* : \mathcal{B} \rightarrow \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ \mathcal{B} .
- (ii) If $f^* : \mathcal{A} \rightarrow \mathcal{B}$ generates its codomain under colimits, then so does $g_0^* : \mathcal{A}' \rightarrow \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ \mathcal{B} .
- (iii) If $q^* : \mathcal{B} \rightarrow \mathcal{B}'$ generates its codomain under colimits, then so does $b^* : \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'}$ $\mathcal{B} \rightarrow \mathcal{B}'$.
- (iv) If p^* , f^* and $q^* f^* \simeq g^* p^*$ are compact, then so is a^* .
- (v) If q^* is compact, then so is c^* .

- (vi) If f^* , p^* , and q^* are compact, then so is b^* .
- (vii) If p^* and f^* are compact, then so are q_0^* and g_0^* .

Proof.

- (i) If the essential image of p^* generates \mathcal{A}' under colimits, then it follows from Remark 3.2.2 that \mathcal{C} is generated under colimits by objects of the form $(p^*(X), 0, 0)$ and $(0, Y, 0)$ as X and Y vary among objects of \mathcal{A} and \mathcal{B} , respectively. In particular, the Verdier quotient $\mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}$ is generated under colimits by the images of such objects under the quotient functor $a^* : \mathcal{C} \rightarrow \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}$, i.e., by objects of the forms $g_0^* p^*(X)$ and $q_0^*(Y)$. But the commutativity of Δ_0 (3.2.c) provides isomorphisms

$$g_0^* p^*(X) \simeq q_0^* f^*(X)$$

in $\mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}$ for every $X \in \mathcal{A}$.

- (ii) Similar to (i).
- (iii) By assumption, \mathcal{B}' is generated under colimits by the essential image of $q^* \simeq b^* q_0^*$. Hence b^* also generates \mathcal{B}' under colimits.
- (iv) Since a^* generates under colimits, it will suffice to show that $a_* a^*$ preserves filtered colimits. Using the cofibre sequence of Remark 3.2.7, we reduce to showing that d^1 preserves filtered colimits. This follows immediately from the description of Remark 3.2.3 (in view of the assumptions).
- (v) Note that the right adjoint $c_* : \mathcal{B}' \rightarrow \mathcal{C}$ is given by the formula $Y' \mapsto (0, \Sigma q_*(Y'), 0)$.
- (vi) Recall that $a_* : \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B} \rightarrow \mathcal{C}$ is conservative (as a^* is essentially surjective) and colimit-preserving by (iii). Therefore b_* preserves colimits if and only if $a_* b_* \simeq c_*$ does. This is true under the assumptions by (iv).
- (vii) By the definition of q_0^* and g_0^* , this follows immediately from (iii).

□

3.3. Characterization of Milnor squares. The following result shows that every Milnor square Δ satisfying base change is isomorphic to one coming from the \odot -construction. More precisely, it is an adjustment of the square Δ_0 (3.2.c) along the canonical functor $b^* : \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B} \rightarrow \mathcal{B}'$, which is an equivalence.

Theorem 3.3.1. *Let Δ be a Milnor square of presentable stable ∞ -categories of the form*

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}'. \end{array}$$

Then we have:

- (i) The square Δ_0 (3.2.c) satisfies base change.
- (ii) If Δ satisfies base change, then the canonical functor

$$b^* : \mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B} \rightarrow \mathcal{B}'$$

is an equivalence. In particular, there is an isomorphism of squares $\Delta \simeq \Delta_0$.

Lemma 3.3.2. *Let Δ be a commutative square of presentable stable ∞ -categories and colimit-preserving functors as above. Let ϕ_{q_0} denote the fibre of the unit η_{q_0} , and similarly for ϕ_q . Then there is a canonical isomorphism $f^* f_* \phi_q \simeq \phi_{q_0}$ such that the diagram*

$$\begin{array}{ccc} f^* f_* \phi_q & \xrightarrow{f^* f_* \text{can}} & f^* f_* \\ \parallel & & \downarrow \varepsilon_f \\ \phi_{q_0} & \xrightarrow{\text{can}} & \text{id} \end{array}$$

commutes, where can denotes the canonical inclusion of the respective fibre.

Proof. Precomposing the cofibre sequence of Remark 3.2.7 with the inclusion $(0, \text{id}, 0) : \mathcal{B} \rightarrow \mathcal{C}$ and projecting the result back to \mathcal{B} yields the cofibre sequence

$$f^* d^l(0, \text{id}, 0) \rightarrow \text{id} \xrightarrow{\eta_{q_0}} q_{0,*} q_0^*$$

of natural transformations $\mathcal{B} \rightarrow \mathcal{B}$. Now apply the description of d^l given in Remark 3.2.3. \square

Lemma 3.3.3. *Let Δ be a commutative square of presentable stable ∞ -categories and colimit-preserving functors as above. If Δ is precartesian, then the exchange transformation for Δ_0 induces an isomorphism*

$$f^* p_* p^* \xrightarrow{\text{Ex}_{\Delta_0} p^*} q_{0,*} g_0^* p^* \simeq q_{0,*} q_0^* f^*.$$

In particular, if Δ is a Milnor square, then Δ_0 satisfies base change.

Proof. Note that the second claim follows from the first in view of the fact that p^* generates \mathcal{A}' under colimits and p^* and q_0^* are compact (Lemma 3.2.8). For the main claim, by construction of the exchange transformation, we have a commutative square

$$\begin{array}{ccc} f^* & \xrightarrow{\eta_p} & f^* p_* p^* \\ \parallel & & \downarrow \\ f^* & \xrightarrow{\eta_{q_0}} & q_{0,*} q_0^* f^* \end{array} \quad (3.3.a)$$

Taking fibres horizontally, we obtain by Lemma 3.3.2 the following commutative diagram:

$$\begin{array}{ccccc} f^* \phi_p & \xrightarrow{\text{can}} & f^* & \xrightarrow{\eta_p} & f^* p_* p^* \\ \downarrow & & \downarrow f^* \eta_f & & \downarrow \\ f^* f_* \phi_q f^* & \xrightarrow{\text{can}} & f^* f_* f^* & & \\ \parallel & & \downarrow \varepsilon_f f^* & & \downarrow \\ \phi_{q_0} f^* & \xrightarrow{\text{can}} & f^* & \xrightarrow{\eta_{q_0}} & q_{0,*} q_0^* f^* \end{array}$$

The upper vertical left-hand arrow is invertible as soon as Δ is precartesian, i.e., when the right-hand square below is cartesian:

$$\begin{array}{ccccc} \phi_p & \xrightarrow{\text{can}} & \text{id} & \xrightarrow{\eta_p} & p_* p^* \\ \downarrow & & \downarrow \eta_f & & \downarrow \eta_g \\ f_* \phi_q f^* & \xrightarrow{\text{can}} & f_* f^* & \xrightarrow{\eta_q} & f_* q_* q^* f^* \simeq p_* g_* g^* p^* \end{array}$$

Thus in that case we find that (3.3.a) is cartesian. \square

Proof of Theorem 3.3.1. The first claim follows from Lemma 3.3.3. For the second, assume that Δ satisfies base change. Since Δ is the adjustment of Δ_0 by b^* , and since b^* is compact by Lemma 3.2.8, this implies by Corollary 3.1.3 that b^* is fully faithful. By Lemma 3.2.8, b^* generates \mathcal{B}' under colimits. Since $\mathcal{A}' \circlearrowleft_{\mathcal{A}}^{\mathcal{B}'}$ \mathcal{B} is cocomplete, it now follows that b^* is an equivalence. \square

3.4. Categorical Milnor excision. Let E be a functor on the ∞ -category of small stable ∞ -categories with values in a stable ∞ -category. Recall that E is called a *localizing invariant* if it sends exact sequences to exact triangles. See e.g. [BGT13], except that we do not assume that E commutes with filtered colimits.

Example 3.4.1. The main example we have in mind is nonconnective algebraic K-theory, as defined in [BGT13, Sect. 9] or via the generalized Bass–Thomason–Trobaugh construction as in [CK20, Thm. C]. We denote it simply by K .

Notation 3.4.2. Let E be a localizing invariant. For any compactly generated stable ∞ -category \mathcal{C} , we set for convenience

$$E(\mathcal{C}) := E(\mathcal{C}^\omega)$$

where \mathcal{C}^ω denotes the full subcategory of compact objects.

Theorem 3.4.3. *Let E be a localizing invariant. Let Δ be a Milnor square of compactly generated stable ∞ -categories of the form*

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array}$$

If Δ satisfies base change, then the induced square $E(\Delta)$

$$\begin{array}{ccc} E(\mathcal{A}) & \xrightarrow{f^*} & E(\mathcal{B}) \\ \downarrow p^* & & \downarrow q^* \\ E(\mathcal{A}') & \xrightarrow{g^*} & E(\mathcal{B}') \end{array}$$

is cartesian.

Theorem 3.4.3 immediately follows, in view of Theorem 3.3.1, from the following statement:

Proposition 3.4.4. *Let the notation be as in Theorem 3.4.3. Then the induced square $E(\Delta_0)$*

$$\begin{array}{ccc} E(\mathcal{A}) & \xrightarrow{f^*} & E(\mathcal{B}) \\ \downarrow p^* & & \downarrow q^* \\ E(\mathcal{A}') & \xrightarrow{g_0^*} & E(\mathcal{A}' \circlearrowleft_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}) \end{array}$$

is cartesian.

Proof. First note that the Verdier localization sequence (see Remark 3.2.7)

$$\mathcal{A} \xrightarrow{d_*} \mathcal{C} \xrightarrow{a^*} \mathcal{A}' \circlearrowleft_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}$$

induces a cofibre sequence

$$E(\mathcal{A}) \xrightarrow{d_*} E(\mathcal{C}) \xrightarrow{a^*} E(\mathcal{A}' \circlearrowleft_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}).$$

Similarly, E sends the Verdier localization sequence $\mathcal{B} \rightarrow \mathcal{C} \rightarrow \mathcal{A}'$ (Remark 3.2.2) to a cofibre sequence. Moreover, both functors in this sequence have *compact* right adjoints which hence yield a splitting

$$E(\mathcal{C}) \simeq E(\mathcal{A}') \oplus E(\mathcal{B})$$

given by the projections $\mathcal{C} \rightarrow \mathcal{A}'$ and $\mathcal{C} \rightarrow \mathcal{B}$. Under this isomorphism, the map $d_* : E(\mathcal{A}) \rightarrow E(\mathcal{C})$ is induced by the maps $p^* : E(\mathcal{A}) \rightarrow E(\mathcal{A}')$ and $f^* : E(\mathcal{A}) \rightarrow E(\mathcal{B})$, and the map $a^* : E(\mathcal{C}) \rightarrow E(\mathcal{A}' \circledast_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B})$ splits thanks to Remark 3.2.6 as

$$a^* : E(\mathcal{C}) \simeq E(\mathcal{A}') \oplus E(\mathcal{B}) \xrightarrow{g_0^* - q_0^*} E(\mathcal{A}' \circledast_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}).$$

Thus we end up with a cofibre sequence of the form

$$E(\mathcal{A}) \xrightarrow{(p^*, f^*)} E(\mathcal{A}') \oplus E(\mathcal{B}) \xrightarrow{g_0^* - q_0^*} E(\mathcal{A}' \circledast_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}).$$

Recall that this is equivalent to the assertion that $E(\Delta_0)$ is cartesian. \square

3.5. Categorical pro-Milnor excision. We now turn our attention to the behaviour of localizing invariants with respect to pro-Milnor squares (Definition 1.4.1). Under some strong assumptions, we will be able to show a pro-variant of Theorem 3.4.3 (see Theorem 3.5.11).

We begin with some preliminary considerations on isomorphisms of pro-systems in $\text{Pro}(\text{Pres}_{\mathcal{C}})$.

Definition 3.5.1. Let $\{f_n^* : \mathcal{C}_n \rightarrow \mathcal{D}_n\}_n$ be a cofiltered system in $\text{Pres}_{\mathcal{C}}$. We say that $\{f_n^*\}_n$ is a *pro-equivalence* if it induces an isomorphism in $\text{Pro}(\text{Pres}_{\mathcal{C}})$.

While a general criterion for pro-equivalences seems out of reach, we will show that there is a large enough supply for our purposes. First note the following class of examples:

Example 3.5.2. Let $\{\phi_n : A_n \rightarrow B_n\}_n$ be a cofiltered system of homomorphisms of connective bounded above \mathcal{E}_1 -rings. If $\{\phi_n\}_n$ induces an isomorphism of underlying pro-spaces, then it follows from [LT19, Lem. 2.28] that it induces an isomorphism of pro-objects in the ∞ -category of \mathcal{E}_1 -rings. In particular, by functoriality of the construction $R \mapsto \text{LMod}_R$ (as a functor from the ∞ -category of \mathcal{E}_1 -rings to $\text{Pres}_{\mathcal{C}}$), we deduce that the extension of scalars functors

$$\{\phi_n^* : \text{LMod}_{A_n} \rightarrow \text{LMod}_{B_n}\}_n$$

define a pro-equivalence.

Example 3.5.3. Since passage to underlying pro-objects preserves finite limits and colimits, it follows that pro-equivalences are closed under finite (co)limits.

We also have closure under certain colocalizations (and a dual statement for localizations that we leave to the reader to formulate):

Lemma 3.5.4. *Suppose given a cofiltered system of commutative squares in $\text{Pres}_{\mathcal{C}}$*

$$\begin{array}{ccc} \mathcal{C}'_n & \xrightarrow{i_{n,*}} & \mathcal{C}_n \\ g_n^* \downarrow & & \downarrow f_n^* \\ \mathcal{D}'_n & \xrightarrow{i_{n,*}} & \mathcal{D}_n \end{array}$$

where $i_{n,*}$ is fully faithful for every n , with right adjoint $i_n^!$. Assume that for every n , the square is horizontally right-adjointable: that is, the base change transformation

$$g_n^* i_n^! \rightarrow i_n^! f_n^*$$

is invertible. If f^* is a pro-equivalence, then so is g^* .

Proof. By assumption, there exists a morphism $v^* : \{\mathcal{D}_n\}_n \rightarrow \{\mathcal{C}_n\}_n$ in $\text{Pro}(\text{Pres}_c)$ which is inverse to $f^* = \{f_n^*\}_n$. Let $u^* : \{\mathcal{D}'_n\}_n \rightarrow \{\mathcal{C}'_n\}_n$ be the morphism in $\text{Pro}(\text{Pres})$ defined as the composite $u^* = i^! v^* i_*$. More explicitly, we may represent v^* by the following data:

- (*) For every index n , there exists an index $n' > n$, a compact colimit-preserving functor $v_n^* : \mathcal{D}_{n'} \rightarrow \mathcal{C}_n$, isomorphisms

$$f_n^* v_n^* \simeq \text{tr}^* : \mathcal{D}_{n'} \rightarrow \mathcal{D}_n, \quad v_n^* f_{n'}^* \simeq \text{tr}^* : \mathcal{C}_{n'} \rightarrow \mathcal{C}_n,$$

and a homotopy coherent system of compatibilities between this data, where we write tr^* for the transition functors.

In terms of such a choice, the morphism u^* is represented by the colimit-preserving functors

$$u_n^* : \mathcal{D}'_{n'} \xrightarrow{i_*} \mathcal{D}_{n'} \xrightarrow{v_n^*} \mathcal{C}_n \xrightarrow{i^!} \mathcal{C}'_n$$

where we omit some decorations for simplicity. Note that we have canonical isomorphisms of functors

$$\begin{aligned} g_n^* u_n^* &\simeq g_n^* i^! v_n^* i_* \simeq i^! f_n^* v_n^* i_* \simeq i^! \text{tr}^* i_* \simeq i^! i_* \text{tr}^* \simeq \text{tr}^* \\ u_n^* g_{n'}^* &\simeq i^! v_n^* i_* g_{n'}^* \simeq i^! v_n^* f_{n'}^* i_* \simeq i^! \text{tr}^* i_* \simeq i^! i_* \text{tr}^* \simeq \text{tr}^*, \end{aligned}$$

where we have used the assumptions that i_* and $i^!$ commute with the vertical arrows. Note moreover that u_n^* is compact: since $g_{n'}^*$ generates under colimits, this follows from the fact that $u_n^* g_{n'}^* \simeq \text{tr}^*$ is compact. Thus the morphism u^* is contained in the subcategory $\text{Pro}(\text{Pres}_c)$ and defines an inverse to $g^* = \{g_n^*\}_n$ in the latter. \square

In the *projectively* generated case (see [HTT, Def. 5.5.8.23]), and when the mapping pro-spaces are uniformly bounded, we can prove the following criterion for pro-equivalence.

Lemma 3.5.5. *Let $\{\mathcal{C}_n\}_n$ and $\{\mathcal{D}_n\}_n$ be cofiltered systems in Pres_c and let $\{f_n^* : \mathcal{C}_n \rightarrow \mathcal{D}_n\}_n$ be a cofiltered system of colimit-preserving compact projective³ functors which generate their codomains under colimits. Consider the following conditions:*

- (i) For every n there exists a small set of compact projective generators $\{C_n^\alpha\}_\alpha$ of \mathcal{C}_n such that every transition functor $\mathcal{C}_n \rightarrow \mathcal{C}_m$ sends $C_n^\alpha \mapsto C_m^\alpha$ for every α and every $m < n$.
- (ii) For every pair of indices α and β , the map

$$\{\text{Maps}_{\mathcal{C}_n}(C_n^\alpha, C_n^\beta)\}_n \rightarrow \{\text{Maps}_{\mathcal{D}_n}(f_n^*(C_n^\alpha), f_n^*(C_n^\beta))\}_n$$

is an isomorphism of pro-spaces.

- (iii) There exists an integer $c \geq 0$ such that the pro-spaces

$$\{\text{Maps}_{\mathcal{C}_n}(C_n^\alpha, C_n^\beta)\}_n$$

are c -truncated for all α and β .

Then $\{f_n^*\}_n$ induces a pro-equivalence.

Proof. For every n , consider the full subcategory $\mathcal{A}_n \subseteq \mathcal{C}_n$ spanned by finite direct sums of the objects C_n^α (as α varies). Then \mathcal{A}_n is an additive ∞ -category for which the inclusion $\mathcal{A}_n \subseteq \mathcal{C}_n$ extends to an equivalence $\mathcal{P}_\Sigma(\mathcal{A}_n) \simeq \mathcal{C}_n$ by [HTT, Prop. 5.5.8.25]. Let $C_n \in \mathcal{C}_n$ be the direct sum of the objects C_n^α (as α varies), and let $\mathcal{A}_n^+ \subseteq \mathcal{C}_n$ be the full subcategory generated by C_n

³A functor $f^* : \mathcal{C} \rightarrow \mathcal{D}$ is called compact projective if its right adjoint preserves sifted colimits.

under finite direct sums and direct summands. By [SAG, Ex. C.1.5.11] this is an idempotent-complete additive ∞ -category equipped with an equivalence $\mathcal{P}_\Sigma(\mathcal{A}_n^+) \simeq \mathrm{LMod}_{A_n}^{\mathrm{cn}}$, where A_n is the connective endomorphism algebra of C_n (as in [SAG, Rem. C.2.1.9]). Then we have inclusions $\mathcal{A}_n \subseteq \mathcal{A}_n^+$ which are closed under finite direct sums and hence give rise by [HTT, Prop. 5.5.8.15] to fully faithful colimit-preserving functors

$$i_* : \mathcal{C}_n \simeq \mathcal{P}_\Sigma(\mathcal{A}_n) \rightarrow \mathcal{P}_\Sigma(\mathcal{A}_n^+) \quad (3.5.a)$$

whose right adjoints $i^!$ are restriction along $\mathcal{A}_n \subseteq \mathcal{A}_n^+$. Similarly let $D_n^\alpha \in \mathcal{D}_n$ be the images of C_n^α , $\mathcal{B}_n \subseteq \mathcal{D}_n$ the additive subcategory they generate, $D_n \in \mathcal{D}_n$ their direct sum over α , and B_n the connective endomorphism algebra of D_n . Since the functors f_n^* are compact projective and generate under colimits, the objects D_n^α form a small set of compact projective generators of \mathcal{D}_n . Thus we similarly get functors $i_* : \mathcal{D}_n \rightarrow \mathrm{LMod}_{B_n}^{\mathrm{cn}}$ fitting into commutative squares

$$\begin{array}{ccc} \mathcal{C}_n & \xrightarrow{i_*} & \mathrm{LMod}_{A_n}^{\mathrm{cn}} \\ \downarrow f_n^* & & \downarrow \phi_n^* \\ \mathcal{D}_n & \xrightarrow{i_*} & \mathrm{LMod}_{B_n}^{\mathrm{cn}}, \end{array}$$

where ϕ_n^* is extension of scalars along the homomorphism $\phi_n : A_n \rightarrow B_n$ induced by f_n^* . Since $\{A_n\}_n$ and $\{B_n\}_n$ are bounded above by assumption, we see by Example 3.5.2 that the functors ϕ_n^* induce a pro-equivalence. To conclude it remains to check the criteria of Lemma 3.5.4 for the above square.

The functors (3.5.a) preserve compact projective objects by construction, so we see that i_* is compact projective. It remains to show that $i^!$ commutes with the vertical arrows. Since all functors in the square commute with sifted colimits, it will suffice to evaluate on the object $A_n \in \mathrm{LMod}_{A_n}^{\mathrm{cn}}$. By construction, the horizontal arrows send

$$\begin{aligned} A_n &\mapsto C_n, \\ B_n &\mapsto D_n, \end{aligned}$$

whence the claim. Thus the conditions of Lemma 3.5.4 are satisfied, and $\{f_n^*\}_n$ is a pro-equivalence. \square

Under the conditions of Lemma 3.5.5, the ∞ -categories \mathcal{C}_n and \mathcal{D}_n will never be stable, but only prestable. Nevertheless, by stabilization, we may deduce a following variant of Lemma 3.5.5 for certain presentable stable ∞ -categories.

Definition 3.5.6. Let $\{\mathcal{C}_i\}_{i \in \mathcal{J}}$ be a diagram of presentable stable ∞ -categories and compact colimit-preserving functors indexed by a small ∞ -category \mathcal{J} . Suppose given full subcategories $(\mathcal{C}_i)_{\geq 0} \subseteq \mathcal{C}_i$ for every $i \in \mathcal{J}$, closed under colimits and extensions, and collections $\{C_i^\alpha\}_{\alpha \in S}$ of compact projective objects of $(\mathcal{C}_i)_{\geq 0}$ indexed by some small set S . We will say that the collections $\{C_i^\alpha\}_\alpha$ form a set of *projective generators* for $\{\mathcal{C}_i\}_i$ if the following conditions hold:

- (i) For every $i \in \mathcal{J}$, the objects $\{C_i^\alpha\}_{\alpha \in S}$ generate $(\mathcal{C}_i)_{\geq 0}$ under colimits and extensions, and the objects $\{\Sigma^{-n}(C_i^\alpha)\}_{n \geq 0, \alpha}$ generate \mathcal{C}_i under colimits.
- (ii) For every morphism $i \rightarrow j$ in \mathcal{J} , the induced functor $u_{i,j}^* : \mathcal{C}_i \rightarrow \mathcal{C}_j$ sends $C_i^\alpha \mapsto C_j^\alpha$ for every $\alpha \in S$, and its right adjoint $u_{i,j,*}$ sends $(\mathcal{C}_j)_{\geq 0}$ to $(\mathcal{C}_i)_{\geq 0}$.

Remark 3.5.7. For $\mathcal{J} = \Delta^0$, Definition 3.5.6 is essentially a presentable version of the notion of weighted ∞ -category as discussed in Subsect. 5.1. For $\mathcal{J} = \Delta^1$, it corresponds to a weight-exact functor which generates under colimits.

Remark 3.5.8. Using [HA, Prop. 1.4.4.11, Lem. 7.2.2.6] we can recast Definition 3.5.6 in terms of t-structures: there are t-structures on the \mathcal{C}_i —accessible, right-complete, and compatible with filtered colimits—with connective parts $(\mathcal{C}_i)_{\geq 0}$ generated under colimits and extensions by $\{C_i^\alpha\}_\alpha$, such that the functors $u_{i,j}^*$ and their right adjoints are right t-exact.

Remark 3.5.9. In case \mathcal{J} has an initial object 0, we will only specify the generators $C_0^\alpha \in \mathcal{C}_0$, so that $C_i^\alpha \in \mathcal{C}_i$ are implicitly defined as the images by the functor $\mathcal{C}_0 \rightarrow \mathcal{C}_i$ for every $i \in \mathcal{J}$. If we have a cofiltered system of \mathcal{J} -indexed diagrams $\{\mathcal{C}_{n,i}\}_{n \in \Lambda, i \in \mathcal{J}}$, where Λ is cofiltered, then projective generators of $\{\mathcal{C}_{n,i}\}_{n,i}$ (as a $\Lambda \times \mathcal{J}$ -indexed diagram) will be similarly specified by a small set S and a collection of objects $\{C_{n,0}^\alpha\}_{\alpha \in S}$.

We can now reformulate Lemma 3.5.5 as follows:

Corollary 3.5.10. *Suppose given a cofiltered system $\{f_n^* : \mathcal{C}_n \rightarrow \mathcal{D}_n\}_n$ of presentable stable ∞ -categories and compact colimit-preserving functors. Suppose there exists a small set S and objects $\{C_n^\alpha\}_{\alpha \in S}$ of \mathcal{C}_n for every n , which projectively generate $\{f_n^*\}_n$ as a cofiltered system of Δ^1 -indexed diagrams. Then $\{f_n^*\}_n$ is a pro-equivalence in case the following conditions hold:*

- (i) For every pair α, β , the map

$$\{\mathrm{Maps}_{\mathcal{C}_n}(C_n^\alpha, C_n^\beta)\}_n \rightarrow \{\mathrm{Maps}_{\mathcal{D}_n}(f_n^*(C_n^\alpha), f_n^*(C_n^\beta))\}_n$$

is an isomorphism of pro-spaces.

- (ii) There exists an integer $c \geq 0$ such that the mapping pro-spaces

$$\{\mathrm{Maps}_{\mathcal{C}_n}(C_n^\alpha, C_n^\beta)\}_n$$

are c -truncated for all α, β .

We now prove our categorical pro-excision statement.

Theorem 3.5.11. *Let E be a localizing invariant. Let Δ be a pro-Milnor square of presentable stable ∞ -categories which can be represented by a cofiltered system $\{\Delta_n\}_n$ of levelwise Milnor squares*

$$\begin{array}{ccc} \mathcal{A}_n & \xrightarrow{f_n^*} & \mathcal{B}_n \\ \downarrow p_n^* & & \downarrow q_n^* \\ \mathcal{A}'_n & \xrightarrow{g_n^*} & \mathcal{B}'_n \end{array}$$

satisfying the following conditions:

- (i) There exists a small set S and objects $\{A_n^\alpha\}_{\alpha \in S}$ of \mathcal{A}_n for every n , which projectively generate $\{\Delta_n\}_n$ as a cofiltered system of $(\Delta^1 \times \Delta^1)$ -indexed diagrams.
- (ii) There is an integer $c \geq 0$ such that for all α, β , the mapping pro-space

$$\{\mathrm{Maps}_{\mathcal{B}'_n}(g_n^* p_n^*(A_n^\alpha), g_n^* p_n^*(A_n^\beta))\}_n$$

is c -truncated.

If Δ satisfies pro-base change, then the induced square of pro-objects

$$\begin{array}{ccc} \{E(\mathcal{A}_n)\}_n & \xrightarrow{f_n^*} & \{E(\mathcal{B}_n)\}_n \\ \downarrow p_n^* & & \downarrow q_n^* \\ \{E(\mathcal{A}'_n)\}_n & \xrightarrow{g_n^*} & \{E(\mathcal{B}'_n)\}_n \end{array}$$

is cartesian.

We will need the following lemma.

Lemma 3.5.12. *Suppose given a precartesian square of presentable stable ∞ -categories and compact colimit-preserving functors*

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f^*} & \mathcal{B} \\ \downarrow p^* & & \downarrow q^* \\ \mathcal{A}' & \xrightarrow{g^*} & \mathcal{B}' \end{array}$$

Assume that the square is projectively generated by a collection $\{A^\alpha\}_\alpha$ of objects of \mathcal{A} . Then the functors $q_0^* : \mathcal{B} \rightarrow \mathcal{A}' \circlearrowleft_{\mathcal{A}'}^{\mathcal{B}'} \mathcal{B}$ and $b^* : \mathcal{A}' \circlearrowleft_{\mathcal{A}'}^{\mathcal{B}'} \mathcal{B} \rightarrow \mathcal{B}'$ are both projectively generated (as Δ^1 -indexed diagrams). In particular, the collection $\{q_0^* f^*(A^\alpha)\}_\alpha$ forms a set of projective generators of the \circlearrowleft -construction $\mathcal{A}' \circlearrowleft_{\mathcal{A}'}^{\mathcal{B}'} \mathcal{B}$.

Proof. We consider the t-structures defined in Remark 3.5.8. To simplify the notation set $\mathcal{Q} := \mathcal{A}' \circlearrowleft_{\mathcal{A}'}^{\mathcal{B}'} \mathcal{B}$. By [HA, Prop. 1.4.4.11] there is a t-structure whose connective part $\mathcal{Q}_{\geq 0}$ is the cocomplete stable subcategory generated by $q_0^*(\mathcal{B}_{\geq 0})$.

It follows from the assumption that each of the functors in the given square generates under colimits, and is compact projective (the latter by [HA, Lem. 7.2.2.6], since it has right t-exact right adjoint by assumption). Hence by Lemma 3.2.8 the functors q_0^* and b^* also generate under colimits and are compact. It remains to show that are right t-exact, and have right t-exact right adjoints. The functor q_0^* is right t-exact by construction. Let us show $q_{0,*}$ is right t-exact. Since $q_0^* f^*$ generates under colimits and is right t-exact, it will suffice to show that $q_{0,*} q_0^* f^*$ is right t-exact. Since Δ is precartesian, Lemma 3.3.3 implies that the latter functor is identified with the right t-exact functor $f^* p_* p^*$.

Since q_0^* generates under colimits and $b^* q_0^* \simeq q^*$ is right t-exact, it follows that $b^* : \mathcal{Q} \rightarrow \mathcal{B}'$ is also right t-exact. Finally, using the fact that $q_{0,*}$ is conservative and t-exact, it follows from the fact that $q_{0,*} b_* \simeq q_*$ is right t-exact that b_* itself is right t-exact. \square

The following key lemma shows that any pro-Milnor square as in Theorem 3.5.11 can be levelwise represented by squares coming from the \circlearrowleft -construction.

Lemma 3.5.13. *Let Δ be a pro-Milnor square as in Theorem 3.5.11. If Δ satisfies pro-base change, then there exists an isomorphism*

$$\{\Delta_{n,0}\}_n \rightarrow \{\Delta_n\}_n$$

of commutative squares in $\text{Pro}(\text{Pres}_c)$, where every $\Delta_{n,0}$ is obtained from Δ_n via the \circlearrowleft -construction.

Proof. Note that by Theorem 3.3.1(i), the induced square $\Delta_{n,0}$

$$\begin{array}{ccc} \mathcal{A}_n & \longrightarrow & \mathcal{B}_n \\ \downarrow & & \downarrow \\ \mathcal{A}'_n & \longrightarrow & \mathcal{A}'_n \circlearrowleft_{\mathcal{A}'_n}^{\mathcal{B}'_n} \mathcal{B}_n \end{array}$$

satisfies base change for every n . To simplify the notation, set $\mathcal{Q}_n := \mathcal{A}'_n \circlearrowleft_{\mathcal{A}'_n}^{\mathcal{B}'_n} \mathcal{B}_n$. It remains only to show that the functors

$$b_n^* : \mathcal{Q}_n \rightarrow \mathcal{B}'_n$$

define a pro-equivalence as n varies. Since $q_n^* \simeq b_n^* q_{n,0}^*$ generates \mathcal{B}'_n under colimits, so does b_n^* . By Lemma 3.5.12, the functor b_n^* is projectively generated for every n (by the objects $B_n^\alpha = f_n^*(A_n^\alpha)$). Thus in order to apply Corollary 3.5.10 it will suffice to show that b_n^* induce isomorphisms on mapping pro-spaces. For this we need a pro-version of Corollary 3.1.3. By Lemma 3.1.2 we have for every n a commutative square of natural transformations

$$\begin{array}{ccc} f_n^* p_{n,*} & \xrightarrow{\text{Ex}_{\Delta_{n,0}}} & q_{n,0,*} g_{n,0}^* \\ \parallel & & \downarrow \eta_{b_n} \\ f_n^* p_{n,*} & \xrightarrow{\text{Ex}_{\Delta_n}} & q_{n,*} g_n^* \end{array}$$

By Lemma 3.3.3, the upper arrow is invertible for every n . Evaluating at the object $A_n'^\alpha := p_n^*(A_n^\alpha) \in \mathcal{A}'_n$ for any α , applying $\text{Maps}_{\mathcal{B}_n}(B_n^\beta, -)$ for any β , and passing to pro-systems yields a commutative diagram of pro-spaces

$$\begin{array}{ccc} \{\text{Maps}_{\mathcal{B}_n}(B_n^\beta, f_n^* p_{n,*}(A_n'^\alpha))\}_n & \xrightarrow{\text{Ex}_{\Delta_{n,0}}} & \{\text{Maps}_{\mathcal{B}_n}(B_n^\beta, q_{n,0,*} g_{n,0}^*(A_n'^\alpha))\}_n \\ \parallel & & \downarrow \\ \{\text{Maps}_{\mathcal{B}_n}(B_n^\beta, f_n^* p_{n,*}(A_n'^\alpha))\}_n & \xrightarrow{\text{Ex}_{\Delta_n}} & \{\text{Maps}_{\mathcal{B}_n}(B_n^\beta, q_{n,*} g_n^*(A_n'^\alpha))\}_n \end{array}$$

where the upper arrow is invertible in $\text{Pro}(\text{Spc})$. The claim is that the right-hand vertical arrow is invertible in $\text{Pro}(\text{Spc})$. Since Δ satisfies pro-base change, it can be represented by some cofiltered system $\{\Delta'_n\}_n$ for which Δ'_n satisfies base change for every n . Choose an isomorphism $\{\Delta_n\}_n \simeq \{\Delta'_n\}_n$ (of squares in $\text{Pro}(\text{Pres}_c)$); re-indexing if necessary, we may assume that it is induced by levelwise morphisms $\Delta_n \rightarrow \Delta'_n$. By functoriality, such an isomorphism gives rise to an identification between the lower arrow in the diagram above and the analogous construction $\{\text{Ex}_{\Delta'_n}\}_n$ for the squares Δ'_n . Since $\text{Ex}_{\Delta'_n}$ is invertible for every n , we conclude that the lower arrow above is also invertible. \square

Proof of Theorem 3.5.11. Combine Lemma 3.5.13 and Proposition 3.4.4. \square

4. LOCALIZING INVARIANTS OF ALGEBRAIC STACKS

Definition 4.0.1. Let \mathcal{X} be a derived algebraic stack which is perfect in the sense of Subsect. A.3. For instance, suppose \mathcal{X} is *ANS* (affine diagonal and nice stabilizers, see Definition A.1.2 and Theorem A.3.2). Then we set

$$E(\mathcal{X}) := E(\mathbf{D}(\mathcal{X})) \simeq E(\text{Perf}(\mathcal{X}))$$

for any localizing invariant E . For example, we have the nonconnective algebraic K-theory spectrum $\mathbf{K}(\mathcal{X})$.

Remark 4.0.2. Let \mathcal{X} and \mathcal{Y} be perfect derived algebraic stacks. Then for any morphism $f : \mathcal{X} \rightarrow \mathcal{Y}$, the functor $f^* : \mathbf{D}(\mathcal{Y}) \rightarrow \mathbf{D}(\mathcal{X})$ preserves perfect complexes and is therefore compact. In particular, there is an induced map

$$f^* : E(\mathcal{Y}) \rightarrow E(\mathcal{X})$$

for any localizing invariant E .

4.1. **Étale excision.** See Subsect. A.3 for the notion of perfectness of stacks.

Theorem 4.1.1. *Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a representable morphism of perfect derived algebraic stacks. Suppose there exists a closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ with quasi-compact open complement such that f is an isomorphism infinitely near \mathcal{Z} , i.e., the induced morphism $\mathcal{X}'_{f^{-1}(\mathcal{Z})} \rightarrow \mathcal{X}_{\mathcal{Z}}$ is invertible. Then for every localizing invariant E , the induced square*

$$\begin{array}{ccc} E(\mathcal{X}) & \xrightarrow{j^*} & E(\mathcal{X} \setminus \mathcal{Z}) \\ \downarrow f^* & & \downarrow \\ E(\mathcal{X}') & \longrightarrow & E(\mathcal{X}' \setminus f^{-1}(\mathcal{Z})) \end{array}$$

is cartesian.

Proof. Apply the criterion of [Tam18, Thm. 18] (cf. [Hoy18, Cor. 13]) to the cartesian square in Theorem 2.2.3. Let $j' : \mathcal{U}' \rightarrow \mathcal{X}'$ denote the open immersion complementary to $f^{-1}(\mathcal{Z})$. Fully faithfulness of the functor $j'_* : \mathbf{D}(\mathcal{U}') \rightarrow \mathbf{D}(\mathcal{X}')$, i.e., invertibility of the unit $\text{id} \rightarrow j'^* j'_*$, follows from the base change formula for the self-intersection $\mathcal{U}' \times_{\mathcal{X}'}^{\mathbf{R}} \mathcal{U}' \simeq \mathcal{U}'$. \square

Specializing to étale neighbourhoods gives:

Corollary 4.1.2 (Étale excision). *Suppose given a Nisnevich square of derived algebraic stacks of the form*

$$\begin{array}{ccc} \mathcal{U}' & \xrightarrow{j'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{U} & \xrightarrow{j} & \mathcal{X} \end{array}$$

where \mathcal{X} and \mathcal{X}' are perfect. Then for every localizing invariant E , the induced square

$$\begin{array}{ccc} E(\mathcal{X}) & \xrightarrow{j^*} & E(\mathcal{U}) \\ \downarrow f^* & & \downarrow \\ E(\mathcal{X}') & \longrightarrow & E(\mathcal{U}') \end{array}$$

is cartesian.

Remark 4.1.3. Note that for affine Nisnevich squares (i.e., squares as above where f is affine), Corollary 4.1.2 can be deduced alternatively from Corollary 2.2.4 and Theorem 3.4.3. This weaker statement would in fact suffice for our purposes in this paper.

Remark 4.1.4. Corollary 4.1.2 also holds for squares as above where f is not assumed representable. This can also be deduced from [Tam18, Thm. 18], e.g. using Nisnevich descent for the presheaf of ∞ -categories $\mathcal{X} \mapsto \mathbf{D}(\mathcal{X})$ and [Kha19, Thm. 2.2.7].

Remark 4.1.5. By [Kha19, Thm. 2.2.7], Corollary 4.1.2 implies that the presheaf of spectra $\mathcal{X} \mapsto E(\mathcal{X})$ satisfies descent for the Grothendieck topology generated by Nisnevich squares. The discussion of [HK19, Prop. 2.8] goes through *mutatis mutandis* in the derived setting to show that this topology is generated by a stacky variant of the usual notion of Nisnevich covers.

4.2. **Formal Milnor and finite excision.**

Theorem 4.2.1. *Suppose given a square of bounded derived algebraic stacks*

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X}. \end{array}$$

with \mathcal{X} noetherian ANS. Assume that i is a closed immersion, f is affine, the square is cartesian on classical truncations, and the formally completed square (2.3.a) is cocartesian. Then for any localizing invariant E , the induced square

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \longrightarrow & \widehat{E}(\mathcal{X}_Z^\wedge) \\ \downarrow & & \downarrow \\ \{E(\mathcal{X}')\} & \longrightarrow & \widehat{E}(\mathcal{X}'_{Z'}) \end{array}$$

is pro-cartesian.

Proof. By Corollary 4.1.2, Theorem A.1.8, Proposition A.1.9 and Theorem A.3.2, we may reduce to the case $\mathcal{X} = [X/G]$, where G is an embeddable nice group scheme over an affine scheme S , and X is an affine derived S -scheme with G -action. Now we may apply the criterion of Theorem 3.5.11 to the square

$$\begin{array}{ccc} \{\mathbf{D}(\mathcal{X})\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}_Z^\wedge) \\ \downarrow & & \downarrow \\ \{\mathbf{D}(\mathcal{X}')\} & \longrightarrow & \widehat{\mathbf{D}}(\mathcal{X}'_{Z'}) \end{array}$$

which is pro-Milnor and satisfies pro-base change by Theorem 2.4.1. The assumptions of Theorem 3.5.11 are verified by Remark 2.2.1 and Proposition A.3.4. \square

Corollary 4.2.2 (Formal Milnor excision). *Suppose given a Milnor square of noetherian algebraic stacks of the form (2.1.b). Then for any localizing invariant E , the induced square*

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \xrightarrow{\widehat{i}^*} & \widehat{E}(\mathcal{X}_Z^\wedge) \\ \downarrow f^* & & \downarrow (f^\wedge)^* \\ \{E(\mathcal{X}')\} & \xrightarrow{\widehat{i}'^*} & \widehat{E}(\mathcal{X}'_{Z'}). \end{array}$$

is pro-cartesian.

Corollary 4.2.3 (Formal finite excision). *Suppose given a finite cdh square of bounded derived stacks*

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

with \mathcal{X} noetherian ANS. Then for any localizing invariant E , the induced square

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \longrightarrow & \widehat{E}(\mathcal{X}_Z^\wedge) \\ \downarrow & & \downarrow \\ \{E(\mathcal{X}')\} & \longrightarrow & \widehat{E}(\mathcal{X}'_{Z'}) \end{array}$$

is pro-cartesian.

Corollary 4.2.4 (Formal nil-excision). *Let \mathcal{X} be a bounded noetherian ANS derived stack with classical truncation \mathcal{X}_{cl} , and let $\mathcal{Z} \hookrightarrow \mathcal{X}$ be a closed immersion with 0-truncated quasi-compact open complement. Then for any localizing invariant E , the induced square*

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \longrightarrow & \widehat{E}(\mathcal{X}_{\mathcal{Z}}^{\wedge}) \\ \downarrow & & \downarrow \\ \{E(\mathcal{X}_{\text{cl}})\} & \longrightarrow & \widehat{E}((\mathcal{X}_{\text{cl}})_{\mathcal{Z}_{\text{cl}}}^{\wedge}) \end{array}$$

is pro-cartesian.

4.3. Derived blow-ups. Given a *quasi-smooth* closed immersion of derived algebraic stacks, one can form the derived blow-up $\widetilde{\mathcal{X}}$ in the sense of [KR18a, 4.1.6], which fits in a commutative square

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{i_{\mathcal{D}}} & \widetilde{\mathcal{X}} \\ \downarrow q & & \downarrow p \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X}, \end{array}$$

where q is the projection of the projectivized normal bundle $\mathcal{E} = \mathbf{P}(\mathcal{N}_{\mathcal{Z}/\mathcal{X}})$ and $i_{\mathcal{D}}$ is a *virtual Cartier divisor*, i.e., a quasi-smooth closed immersion of virtual codimension 1. See [KR18a] for background on quasi-smoothness and virtual Cartier divisors. This square is not homotopy cartesian, but it is an abstract blow-up square in the sense of Definition 2.1.3. The following result was proven in [Kha20, Thm. A]:

Theorem 4.3.1. *Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a quasi-smooth closed immersion of derived algebraic stacks. Then for every localizing invariant E , there is a cartesian square*

$$\begin{array}{ccc} E(\mathcal{X}) & \xrightarrow{i^*} & E(\mathcal{Z}) \\ \downarrow p^* & & \downarrow q^* \\ E(\widetilde{\mathcal{X}}) & \longrightarrow & E(\mathcal{D}). \end{array}$$

In this subsection, we derive a formal version of this statement. This is a special case of formal proper excision.

Proposition 4.3.2. *Let the notation be as above and assume that \mathcal{X} is a noetherian ANS stack. Then the square*

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \longrightarrow & \widehat{E}(\mathcal{X}_{\mathcal{Z}}^{\wedge}) \\ \downarrow & & \downarrow \\ \{E(\widetilde{\mathcal{X}})\} & \longrightarrow & \widehat{E}(\widetilde{\mathcal{X}}_{\mathcal{D}}^{\wedge}) \end{array}$$

is pro-cartesian.

Remark 4.3.3. Note that if \mathcal{X} is ANS, then so is the derived blow-up $\widetilde{\mathcal{X}}$ by Lemma A.1.7.

Remark 4.3.4. If \mathcal{X} has the resolution property, the quasi-smooth closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ can be realized as the derived zero locus of some section s of a vector bundle \mathcal{V} on \mathcal{X} (Remark A.2.3). For every integer n , let $\mathcal{Z}(n) \rightarrow \mathcal{X}$ denote the derived zero locus of $s^{\otimes n}$. By Lemma B.1.5, the formal completion of \mathcal{X} can then be represented by

$$\mathcal{X}^{\wedge} \simeq \{\mathcal{Z}(n)\}_n.$$

Similarly, it follows from Remark B.1.2 that the formal completion of the derived blow-up can be represented by

$$\tilde{\mathcal{X}}^\wedge \simeq \{\mathcal{Z}'(n)\}_n,$$

where $\mathcal{Z}'(n)$ denotes the derived base change $\mathcal{Z}(n) \times_{\mathcal{X}} \tilde{\mathcal{X}}$.

Construction 4.3.5. Assume that \mathcal{X} has the resolution property and let $\mathcal{Z}(n)$ be as in Remark 4.3.4. The derived blow-up square defining $\tilde{\mathcal{X}}$ is the derived base change of the classical blow-up square

$$\begin{array}{ccc} \mathcal{D}_{\mathcal{V}} & \longrightarrow & \tilde{\mathcal{V}} \\ \downarrow & & \downarrow \\ \mathcal{X} & \xrightarrow{0} & \mathcal{V} \end{array}$$

of the zero section $0 : \mathcal{X} \rightarrow \mathcal{V}$. Let $\mathcal{X}(n)$ denote the n th infinitesimal thickening of the latter. Write $\mathcal{D}_{\mathcal{V}}(n)$ for the (classical) fibre product of $\mathcal{X}(n)$ and $\tilde{\mathcal{V}}$ over \mathcal{V} (which is an effective Cartier divisor in $\tilde{\mathcal{V}}$) and let $\mathcal{Z}'_{\mathcal{V}}(n)$ denote the derived fiber product. The two commutative squares

$$\begin{array}{ccc} \mathcal{D}_{\mathcal{V}}(n) & \longrightarrow & \tilde{\mathcal{V}} \\ \downarrow & & \downarrow \\ \mathcal{X}(n) & \longrightarrow & \mathcal{V}, \end{array} \quad \begin{array}{ccc} \mathcal{Z}'_{\mathcal{V}}(n) & \longrightarrow & \tilde{\mathcal{V}} \\ \downarrow & & \downarrow \\ \mathcal{X}(n) & \longrightarrow & \mathcal{V} \end{array} \quad (4.3.a)$$

define by derived base change to \mathcal{X} the squares

$$\begin{array}{ccc} \mathcal{D}(n) & \longrightarrow & \tilde{\mathcal{X}} \\ \downarrow & & \downarrow \\ \mathcal{Z}(n) & \longrightarrow & \mathcal{X}, \end{array} \quad \begin{array}{ccc} \mathcal{Z}'(n) & \longrightarrow & \tilde{\mathcal{X}} \\ \downarrow & & \downarrow \\ \mathcal{Z}(n) & \longrightarrow & \mathcal{X}. \end{array} \quad (4.3.b)$$

The right-hand square is homotopy cartesian. The left-hand square is a derived blow-up square for $n = 1$, and otherwise is a thickening of the latter.⁴

The ind-stack $\{\mathcal{D}(n)\}$ gives another presentation of $\tilde{\mathcal{X}}^\wedge$:

Lemma 4.3.6. *If \mathcal{X} is a noetherian algebraic stack, then the morphisms $\mathcal{D}(n) \rightarrow \mathcal{Z}'(n)$ induce an isomorphism*

$$\{\mathcal{D}(n)\}_n \rightarrow \{\mathcal{Z}'(n)\}_n \simeq \tilde{\mathcal{X}}^\wedge$$

of ind-stacks over $\tilde{\mathcal{X}}$.

Proof. By derived base change, it will suffice to show the claim for the analogous constructions over $\tilde{\mathcal{V}}$. Note that both squares in (4.3.a) consist of classical stacks, with the sole exception of $\mathcal{Z}'_{\mathcal{V}}(n)$. In fact, the morphism $\mathcal{D}_{\mathcal{V}}(n) \rightarrow \mathcal{Z}'_{\mathcal{V}}(n)$ exhibits the domain as the classical truncation of the codomain. Thus the morphism

$$\{\mathcal{D}_{\mathcal{V}}(n)\}_n \rightarrow \{\mathcal{Z}'_{\mathcal{V}}(n)\}_n$$

over $\tilde{\mathcal{V}}$ can be identified with the morphism from the classical formal completion of $\tilde{\mathcal{V}}$ (in the classical stack $\mathcal{D}_{\mathcal{V}}$) to the formal completion in the sense of Definition B.1.1, and is invertible by Remark B.1.6. \square

⁴In fact, $\mathcal{D}(n)$ can be described as the n -fold sum $n\mathcal{D}$ of the virtual Cartier divisor \mathcal{D} , but we do not need this here.

Notation 4.3.7. To simplify the notation, we will compress commutative squares of ind-stacks of the form

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow \\ \mathcal{Z} & \longrightarrow & \mathcal{X}, \end{array}$$

where the horizontal arrows can be represented by levelwise closed immersions, into morphisms of pairs

$$(\mathcal{X}', \mathcal{Z}') \rightarrow (\mathcal{X}, \mathcal{Z}).$$

We will say that such a morphism is an *E-equivalence* if it induces an isomorphism

$$\widehat{E}(\mathcal{X}, \mathcal{Z}) \rightarrow \widehat{E}(\mathcal{X}', \mathcal{Z}'),$$

where $\widehat{E}(\mathcal{X}, \mathcal{Z})$ is defined as the fibre of $\widehat{E}(\mathcal{X}) \rightarrow \widehat{E}(\mathcal{Z})$.

To show Proposition 4.3.2, we will need to compare the pairs

$$\{(\widetilde{\mathcal{X}}, \mathcal{Z}'(n))\}_n \quad \text{and} \quad \{(\mathcal{X}, \mathcal{Z}(n))\}_n,$$

up to *E*-equivalence. By Theorem 4.3.1, we have *E*-equivalence of the pair $\{(\mathcal{X}, \mathcal{Z}(n))\}_n$ with its derived blow up $\{(\widetilde{\mathcal{X}}(n), \mathcal{D}(n, n))\}_n$, which is *E*-equivalent to the pair $\{(\widetilde{\mathcal{X}}(n), \mathcal{Z}'(n, n))\}_n$ by Lemma 4.3.6, where $\mathcal{Z}'(n, n)$ is the derived pullback of $\mathcal{Z}(n)$ in $\widetilde{\mathcal{X}}(n)$. The goal is therefore to understand the relation between $\{(\widetilde{\mathcal{X}}(n), \mathcal{Z}'(n, n))\}_n$ and $\{(\widetilde{\mathcal{X}}, \mathcal{Z}'(n))\}_n$. For this purpose it will be convenient to introduce the following bi-indexed construction.

Construction 4.3.8. For every pair of natural numbers n and k , consider the following two commutative squares:

$$\begin{array}{ccc} \mathcal{D}_{\mathcal{V}}(n, k) & \longrightarrow & \widetilde{\mathcal{V}}(k) \\ \downarrow & & \downarrow \\ \mathcal{X}(n) & \longrightarrow & \mathcal{V}, \end{array} \quad \begin{array}{ccc} \mathcal{Z}'_{\mathcal{V}}(n, k) & \longrightarrow & \widetilde{\mathcal{V}}(k) \\ \downarrow & & \downarrow \\ \mathcal{X}(n) & \longrightarrow & \mathcal{V} \end{array} \quad (4.3.c)$$

Here $\widetilde{\mathcal{V}}(k)$ is the blow-up of \mathcal{V} centred in $\mathcal{X}(k)$. The left-hand square is classically cartesian and the right-hand square is homotopy cartesian. As above, all stacks are underived except for $\mathcal{Z}'_{\mathcal{V}}(n, k)$.

Now by derived base change to \mathcal{X} we get the squares

$$\begin{array}{ccc} \mathcal{D}(n, k) & \longrightarrow & \widetilde{\mathcal{X}}(k) \\ \downarrow & & \downarrow \\ \mathcal{Z}(n) & \longrightarrow & \mathcal{X}, \end{array} \quad \begin{array}{ccc} \mathcal{Z}'(n, k) & \longrightarrow & \widetilde{\mathcal{X}}(k) \\ \downarrow & & \downarrow \\ \mathcal{Z}(n) & \longrightarrow & \mathcal{X}. \end{array} \quad (4.3.d)$$

Note that we have

$$\mathcal{D}(n, 1) = \mathcal{D}(n), \quad \mathcal{Z}'(n, 1) = \mathcal{Z}'(n)$$

for every n . We regard $\mathcal{D}(n, k)$ and $\mathcal{Z}'(n, k)$ as ind-stacks indexed by the poset of pairs $(n, k) \in \mathbf{N} \times \mathbf{N}$, where $(n, k) \leq (n', k')$ iff $n \leq n'$ and $k \leq k'$.

Proof of Proposition 4.3.2. By Corollary 4.1.2, Theorem A.1.8, Proposition A.1.9 and Theorem A.3.2, we may reduce to the case $\mathcal{X} = [X/G]$, where G is an embeddable nice group scheme over an affine scheme S , and X is an affine S -scheme with G -action. Then by Proposition A.2.5, \mathcal{X} has the resolution property so we are in the situation of Remark 4.3.4 and Construction 4.3.5.

We need to show that the morphism of pairs

$$\{(\tilde{\mathcal{X}}, \mathcal{Z}'(n))\}_n \rightarrow \{(\mathcal{X}, \mathcal{Z}(n))\}_n \quad (4.3.e)$$

is an E -equivalence. Passing to the bi-indexed system constructed in Construction 4.3.8, note that for $k = n$, the square

$$\begin{array}{ccc} \mathcal{D}(n, n) & \longrightarrow & \tilde{\mathcal{X}}(n) \\ \downarrow & & \downarrow \\ \mathcal{Z}(n) & \longrightarrow & \mathcal{X} \end{array}$$

is a derived blow-up square and therefore by Theorem 4.3.1,

$$\{(\tilde{\mathcal{X}}(n), \mathcal{D}(n, n))\}_n \rightarrow \{(\mathcal{X}, \mathcal{Z}(n))\}_n$$

induces a levelwise E -equivalence. For each k , Lemma 4.3.6 gives an identification of ind-stacks

$$\{\mathcal{D}(n, k)\}_n \rightarrow \{\mathcal{Z}'(n, k)\}_n,$$

which induces an equivalence

$$\{\mathcal{D}(n, n)\}_n \rightarrow \{\mathcal{Z}'(n, n)\}_n.$$

Therefore we have an E -equivalence

$$\{(\tilde{\mathcal{X}}(n), \mathcal{Z}'(n, n))\}_n \simeq \{(\tilde{\mathcal{X}}(n), \mathcal{D}(n, n))\}_n \rightarrow \{(\mathcal{X}, \mathcal{Z}(n))\}_n. \quad (4.3.f)$$

The inclusion of posets $\Delta_{\mathbf{N}} \subseteq \mathbf{N} \times \mathbf{N} \supseteq \mathbf{N} \times \{1\}$, induces morphisms

$$\{(\tilde{\mathcal{X}}(m), \mathcal{Z}'(m, m))\}_m \rightarrow \{(\tilde{\mathcal{X}}(k), \mathcal{Z}'(n, k))\}_{n,k} \leftarrow \{(\tilde{\mathcal{X}}(n), \mathcal{Z}'(n, 1))\}_n \simeq \{(\tilde{\mathcal{X}}(n), \mathcal{Z}'(n))\}_n,$$

where the left arrow is an equivalence by cofinality and we shall see that the right arrow is an equivalence by finite formal excision. As one can see by base change from \mathcal{V} , there is for every k a *finite* morphism

$$\tilde{\mathcal{X}} \rightarrow \tilde{\mathcal{X}}(k)$$

which fits in a commutative diagram

$$\begin{array}{ccc} \mathcal{Z}'(n) & \longrightarrow & \tilde{\mathcal{X}} \\ \downarrow & & \downarrow \\ \mathcal{Z}'(n, k) & \longrightarrow & \tilde{\mathcal{X}}(k) \\ \downarrow & & \downarrow \\ \mathcal{Z}(n) & \longrightarrow & \mathcal{X} \end{array}$$

where the squares are homotopy cartesian for each n, k . This provides a factorization

$$\{(\tilde{\mathcal{X}}, \mathcal{Z}'(n))\}_n \rightarrow \{(\tilde{\mathcal{X}}(n), \mathcal{Z}'(n, n))\}_n \rightarrow \{(\mathcal{X}, \mathcal{Z}(n))\}_n, \quad (4.3.g)$$

where the first arrow is an E -equivalence by finite excision (Corollary 4.2.3) and the second arrow is an E -equivalence by (4.3.f). \square

4.4. Formal proper excision. The following is the main result of this section. It generalizes Corollary 4.2.3 and Proposition 4.3.2 to arbitrary proper cdh squares. The proof is essentially the same as that in [Kha20, 5.3.4].

Theorem 4.4.1. *Suppose given a proper cdh square of algebraic stacks*

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

with \mathcal{X} noetherian and ANS. Then for any localizing invariant E , the induced square

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \longrightarrow & \widehat{E}(\mathcal{X}^\wedge_{\mathcal{Z}}) \\ \downarrow & & \downarrow \\ \{E(\mathcal{X}')\} & \longrightarrow & \widehat{E}(\mathcal{X}'^\wedge_{\mathcal{Z}'}) \end{array}$$

is pro-cartesian.

Proof. Let us first demonstrate the claim in the case where f is the projection of the blow-up $\mathcal{X}' = \mathrm{Bl}_{\mathcal{Z}} \mathcal{X}$. By Nisnevich descent (Corollary 4.1.2), Theorem A.1.8 and Proposition A.1.9, we may assume that $\mathcal{X} = [X/G]$ where G is an embeddable nice group scheme over an affine scheme S , and X is an affine derived S -scheme with G -action. Since \mathcal{X} has the resolution property (Proposition A.2.5), $i : \mathcal{Z} \rightarrow \mathcal{X}$ is the classical truncation of a quasi-smooth closed immersion $\widetilde{\mathcal{Z}} \rightarrow \mathcal{X}$ (Construction A.2.2). Let $\widetilde{\mathcal{X}} \rightarrow \mathcal{X}$ denote the derived blow-up of the latter. Since the formal completion \mathcal{X}^\wedge only depends on the classical truncation $\widetilde{\mathcal{Z}}_{\mathrm{cl}} \simeq \mathcal{Z}$, the square in question factors as in the diagram below:

$$\begin{array}{ccc} \{E(\mathcal{X})\} & \longrightarrow & \widehat{E}(\mathcal{X}^\wedge) \\ \downarrow & & \downarrow \\ \{E(\widetilde{\mathcal{X}})\} & \longrightarrow & \widehat{E}(\widetilde{\mathcal{X}}^\wedge) \\ \downarrow & & \downarrow \\ \{E(\mathcal{X}')\} & \longrightarrow & \widehat{E}(\mathcal{X}'^\wedge). \end{array}$$

The upper square is cartesian by Proposition 4.3.2. Since $\mathcal{X}' \rightarrow \widetilde{\mathcal{X}}$ is a closed immersion which is an isomorphism away from \mathcal{Z} , the lower square is also cartesian by Corollary 4.2.3. This concludes the proof in the case of a blow-up square.

Now consider the case of an arbitrary proper morphism f . By Corollary 4.2.3 it is safe to replace \mathcal{X} by the schematic closure of $\mathcal{X} \setminus \mathcal{Z}$ and thereby assume that $\mathcal{X} \setminus \mathcal{Z}$ is schematically dense in \mathcal{X} and $\mathcal{X}' \setminus \mathcal{Z}'$ is schematically dense in \mathcal{X}' . In this case Rydh's stacky generalization of Raynaud–Gruson (see [Ryd17], [HK19, Cor. 2.4]) to $f : \mathcal{X}' \rightarrow \mathcal{X}$ yields a proper morphism $f' : \mathcal{X}'' \rightarrow \mathcal{X}'$ such that $f \circ f' : \mathcal{X}'' \rightarrow \mathcal{X}$ is a sequence of $(\mathcal{X} \setminus \mathcal{Z})$ -admissible blow-ups. In particular, f' sits in a proper cdh square Q' over the original square Q . Applying the construction again to f' yields a third proper cdh square Q'' over Q' . Then it suffices to show the claim for the two squares $Q' \circ Q$ and $Q'' \circ Q'$, so we have reduced to the case where f is a sequence of $(\mathcal{X} \setminus \mathcal{Z})$ -admissible blow-ups. By induction, we may as well assume it is a $(\mathcal{X} \setminus \mathcal{Z})$ -admissible blow-up. Using Corollary 4.2.3 and the same argument as in [KST17, Claim 5.3], one finally reduces to the case of a blow-up considered above. \square

Remark 4.4.2. Theorem 4.4.1 also holds “with supports” in any closed substack $\mathcal{Y} \subseteq \mathcal{X}$. That is, one can replace $E(\mathcal{X})$ with $E(\mathcal{X} \text{ on } \mathcal{Y})$, $\widehat{E}(\mathcal{X}_z^\wedge)$ with $\widehat{E}(\mathcal{X}_z^\wedge \text{ on } \mathcal{Y}_{y \cap z}^\wedge)$, etc. Here $E(\mathcal{X} \text{ on } \mathcal{Y})$ is E applied to the kernel of the restriction $\text{Perf}(\mathcal{X}) \rightarrow \text{Perf}(\mathcal{X} \setminus \mathcal{Y})$ as in [TT07]. Note that since E is localizing, there are exact triangles

$$E(\mathcal{X} \text{ on } \mathcal{Y}) \rightarrow E(\mathcal{X}) \rightarrow E(\mathcal{X} \setminus \mathcal{Y}).$$

Therefore, the “with supports” variant of Theorem 4.4.1 follows immediately from the “without supports” one.

5. NEGATIVE K-THEORY

5.1. Nil-invariance of negative K-groups. In this subsection we prove a nil-invariance result for sufficiently negative K-groups of ANS stacks (Corollary 5.1.4).

We make use of the formalism of weight structures of Bondarko. Let \mathcal{C} be an additive ∞ -category which is projectively generated in the sense of [HTT, Def. 5.5.8.23]. Then the full subcategory \mathcal{A} of compact projective objects is an idempotent-complete additive ∞ -category for which the inclusion $\mathcal{A} \subset \mathcal{C}$ extends to an equivalence $\mathcal{P}_\Sigma(\mathcal{A}) \simeq \mathcal{C}$ by [HTT, Prop. 5.5.8.25]. By [SAG, Prop. C.1.5.7], \mathcal{C} is prestable and thus embeds fully faithfully into its stabilization $\text{Spt}(\mathcal{C})$. In this situation, the full subcategory $\mathcal{D} = \text{Spt}(\mathcal{C})^\omega$ of compact objects admits a *weight structure* in the sense of [Bon10]. This weight structure is bounded, its heart $\mathcal{D}^{w=0}$ is \mathcal{A} , and the subcategory $\mathcal{D}^{w \geq 0} \subseteq \mathcal{D}$ of connective objects is \mathcal{C}^ω . Moreover, every bounded weight structure arises in this manner: in fact, the ∞ -category of weighted ∞ -categories and weight-exact functors is equivalent to the ∞ -category of idempotent-complete additive ∞ -categories by [Sos19, Prop. 3.3] (see also [Sos21, Props. 3.1.4, 3.1.5]). We refer the reader to [Sos19, 1.3], [Sos21, 3.1], or [ES20] for an ∞ -categorical account of the theory of weight structures, originally developed in [Bon10].

Example 5.1.1. Let A be a connective \mathcal{E}_∞ -ring. Then the ∞ -category Mod_A^{cn} of connective A -modules is projectively generated by A . Thus there exists a canonical weight structure on the stable ∞ -category Perf_A whose heart is the full subcategory of finite projective A -modules.

The next example, a slight generalization of [Sos21, Theorem 3.4.3], will play in an important role in what follows.

Example 5.1.2. Let R be a commutative ring, G an embeddable linearly reductive group scheme over R , and A a derived commutative ring over R . Then by Proposition A.3.4, the ∞ -category $\mathbf{D}([\text{Spec}(A)/G])_{\geq 0}$ of connective G -equivariant A -modules is projectively generated and there exists a canonical weight structure on $\mathbf{D}([\text{Spec}(A)/G])$ whose heart is the full subcategory spanned by objects of the form $p^*(\mathcal{E})$, where $p: [\text{Spec}(A)/G] \rightarrow BG$ is the projection and $\mathcal{E} \in \mathbf{D}(BG)$ is a finite projective G -equivariant R -module.

Proposition 5.1.3. *Let R be a commutative ring and G an embeddable linearly reductive group scheme over R . For any G -equivariant nilpotent extension $A \rightarrow B$ of connective \mathcal{E}_∞ -algebras over R with G -actions (i.e., a π_0 -surjection with nilpotent kernel), the induced map*

$$\mathbf{K}_{-n}([\text{Spec}(A)/G]) \rightarrow \mathbf{K}_{-n}([\text{Spec}(B)/G])$$

is an isomorphism for every $n \geq 0$.

Proof. Example 5.1.2 together with Lemma A.2.6 shows that the morphism $A \rightarrow B$ induces an equivalence of the homotopy categories of the heart of the weight structures on the ∞ -categories $\mathbf{D}([\text{Spec}(A)/G])$ and $\mathbf{D}([\text{Spec}(B)/G])$. The required isomorphism on negative K-groups therefore follows from [Sos19, Theorem 4.3]. \square

This further implies the following nil-invariance statement for a large class of stacks, which will be crucial for the proof of Weibel’s conjecture.

Corollary 5.1.4. *Let \mathcal{X} be a ANS derived stack of Nisnevich cohomological dimension d . Then for any surjective closed immersion $i : \mathcal{X}' \rightarrow \mathcal{X}$, the map*

$$i^* : K_{-n}(\mathcal{X}) \rightarrow K_{-n}(\mathcal{X}')$$

is an isomorphism for every $n > d$.

Proof. By Corollary 4.1.2, we may regard $K(-)$ and $K(- \times_{\mathcal{X}} \mathcal{X}')$ as Nisnevich sheaves of spectra on the small étale site of \mathcal{X} . Then the fibre \mathcal{F} of the morphism $K(-) \rightarrow K(- \times_{\mathcal{X}} \mathcal{X}')$ is also a Nisnevich sheaf. The claim is that $\pi_{-n}(\mathcal{F}(\mathcal{X})) = 0$ for $n > d$. Considering the descent spectral sequence

$$H_{\text{Nis}}^p(\mathcal{X}, \pi_q^{\text{Nis}}(\mathcal{F})) \Rightarrow \pi_{q-p}(\mathcal{F}(\mathcal{X})),$$

it will suffice to show that the left-hand side is trivial for $q < 0$ and for $p > d$. By Theorem A.1.8 and Proposition A.1.9, there exists an embeddable linearly reductive group scheme G over an affine scheme S , and affine derived schemes U_i over S with G -action, together with étale morphisms $[U_i/G] \rightarrow \mathcal{X}$ which generate a Nisnevich covering. By Proposition 5.1.3, the spectrum $\mathcal{F}([U_i/G])$ is connective for all i . Hence the homotopy sheaves $\pi_q^{\text{Nis}}(\mathcal{F})$ vanish for $q < 0$. \square

5.2. Killing by blow-ups. The killing lemma, proven in [KS17, Prop. 5], says that for any negative K-theory class one can find a suitable (sequence of) blow-ups along which the inverse image vanishes. It was generalized to stacks, using Rydh’s generalization of Raynaud–Gruson flatification, in [HK19, Prop. 7.3]. The following is a “with supports” variant of the killing lemma, which we will require for our proof of the Weibel conjecture.

Proposition 5.2.1. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a smooth morphism of finite type where \mathcal{Y} is a reduced noetherian ANS stack and \mathcal{X} satisfies the resolution property. Let $\mathcal{Z} \subseteq \mathcal{X}$ be a closed substack. Then for any integer $i > 0$ and any class $\alpha \in K_{-i}(\mathcal{X}$ on $\mathcal{Z})$, there exists a sequence of blow-ups $q : \mathcal{Y}' \rightarrow \mathcal{Y}$ with nowhere dense centres such that $q_{\mathcal{X}}^*(\alpha) = 0$ in $K_{-i}(\mathcal{X}'$ on $\mathcal{Z}')$, where $\mathcal{X}' := \mathcal{X} \times_{\mathcal{Y}} \mathcal{Y}'$, $\mathcal{Z}' := \mathcal{Z} \times_{\mathcal{X}} \mathcal{X}'$, and $q_{\mathcal{X}} : \mathcal{X}' \rightarrow \mathcal{X}$ is the projection.*

The proof will use the following standard lemma. For a noetherian algebraic stack \mathcal{X} , let $\text{Coh}^b(\mathcal{X})$ be the derived ∞ -category of coherent complexes on \mathcal{X} (see e.g. [Kha22, Defn. 1.5] where it is denoted $\mathbf{D}_{\text{coh}}(\mathcal{X})$). Given a closed substack \mathcal{Z} , $\text{Coh}^b(\mathcal{X}$ on $\mathcal{Z})$ denotes the full subcategory spanned by complexes supported set-theoretically on $|\mathcal{Z}|$.

Lemma 5.2.2. *Let \mathcal{X} be a noetherian algebraic stack and \mathcal{Z} a closed substack. Then for any open immersion $j : \mathcal{U} \rightarrow \mathcal{X}$, the restriction functor*

$$j^* : \text{Coh}^b(\mathcal{X} \text{ on } \mathcal{Z}) \rightarrow \text{Coh}^b(\mathcal{U} \text{ on } \mathcal{Z} \cap \mathcal{U})$$

is essentially surjective.

Proof. Since j^* is t-exact and the t-structures are bounded, it is enough to show that it induces an essentially surjective functor on the hearts. That is, it is enough to show that every coherent sheaf \mathcal{F} on \mathcal{U} supported on $\mathcal{Z} \cap \mathcal{U}$ extends to a coherent sheaf $\tilde{\mathcal{F}}$ on \mathcal{X} supported on \mathcal{Z} . Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ and $i_{\mathcal{U}} : \mathcal{Z} \cap \mathcal{U} \rightarrow \mathcal{U}$ denote the inclusions. We may write $\mathcal{F} \simeq i_{\mathcal{U},*}(\mathcal{G})$ for some coherent sheaf \mathcal{G} on $\mathcal{Z} \cap \mathcal{U}$. Now \mathcal{G} can be extended to a coherent sheaf $\tilde{\mathcal{G}}$ on \mathcal{Z} (see e.g. [LMB18, Cor. 15.5]). By the base change formula, $i_{\mathcal{U},*}(\tilde{\mathcal{G}})|_{\mathcal{U}} \simeq i_{\mathcal{U},*}(\tilde{\mathcal{G}}|_{\mathcal{Z} \cap \mathcal{U}}) \simeq \mathcal{F}$. That is, $\tilde{\mathcal{F}} = i_{\mathcal{X},*}(\tilde{\mathcal{G}})$ is an extension of \mathcal{F} as desired. \square

Proof of Proposition 5.2.1. By inductive application of the Bass fundamental theorem (see e.g. [Kha22, Thm. 2.21], which holds with supports just as in [TT07, Thm. 7.5]), there is a canonical surjection

$$\mathrm{Coker}(\mathrm{K}_0(\mathcal{X} \times \mathbf{A}^i \text{ on } \mathcal{Z} \times \mathbf{A}^i) \xrightarrow{j^*} \mathrm{K}_0(\mathcal{X} \times \mathbf{G}_m^i \text{ on } \mathcal{Z} \times \mathbf{G}_m^i)) \rightarrow \mathrm{K}_{-i}(\mathcal{X} \text{ on } \mathcal{Z}).$$

Therefore it suffices to show that for any $\alpha \in \mathrm{K}_0(\mathcal{X} \times \mathbf{G}_m^i \text{ on } \mathcal{Z} \times \mathbf{G}_m^i)$, there exists a sequence of blow-ups $q: \mathcal{Y}' \rightarrow \mathcal{Y}$ with nowhere dense centers such that the image of α in $\mathrm{K}_0(\mathcal{X}' \times \mathbf{G}_m^i \text{ on } \mathcal{Z}' \times \mathbf{G}_m^i)$ lifts to a class in $\mathrm{K}_0(\mathcal{X} \times \mathbf{A}^i \text{ on } \mathcal{Z} \times \mathbf{A}^i)$. By definition, we may write $\alpha = [\mathcal{F}]$ where \mathcal{F} is a perfect complex on $\mathcal{X} \times \mathbf{G}_m^i$ supported on $\mathcal{Z} \times \mathbf{G}_m^i$.

By Lemma 5.2.2, we can extend \mathcal{F} to some $\mathcal{E} \in \mathrm{Coh}^b(\mathcal{X} \times \mathbf{A}^i \text{ on } \mathcal{Z} \times \mathbf{A}^i)$. Since $\mathcal{X} \times \mathbf{A}^i$ has the resolution property (as it is affine over \mathcal{X}), we may assume that \mathcal{E} is represented by a chain complex \mathcal{E}_\bullet of finite locally free sheaves with $\mathcal{E}_n = 0$ for $n \ll 0$. Since its restriction $\mathcal{F} \simeq j^*(\mathcal{E})$ is perfect, say of Tor-amplitude $\leq a$, we may replace \mathcal{E} by its truncation $\tau_{\leq a}(\mathcal{E})$ (which still restricts to \mathcal{F}) so that it may be represented by a chain complex \mathcal{E}_\bullet with the following properties:

- (1) $\mathcal{E}_n = 0$ for $n \ll 0$ or $n > a$;
- (2) \mathcal{E}_n is finite locally free for all $n < a$;
- (3) \mathcal{E}_a is coherent and $j^*(\mathcal{E}_a)$ is finite locally free.

By Rydh's stacky generalization of Raynaud–Gruson (see [Ryd17, Thm. 4.2]), we can argue as in the proof of [HK19, Prop. 7.3] to produce a sequence of blow-ups $q: \mathcal{Y}' \rightarrow \mathcal{Y}$ such that the strict transform \mathcal{E}'_a of \mathcal{E}_a on $\mathcal{X}' \times \mathbf{A}^i$ is flat over \mathcal{X}' . For every n , the strict transform \mathcal{E}'_n of \mathcal{E}_n on $\mathcal{X}' \times \mathbf{A}^i$ is the cokernel of the inclusion $\mathcal{H}_{\mathcal{D} \times_{\mathcal{Y}} \mathcal{X} \times \mathbf{A}^i}^0(q_{\mathcal{X}}^*(\mathcal{E}_n)) \hookrightarrow q_{\mathcal{X}}^*(\mathcal{E}_n)$, where $\mathcal{D} \subseteq \mathcal{Y}' \times \mathbf{A}^i$ is the exceptional divisor. Thus we may regard \mathcal{E}'_\bullet as a chain complex with the following properties:

- (a) \mathcal{E}'_a is of finite tor-amplitude on $\mathcal{X}' \times \mathbf{A}^i$, since it is flat over \mathcal{Y}' (see e.g. [HK19, Lem. 7.2]).
- (b) For every $n < a$ we have $\mathcal{E}'_n = q_{\mathcal{X}}^*(\mathcal{E}_n)$, since \mathcal{E}_n is already flat over $\mathcal{X}' \times \mathbf{A}^i$.
- (c) For every n , we have $\mathcal{E}'_n|_{\mathcal{X}' \times \mathbf{G}_m^i} = q_{\mathcal{X}}^*(\mathcal{F}_n)$, because \mathcal{F}_n is already flat over \mathcal{X}' .

Since each term of \mathcal{E}'_\bullet is of finite Tor-amplitude on $\mathcal{X}' \times \mathbf{A}^i$, \mathcal{E}'_\bullet represents a perfect complex $\mathcal{E}' \in \mathrm{Perf}(\mathcal{X}' \times \mathbf{A}^i)$. Since the chain complex $q_{\mathcal{X}}^*(\mathcal{E}_\bullet)$ has homology supported on $\mathcal{Z}' \times \mathbf{A}^i$, the same holds for its quotient \mathcal{E}'_\bullet , hence in particular $\mathcal{E}' \in \mathrm{Perf}(\mathcal{X}' \times \mathbf{A}^i \text{ on } \mathcal{Z}' \times \mathbf{A}^i)$. Finally, since $\mathcal{E}'|_{\mathcal{X}' \times \mathbf{G}_m^i} \simeq q_{\mathcal{X}}^*(\mathcal{F})$ in $\mathrm{Perf}(\mathcal{X}' \times \mathbf{G}_m^i \text{ on } \mathcal{Z}' \times \mathbf{G}_m^i)$, we find that the class $[\mathcal{E}'] \in \mathrm{K}_0(\mathcal{X}' \times \mathbf{A}^i \text{ on } \mathcal{Z}' \times \mathbf{A}^i)$ lifts $q_{\mathcal{X}}^*(\alpha)$ as claimed. \square

5.3. Weibel's conjecture (I).

Theorem 5.3.1. *Let \mathcal{X} be a noetherian ANS stack of fppf-covering dimension d (see Definition A.4.1). Then the negative K-groups $\mathrm{K}_{-i}(\mathcal{X} \text{ on } \mathcal{Y})$ vanish for all $i > d$.*

Lemma 5.3.2. *Let \mathcal{X} be a reduced noetherian ANS stack with the resolution property of fppf-covering dimension d (see Definition A.4.1). Then for any closed substack $\mathcal{Y} \subseteq \mathcal{X}$, the negative K-groups $\mathrm{K}_{-i}(\mathcal{X} \text{ on } \mathcal{Y})$ vanish for all $i > d$.*

Proof. We argue by induction on d . For any element $\gamma \in \mathrm{K}_{-i}(\mathcal{X} \text{ on } \mathcal{Y})$, there exists by Proposition 5.2.1 a sequence of blow-ups $f: \mathcal{X}' \rightarrow \mathcal{X}$ with nowhere dense centers such that $f^*(\gamma) = 0$ in

$K_{-i}(\mathcal{X}' \text{ on } f^{-1}(\mathcal{Y}))$. This fits in a proper cdh square

$$\begin{array}{ccc} \mathcal{Z}' & \longrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow f \\ \mathcal{Z} & \longrightarrow & \mathcal{X} \end{array}$$

where $\mathcal{Z} \subseteq \mathcal{X}$ is any nowhere dense closed substack for which f is an isomorphism over $\mathcal{X} \setminus \mathcal{Z}$. Let $\mathcal{Z}(n)$ and $\mathcal{Z}'(n)$ denote the n th infinitesimal thickenings of \mathcal{Z} and \mathcal{Z}' , respectively. By Theorem 4.4.1 (and Remark 4.4.2) and Remark B.1.6, we get a long exact sequence

$\cdots \rightarrow \{K_{-i+1}(\mathcal{Z}'(n)) \text{ on } f^{-1}(\mathcal{Y})\}_n \rightarrow K_{-i}(\mathcal{X} \text{ on } \mathcal{Y}) \rightarrow \{K_{-i}(\mathcal{Z}(n)) \text{ on } \mathcal{Y}\}_n \oplus K_{-i}(\mathcal{X}' \text{ on } f^{-1}(\mathcal{Y})) \rightarrow \cdots$ of pro-abelian groups. Now $\mathcal{Z}(n)$ and $\mathcal{Z}'(n)$ are reduced noetherian ANS stacks satisfying the resolution property (Lemma A.1.7 and Lemma A.2.4) and of fppf-covering dimension $< d$, so $\{K_{-i+1}(\mathcal{Z}'(n)) \text{ on } f^{-1}(\mathcal{Y})\}_n$ and $\{K_{-i}(\mathcal{Z}(n)) \text{ on } \mathcal{Y}\}_n$ both vanish by the induction hypothesis. It follows that $f^* : K_{-i}(\mathcal{X}) \rightarrow K_{-i}(\mathcal{X}')$ is injective and hence that $\gamma = 0$. \square

Proof of Theorem 5.3.1. We again argue by induction on d . Suppose $d = 0$. By Corollary 4.1.2 and Theorem A.3.2, there is a convergent Nisnevich-descent spectral sequence:

$$H_{\text{Nis}}^p(\mathcal{X}, \pi_q^{\text{Nis}}(\mathbb{K})) \Rightarrow K_{q-p}(\mathcal{X}),$$

where $\pi_q^{\text{Nis}}(\mathbb{K})$ denotes the Nisnevich sheaf associated with K_q . It follows from Proposition A.4.4 and the previous case that $H_{\text{Nis}}^p(\mathcal{X}, \pi_q^{\text{Nis}}(\mathbb{K}))$ vanishes for all $q < 0$ and $p > 0$ and therefore also $K_{-i}(\mathcal{X}) = 0$ for $i > 0$.

Now suppose $d \geq 1$. By Theorem A.1.8 and Proposition A.2.5, there is a finite sequence of open immersions $\emptyset = \mathcal{U}_0 \hookrightarrow \mathcal{U}_1 \hookrightarrow \cdots \hookrightarrow \mathcal{U}_n = \mathcal{X}$, and Nisnevich squares

$$\begin{array}{ccc} \mathcal{W}_j & \longrightarrow & \mathcal{V}_j \\ \downarrow & & \downarrow \\ \mathcal{U}_{j-1} & \longrightarrow & \mathcal{U}_j, \end{array}$$

where each \mathcal{W}_j and \mathcal{V}_j satisfy the resolution property and have fppf-covering dimension $\leq d$. We prove by induction on j that for $i > d$, $K_{-i}(\mathcal{U}_j)$ vanishes. For $j = 0$, this follows from the previous case. By Proposition A.4.4 and Corollary 5.1.4, we may assume that \mathcal{U}_j is reduced. Choose $\gamma \in K_{-i}(\mathcal{U}_j)$. By induction on j , and the previous case for stacks with resolution property, the groups $K_{-l}(\mathcal{W}_j)$, $K_{-l}(\mathcal{V}_j)$ and $K_{-l}(\mathcal{U}_{j-1})$ vanish for all $l > d$. By Nisnevich descent (Corollary 4.1.2), we have a long exact sequence:

$$\cdots \rightarrow K_{-i+1}(\mathcal{W}_j) \xrightarrow{\partial} K_{-i}(\mathcal{U}_j) \rightarrow K_{-i}(\mathcal{U}_{j-1}) \oplus K_{-i}(\mathcal{V}_j) \rightarrow \cdots.$$

By induction hypothesis on j , we deduce that $\gamma = \partial(\alpha)$ for some $\alpha \in K_{-i+1}(\mathcal{W}_j)$. By applying the killing lemma (Proposition 5.2.1) to the étale morphism $\mathcal{W}_j \rightarrow \mathcal{U}_j$, we can find a sequence of blow-ups $f : \mathcal{U}'_j \rightarrow \mathcal{U}_j$ with nowhere dense centers such that for the induced map $f_W : \mathcal{W}'_j := \mathcal{U}'_j \times_{\mathcal{U}_j} \mathcal{W}_j \rightarrow \mathcal{W}_j$, $f_W^*(\alpha) = 0$ in $K_{-i+1}(\mathcal{W}'_j)$. Since \mathcal{U}'_j is again a noetherian ANS stack (Lemma A.1.7), we conclude that $f^*(\gamma) = f^*(\partial(\alpha)) = \partial(f_W^*(\alpha)) = 0$. Now as in the first case, by using Theorem 4.4.1 and the induction hypothesis on d , we conclude that $\gamma = 0$. \square

5.4. Weibel's conjecture (II).

Theorem 5.4.1. *Let \mathcal{X} be a noetherian ANS stack of smooth-covering dimension d (see Definition A.4.1). Then for any vector bundle $\pi : \mathcal{E} \rightarrow \mathcal{X}$, the map*

$$\pi^* : K_{-i}(\mathcal{X}) \rightarrow K_{-i}(\mathcal{E})$$

is an isomorphism for all $i \geq d$.

Proof. The zero section induces a retraction of $\pi^* : K(\mathcal{X}) \rightarrow K(\mathcal{E})$, so it is enough to show that π^* is surjective. If \mathcal{F} denotes the cofiber of the morphism $\pi^* : K(-) \rightarrow K(- \times_{\mathcal{X}} \mathcal{E})$ on the small étale site of \mathcal{X} , then it suffices to show that $\mathcal{F}_{-i}(\mathcal{X})$ vanishes for all $i \geq d$. By Proposition 5.1.3, \mathcal{F}_{-i} is nil-invariant for all $i \geq 0$ for any $[X/G]$, where X is an affine scheme and G is an embeddable linearly reductive group scheme. Therefore by Corollary 4.1.2, Theorem A.1.8 and Proposition A.1.9, \mathcal{F}_{-i} is nil-invariant for all $i \geq d$ (arguing as in the proof of Corollary 5.1.4). Thus we can assume that \mathcal{X} is reduced.

Suppose $d = 0$. Then there exists a smooth surjection $u : X \rightarrow \mathcal{X}$ by a 0-dimensional noetherian scheme X . Since \mathcal{X} is reduced, X is also reduced and hence regular. Since perfectness is fppf-local, we find that every cohomologically bounded pseudocoherent complex on \mathcal{X} is perfect (see e.g. the proof of [HK19, Lem. 5.6]), and the same for \mathcal{E} . Thus the map $\pi^* : K(\mathcal{X}) \rightarrow K(\mathcal{E})$ is identified with the inverse image $\pi^* : G(\mathcal{X}) \rightarrow G(\mathcal{E})$, which is invertible by [Kha22, Thm. 3.5]. Now assume $d > 0$ and the statement is known for smooth-covering dimension $< d$.

Case 1: \mathcal{X} has the resolution property. If \mathcal{X} has the resolution property, then by Lemma A.1.7, Theorem A.3.2 and Lemma A.2.4, \mathcal{E} is a perfect stack with the resolution property. By Proposition 5.2.1, for any $\gamma \in K_{-d}(\mathcal{E})$, there exists a sequence of blow-ups $f : \mathcal{X}' \rightarrow \mathcal{X}$ with nowhere dense centers such that γ goes to 0 in $K_{-d}(\mathcal{X}' \times_{\mathcal{X}} \mathcal{E})$. Choose $\mathcal{Z} \subseteq \mathcal{X}$ a nowhere dense closed substack such that f is an isomorphism over $\mathcal{X} \setminus \mathcal{Z}$ and let $\mathcal{Z}' = \mathcal{Z} \times_{\mathcal{X}} \mathcal{X}'$. The n th infinitesimal thickenings $\mathcal{Z}(n)$ satisfy the induction hypothesis by Lemmas A.1.7 and A.2.4. Combining formal proper excision (Theorem 4.4.1) with Remark B.1.6, the isomorphism $K_{-d}(\mathcal{E} \times_{\mathcal{X}} \mathcal{Z}(n)) \simeq K_{-d}(\mathcal{Z}(n)) \simeq 0$ (by induction hypothesis), and Theorem 5.3.1, we get a commutative diagram with exact rows:

$$\begin{array}{ccccc} \{K_{-d+1}(\mathcal{Z}'(n))\}_n & \longrightarrow & K_{-d}(\mathcal{X}) & \longrightarrow & K_{-d}(\mathcal{X}') \\ \downarrow \pi_{\mathcal{Z}'}^* & & \downarrow \pi^* & & \downarrow \pi_{\mathcal{Z}}^* \\ \{K_{-d+1}(\mathcal{E} \times_{\mathcal{X}} \mathcal{Z}'(n))\}_n & \longrightarrow & K_{-d}(\mathcal{E}) & \longrightarrow & K_{-d}(\mathcal{E} \times_{\mathcal{X}} \mathcal{X}') \end{array}$$

Since $\pi_{\mathcal{Z}'}^*$ is also an isomorphism by induction hypothesis, this implies that γ is in the image of π^* .

Case 2: \mathcal{X} is arbitrary. In general, there exists by Theorem A.1.8 and Proposition A.2.5 a finite sequence of open immersions $\emptyset = \mathcal{U}_0 \hookrightarrow \mathcal{U}_1 \hookrightarrow \dots \hookrightarrow \mathcal{U}_n = \mathcal{X}$ and Nisnevich squares

$$\begin{array}{ccc} \mathcal{W}_j & \longrightarrow & \mathcal{V}_j \\ \downarrow & & \downarrow p \\ \mathcal{U}_{j-1} & \longrightarrow & \mathcal{U}_j, \end{array}$$

where each \mathcal{V}_j satisfies the resolution property. We prove by induction on j that for $i \geq d$, $\mathcal{F}_{-i}(\mathcal{U}_j)$ vanishes. For $j = 0$, this follows from Case 1. Assuming the groups $\mathcal{F}_{-i}(\mathcal{U}_{j-1})$ vanish for all $l \geq d$, we will show that $\mathcal{F}_{-i}(\mathcal{U}_j)$ also vanishes.

Choose $\gamma \in \mathcal{F}_{-i}(\mathcal{U}_j)$ for some $i \geq d$. We will show that $\gamma = 0$. By localization for \mathcal{F} , we get an exact sequence of homotopy groups

$$\mathcal{F}_{-i}(\mathcal{U}_j \text{ on } \mathcal{Z}_j) \rightarrow \mathcal{F}_{-i}(\mathcal{U}_j) \rightarrow \mathcal{F}_{-i}(\mathcal{U}_{j-1}),$$

where \mathcal{Z}_j denotes the (reduced) complement of the open substack $\mathcal{U}_{j-1} \subseteq \mathcal{U}_j$. Since $\mathcal{F}_{-i}(\mathcal{U}_{j-1})$ vanishes by induction hypothesis, γ lifts to a class $\mathcal{F}_{-i}(\mathcal{U}_j \text{ on } \mathcal{Z}_j)$ (which we also denote γ). Let $\tilde{\gamma}$ denote its image by $p^* : \mathcal{F}_{-i}(\mathcal{U}_j \text{ on } \mathcal{Z}_j) \rightarrow \mathcal{F}_{-i}(\mathcal{V}_j \text{ on } p^{-1}(\mathcal{Z}_j))$.

By applying Proposition 5.2.1⁵ to $\tilde{\gamma}$ and the smooth morphism $\mathcal{V}_j \rightarrow \mathcal{U}_j$, we can find a sequence of blow-ups $q : \mathcal{U}'_j \rightarrow \mathcal{U}_j$ with nowhere dense centres such that $\tilde{\gamma}$ vanishes after inverse image along the base change $qv : \mathcal{V}'_j := \mathcal{U}'_j \times_{\mathcal{U}_j} \mathcal{V}_j \rightarrow \mathcal{V}_j$. The cartesian square

$$\begin{array}{ccc} \mathcal{V}'_j & \xrightarrow{qv} & \mathcal{V}_j \\ \downarrow & & \downarrow p \\ \mathcal{U}'_j & \xrightarrow{q} & \mathcal{U}_j \end{array}$$

gives rise to a commutative square

$$\begin{array}{ccc} \mathcal{F}_{-i}(\mathcal{U}_j \text{ on } \mathcal{Z}_j) & \longrightarrow & \mathcal{F}_{-i}(\mathcal{V}_j \text{ on } \mathcal{Z}_j) \\ \downarrow q^* & & \downarrow q_v^* \\ \mathcal{F}_{-i}(\mathcal{U}'_j \text{ on } \mathcal{Z}_j) & \longrightarrow & \mathcal{F}_{-i}(\mathcal{V}'_j \text{ on } \mathcal{Z}_j), \end{array}$$

where the horizontal arrows are invertible by excision (and we implicitly base change \mathcal{Z}_j where necessary). Therefore, we get that $q^*(\gamma)$ vanishes in $\mathcal{F}_{-i}(\mathcal{U}'_j \text{ on } \mathcal{Z}_j)$ and in particular in $\mathcal{F}_{-i}(\mathcal{U}'_j)$.

By construction, q is an isomorphism over $\mathcal{U}_j \setminus \mathcal{D}$ for some nowhere dense closed substack $\mathcal{D} \subseteq \mathcal{U}_j$. Let $\mathcal{D}' = \mathcal{D} \times_{\mathcal{U}_j} \mathcal{U}'_j$. Using Theorem 4.4.1, we have an exact sequence of pro-abelian groups

$$\{\mathcal{F}_{-i+1}(\mathcal{D}'(n))\}_n \rightarrow \mathcal{F}_{-i}(\mathcal{U}_j) \rightarrow \mathcal{F}_{-i}(\mathcal{U}'_j) \oplus \{\mathcal{F}_{-i}(\mathcal{D}(n))\}_n,$$

where $\mathcal{F}_{-i+1}(\mathcal{D}'(n))$ and $\mathcal{F}_{-i}(\mathcal{D}(n))$ vanish as they satisfy the induction hypothesis on d (by Lemmas A.1.7 and A.2.4). But since $q^*(\gamma)$ vanishes in $\mathcal{F}_{-i}(\mathcal{U}'_j)$, we have $\gamma = 0$ in $\mathcal{F}_{-i}(\mathcal{U}_j)$ as desired. \square

Remark 5.4.2. The argument in the proof of Theorem 5.4.1 can also be used to generalize Theorem 5.3.1 to any stack \mathcal{X} that is *smooth* and affine over a noetherian ANS stack of fppf-covering dimension d .

APPENDIX A. ALGEBRAIC STACKS

A.1. ANS stacks.

Definition A.1.1. Let G be an affine fppf group scheme over an affine scheme S .

- (i) We say that G is *linearly reductive* if direct image along the morphism $BG \rightarrow S$ is t-exact (i.e., it is cohomologically affine).
- (ii) We say that G is *nice* if it is an extension of a finite étale group scheme, of order prime to the characteristics of S , by a group scheme of multiplicative type.
- (iii) We say that G is *embeddable* if it is a closed subgroup of $\mathrm{GL}_S(\mathcal{E})$ for some finite locally free sheaf \mathcal{E} on S .

Nice group schemes are linearly reductive by [AHR19, Rem. 2.2].

Definition A.1.2. A derived algebraic stack \mathcal{X} is called *ANS* if it has affine diagonal and nice stabilizers.

Example A.1.3. In characteristic zero any reductive group G (such as $\mathrm{GL}_{n,S}$) is linearly reductive. In characteristic $p > 0$, any linearly reductive group is nice [AHR19, Thm. 18.9].

⁵To be precise, we use the statement for \mathcal{F} in place of K , which holds since there is a natural splitting $K(- \times_{\mathcal{X}} \mathcal{E}) \simeq K(-) \oplus \mathcal{F}(-)$.

Example A.1.4. Let G be a finite étale group scheme over a field k . If G has order prime to the characteristic of k , then G is nice and embeddable. It follows that any separated Deligne–Mumford stack over k is ANS as long as it is tame (i.e., has all stabilizers of order prime to the characteristic).

Example A.1.5. Any algebraic stack with affine diagonal that is tame in the sense of [AOV08, Def. 3.1] is ANS. This generalizes Example A.1.4.

Example A.1.6. Tori are embeddable group schemes of multiplicative type (hence nice). Thus if T is a torus over an affine scheme S acting on an algebraic space X over S with affine diagonal, then the quotient $[X/T]$ is ANS. (However, it is typically not tame in the sense of [AOV08].)

Lemma A.1.7. *Let \mathcal{X} be an ANS derived stack. Let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a representable morphism with affine diagonal. Then \mathcal{X}' is ANS.*

Proof. Since f is representable, the stabilizers of \mathcal{X}' are subgroups of those of \mathcal{X} . □

The following is the main result of [AHHLR] in the classical case. The generalization to derived stacks is immediate.

Theorem A.1.8 (Alper–Hall–Halpern–Leistner–Rydh). *Let \mathcal{X} be an ANS derived stack. Then there exists a finite sequence of open immersions*

$$\emptyset = \mathcal{U}_0 \hookrightarrow \mathcal{U}_1 \hookrightarrow \dots \hookrightarrow \mathcal{U}_n = \mathcal{X},$$

an embeddable nice group scheme G over an affine scheme S , and Nisnevich squares

$$\begin{array}{ccc} \mathcal{W}_i & \longrightarrow & \mathcal{V}_i \\ \downarrow & & \downarrow \\ \mathcal{U}_{i-1} & \longrightarrow & \mathcal{U}_i \end{array}$$

where \mathcal{V}_i is étale and affine over \mathcal{U}_i and quasi-affine over BG .

Proof. This follows by combining [AHHLR, Thm. 6.3]⁶ with [HK19, Prop. 2.9]. See [KR21, Thm. 2.12] for details. □

Proposition A.1.9. *Let $\mathcal{X} = [X/G]$ be the quotient of a quasi-compact separated derived algebraic space X with action of a nice group scheme G over an affine scheme S . Then \mathcal{X} admits a scallop decomposition of the form $(\mathcal{U}_i, \mathcal{V}_i, u_i)_i$, where \mathcal{V}_i is of the form $[V_i/G]$ for some affine derived schemes V_i over S with G -action, and u_i is an affine morphism for each i .*

Proof. By generalized Sumihiro (see [KR21, Theorem 2.14]), \mathcal{X} admits an affine Nisnevich cover $u : \mathcal{V} \rightarrow \mathcal{X}$ where \mathcal{V} is of the form $[V/G]$ with V an affine scheme over S with G -action. The desired scallop decomposition is obtained by a G -equivariant version of the construction in the proof of [RG71, Lem. 5.7.5] or [SAG, Prop. 3.2.2.4], which goes through *mutatis mutandis*:

For every $i \geq 0$, define $\mathcal{U}^i \subseteq \mathcal{X}$ as the substack of points where the fibre of u has $\geq i$ geometric points. We have $\mathcal{U}^1 = \mathcal{X}$ (since u is surjective) and $\mathcal{U}^{n+1} = \emptyset$ for some large enough n (since \mathcal{X} is quasi-compact). This gives a finite filtration of \mathcal{X} by quasi-compact opens $\mathcal{U}_i := \mathcal{U}^{n+1-i}$.

Consider the fibre powers V^i of V over X and $\mathcal{V}^i = [V^i/G]$ of \mathcal{V} over \mathcal{X} , respectively. Since u is affine, so is each V^i . Since $V \rightarrow X$ is affine and étale, the “big diagonal” $\Delta^i \subseteq V^i$ is an open and

⁶See [AHR19, Cor. 17.3] and [AOV08, Thm. 3.2] for documented special cases of this result.

closed subscheme. The permutation action of the symmetric group Σ_i on V^i is free away from Δ^i , and commutes with the factorwise G -action on V^i . Thus we can write

$$\mathcal{V}_i := [(V^i \setminus \Delta^i)/G \times \Sigma_i] \simeq [W_i/G],$$

where $W_i = [(V^i \setminus \Delta^i)/\Sigma_i]$, as a quotient of an affine scheme by a free action of a finite group, is an affine scheme. Now one checks, exactly as in [SAG, Prop. 3.2.2.4], that the canonical morphisms $\mathcal{V}_i \rightarrow \mathcal{X}$ factor through affine étale morphisms $u_i : \mathcal{V}_i \rightarrow \mathcal{U}_i$, and that the resulting construction $(\mathcal{U}_i, \mathcal{V}_i, u_i)_i$ is indeed a scallop decomposition. \square

A.2. The resolution property.

Definition A.2.1. Let \mathcal{X} be a derived algebraic stack. We say that \mathcal{X} has the *resolution property* if for every discrete coherent sheaf \mathcal{F} of finite type on \mathcal{X} , there exists a finite locally free sheaf \mathcal{E} and a surjection $\mathcal{E} \twoheadrightarrow \mathcal{F}$.

The following construction is one of the pleasant consequences of the resolution property.

Construction A.2.2. Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a closed immersion of derived stacks. If i is almost of finite presentation (e.g. \mathcal{X} is noetherian), then the ideal $\mathcal{J} \subseteq \pi_0(\mathcal{O}_{\mathcal{X}})$ defining \mathcal{Z}_{cl} in \mathcal{X}_{cl} is of finite type. Thus if \mathcal{X} admits the resolution property, there exists a surjection $\mathcal{E} \twoheadrightarrow \mathcal{J}$ from a finite locally free sheaf \mathcal{E} on \mathcal{X} . The induced morphism $s : \mathcal{E} \rightarrow \mathcal{J} \rightarrow \mathcal{O}_{\mathcal{X}}$ can be viewed as a section of the vector bundle

$$\mathbf{V}_{\mathcal{X}}(\mathcal{E}) = \text{Spec}_{\mathcal{X}}(\text{Sym}_{\mathcal{O}_{\mathcal{X}}}(\mathcal{E})),$$

and its derived zero locus defines a quasi-smooth closed immersion $\tilde{i} : \tilde{\mathcal{Z}} \rightarrow \mathcal{X}$ whose 0-truncation is $i : \mathcal{Z} \rightarrow \mathcal{X}$ and which fits in a homotopy cartesian square

$$\begin{array}{ccc} \tilde{\mathcal{Z}} & \xrightarrow{\tilde{i}} & \mathcal{X} \\ \downarrow & & \downarrow s \\ \mathcal{X} & \xrightarrow{0} & \mathbf{V}_{\mathcal{X}}(\mathcal{E}). \end{array}$$

Repeating this construction with the section $s^{\otimes n}$, for any $n > 0$, gives a tower of infinitesimal thickenings

$$\mathcal{Z} \hookrightarrow \tilde{\mathcal{Z}} = \tilde{\mathcal{Z}}(1) \hookrightarrow \tilde{\mathcal{Z}}(2) \hookrightarrow \dots$$

Remark A.2.3. If \mathcal{X} is a derived stack with the resolution property, then any quasi-smooth closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ fits in a homotopy cartesian square

$$\begin{array}{ccc} \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \\ \downarrow & & \downarrow s \\ \mathcal{X} & \xrightarrow{0} & \mathbf{V}_{\mathcal{X}}(\mathcal{E}) \end{array}$$

where \mathcal{E} is a finite locally free sheaf on \mathcal{X} . This follows by a variant of the proof of [KR18a, Prop. 2.3.8].

We discuss some examples of derived stacks with the resolution property. First, recall that in the classical setting, the property is stable under affine morphisms:

Lemma A.2.4. *Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a quasi-affine morphism of (classical) algebraic stacks. If \mathcal{Y} has the resolution property, then so does \mathcal{X} .*

Proof. See [HR17, Lem. 7.1] or [Gro17, Prop. 1.8(v)]. \square

The classifying stack BG has the resolution property for embeddable linearly reductive group schemes G , so one finds that affine quotient stacks $[X/G]$ also admit the resolution property (see [AHR19, Rmk. 2.5]). We prove the following derived generalization of this statement:

Proposition A.2.5. *Let S be an affine scheme, G an embeddable linearly reductive group scheme over S , and X a derived affine S -scheme with G -action. Then the derived stack $\mathcal{X} = [X/G]$ admits the resolution property.*

The proof of Proposition A.2.5 will require the following lemma. We write $\mathbf{D}_{\text{lfr}}(\mathcal{Y})$ for the full subcategory of $\mathbf{D}(\mathcal{Y})$ spanned by the finite locally free sheaves, for any derived algebraic stack \mathcal{Y} .

Lemma A.2.6. *Let the notation be as in Proposition A.2.5. For any integer $n \geq 0$, let $i : \tau_{\leq n}(\mathcal{X}) \rightarrow \mathcal{X}$ be the inclusion of the n -truncation. Then we have:*

- (i) *For any finite locally free $\mathcal{E} \in \mathbf{D}_{\text{lfr}}(\mathcal{X})$ and any $\mathcal{F} \in \mathbf{D}(\mathcal{X})$, the canonical map of n -truncated spaces*

$$\tau_{\leq n} \text{Maps}(\mathcal{E}, \mathcal{F}) \rightarrow \text{Maps}(\tau_{\leq n}(\mathcal{E}), \tau_{\leq n}(\mathcal{F}))$$

is invertible.

- (ii) *The induced functor of $(n+1)$ -categories*

$$i^* : \tau_{\leq n+1} \mathbf{D}_{\text{lfr}}(\mathcal{X}) \rightarrow \mathbf{D}_{\text{lfr}}(\tau_{\leq n}(\mathcal{X}))$$

is an equivalence.

In particular, the functor of ordinary categories

$$\mathbf{h} \mathbf{D}_{\text{lfr}}(\mathcal{X}) \rightarrow \mathbf{D}_{\text{lfr}}(\mathcal{X}_{\text{cl}})$$

is an equivalence (where \mathbf{h} denotes the homotopy category).

Proof. Note that the map in the first claim is induced by the morphism in $\mathbf{D}(\mathcal{X})$

$$\tau_{\leq n} \underline{\text{Hom}}_{\mathcal{O}_{\mathcal{X}}}(\mathcal{E}, \mathcal{F}) \rightarrow i_* \underline{\text{Hom}}_{\mathcal{O}_{\tau_{\leq n}(\mathcal{X})}}(\tau_{\leq n}(\mathcal{E}), \tau_{\leq n}(\mathcal{F})), \quad (\text{A.2.a})$$

where $\underline{\text{Hom}}$ denotes the internal Hom, by applying in succession the functors of direct image along $f : \mathcal{X} \rightarrow BG$ (which is t-exact since f is affine), direct image along $BG \rightarrow S$ (which is t-exact since G is linearly reductive), and (derived) global sections (which is t-exact since S is affine). Therefore it will suffice to show that (A.2.a) is invertible. By fpqc descent, this can be checked after inverse image along the smooth surjection $X \rightarrow \mathcal{X}$. Since i is representable, i_* satisfies base change and we are thus reduced to the affine case, which is well-known (see e.g. [Kha17, Claim 4.3]).

Consider now claim (ii). By (i) the functor in question is fully faithful (on finite locally frees). For essentially surjectivity, we may assume $n = 0$ (so that $\tau_{\leq n}(\mathcal{X}) = \mathcal{X}_{\text{cl}}$). Since BG has the resolution property [AHR19, Rmk. 2.5], Lemma A.2.4 implies that for every finite locally free sheaf $\mathcal{E} \in \mathbf{D}(\mathcal{X}_{\text{cl}})$, there exists a finite locally free sheaf $\mathcal{F} \in \mathbf{D}(BG)$ and a surjection $g^*(\mathcal{F}) \rightarrow \mathcal{E}$ where $g : \mathcal{X}_{\text{cl}} \rightarrow BG$. Certainly $g^*(\mathcal{F}) \in \mathbf{D}(\mathcal{X}_{\text{cl}})$ lifts to $f^*(\mathcal{F}) \in \mathbf{D}(\mathcal{X})$, so we are reduced to show that if $\mathcal{E} \twoheadrightarrow \mathcal{F}$ is surjection of locally free sheaves on \mathcal{X}_{cl} and $\mathcal{E} = i^* \tilde{\mathcal{E}}$ for some locally free $\tilde{\mathcal{E}} \in \mathbf{D}(\mathcal{X})$, then \mathcal{F} also extends to a locally free sheaf $\tilde{\mathcal{F}}$ on \mathcal{X} . Since G is linearly reductive, \mathcal{F} is projective by [Hoy17, Lem. 2.17] and the surjection $\mathcal{E} \twoheadrightarrow \mathcal{F}$ splits. The resulting map $e_0 : \mathcal{E} \twoheadrightarrow \mathcal{F} \rightarrow \mathcal{E}$ is an idempotent with image \mathcal{F} . By claim (i), we can extend e_0 to an idempotent endomorphism e of $\tilde{\mathcal{E}}$. If we set

$$\begin{aligned} \tilde{\mathcal{F}} &:= \varinjlim (\tilde{\mathcal{E}} \xrightarrow{e} \tilde{\mathcal{E}} \xrightarrow{e} \dots), \\ \tilde{\mathcal{F}}_1 &:= \varinjlim (\tilde{\mathcal{E}} \xrightarrow{\text{id}-e} \tilde{\mathcal{E}} \xrightarrow{\text{id}-e} \dots), \end{aligned}$$

then the induced morphism $\phi : \widetilde{\mathcal{E}} \rightarrow \widetilde{\mathcal{F}} \oplus \widetilde{\mathcal{F}}_1$ is invertible. Indeed, by fpqc descent we may pull back to X and thereby assume \mathcal{X} is affine, in which case it is clear that ϕ is an isomorphism on homotopy groups. In particular, it follows that $\widetilde{\mathcal{F}}$ is locally free, and clearly $i^*(\widetilde{\mathcal{F}}) \simeq \mathcal{F}$ by construction. \square

Proof of Proposition A.2.5. By [AHR19, Rmk. 2.5], the classical truncation $[X_{\text{cl}}/G]$ has the resolution property. Hence the claim follows from Lemma A.2.6. \square

A.3. Compact generation.

Definition A.3.1. A derived algebraic stack \mathcal{X} is *perfect* if the stable ∞ -category $\mathbf{D}(\mathcal{X})$ is compactly generated by its full subcategory $\text{Perf}(\mathcal{X})$ of perfect complexes.

In this subsection we prove the following result, which is a derived generalization of [AHR19, Prop. 14.1].

Theorem A.3.2. *Let \mathcal{X} be an ANS derived stack. Then \mathcal{X} is perfect.*

Lemma A.3.3. *Let G be an embeddable linearly reductive group scheme over an affine scheme S . Then the classifying stack BG is perfect. Moreover, $\mathbf{D}(BG)$ is compactly generated by finite representations of G (i.e., finite projectives).*

Proof. Since G is affine, BG has affine diagonal. By Proposition A.2.5, BG has the resolution property. Hence the claim follows from [HR17, Prop. 8.4]. \square

The following shows that, for nice enough quotients of affine derived schemes, the derived ∞ -category is not only compactly generated, but projectively generated in the sense of Definition 3.5.6.

Proposition A.3.4. *Let G be an embeddable linearly reductive group scheme over an affine scheme S . Then for any affine derived scheme X with G -action, the quotient stack $[X/G]$ is perfect and even crisp in the sense of [HR17]. Moreover, if $p : [X/G] \rightarrow BG$ is the projection, then the collection of objects $\{p^*(\mathcal{E})\}$ forms a small set of compact projective generators for $\mathbf{D}([X/G])_{\geq 0}$, as $\mathcal{E} \in \mathbf{D}(BG)$ varies over finite locally free G -modules on S .*

Proof. Since p is affine, the functor $p^* : \mathbf{D}(BG) \rightarrow \mathbf{D}([X/G])$ is compact and generates its codomain under colimits (2.2.1). This already implies that $\mathbf{D}([X/G])$ is compactly generated by the objects of the stated form.

We now demonstrate the stronger property of crispness. By Proposition A.2.5, $[X/G]$ admits the resolution property. Since $[X/G]$ has affine diagonal, [HR17, Prop. 8.4] then implies that $[X/G]$ is crisp. Note that *loc. cit.* only discusses classical stacks, but we only need the argument from the last paragraph of the proof, which immediately generalizes to the derived setting.

Finally let us show that the object $p^*(\mathcal{E}) \in \mathbf{D}([X/G])_{\geq 0}$ is projective for every finite representation $\mathcal{E} \in \mathbf{D}(BG)$. By [HA, Lem. 7.2.2.6] it will suffice to show that the functor $\mathcal{M}aps(p^*(\mathcal{E}), -) : \mathbf{D}(BG) \rightarrow \text{Spt}$ is t-exact, where $\mathcal{M}aps(-, -)$ denotes the mapping spectrum functor in the stable ∞ -category $\mathbf{D}(BG)$. We have a canonical isomorphism

$$\mathcal{M}aps(p^*(\mathcal{E}), -) \simeq \Gamma(S, (p_* \underline{\text{Hom}}(\mathcal{E}, -))^G).$$

Note that $\underline{\text{Hom}}(\mathcal{E}, -)$ is t-exact because \mathcal{E} is projective in $\mathbf{D}(S)$, p_* is t-exact since p is affine, the G -invariants functor $(-)^G$ is identified with the direct image functor $\mathbf{D}(BG) \rightarrow \mathbf{D}(S)$ and

hence is t-exact because G is linearly reductive, and the (derived) global sections functor $\Gamma(S, -)$ is t-exact since S is affine. \square

Proof of Theorem A.3.2. By Theorem A.1.8, we can in particular find an affine étale surjection onto \mathcal{X} from a finite coproduct of quotient stacks of the form $[X/G]$, where G is a nice group scheme over an affine scheme S and X is an affine derived S -scheme with G -action. It will now suffice to show that the property of crispness can be detected by affine étale surjections. Indeed, we note that the proof given in [HR17, Thm. C] for the classical case generalizes to our setting, following Example 9.4 of *loc. cit.* \square

A.4. Dimension of algebraic stacks. Recall several useful notions of dimension from [HK19].

Definition A.4.1. Let \mathcal{X} be a noetherian stack.

- (i) The *Krull dimension* $\dim(\mathcal{X})$ of \mathcal{X} is the dimension of the underlying topological space $|\mathcal{X}|$.
- (ii) The *blow-up dimension* $\text{bl dim}(\mathcal{X})$ is the supremum over integers $n \geq 0$ for which there exists a sequence of maps

$$\mathcal{X}_n \rightarrow \cdots \rightarrow \mathcal{X}_0 = \mathcal{X}$$

such that \mathcal{X}_i is a nonempty nowhere dense closed substack in an iterated blow-up of \mathcal{X}_{i-1} for all $i > 0$. If \mathcal{X} is empty, then $\text{bl dim}(\mathcal{X}) = -1$ by convention.

- (iii) The *fppf-covering dimension* $\text{cov dim}_{\text{fppf}}(\mathcal{X})$ is the minimal integer $-1 \leq n \leq \infty$ such that there exists an fppf morphism $X \rightarrow \mathcal{X}$ where X is a noetherian scheme of Krull dimension n .
- (iv) The *covering dimension* $\text{cov dim}_{\text{sm}}(\mathcal{X})$ (more precisely, *smooth-covering dimension*) is the minimal integer $-1 \leq n \leq \infty$ such that there exists a smooth surjection $X \rightarrow \mathcal{X}$ where X is a noetherian scheme of Krull dimension n .

Remark A.4.2. In general, one has the inequalities $\dim(\mathcal{X}) \leq \text{bl dim}(\mathcal{X}) \leq \text{cov dim}_{\text{fppf}}(\mathcal{X}) \leq \text{cov dim}_{\text{sm}}(\mathcal{X})$. For quasi-Deligne–Mumford stacks these are all equal to the usual dimension as defined in [Stacks, Tag 0AFL]. See [HK19, Lemma 7.8].

Example A.4.3. Let G be an fppf group scheme over an algebraic space S . If G acts on an algebraic space X over S , then the quotient stack $\mathcal{X} = [X/G]$ is of fppf-covering dimension $\leq \dim(X)$.

In this subsection we will show that the Nisnevich cohomological dimension of an algebraic stack \mathcal{X} , which we denote by $\text{cd}_{\text{Nis}}(\mathcal{X})$, is bounded by its fppf-covering dimension:

Proposition A.4.4. *Let \mathcal{X} be a noetherian stack and let \mathcal{F} be a sheaf of abelian groups on the Nisnevich site of \mathcal{X} . Then*

$$\mathbf{H}_{\text{Nis}}^i(\mathcal{X}, \mathcal{F}) = 0$$

for all $i > \text{cov dim}_{\text{fppf}}(\mathcal{X})$. In other words, $\text{cd}_{\text{Nis}}(\mathcal{X}) \leq \text{cov dim}_{\text{fppf}}(\mathcal{X})$.

Recall that the Nisnevich topology on the category of noetherian algebraic stacks is generated by a cd-structure [HK19, Sect. 2C], which is clearly complete and regular. To establish Proposition A.4.4 we will show that this cd-structure is bounded with respect to a density structure.

Construction A.4.5. For any noetherian algebraic stack \mathcal{X} , let $\text{Stk}_{/\mathcal{X}}^{\text{DM}}$ be the category of algebraic stacks \mathcal{Y} over \mathcal{X} for which the structural morphism $\mathcal{Y} \rightarrow \mathcal{X}$ is representable by Deligne–Mumford stacks. Choose an fppf covering $\mathcal{S} \rightarrow \mathcal{X}$ from a Deligne–Mumford stack \mathcal{S} . We define a density structure $D_i^{\mathcal{S}}(-)$ in the sense of [Voe10a, Definition 2.20] on the category $\text{Stk}_{/\mathcal{X}}^{\text{DM}}$. For

$\mathcal{Y} \in \text{Stk}_{/\mathcal{X}}^{\text{DM}}$, let $q : \mathcal{S}_{\mathcal{Y}} \rightarrow \mathcal{Y}$ be the fppf covering of \mathcal{Y} given by the base change of $\mathcal{S} \rightarrow \mathcal{X}$ along \mathcal{Y} . For $i \geq 0$, let $D_i^{\mathcal{S}}(\mathcal{Y})$ denote the class of open substacks $\mathcal{U} \rightarrow \mathcal{Y}$ such that the open substack $\mathcal{U} \times_{\mathcal{Y}} \mathcal{S}_{\mathcal{Y}} \rightarrow \mathcal{S}_{\mathcal{Y}}$ defines an element of the class $D_i(\mathcal{S}_{\mathcal{Y}})$, where $D_*(-)$ denotes the density structure on the category of Deligne–Mumford stacks defined in [KØ12, Definition 4.4]. It is easy to check that the classes $D_i^{\mathcal{S}}(-)$ define a density structure which is locally of finite dimension and the dimension of \mathcal{X} with respect to the density structure is equal to the Krull dimension $\dim(\mathcal{S})$.

Recall that for a Deligne–Mumford stack \mathcal{X} and $i \geq 0$, $D_i(\mathcal{X})$ is the collection of open substacks $\mathcal{U} \rightarrow \mathcal{X}$ such that for every irreducible, reduced closed substack \mathcal{Z} of \mathcal{X} with $\mathcal{Z} \times_{\mathcal{X}} \mathcal{U}$ the empty stack, there exists a sequence $\mathcal{Z} = \mathcal{Z}_0 \not\subset \mathcal{Z}_1 \not\subset \dots \not\subset \mathcal{Z}_i$ of irreducible, reduced closed substacks of \mathcal{X} .

Lemma A.4.6. *Let \mathcal{X} be a Deligne–Mumford stack and $\mathcal{U} \in D_i(\mathcal{X})$. For any open substack \mathcal{V} of \mathcal{X} , we have $\mathcal{U} \cap \mathcal{V} \in D_i(\mathcal{V})$.*

Proof. By [KØ12, Lemma 4.5] it suffices to show that $|\mathcal{U} \cap \mathcal{V}| \in D_i(|\mathcal{V}|)$. Now we can use the same argument as in the proof of [Voe10b, Lemma 2.5]. \square

Proposition A.4.7. *The Nisnevich cd-structure on the category $\text{Stk}_{/\mathcal{X}}^{\text{DM}}$ is bounded with respect to the density structure $D_*^{\mathcal{S}}(-)$.*

Proof. We need to show that every Nisnevich square is reducing with respect to the density structure. Consider a Nisnevich square Q in $\text{Stk}_{/\mathcal{X}}^{\text{DM}}$:

$$\begin{array}{ccc}
 \mathcal{W} & \xleftarrow{j_{\mathcal{V}}} & \mathcal{V} \\
 \downarrow & & \downarrow p \\
 \mathcal{U} & \xleftarrow{j} & \mathcal{Y},
 \end{array} \tag{A.4.a}$$

where p is étale, j is an open immersion and the induced morphism $p^{-1}(\mathcal{Y} \setminus \mathcal{U}) \rightarrow (\mathcal{Y} \setminus \mathcal{U})$ is invertible. Choose $\mathcal{W}_0 \in D_{i-1}^{\mathcal{S}}(\mathcal{W})$, $\mathcal{U}_0 \in D_i^{\mathcal{S}}(\mathcal{U})$ and $\mathcal{V}_0 \in D_i^{\mathcal{S}}(\mathcal{V})$. To show that the above square Q is reducing with respect to the density structure, we need to prove that there exists a Nisnevich square Q' in $\text{Stk}_{/\mathcal{X}}^{\text{DM}}$

$$\begin{array}{ccc}
 \mathcal{W}' & \xleftarrow{\quad} & \mathcal{V}' \\
 \downarrow & & \downarrow \\
 \mathcal{U}' & \xleftarrow{\quad} & \mathcal{Y}',
 \end{array} \tag{A.4.b}$$

and a morphism $Q' \rightarrow Q$ such that $\mathcal{W}' \rightarrow \mathcal{W}$ factors through $\mathcal{W}' \rightarrow \mathcal{W}_0$, $\mathcal{U}' \rightarrow \mathcal{U}$ through $\mathcal{U}' \rightarrow \mathcal{U}_0$, $\mathcal{V}' \rightarrow \mathcal{V}$ through $\mathcal{V}' \rightarrow \mathcal{V}_0$, and $\mathcal{Y}' \in D_i^{\mathcal{S}}(\mathcal{Y})$ (see [Voe10a, Definition 2.21]). Applying Lemma A.4.8 to the morphism $j \amalg p$, we can find $\mathcal{Y}_0 \in D_i^{\mathcal{S}}(\mathcal{Y})$ such that $j^{-1}(\mathcal{Y}_0) \subseteq \mathcal{U}_0$ and $p^{-1}(\mathcal{Y}_0) \subseteq \mathcal{V}_0$. Therefore by base changing (A.4.a) along $\mathcal{Y}_0 \hookrightarrow \mathcal{Y}$ and then replacing \mathcal{Y} by \mathcal{Y}_0 we are reduced to the case when $\mathcal{U} = \mathcal{U}_0$ and $\mathcal{V} = \mathcal{V}_0$ in (A.4.a). Note that $\mathcal{W}_0 \times_{\mathcal{Y}} \mathcal{Y}_0$ is in $D_{i-1}^{\mathcal{S}}(\mathcal{U}_0 \times_{\mathcal{Y}} \mathcal{V}_0)$ by Lemma A.4.6.

Let $\mathcal{Z} = \mathcal{W} \setminus \mathcal{W}_0$ and $\mathcal{C} = \mathcal{Y} \setminus \mathcal{U}$ and set $\mathcal{W}' = \mathcal{W}_0$, $\mathcal{U}' = \mathcal{U}$, $\mathcal{V}' = \mathcal{V} \setminus \text{cl}_{\mathcal{V}}(\mathcal{Z})$ and $\mathcal{Y}' = \mathcal{Y} \setminus (\mathcal{C} \cap \text{cl}_{\mathcal{Y}}(p \circ j_{\mathcal{V}}(\mathcal{Z})))$ in (A.4.b) to obtain the Nisnevich square Q' with a natural morphism to Q given by inclusions. Now $\mathcal{Y}' \times_{\mathcal{Y}} \mathcal{S}_{\mathcal{Y}} \rightarrow \mathcal{S}_{\mathcal{Y}} \in D_i(\mathcal{S}_{\mathcal{Y}})$ by the proof of [KØ12, Proposition 4.9]. Therefore $Q' \rightarrow Q$ satisfies all the required properties. \square

Lemma A.4.8. *Let $f : \mathcal{W} \rightarrow \mathcal{Y}$ be an étale surjection of noetherian stacks. Then for any $i \geq 0$ and $\mathcal{W}_0 \in D_i^{\mathcal{S}}(\mathcal{W})$ there exists $\mathcal{Y}_0 \in D_i^{\mathcal{S}}(\mathcal{Y})$ such that $f^{-1}(\mathcal{Y}_0) \subset \mathcal{W}_0$.*

Proof. Let $\mathcal{S}_{\mathcal{W}} = \mathcal{S} \times_{\mathcal{X}} \mathcal{W}$, $\mathcal{S}_{\mathcal{W}_0} = \mathcal{S} \times_{\mathcal{X}} \mathcal{W}_0$ and let $f_S : \mathcal{S}_{\mathcal{W}} \rightarrow \mathcal{S}_{\mathcal{Y}}$ denote the base change of $f : \mathcal{W} \rightarrow \mathcal{Y}$ along the fppf covering $q : \mathcal{S}_{\mathcal{Y}} \rightarrow \mathcal{Y}$ and $\tilde{q} : \mathcal{S}_{\mathcal{W}} \rightarrow \mathcal{W}$ denote the base change of q along f . Then by [KØ12, Lemma 4.7] applied to f_S there exists $\tilde{\mathcal{Y}} \in D_i(\mathcal{S}_{\mathcal{Y}})$ such that $f_S^{-1}(\tilde{\mathcal{Y}}) \subset \mathcal{S}_{\mathcal{W}_0}$. Let $\mathcal{Y}_0 = q(\tilde{\mathcal{Y}})$, then $\mathcal{Y}_0 \in D_i(\mathcal{Y})$ since q is open (see [LMB18, Proposition 5.6]), $\tilde{\mathcal{Y}} \subset \mathcal{Y}_0 \times_{\mathcal{Y}} \mathcal{S}_{\mathcal{Y}}$ and $\tilde{\mathcal{Y}} \in D_i(\mathcal{S}_{\mathcal{Y}})$. Moreover $f^{-1}(\mathcal{Y}_0) = f^{-1}(q(\tilde{\mathcal{Y}})) \subset \tilde{q}(f_S^{-1}(\tilde{\mathcal{Y}})) \subset \tilde{q}(\mathcal{S}_{\mathcal{W}_0}) = \mathcal{W}_0$. \square

APPENDIX B. FORMAL STACKS

B.1. Formal completion. We briefly review some elements of formal derived algebraic geometry. Good references are [HLP14, Sect. 2.1] and [GR14]. The following definition is [HLP14, Def. 2.1.1].

Definition B.1.1 (Formal completion). Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a closed immersion of derived stacks with quasi-compact open complement. The *formal completion* of \mathcal{X} in \mathcal{Z} is the derived prestack $\mathcal{X}_{\mathcal{Z}}^{\wedge}$ whose R -points, for any derived commutative ring R , are the R -points $x : \mathrm{Spec}(R) \rightarrow \mathcal{X}$ which factor *set-theoretically* through the underlying topological space $|\mathcal{Z}| \subseteq |\mathcal{X}|$. By definition, $i : \mathcal{Z} \rightarrow \mathcal{X}$ factors as

$$i : \mathcal{Z} \rightarrow \mathcal{X}_{\mathcal{Z}}^{\wedge} \xrightarrow{\hat{i}} \mathcal{X},$$

where the first arrow induces an isomorphism on reductions, and the second arrow is a monomorphism. When there is no risk of ambiguity, we will write simply \mathcal{X}^{\wedge} for $\mathcal{X}_{\mathcal{Z}}^{\wedge}$.

Remark B.1.2. Note that the formal completion $\mathcal{X}_{\mathcal{Z}}^{\wedge}$ only depends on the underlying topological space $|\mathcal{Z}|$. In particular, if we have a commutative square

$$\begin{array}{ccc} \mathcal{Z}' & \xrightarrow{i'} & \mathcal{X}' \\ \downarrow g & & \downarrow f \\ \mathcal{Z} & \xrightarrow{i} & \mathcal{X} \end{array}$$

where $|\mathcal{Z}'|$ is the set-theoretic inverse image $f^{-1}(|\mathcal{Z}|)$, then the formal completion \mathcal{X}'^{\wedge} is the derived base change of \mathcal{X}^{\wedge} . That is, we have a homotopy cartesian square

$$\begin{array}{ccc} \mathcal{X}'_{\mathcal{Z}'}^{\wedge} & \hookrightarrow & \mathcal{X}' \\ \downarrow f^{\wedge} & & \downarrow f \\ \mathcal{X}_{\mathcal{Z}}^{\wedge} & \hookrightarrow & \mathcal{X}. \end{array}$$

Remark B.1.3. The formal completion of a derived algebraic stack \mathcal{X} along a closed immersion $i : \mathcal{Z} \rightarrow \mathcal{X}$ is always an *ind-algebraic stack*. More precisely, one has the following canonical isomorphism (see [GR14, Prop. 6.5.5]):

$$\{\tilde{\mathcal{Z}}\}_{\tilde{\mathcal{Z}} \rightarrow \mathcal{X}} \rightarrow \mathcal{X}_{\mathcal{Z}}^{\wedge},$$

where the source is the ind-system indexed by the filtered ∞ -category of closed immersions $\tilde{\mathcal{Z}} \rightarrow \mathcal{X}$ that induce an isomorphism $\tilde{\mathcal{Z}}_{\mathrm{red}} \simeq \mathcal{Z}_{\mathrm{red}}$ on reductions. Note that the transition morphisms are surjective closed immersions.

In the affine case, we can give the following more familiar (but less canonical) description.

Example B.1.4. Let A be a derived commutative ring and $I \subseteq \pi_0(A)$ an ideal, and consider the formal completion of $\mathrm{Spec}(A)^{\wedge}$ in the vanishing locus of I . Choosing generators f_1, \dots, f_m for the ideal I , there is an equivalence

$$\mathrm{Spec}(A)^{\wedge} \simeq \{\mathrm{Spec}(A//\langle f_1^n, \dots, f_m^n \rangle)\}_n.$$

See [SAG, Proof of Prop. 8.1.2.1] or [HLP14, Prop. 2.1.2].

If A is an ordinary commutative ring and is noetherian, then we recover the classical formal completion, i.e., the formal spectrum of A as an I -adic ring:

$$\mathrm{Spec}(A)^\wedge \simeq \{\mathrm{Spec}(A/(f_1^n, \dots, f_m^n))\}_n.$$

See [SAG, Lem. 17.3.5.7], [HLP14, Prop. 2.1.4], or [GR14, Proof of Prop. 6.8.2].

More generally in the presence of the resolution property, we have:

Lemma B.1.5. *Let $i : \mathcal{Z} \rightarrow \mathcal{X}$ be a closed immersion almost of finite presentation between derived algebraic stacks for which \mathcal{X} has the resolution property. Let $\tilde{\mathcal{Z}}(n)$ be as in Construction A.2.2. Then there is an isomorphism of ind-stacks*

$$\mathcal{X}_{\mathcal{Z}}^\wedge \simeq \{\tilde{\mathcal{Z}}(n)\}.$$

Proof. This is a simple cofinality argument using the description of the formal completion given in Remark B.1.3. \square

Remark B.1.6. If \mathcal{X} is a *classical* stack, then the formal completion (in any closed substack) in the sense of Definition B.1.1 coincides with the classical formal completion, as long as \mathcal{X} is *noetherian*. This follows from Example B.1.4, see [HLP14, Cor. 2.1.5].

For the reader's convenience, we spell out Lemma B.1.5 in the equivariant (quotient stack) case.

Construction B.1.7. Let G be a group scheme over a commutative ring R , and let A be a derived commutative R -algebra with G -action. Let M be a locally free G -equivariant A -module and $s : M \rightarrow A$ a homomorphism of G -equivariant A -modules. The *derived quotient* of A by s is the G -equivariant A -algebra formed by attaching a cell $s \simeq 0$, i.e., by the cocartesian square in G -equivariant derived commutative rings

$$\begin{array}{ccc} \mathrm{Sym}_A(M) & \xrightarrow{0} & A \\ \downarrow s & & \downarrow \\ A & \longrightarrow & A//s, \end{array}$$

where the map 0 is induced by adjunction from the zero map $M \rightarrow A$, and similarly for s .

Similarly, for any $n > 0$, we also write $A//s^n$ for the same construction where s is replaced by $s^{\otimes n} : M^{\otimes n} \rightarrow A^{\otimes n} \simeq A$ (the n -fold derived tensor product taken over A).

Lemma B.1.8. *Let G be a group scheme over a commutative ring R and $A \twoheadrightarrow B$ a homomorphism of G -equivariant derived commutative R -algebras which is surjective on π_0 . Assume that the quotient stack $[\mathrm{Spec}(\pi_0(A))/G]$ admits the resolution property, so that there exists a locally free G -equivariant A -module M and $s : M \rightarrow A$ whose image on π_0 is equal to the kernel of $\pi_0(A) \rightarrow \pi_0(B)$. Then we have the following presentation of the formal completion:*

$$[\mathrm{Spec}(A)/G]^\wedge \simeq \{[\mathrm{Spec}(A//s^n)/G]\}_{n>0}.$$

Lemma B.1.9. *Let G be a group scheme over a commutative ring R and $A \twoheadrightarrow B$ a surjective homomorphism of G -equivariant commutative R -algebras with kernel I . Assume that the quotient stack $[\mathrm{Spec}(A)/G]$ admits the resolution property, so that there exists a locally free G -equivariant A -module M and $s : M \rightarrow A$ whose image is I . Then we have the following presentation of the formal completion:*

$$[\mathrm{Spec}(A)/G]^\wedge \simeq \{[\mathrm{Spec}(A/I^n)/G]\}_{n>0}.$$

B.2. Quasi-coherent sheaves. For formal stacks (or more generally ind-stacks) such as the formal completion, there is a natural pro- ∞ -categorical refinement of the stable ∞ -category of quasi-coherent sheaves.

Construction B.2.1. If \mathcal{C} denotes the ∞ -category of derived algebraic stacks and representable morphisms, then by Remark 2.2.1 the assignment $\mathcal{X} \mapsto \mathbf{D}(\mathcal{X})$ can be regarded as a functor $\mathcal{C}^{\text{op}} \rightarrow \text{Pres}_c$ valued in the ∞ -category of presentable ∞ -categories and compact colimit-preserving functors. Now consider its Ind-extension

$$\text{Ind}(\mathcal{C})^{\text{op}} \simeq \text{Pro}(\mathcal{C}^{\text{op}}) \rightarrow \text{Pro}(\text{Pres}_c).$$

Any derived ind-algebraic stack \mathcal{X} can be regarded an ind-object in \mathcal{C} and hence gives rise to a canonical pro- ∞ -category which we denote $\widehat{\mathbf{D}}(\mathcal{X})$.

Example B.2.2. Let \mathcal{X} be a derived algebraic stack. Then $\widehat{\mathbf{D}}(\mathcal{X})$ is the constant pro- ∞ -category $\{\mathbf{D}(\mathcal{X})\}$.

Example B.2.3. If \mathcal{X} is a derived ind-algebraic stack represented by a filtered system $\{\mathcal{X}_n\}_n$, then $\widehat{\mathbf{D}}(\mathcal{X})$ is represented by the cofiltered system $\{\mathbf{D}(\mathcal{X}_n)\}_n$.

Example B.2.4. Let A be a derived commutative ring and $I \subseteq \pi_0(A)$ an ideal, and consider the formal completion $\text{Spec}(A)^\wedge$ in the vanishing locus of I . Then choosing generators f_1, \dots, f_m for the ideal I , we have an equivalence

$$\widehat{\mathbf{D}}(\text{Spec}(A)^\wedge) \simeq \{\mathbf{D}(A/(f_1^n, \dots, f_m^n))\}_n$$

by Example B.1.4. If A is a noetherian commutative ring, then we have also

$$\widehat{\mathbf{D}}(\text{Spec}(A)^\wedge) \simeq \{\mathbf{D}(A/(f_1^n, \dots, f_m^n))\}_n$$

again by Example B.1.4.

APPENDIX C. WEAK PRO-MILNOR SQUARES

In this section we continue to use the language of weight structures as in Sect. 5. For convenience, the term *weighted ∞ -category* will refer to an essentially small stable ∞ -category with a bounded weight structure. We write $\mathcal{C}^{w=0}$ for the weight-heart of a weighted ∞ -category \mathcal{C} .

C.1. Connected invariants.

Definition C.1.1.

- (i) We say that a weight-exact functor $f : \mathcal{C} \rightarrow \mathcal{D}$ between weighted ∞ -categories is *thickly surjective* if every object $Y \in \mathcal{D}^{w=0}$ is a direct summand of $f(X)$ for some object $X \in \mathcal{C}^{w=0}$. In other words, if the induced functor f^* on Ind-completions generates its codomain under colimits.
- (ii) Let \mathcal{C} and \mathcal{D} be weighted ∞ -categories and $k \geq 0$ an integer. A thickly surjective weight-exact functor $f : \mathcal{C} \rightarrow \mathcal{D}$ is *k -connective* if it induces k -connective maps

$$\text{Maps}_{\mathcal{C}^{w=0}}(X, Y) \rightarrow \text{Maps}_{\mathcal{D}^{w=0}}(f(X), f(Y))$$

for all X and Y in $\mathcal{C}^{w=0}$. In other words, if the induced functor of $(k+1)$ -categories

$$\tau_{\leq k+1}(\mathcal{C}) \rightarrow \tau_{\leq k+1}(\mathcal{D})$$

is fully faithful (and hence an equivalence) when restricted to the weight-hearts.

Example C.1.2. If $A \rightarrow B$ is a k -connective map of connective \mathcal{E}_∞ -rings, then the induced functor $\mathrm{Perf}_A \rightarrow \mathrm{Perf}_B$ is k -connective with respect to the weight structures of Example 5.1.1.

The following definition is a variant of [LT19, Def. 2.5].

Definition C.1.3. Let E be a spectrum-valued functor on the ∞ -category of small stable ∞ -categories. We say that E is *connected* if for any k -connective functor $\mathcal{C} \rightarrow \mathcal{D}$ of weighted ∞ -categories, the induced map of spectra $E(\mathcal{C}) \rightarrow E(\mathcal{D})$ is $(k+1)$ -connective.

Example C.1.4. Connective K-theory is an example of a connected invariant. This follows from [Fon18, Cor. 5.16] and the fact that plus-construction sends k -connected maps of spaces into $k+1$ -connected maps (cf. [LT19, Lem. 2.4]). Moreover, nonconnective K-theory is also an example by [Sos19, Thm. 4.3].

Remark C.1.5. Let E be a localizing invariant. If E is connected, then it is 1-connective in the sense of [LT19, Def. 2.5]. The converse holds if E commutes with filtered colimits. If E does not commute with filtered colimits, one can still show a weaker statement: if E is 2-connective in the sense of [LT19], then it is connected. This follows from the fact that any weighted ∞ -category can be functorially realized as the kernel of a weight-exact localization functor $\mathrm{Perf}_{A(\mathcal{C})} \rightarrow \mathrm{Perf}_{B(\mathcal{C})}$ for some map of connective \mathcal{E}_1 -rings $A(\mathcal{C}) \rightarrow B(\mathcal{C})$; see the proof of Lemma 3.5.5. (See [Bon10, Sect. 8.1] or [BS16] for the theory of weight-exact localizations.) Any n -connective functor $\mathcal{C} \rightarrow \mathcal{D}$ clearly induces an n -connective map on $A(\mathcal{C}) \rightarrow A(\mathcal{D})$, and it also induces an n -connective map $B(\mathcal{C}) \rightarrow B(\mathcal{D})$ by the universal properties of localization and of truncation.

C.2. Weak pro-Milnor squares. The starting point for our definition of weak pro-Milnor squares is the following classical definition (see e.g. [AM69, Sect. 4]):

Definition C.2.1. Let $\{f_n : X_n \rightarrow Y_n\}_n$ be a morphism of cofiltered systems of spectra. We say that f is *pro- k -connective* if the induced map

$$\{\tau_{\leq k}(X_n)\}_n \rightarrow \{\tau_{\leq k}(Y_n)\}_n$$

is an isomorphism in $\mathrm{Pro}(\mathrm{Spt})$ for every k . It is a *weak pro-equivalence* if it is pro- k -connective for all k .

Note that the same definition makes sense for the ∞ -category of \mathcal{E}_1 -rings, for instance, in place of Spt . The following can be viewed as a many-object generalization (see Example 5.1.1). A *weighted pro- ∞ -category* is a pro-object in the ∞ -category of weighted ∞ -categories and weight-exact functors.

Definition C.2.2. Let $\{f_n : \mathcal{C}_n \rightarrow \mathcal{D}_n\}_n$ be a cofiltered system of weight-exact functors between weighted ∞ -categories and $k \geq 0$ an integer. We say that it is *pro- k -connective* if the induced morphism of pro- ∞ -categories

$$\{\tau_{\leq k+1}(\mathcal{C}_n^{w=0})\}_n \rightarrow \{\tau_{\leq k+1}(\mathcal{D}_n^{w=0})\}_n$$

is invertible. If it is k -connective for all k , then it is called a *weak pro-equivalence* of weighted pro- ∞ -categories.

This enables us to define weak pro-Milnor squares of weighted ∞ -categories.

Definition C.2.3. Let $\{\Delta_n\}_n$ be a cofiltered system of commutative squares of weighted ∞ -categories and weight-exact functors of the form

$$\begin{array}{ccc} \mathcal{A}_n & \xrightarrow{f_n} & \mathcal{B}_n \\ \downarrow p_n & & \downarrow q_n \\ \mathcal{A}'_n & \xrightarrow{g_n} & \mathcal{B}'_n. \end{array}$$

For every n , let $\mathcal{A}_n^+ \subseteq \mathcal{A}'_n \times_{\mathcal{B}'_n} \mathcal{B}_n$ denote the full subcategory of the pullback (taken in the ∞ -category of weighted ∞ -categories) generated under finite colimits, finite limits, and retracts by the essential image of $\mathcal{A}_n \rightarrow \mathcal{A}'_n \times_{\mathcal{B}'_n} \mathcal{B}_n$. Note that \mathcal{A}_n^+ inherits a weight structure from $\mathcal{A}'_n \times_{\mathcal{B}'_n} \mathcal{B}_n$. We say that Δ is *k-pro-precartesian* if the functors $\mathcal{A}_n \rightarrow \mathcal{A}_n^+$ induce a pro- k -connective functor on weighted pro- ∞ -categories. We say it is *weakly pro-precartesian* if it is k -pro-precartesian for all k .

Definition C.2.4. Let $\{\Delta_n\}_n$ be a cofiltered system of commutative squares of weighted ∞ -categories and weight-exact functors as above. We say that $\{\Delta_n\}_n$ is *k-pro-Milnor* if it is pro-precartesian and each of the functors $f_n^*, g_n^*, p_n^*, q_n^*$ is thickly surjective. It is *weakly pro-Milnor* if it is k -pro-Milnor for all k .

Construction C.2.5. Suppose given a commutative square

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f} & \mathcal{B} \\ \downarrow p & & \downarrow q \\ \mathcal{A}' & \xrightarrow{g} & \mathcal{B}' \end{array}$$

of weighted ∞ -categories. Write $\widehat{\mathcal{A}}, \widehat{\mathcal{B}}$, etc. for the Ind-completions and consider the \odot -construction $\widehat{\mathcal{Q}} := \widehat{\mathcal{A}'} \odot_{\widehat{\mathcal{A}}}^{\widehat{\mathcal{B}'}} \widehat{\mathcal{B}}$. By Lemma 3.5.12 there is a weight structure on $\mathcal{Q} := \widehat{\mathcal{Q}}^\omega$ such that all the functors in the induced square

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{f} & \mathcal{B} \\ \downarrow p & & \downarrow q_0 \\ \mathcal{A}' & \xrightarrow{g_0} & \mathcal{Q}, \end{array}$$

as well as the functor $b : \mathcal{Q} \rightarrow \mathcal{B}'$, are weight-exact. We call \mathcal{Q} the *weighted \odot -construction*, and denote it by

$$\mathcal{A}' \odot_{\mathcal{A}}^{\mathcal{B}'} \mathcal{B}.$$

Definition C.2.6. Let $\{\Delta_n\}_n$ be a cofiltered system of commutative squares of weighted ∞ -categories and weight-exact functors of the form

$$\begin{array}{ccc} \mathcal{A}_n & \xrightarrow{f_n} & \mathcal{B}_n \\ \downarrow p_n & & \downarrow q_n \\ \mathcal{A}'_n & \xrightarrow{g_n} & \mathcal{B}'_n. \end{array} \tag{C.2.a}$$

If Δ is a pro- k -Milnor square, then we say it *pro- k -base change* if the functors $\mathcal{A}'_n \odot_{\mathcal{A}_n}^{\mathcal{B}'_n} \mathcal{B}_n \rightarrow \mathcal{B}'_n$ induce a pro- k -connective functor on weighted pro- ∞ -categories. If Δ is a weak pro-Milnor square, then we say it *weakly satisfies pro-base change* if satisfies pro- k -base change for all k .

C.3. Weak pro-excision. For connected invariants, we have the following analogue of Theorem 3.5.11:

Theorem C.3.1 (Weak pro-excision). *Let E be a connected localizing invariant. Suppose given a weak pro-Milnor square of weighted pro- ∞ -categories of the form (C.2.a) weakly satisfying pro-base change. Then the induced square of pro-spectra*

$$\begin{array}{ccc} \{E(\mathcal{A}_n)\}_n & \longrightarrow & \{E(\mathcal{B}_n)\}_n \\ \downarrow & & \downarrow \\ \{E(\mathcal{A}'_n)\}_n & \longrightarrow & \{E(\mathcal{B}'_n)\}_n \end{array}$$

is weakly pro-cartesian. That is, the morphisms $E(\mathcal{A}_n) \rightarrow E(\mathcal{A}'_n) \times_{E(\mathcal{B}'_n)} E(\mathcal{B}_n)$ induce a weak pro-equivalence of pro-spectra.

Lemma C.3.2. *Let $\{f_n : \mathcal{C}_n \rightarrow \mathcal{D}_n\}_n$ be a cofiltered system of weight-exact functors between idempotent-complete weighted ∞ -categories. If it is pro- k -connective, then it is isomorphic to a pro-system of levelwise k -connective functors.*

Proof. For every n , consider the commutative square

$$\begin{array}{ccc} \mathcal{C}_n & \xrightarrow{f_n} & \mathcal{D}_n \\ \downarrow & & \downarrow \\ \tau_{\leq k+1}(\mathcal{C}_n) & \xrightarrow{f_n} & \tau_{\leq k+1}(\mathcal{D}_n) \end{array}$$

By assumption, the lower arrow induces an isomorphism of pro-objects as n varies. Since passage to underlying pro-objects commutes with finite limits, it follows that the base changes $\mathcal{E}_n \rightarrow \mathcal{D}_n$ also induce an isomorphism of pro-objects. Thus it will suffice to show that for every n , the induced functor $f'_n : \mathcal{C}_n \rightarrow \mathcal{E}_n$ is k -connective. By construction of \mathcal{E}_n , the functor $\mathcal{E}_n \rightarrow \tau_{\leq k+1}(\mathcal{C}_n)$ induces an equivalence on homotopy categories of the weight-hearts. Since the same holds for $\mathcal{C}_n \rightarrow \tau_{\leq k+1}(\mathcal{C}_n)$, it follows that $\mathcal{C}_n \rightarrow \mathcal{E}_n$ is thickly surjective.

Now for any two objects X and Y in $\mathcal{C}_n^{w=0}$, consider the commutative triangle

$$\begin{array}{ccc} \mathrm{Maps}_{\mathcal{C}_n}(X, Y) & \longrightarrow & \tau_{\leq k} \mathrm{Maps}_{\mathcal{E}_n}(f'_n(X), f'_n(Y)) \\ & \searrow & \downarrow \\ & & \tau_{\leq k} \mathrm{Maps}_{\mathcal{C}_n}(X, Y). \end{array}$$

Note that the vertical and diagonal maps have fibres

$$\tau_{\geq k+1} \mathrm{Maps}_{\mathcal{D}_n}(f_n(X), f_n(Y)) \text{ and } \tau_{\geq k+1} \mathrm{Maps}_{\mathcal{C}_n}(X, Y),$$

respectively. These are both $(k+1)$ -connective, so it follows from the octahedral axiom that the horizontal map also has k -connective fibre. \square

Corollary C.3.3. *Let E be a connected invariant. Then E sends pro- k -connective maps of idempotent-complete weighted ∞ -categories to $(k+1)$ -connective maps of pro-spectra. In particular, it sends weak pro-equivalences to weak pro-equivalences.*

Proof of Theorem C.3.1. Since $\{\Delta_n\}_n$ is weakly pro-Milnor, it is weakly pro-equivalent to the pro-system induced by the squares

$$\begin{array}{ccc} \mathcal{A}_n & \longrightarrow & \mathcal{B}_n \\ \downarrow & & \downarrow \\ \mathcal{A}'_n & \longrightarrow & \mathcal{A}'_n \circlearrowleft_{\mathcal{A}'_n}^{\mathcal{B}'_n} \mathcal{B}_n. \end{array}$$

By Theorem 3.4.3, E sends the above square to a cartesian square of spectra for every n . Thus the claim follows from Corollary C.3.3. \square

REFERENCES

- [AHHLR] J. Alper, J. Hall, D. Halpern-Leistner, and D. Rydh, *Artin algebraization for pairs and applications to the local structure of stacks and Ferrand pushouts*, in preparation
- [AHR19] J. Alper, J. Hall, and D. Rydh, *The étale local structure of algebraic stacks*, arXiv e-prints (2019), preprint arXiv:1912.06162
- [AM69] M. Artin and B. Mazur, *Étale homotopy*, vol. 100, Springer, Cham, 1969
- [AOV08] D. Abramovich, M. Olsson, and A. Vistoli, *Tame stacks in positive characteristic*, Ann. Inst. Fourier **58** (2008), no. 4, pp. 1057–1091
- [Art70] M. Artin, *Algebraization of formal moduli. II: Existence of modifications*, Ann. Math. (2) **91** (1970), pp. 88–135
- [BGT13] A. J. Blumberg, D. Gepner, and G. Tabuada, *A universal characterization of higher algebraic K-theory*, Geometry & Topology **17** (2013), no. 2, pp. 733–838
- [Bon10] M. Bondarko, *Weight structures vs. t-structures; weight filtrations, spectral sequences, and complexes (for motives and in general)*, K-theory **6** (2010), no. 3, pp. 387–504
- [BS16] M. Bondarko and V. Sosnilo, *Non-commutative localizations of additive categories and weight structures; applications to birational motives*, Journal of the Institute of Mathematics of Jussieu **17** (2016), no. 4, p. 785–821, <http://dx.doi.org/10.1017/S1474748016000207>
- [CK20] D.-C. Cisinski and A. A. Khan, *\mathbf{A}^1 -homotopy invariance in spectral algebraic geometry*, preprint arXiv:1705.03340
- [EHIK20] E. Elmanto, M. Hoyois, R. Iwasa, and S. Kelly, *Milnor excision for motivic spectra*, preprint arXiv:2004.12098
- [ES20] E. Elmanto and V. Sosnilo, *On nilpotent extensions of ∞ -categories and the cyclotomic trace*, preprint arXiv:2010.09155
- [Fer03] D. Ferrand, *Conducteur, descente et pincement*, Bull. Soc. Math. Fr. **131** (2003), no. 4, pp. 553–585
- [Fon18] E. E. Fontes, *Weight structures and the algebraic K-theory of stable ∞ -categories*, preprint arXiv:1812.09751
- [GH06] T. Geisser and L. Hesselholt, *Bi-relative algebraic K-theory and topological cyclic homology*, Invent. Math. **166** (2006), no. 2, pp. 359–395
- [GR14] D. Gaitsgory and N. Rozenblyum, *DG indschemes*, Contemporary Mathematics (2014), p. 139–251, <http://dx.doi.org/10.1090/conm/610/12080>
- [GR17] D. Gaitsgory and N. Rozenblyum, *A study in derived algebraic geometry. Volume I: Correspondences and duality*, vol. 221, Providence, RI: American Mathematical Society (AMS), 2017
- [Gro17] P. Gross, *Tensor generators on schemes and stacks*, Algebraic Geometry (2017), p. 501–522, <http://dx.doi.org/10.14231/ag-2017-026>
- [HK19] M. Hoyois and A. Krishna, *Vanishing theorems for the negative K-theory of stacks*, Annals of K-Theory **4** (2019), no. 3, p. 439–472, <http://dx.doi.org/10.2140/akt.2019.4.439>
- [HLP14] D. Halpern-Leistner and A. Preygel, *Mapping stacks and categorical notions of properness*, preprint arXiv:1402.3204
- [Hoy17] M. Hoyois, *The six operations in equivariant motivic homotopy theory*, Adv. Math. **305** (2017), pp. 197–279, <https://doi.org/10.1016/j.aim.2016.09.031>
- [Hoy18] M. Hoyois, *K-theory of dualizable categories*, available at <http://www.mathematik.ur.de/hoyois/papers/efimov.pdf>, 2018
- [HR17] J. Hall and D. Rydh, *Perfect complexes on algebraic stacks*, Compositio Mathematica **153** (2017), no. 11, p. 2318–2367, <http://dx.doi.org/10.1112/S0010437X17007394>
- [Ker18] M. Kerz, *On negative algebraic K-groups*, Proceedings of the International Congress of Mathematicians, ICM 2018, Rio de Janeiro, Brazil, August 1–9, 2018. Volume II. Invited lectures, 2018, pp. 163–172

- [Kha17] A. A. Khan, *Pro-systems of K-theory spectra*, Lecture notes, available at <https://www.preschema.com/lecture-notes/kdescent/lect8.pdf>, 2017
- [Kha19] A. A. Khan, *The Morel–Voevodsky localization theorem in spectral algebraic geometry*, *Geom. Topol.* **23** (2019), no. 7, pp. 3647–3685
- [Kha20] A. A. Khan, *Algebraic K-theory of quasi-smooth blow-ups and cdh descent*, *Annales Henri Lebesgue* **3** (2020), pp. 1091–1116, ahl.centre-mersenne.org/item/AHL_2020__3__1091_0/
- [Kha22] A. A. Khan, *K-theory and G-theory of derived algebraic stacks*, *Japanese Journal of Mathematics* (2022)
- [KØ12] A. Krishna and P. A. Østvær, *Nisnevich descent for K-theory of Deligne–Mumford stacks*, *J. K-Theory* **9** (2012), no. 2, pp. 291–331
- [KR18a] A. A. Khan and D. Rydh, *Virtual Cartier divisors and blow-ups*, preprint [arXiv:1802.05702](https://arxiv.org/abs/1802.05702)
- [KR18b] A. Krishna and C. Ravi, *Algebraic K-theory of quotient stacks*, *Annals of K-Theory* **3** (2018), no. 2, p. 207–233, <http://dx.doi.org/10.2140/akt.2018.3.207>
- [KR21] A. A. Khan and C. Ravi, *Generalized cohomology theories for algebraic stacks*, arXiv e-prints (2021), preprint [arXiv:2106.15001](https://arxiv.org/abs/2106.15001)
- [KS17] M. Kerz and F. Strunk, *On the vanishing of negative homotopy K-theory*, *J. Pure Appl. Algebra* **221** (2017), no. 7, pp. 1641–1644
- [KST17] M. Kerz, F. Strunk, and G. Tamme, *Algebraic K-theory and descent for blow-ups*, *Inventiones mathematicae* **211** (2017), no. 2, p. 523–577, <http://dx.doi.org/10.1007/s00222-017-0752-2>
- [LMB18] G. Laumon and L. Moret-Bailly, *Champs algébriques*, vol. 39, Springer, 2018
- [LT19] M. Land and G. Tamme, *On the K-theory of pullbacks*, *Annals of Mathematics* **190** (2019), no. 3, pp. 877–930, <https://www.jstor.org/stable/10.4007/annals.2019.190.3.4>
- [HTT] J. Lurie, *Higher topos theory*, no. 170, Princeton University Press, 2009
- [HA] ———, *Higher algebra*, Preprint, available at www.math.harvard.edu/~lurie/papers/HigherAlgebra.pdf (2017), Version of 2017-09-18
- [SAG] ———, *Spectral algebraic geometry*, Preprint, available at www.math.harvard.edu/~lurie/papers/SAG-rootfile.pdf (2018), Version of 2018-02-03
- [Mil71] J. W. Milnor, *Introduction to algebraic K-theory*, vol. 72, Princeton University Press, Princeton, NJ, 1971
- [MNN17] A. Mathew, N. Naumann, and J. Noel, *Nilpotence and descent in equivariant stable homotopy theory*, *Advances in Mathematics* **305** (2017), p. 994–1084
- [Mor18] M. Morrow, *Pro unitality and pro excision in algebraic K-theory and cyclic homology*, *J. Reine Angew. Math.* **736** (2018), pp. 95–139
- [Rao74] J.-C. Raoult, *Compactification des espaces algébriques*, *C. R. Acad. Sci., Paris, Sér. A* **278** (1974), pp. 867–869
- [RG71] M. Raynaud and L. Gruson, *Critères de platitude et de projectivité. Techniques de "platification" d'un module. (Criterial of flatness and projectivity. Technics of "flatification of a module.)*, *Invent. Math.* **13** (1971), pp. 1–89
- [Ryd11] D. Rydh, *Compactification of tame Deligne–Mumford stacks*, Preprint, available at <https://people.kth.se/~dary/tamecompactification20110517.pdf>, 2011
- [Ryd17] ———, *Equivariant flatification, étalification and compactification*, In preparation (2017)
- [Sos19] V. Sosnilo, *Theorem of the heart in negative K-theory for weight structures*, *Doc. Math.* **24** (2019), pp. 2137–2158
- [Sos21] V. Sosnilo, *Regularity of Spectral Stacks and Discreteness of Weight-Hearts*, *The Quarterly Journal of Mathematics* (2021), <https://doi.org/10.1093/qmath/haab017>, haab017
- [Sta20] J. Stapleton, *Weibel's conjecture for twisted K-theory*, *Annals of K-Theory* **5** (2020), no. 3, pp. 621–637
- [Sus95] A. A. Suslin, *Excision in the integral algebraic K-theory*, Number theory, algebra and algebraic geometry. Collected papers. In honor of the seventieth birthday of Academician Igor Rostislavovich Shafarevich, Moscow: Maik Nauka/Interperiodica Publishing, 1995, pp. 255–279
- [Tam18] G. Tamme, *Excision in algebraic K-theory revisited*, *Compositio Mathematica* **154** (2018), no. 9, p. 1801–1814, <http://dx.doi.org/10.1112/S0010437X18007236>
- [Stacks] The Stacks Project Authors, *Stacks Project*, <https://stacks.math.columbia.edu>, 2020
- [Tho87] R. W. Thomason, *Algebraic K-theory of group scheme actions*, Algebraic topology and algebraic K-theory, Proc. Conf., Princeton, NJ (USA), *Ann. Math. Stud.* 113, 539–563 (1987)., 1987
- [TT07] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, *The Grothendieck Festschrift*, Springer, 2007, pp. 247–435
- [Voe10a] V. Voevodsky, *Homotopy theory of simplicial sheaves in completely decomposable topologies*, *J. Pure Appl. Algebra* **214** (2010), no. 8, pp. 1384–1398, <https://doi.org/10.1016/j.jpaa.2009.11.004>
- [Voe10b] ———, *Unstable motivic homotopy categories in Nisnevich and cdh-topologies*, *J. Pure Appl. Algebra* **214** (2010), no. 8, pp. 1399–1406, <https://doi.org/10.1016/j.jpaa.2009.11.005>

[Wei80] C. A. Weibel, *K-theory and analytic isomorphisms*, Invent. Math. **61** (1980), pp. 177–197

DEPARTMENT MATHEMATIK, LMU MÜNCHEN, THERESIENSTR. 39, 80333 MÜNCHEN, GERMANY

Email address: tom.bachmann@zoho.com

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, NO. 1, SEC. 4, ROOSEVELT ROAD, TAIPEI 106319, TAIWAN

Email address: adeelkhan@gate.sinica.edu.tw

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY

Email address: ravi@mpim-bonn.mpg.de

LABORATORY OF MODERN ALGEBRA AND APPLICATIONS, ST. PETERSBURG STATE UNIVERSITY, 14TH LINE, 29B, 199178 SAINT PETERSBURG, RUSSIA

ST. PETERSBURG DEPARTMENT OF STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, FONTANKA, 27, 191023 SAINT PETERSBURG, RUSSIA

Email address: vsosnilo@gmail.com