



**Manchester
Metropolitan
University**

Venckunas, Tomas and Degens, Hans ORCID logoORCID:
<https://orcid.org/0000-0001-7399-4841> (2022) Genetic polymorphisms of
muscular fitness in young healthy men. PLoS One, 17 (9). e0275179-
e0275179. ISSN 1932-6203

Downloaded from: <https://e-space.mmu.ac.uk/630504/>

Version: Published Version

Publisher: Public Library of Science (PLoS)

DOI: <https://doi.org/10.1371/journal.pone.0275179>

Usage rights: Creative Commons: Attribution 4.0

Please cite the published version


<https://e-space.mmu.ac.uk>

RESEARCH ARTICLE

Genetic polymorphisms of muscular fitness in young healthy men

Tomas Venckunas^{1*}, Hans Degens^{1,2}

1 Institute of Sport Science and Innovations, Lithuanian Sports University, Kaunas, Lithuania, **2** Department of Life Sciences, Musculoskeletal Science and Sports Medicine Research Centre, Institute of Sport, Manchester Metropolitan University, Manchester, United Kingdom

* tomas.venckunas@lsu.lt

Abstract

The effects of genetic polymorphisms on muscle structure and function remain elusive. The present study tested for possible associations of 16 polymorphisms (across ten candidate genes) with fitness and skeletal muscle phenotypes in 17- to 37-year-old healthy Caucasian male endurance ($n = 86$), power/strength ($n = 75$) and team athletes ($n = 60$), and non-athletes ($n = 218$). Skeletal muscle function was measured with eight performance tests covering multiple aspects of muscular fitness. Along with body mass and height, the upper arm and limb girths, and maximal oxygen uptake were measured. Genotyping was conducted on DNA extracted from blood. Of the 16 polymorphisms studied, nine (spanning seven candidate genes and four gene families/signalling pathways) were independently associated with at least one skeletal muscle fitness measure (size or function, or both) measure and explained up to 4.1% of its variation. Five of the studied polymorphisms (activin- and adreno-receptors, as well as myosine light chain kinase 1) in a group of one to three combined with body height, age and/or group explained up to 20.4% of the variation of muscle function. *ACVR1B* (rs2854464) contributed 2.0–3.6% to explain up to 14.6% of limb proximal girths. The G allele (genotypes AG and GG) of the *ACVR1B* (rs2854464) polymorphism was significantly overrepresented among team (60.4%) and power (62.0%) athletes compared to controls (52.3%) and endurance athletes (39.2%), and G allele was also most consistently/frequently associated with muscle size and power. Overall, the investigated polymorphisms determined up to 4.1% of the variability of muscular fitness in healthy young humans.

OPEN ACCESS

Citation: Venckunas T, Degens H (2022) Genetic polymorphisms of muscular fitness in young healthy men. PLoS ONE 17(9): e0275179. <https://doi.org/10.1371/journal.pone.0275179>

Editor: Dalton Müller Pessôa Filho, Universidade Estadual Paulista Julio de Mesquita Filho - Campus de Bauru, BRAZIL

Received: May 11, 2022

Accepted: September 12, 2022

Published: September 27, 2022

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0275179>

Copyright: © 2022 Venckunas, Degens. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The author received no specific funding for this work.

Introduction

As early as during adolescence, low skeletal muscle fitness poses a significant risk for chronic disease independent of cardiovascular fitness and overweight [1]. Although it is generally accepted that genetic variation accounts for >50% of the variability in muscle mass and function, and exercise performance [2–8], little is known about what specific genetic variants contribute to the ability to perform specific motor tasks, such as polymorphisms that predispose to high muscle strength and power. In addition, most studies performed to assess the relationship between polymorphisms and muscle function or exercise capacity lack precision in

Competing interests: The authors have declared that no competing interests exist.

phenotyping the participants, as detailed measurements with a battery of physiological and performance tests is time-consuming and challenging for both the subjects and the investigators. Yet, a study like this may help in “talent selection”, where a good panel of polymorphisms may be used to direct people to sports they are “suited” best.

The ACE and ACTN3 genes are perhaps the most investigated genes and it has been observed that for instance the ACTN3 R-allele was associated with greater muscle power [9, 10]. It is equivocal whether there are such associations with ACE polymorphisms, where some studies do not find them [10] while the others did observe an association with muscle strength [11]. Perhaps the discrepancies between studies are related to differences in the parameters of muscle function that were studied and/or differences between populations, as Nazarov et al. [12] observed a higher prevalence of D-allele carriers in power athletes than endurance athletes and non-athletes.

The myostatin signalling pathway has emerged as one of the strongest candidates to explain the inter-individual variation in skeletal muscle phenotypes [13–15]. Indeed, muscle strength has been associated with polymorphisms in several genes involved in the myostatin signalling pathway [16–19]. For instance, the amount of lean body mass in older women has been found to associate with genetic variation in *ACVR2B* (rs2276541) [20] and the K153R polymorphism (rs1805086) of the myostatin (*MSTN*) gene has been associated with muscle strength in young men and women [21] and in aged women [22], and with muscle power in young men [23]. Another *MSTN* polymorphism (A55T) has been linked to muscle strength recovery after exercise-induced muscle damage in young men [24], and both K153R and A55T polymorphism have been associated with strength training-induced muscle hypertrophy [25]. As Mstn acts via its receptors *ACVR1B* and *ACVR2B*, it is of interest to know whether associations also exist between polymorphisms of the *ACVR1B* and *ACVR2B* genes with muscle size, strength and power.

Despite these reported associations between genotype and muscle mass and function, as illustrated above for the associations of muscle structure and function with genetic polymorphisms, so far studied SNPs have failed to explain even half of the estimated heritability for the trait [3, 26–28]. In fact, most investigated candidate genes have been shown to influence baseline muscle phenotypes as well as their response to training (i.e. muscle plasticity) only to a small extent at best [3, 10, 19, 29–31]. There thus still remains a long way to dissect the genetic factors that affect muscular fitness in humans.

To obtain a better understanding of the associations between genotype and muscle structure and function, in the current study we assessed the association of sixteen common genetic polymorphisms with a wide range of muscle fitness phenotypes in a large cohort of young healthy men varying in their training and fitness characteristics. Among the most investigated candidate genes *ACTN3* and *ACE*, we have also included genes of the myostatin signalling pathway (because of its primary relevance to muscularity), as well as those regulating myofibrillar contractile properties in smooth (*MYLK1*), skeletal (*MYLK2*) and cardiac (*MYLK3*) muscle, and putative modifiers of resting and exercise energy metabolism and muscle blood flow *via* differentiate sensitivity to catecholamine signalling (i.e. genes encoding adrenoreceptors *ADRA2A* and *ADRA2B*). We *hypothesized* that these less investigated candidate genes are at least as important as *ACTN3* and *ACE* for the muscular fitness.

Materials and methods

Subjects

The participants were 17- to 37-year-old Caucasian men from the Genetics and Epigenetics of Lithuanian Athletes from Kaunas (GELAK) cohort recruited as described previously [32, 33]. Potential participants were excluded from taking part in the study if they were known to suffer

Table 1. Anthropometrics, muscular performance and cardiovascular fitness of the participants.

	C	E	P	T
Age (y)	24.5 ± 4.3 (218)	22.3 ± 3.6 (86) ^a	22.7 ± 3.5 (75) ^a	21.5 ± 3.1 (60) ^a
Height (m)	1.80 ± 0.06 (216)	1.79 ± 0.05 (86)	1.81 ± 0.06 (75)	1.87 ± 0.08 (58) ^{a,b,c}
Body mass (kg)	77.4 ± 11.0 (218)	70.5 ± 6.7 (85) ^a	81.5 ± 11.9 (72) ^{a,b}	80.8 ± 9.1 (59) ^b
BMI (kg·m ⁻²)	23.8 ± 2.9 (215)	21.9 ± 1.7 (84) ^a	25.0 ± 2.8 (72) ^{a,b}	23.1 ± 1.5 (58) ^{b,c}
Thigh girth (cm)	54.5 ± 5.6 (217)	53.0 ± 4.0 (86)	57.5 ± 4.7 (75) ^{a,b}	56.2 ± 2.8 (59) ^b
Upper arm girth (cm)	30.9 ± 3.7 (217)	28.0 ± 2.0 (86) ^a	33.7 ± 3.7 (75) ^{a,b}	31.2 ± 2.0 (59) ^{b,c}
Balance (no. of attempts)	8.02 ± 4.50 (216)	6.85 ± 7.14 (85)	7.36 ± 4.73 (73)	6.97 ± 3.34 (58)
Handgrip strength (kg)	68.7 ± 10.1 (216)	61.6 ± 8.7 (84) ^a	70.6 ± 11.4 (73) ^b	68.4 ± 8.8 (58) ^b
Knee extension PT (Nm)	251 ± 47 (215)	240 ± 37 (79) ^a	301 ± 68 (72) ^{a,b}	298 ± 46 (54) ^{a,b}
Knee flexion PT (Nm)	135 ± 25 (215)	134 ± 22 (79)	152 ± 35 (72) ^{a,b}	161 ± 25 (54) ^{a,b}
Vertical jump power (W·kg ⁻¹)	26.5 ± 2.0 (212)	26.3 ± 2.0 (80)	28.4 ± 2.3 (70) ^{a,b}	28.1 ± 1.8 (55) ^{a,b}
Shuttle agility run (s)	20.2 ± 1.3 (211)	19.6 ± 0.9 (85) ^a	19.7 ± 1.0 (73) ^a	19.7 ± 1.4 (50)
30 m run (s)	4.56 ± 0.24 (214)	4.43 ± 0.18 (84) ^a	4.38 ± 0.19 (73) ^{a,b}	4.31 ± 0.15 (58) ^{a,b,c}
Wingate (rev·30 s ⁻¹)	53.0 ± 5.8 (208)	59.6 ± 5.0 (78) ^a	56.6 ± 4.9 (72) ^b	58.3 ± 4.6 (58) ^a
Wingate fatigue resistance (%)	51.5 ± 7.9 (208)	61.6 ± 8.2 (79) ^a	53.8 ± 8.6 (72) ^b	56.2 ± 7.5 (58) ^{a,b}
Pull-ups (no.)	8.5 ± 4.8 (200)	9.6 ± 3.8 (82)	13.0 ± 5.0 (70) ^{a,b}	9.0 ± 4.2 (56) ^c
VO ₂ max (ml·kg ⁻¹ ·min ⁻¹)	50.8 ± 6.0 (186)	64.1 ± 10.2 (74) ^a	51.7 ± 5.8 (67) ^b	54.6 ± 4.9 (48) ^{a,b}

C: Control; E: Endurance athletes; P: Power athletes; T: Team athletes; Values are mean ± SD; between parentheses number of individuals; BMI: body mass index; PT: peak torque; VO₂max: maximal oxygen uptake.

^a: different from control at p ≤ 0.024

^b: different from endurance athletes at p < 0.05

^c: different from power athletes at p < 0.05.

<https://doi.org/10.1371/journal.pone.0275179.t001>

from cardiovascular diseases or hypertension. The vast majority of the athletes were competitive, trained 3–14 times a week during the study, and were selected from the Registers of the Lithuanian Sports Federations. The study was approved by the Lithuanian National Committee for Bioethics, adhered to the guidelines of the declaration of Helsinki, and participants provided written informed consent before participation in the study. Athletes were divided into 3 groups as described previously [33]: endurance athletes (n = 86; distance runners, road cyclists, paddlers and skiers), power/strength athletes (n = 75; track and field sprinters, jumpers, throwers, combat sport athletes and body builders) and team athletes (n = 60; basketball, soccer and volleyball). Non-athletes (n = 218) trained for less than 2 h per week for the last 5 years. Characterization of the participants is provided in [Table 1](#).

Anthropometrics

Anthropometrics were determined before the performance tasks with the participants standing barefoot and wearing only shorts. Body height was measured using a metal stadiometer with the subject not touching any surface, and girths were measured with the flexible strap, both at 0.1 cm resolution. Dominant leg and dominant arm girths have been used for the analysis in this study. Body mass was measured after the overnight fast on a separate visit to the lab with electronic scales (TBF 300; Tanita, Tokyo, Japan) to the nearest 0.1 kg.

Performance measurements

Muscular fitness. Subjects arrived at the lab at least 2 hours after the last meal. Before the tests, subjects warmed-up with 6–8 min of cycling on an electromechanically braked cycle

ergometer (Ergometrics–800S, Ergo Line, Medical Measurement Systems; Bitz, Germany) at ~70 rpm at the power in Watts equal to the body mass in kg, and then performed some light dynamic stretching exercises. The order of the tests was as follows: handgrip strength, Flamingo balance test, isokinetic dynamometry, pull-ups, countermovement jump, 10 x 5 m shuttle run, 30 m sprint run, and Wingate test.

Handgrip strength

With the subject sitting with the hands on the table, handgrip strength at a comfortable grip position was measured with a hydraulic dynamometer (SH5001, Saehan Corporation, Korea) as the best of six attempts (three with each hand). The attempts were separated by ~1 min of rest, and to ensure a good grip of the dynamometer, a towel was used to dry/clean/absorb any sweat from the handle and the palms between the attempts.

Balance. The Flamingo balance test assesses one-leg standing ability on a 50 cm long, 5 cm high and 3 cm wide rigid beam [34]. In brief, the subject had to flex the free leg and grip the back of the foot with his hand of the same side, and after release of the support (and immediate start of the stopwatch by the observing researcher) try to remain on the beam in this position for 1 min. When balance is lost, (subject releases the suspended leg), the stopwatch is stopped and the test resumed without delay by the subject. The number of attempts needed to accumulate a total 60 s standing time is counted and considered inversely related to the static balance ability. Before the test, subjects were given one try with each leg to become familiar with the test and to decide which leg is dominant (more convenient to stand on), and that leg was used to start the test. Then the test was immediately performed on the other leg. For further analyses, the average number of attempts of each leg was used.

Isokinetic dynamometry. Knee extension and flexion torque of the dominant (the right one in most cases) leg was measured with the subject seated comfortably on a dynamometer (Biodex Pro3, USA) as described in more detail in Degens et al. [35]. The peak torques achieved for flexion and extension during three consecutive maximal effort full-range isokinetic flexion–extension contraction cycles at $30^{\circ} \cdot s^{-1}$ were used for further analysis.

Pull-ups. The number of full pull-ups (pronated grip) the subject is capable to perform in one set was measured, with the execution technique supervised (full extension of the arms at the end of eccentric phase, no leg swing to get make use of inertia, chin required to be raised above the bar) by a researcher who was also giving verbal encouragement to perform as many repetitions as possible with rests of no longer than 3 seconds between consecutive pull-ups. One attempt was allowed; subjects were not permitted to release the bar with either hand during the set. The bar was 3 cm in diameter, rigid and the width of the grasp was as preferred by the subject but corresponded to about the width of the shoulders.

As a measure of maximal leg extension *muscle power*, the participants performed three countermovement jumps (arms on the waist) separated by at least 1 min. The best jump was used for further analysis. Jump velocity at take-off (v in $m \cdot s^{-1}$) was calculated as:

$$v = a * t_f / 2$$

where ‘ a ’ is the gravitational acceleration ($9.81 m \cdot s^{-2}$) and ‘ t_f ’ the flight time of the jump. The power was estimated as:

$$W = \text{body mass} * a * v$$

and expressed relative to a subject’s body mass in kg.

To measure agility and *anaerobic power*, the subjects performed a 10 x 5 m shuttle run test according to the protocol of the Eurofit battery [34, 36], followed by a 30-m sprint run from a

standing start. Both running tests were conducted on concrete flooring, and the better of two attempts in the shuttle run and the best of the three attempts in 30-m sprint run were used for subsequent analysis. The time in the shuttle run was measured by an investigator with a stopwatch while it was measured electronically for the 30-m sprint.

The battery of the muscular fitness tests was concluded with a Wingate test to estimate anaerobic power and anaerobic endurance capacity [37]. In brief, the Wingate test was carried out on a mechanically-braked cycle ergometer (Monark 824E, Sweden) and preceded with fast unresisting acceleration to maximal pedaling cadence, after which the brake weight equaling 7.5% of body mass (to the nearest 0.1 kg) was applied at the signal of the subject, to initiate the 30-s all-out effort. Before and during the test, subjects were verbally encouraged to exert maximally during the entire 30 seconds of the test. The number of full pedaling cycles per 30 s was then calculated, as was the power produced per each 5-s interval to derive a fatigue index, which is the percentage of power drop during the test.

Aerobic capacity (cardiorespiratory fitness)

Aerobic capacity was measured with a VO_2max test during a separate visit to the lab [35]. In brief, a ramp treadmill (H/P/Cosmos Sports & Medical GMBH, Germany) protocol of continuous incremental running until exhaustion was applied. After a warm-up of slow running for some minutes to familiarize with the treadmill, participants started the test by jogging at $7 \text{ km}\cdot\text{h}^{-1}$ for 3 min at an initial gradient of 1%, and then the speed of the treadmill belt was increasing by $0.1 \text{ km}\cdot\text{h}^{-1}$ each 6 seconds. The treadmill speed remained constant after it reached $20 \text{ km}\cdot\text{h}^{-1}$, and then the gradient of the treadmill was increased by 0.05% every 6 s from the initial 1%. Throughout the test, breath-by-breath gas analysis was performed using an Oxycon Mobile gas analyzer (Viasys, Germany), and heart rate (HR) was recorded with a HR meter (Polar 810s, Finland). During the later stages of the test, participants were verbally encouraged to keep running as long as they can. VO_2max was calculated as highest average of 20 consecutive seconds.

Genotyping

Venous blood was collected from an antecubital vein. DNA for genotyping was extracted from the venous blood samples using the NucleoSpin Blood kit (Macherey-Nagel, GmbH & Co. KG, Düren, Germany) according to the instructions by the manufacturer [37]. The polymorphisms analysed are listed in Table 2.

ACE I/D (rs4341) genotypes were determined as described in Moran et al. [38]. *ACTN3* R577X (rs1815739), *MSTN* K153R (rs1805086), *ACVR1B* (rs2854464), *ACVR2B* (rs3792527 and rs7372545), *MYLK1* (C49T, rs2700352; H21P, rs28497577; and rs820336), *MYLK2* (rs6060965, rs4911532 and rs6119729), *MYLK3* (rs36471), *ADRA2A* (rs553668) and *ADRA2B* (rs4066772) genotypes were determined using PCR-RFLP described in detail in Moran et al. [38] and below.

Copy number variation

Primer and hydrolysis probe assays (Integrated DNA Technologies Inc., USA) were designed to amplify a region lying in the introns of the genes of interest. The Basic Local Alignment Search Tool (BLAST, National Centre for Biotechnology Information) was used to ensure that the primers and probes were specific to the genes of interest and did not recognise any other sequences. All samples were run in triplicate and each plate contained a non-template control also run in triplicate. A reaction volume of 12.5 μL was used for all reactions, containing 6.25 μL of 2x Type-it CNV Probe PCR Master Mix (Qiagen Ltd, UK), 0.5 μL of 25x Ref assay,

Table 2. Genetic polymorphisms analysed and genotype distribution (%) in C: Control; E: Endurance athletes; P: Power athletes; T: Team athletes.

	Genotype	C	E	P	T
ACE (rs4341)	II	23.2	33.3	15.5	16.4
	ID	50.0	48.1	62.0	52.7
	DD	26.8	18.5	22.5	30.9
MYLK1 (rs2700352)	AA	0.5	1.2	1.4	0.0
	AG	32.0	32.1	26.8	21.8
	GG	67.5	66.7	71.8	78.2
MYLK1 H21P (rs28497577)	GG	82.5	85.2	90.1	81.8
	GT	17.0	13.6	9.9	18.2
	TT	0.5	1.2	0.0	0.0
MYLK1 (rs820336)	CC	8.2	6.2	9.9	7.3
	TC	42.3	42.0	35.2	36.4
	TT	49.5	51.9	54.9	56.4
MYLK2 (rs6060965)	AA	88.7	86.4	90.1	92.7
	AG	11.3	13.6	9.9	7.3
	GG	0.0	0.0	0.0	0.0
MYLK2 (rs4911532)	CC	29.4	35.8	39.4	30.9
	CT	54.1	40.7	45.1	50.9
	TT	16.5	23.5	15.5	18.2
MYLK2 (rs6119729)	CC	55.7	59.3	57.7	54.5
	CT	37.6	28.4	33.8	38.2
	TT	6.7	12.3	8.5	7.3
MYLK3 (rs36471)	AA	66.5	59.3	63.4	60.0
	GA	29.9	34.6	28.2	32.7
	GG	3.6	6.2	8.5	7.3
MYLK3 V180L (rs28407821)	AA	30.9	28.4	26.8	29.1
	CA	46.4	50.6	56.3	47.3
	CC	22.7	21.0	16.9	23.6
MSTN K153R (rs1805086)	CC	0.0	0.0	0.0	0.0
	TC	0.0	2.3	1.3	0.0
	TT	100.0	97.7	98.7	100.0
ACVR1B (rs2854464)	AA	47.7	60.8	38.0 ^b	39.6 ^{a,b}
	GA	39.9	30.4	43.7 ^b	32.1 ^{a,b}
	GG	12.4	8.9	18.3 ^b	28.3 ^{a,b}
ACVR2B (rs3792527)	CC	36.3	27.5	34.8	21.8
	TC	50.0	47.5	39.1	49.1
	TT	13.7	25	26.1	29.1
ACVR2B (rs7372545)	GG	36.5	27.2	28.6	29.6
	GT	36.0	43.2	38.6	37.0
	TT	27.5	29.6	32.9	33.3
ACTN3 R577X (rs1815739)	RR	34.9	35.8	38.0	41.8
	RX	51.6	55.6	52.1	45.5
	XX	13.5	8.6	9.9	12.7
ADRA2A (rs553668)	AA	0.5	1.3	1.5	1.9
	GA	26.3	24.1	14.7	28.3
	GG	73.2	74.7	83.8	69.8
ADRA2B (rs4066772)	II	40.7	27.2	38.0	43.6
	ID	48.5	54.3	39.4	38.2

(Continued)

Table 2. (Continued)

	Genotype	C	E	P	T
	DD	10.8	18.5	22.5	18.2
Genotype score		28.6 ± 2.4 (178)	28.8 ± 2.4 (79)	29.1 ± 2.4 (67)	29.7 ± 2.3 (52)

ACE: Angiotensin converting enzyme; MYLK1: myosin light chain kinase smooth muscle; MYLK2: MYLK skeletal muscle; MYLK3: MYLK cardiac muscle; MSTN: myostatin; ACVR: activin receptors; ACTN3: α -actinin-3; ADRA2: α -2-adrenergic receptor

^a: different from control $P = 0.03$

^b: different from endurance $p \leq 0.037$.

<https://doi.org/10.1371/journal.pone.0275179.t002>

0.5 μ l of 25x assay, 2.75 μ l of nuclease-free water and 2 μ l of DNA normalised to 5 ng/ μ L. All reactions were performed on the same LightCycler 480 (Roche Diagnostics Ltd, UK) system and included activation at 95°C for 5 min followed by 40 cycles of alternative 30-s periods at 95°C and 60°C as recommended by the Type-it CNV probe protocol.

The genotype score was calculated as the sum of the scores for each genotype, where 3 was allocated to the homozygous genotype of a certain gene that according to the literature was expected to correlate with fitness positively, while a score of 1 reflected the homozygous genotype that was expected to correlate with fitness negatively. A score of two was given to the heterozygous genotype.

Statistical analyses

The Shapiro–Wilk test showed that the data were normally distributed. An ANOVA with a Bonferroni-corrected post-hoc test was used to assess differences between groups. Hardy-Weinberg equilibrium was tested for all genotypes with a χ^2 test. A Kruskal-Wallis test was performed to determine differences in the frequency distribution of polymorphisms between groups. Stepwise linear regression was performed with group, age, height and all polymorphisms as factors to assess which polymorphisms correlated with muscle fitness phenotypes. Differences and correlations were considered significant at $p < 0.05$. Phenotypic data is shown as mean \pm SD.

Results

Participant characteristics and indices of fitness

Table 1 shows participant characteristics and fitness parameters. The BMI and limb girths were largest in power athletes and smallest in endurance athletes, while team sport athletes were the tallest ($p < 0.05$). Team and power athletes had large knee extension and flexion torques, jump power and sprint running abilities than endurance athletes and controls (Table 1). Power athletes performed the largest number of pull-ups in 30 s. Anaerobic capacity, represented here as Wingate test average power and agility shuttle run time, was better in all three groups of athletes compared to controls; maximal oxygen uptake (VO_{2max}) and fatigue resistance in the Wingate anaerobic test were highest in endurance athletes.

Genotype frequencies

Table 2 shows the distribution of genotype frequencies of the studied genetic polymorphisms in the three groups. All studied polymorphisms were in Hardy-Weinberg equilibrium in all groups. *ACVR1B* was the only polymorphism which differed between the groups with G allele overrepresented over A allele among team/power athletes compared to controls and

endurance athletes. Endurance athletes had a lower frequency of the GG genotype for the *ACVR1B* (rs2854464) gene than power ($p = 0.037$) and team ($p = 0.004$) athletes. In controls, the GG genotype frequency was also less than in team athletes ($p = 0.03$). There was no significant difference in genotype score between groups.

Genotype effect on muscle size and exercise performance

Of the 16 polymorphisms tested, nine were associated with at least one of the muscular fitness and/or size phenotypes (Table 3).

The polymorphisms associated with muscular size/fitness were of seven genes (Table 3): *ACVR1B* plus *MYLK1* (rs2700352) explained 2.7% of thigh girth, and 3.3% of upper arm girth was explained by *ACVR1B*. For jumping power ($W \cdot kg^{-1}$) the GG genotype associated with larger power ($R^2 = 0.024$; $p = 0.002$). Balance and pull-ups were predicted by the combination of *ADRA2A* and *ACTN3 R577X* for 2% and 1.8%, respectively. Knee extension torque was explained for 4.1% by *ACVR1B*, *MYLK1* (rs2700352), *MYLK1* (rs820336) and *MYLK3* (rs36471), and knee flexion torque for 1.7% by the *MYLK1* (rs2700352) polymorphism. The *MYLK1* H21P polymorphism, together with *ACVR2B* (rs379527) explained 3.1% to the variation in the shuttle agility run performance. The *ACVR2B* (rs379527) explained 4.0% of the variation of the Wingate test fatigue index.

Together with age, group and height as additional factors genotypes could explain up to 20.4% (knee extension torque) of the variation in muscular fitness (Table 4).

Discussion

The current study on young Caucasian men, including athletes from various sports and healthy controls, found that nine of the 16 genetic polymorphisms tested and residing in regions coding for seven genes—myostatin, myostatin signaling receptors Activins 1B and 2B, myosin light chain kinases 1 and 3, alpha-actinin-3, and α -2-adrenergic receptor A—explained up to 4.1% of the variation of muscular fitness phenotypes either separately or when in combination. Activin receptor 1B was the only polymorphism that separated the groups, with the G allele overrepresented over the A allele among team/power athletes compared to controls and endurance athletes. The G allele of this gene was also most consistently associated with muscle size and power.

Multiple genetic polymorphisms have been identified to associate with the status of being a power athlete [30, 39]. However, the contribution of particular genetic markers for muscular fitness appears to be highly dependent on the specific characteristics of the population studied (age, ethnicity, athletic status etc.). For instance, five out of ten tested candidate SNPs were found to be associated with muscle power and sprint running performance in elite soccer players with their importance depending on the pubertal status of the athletes [40]. The polymorphisms associated with muscular fitness in our healthy young men irrespective of being a non-athlete, power, endurance or team athlete, therefore warrants testing for replication on other populations as aged people.

While myostatin (*MSTN*, *GDF-8*) is often included in power-oriented sport polygenic profiling [41], the K153R polymorphism with the strongest effect on muscle mass and strength is rather rare [42]. On its own, the K153R polymorphism explained up to 6.5% of the variance in some parameters of muscular fitness, even though only 3 subjects (two endurance and one power athlete) had the R allele in our cohort of 439 young men. The rarity of R allele in most of the world populations explains why most of the studies are underpowered to confirm phenotypic effects of this polymorphism [43–45]. However, polymorphisms of *ACVR1B* have been reported to modulate the response of muscle size and strength/power to rehabilitation of

Table 3. Correlations between polymorphisms and fitness parameters.

	ACE (rs4341)	MYLK1 (rs2700352)	MYLK1 H21P (rs28497577)	MYLK1 (rs820336)	MYLK2 (rs606965)	MYLK2 (rs4911532)	MYLK2 (rs6119729)	MYLK3 (rs56471)	MYLK3V180L (rs28407821)	MSTNK153R (rs1805086)	ACVR1B (rs2854464)	ACVR2B (rs3792527)	ACVR2B (rs7372545)	ACTN3R577X (rs1815739)	ADRA2A (rs553668)	ADRA2B (rs4066772)
Thigh girth	R = -0.025; p = 0.611	R = 0.168; p = 0.001	R = -0.121; p = 0.015	R = 0.077; p = 0.124	R = -0.097; p = 0.052	R = -0.020; p = 0.694	R = 0.013; p = 0.788	R = -0.032; p = 0.523	R = 0.029; p = 0.558	R = -0.177; p < 0.001	R = 0.127; p = 0.011	R = -0.005; p = 0.922	R = 0.012; p = 0.806	R = -0.031; p = 0.539	R = 0.107; p = 0.035	R = -0.011; p = 0.830
	$R^2_{adj} = 0.027$; p = 0.002	$R^2_{adj} = 0.027$; p = 0.002									$R^2_{adj} = 0.017$; p = 0.007					
Upper arm girth	R = 0.067; p = 0.178	R = 0.102; p = 0.041	R = 0.081; p = 0.104	R = 0.036; p = 0.471	R = -0.030; p = 0.544	R = -0.041; p = 0.412	R = 0.015; p = 0.768	R = 0.016; p = 0.741	R = 0.051; p = 0.308	R = -0.038; p = 0.444	R = 0.175; p < 0.001	R = -0.070; p = 0.165	R = -0.011; p = 0.828	R = -0.004; p = 0.944	R = 0.051; p = 0.316	R = -0.014; p = 0.785
											$R^2_{adj} = 0.033$; p < 0.001					
Vertical jump power (W.kg ⁻¹)	R = 0.002; p = 0.964	R = -0.056; p = 0.265	R = -0.115; p = 0.022	R = 0.133; p = 0.008	R = -0.098; p = 0.051	R = 0.013; p = 0.793	R = -0.002; p = 0.966	R = -0.022; p = 0.659	R = 0.020; p = 0.699	R = -0.249; p < 0.001	R = 0.113; p = 0.026	R = -0.002; p = 0.974	R = -0.071; p = 0.165	R = -0.035; p = 0.487	R = 0.045; p = 0.375	R = -0.033; p = 0.520
											$R^2_{adj} = 0.024$; p = 0.002					
Balance (no. of attempts)	R = 0.060; p = 0.231	R = -0.024; p = 0.635	R = 0.085; p = 0.090	R = -0.095; p = 0.058	R = 0.056; p = 0.262	R = -0.031; p = 0.541	R = -0.033; p = 0.505	R = 0.060; p = 0.231	R = -0.011; p = 0.827	R = 0.255; p < 0.001	R = 0.111; p = 0.824	R = -0.031; p = 0.542	R = 0.019; p = 0.713	R = -0.085; p = 0.091	R = 0.043; p = 0.393	R = -0.049; p = 0.324
														$R^2_{adj} = 0.010$; p = 0.026		
Handgrip strength	R = 0.061; p = 0.222	R = -0.011; p = 0.821	R = 0.011; p = 0.828	R = 0.011; p = 0.833	R = 0.018; p = 0.726	R = -0.035; p = 0.481	R = 0.076; p = 0.131	R = -0.027; p = 0.588	R = 0.006; p = 0.900	R = -0.070; p = 0.165	R = 0.092; p = 0.067	R = -0.056; p = 0.269	R = -0.068; p = 0.177	R = 0.001; p = 0.988	R = 0.038; p = 0.454	R = -0.062; p = 0.220
Knee extension peak torque	R = 0.026; p = 0.607	R = 0.112; p = 0.027	R = 0.043; p = 0.398	R = 0.107; p = 0.035	R = -0.078; p = 0.124	R = -0.003; p = 0.956	R = 0.055; p = 0.283	R = 0.101; p = 0.046	R = 0.049; p = 0.339	R = -0.026; p = 0.614	R = 0.114; p = 0.025	R = -0.007; p = 0.899	R = -0.025; p = 0.628	R = -0.053; p = 0.296	R = 0.033; p = 0.522	R = -0.031; p = 0.542
Knee flexion peak torque	R = 0.017; p = 0.738	R = 0.152; p = 0.003	R = -0.075; p = 0.140	R = 0.064; p = 0.211	R = -0.049; p = 0.339	R = 0.027; p = 0.592	R = 0.012; p = 0.818	R = 0.041; p = 0.420	R = 0.008; p = 0.873	R = 0.032; p = 0.523	R = 0.076; p = 0.139	R = -0.074; p = 0.149	R = -0.027; p = 0.602	R = -0.024; p = 0.645	R = 0.064; p = 0.214	R = 0.016; p = 0.747
Shuttle agility run (s)	R = -0.019; p = 0.713	R = 0.069; p = 0.176	R = 0.102; p = 0.044	R = 0.038; p = 0.457	R = 0.057; p = 0.265	R = -0.077; p = 0.131	R = 0.001; p = 0.982	R = -0.027; p = 0.600	R = 0.035; p = 0.486	R = -0.047; p = 0.354	R = 0.047; p = 0.361	R = 0.149; p = 0.003	R = -0.023; p = 0.660	R = 0.042; p = 0.414	R = 0.054; p = 0.296	R = -0.090; p = 0.075
30 m run (s)	R = -0.026; p = 0.610	R = -0.012; p = 0.809	R = 0.045; p = 0.369	R = -0.030; p = 0.546	R = 0.045; p = 0.368	R = -0.013; p = 0.789	R = -0.003; p = 0.959	R = 0.037; p = 0.466	R = 0.024; p = 0.631	R = -0.064; p = 0.205	R = -0.089; p = 0.079	R = -0.084; p = 0.097	R = 0.017; p = 0.734	R = 0.010; p = 0.844	R = -0.046; p = 0.368	R = -0.008; p = 0.868
Wingate (rev.30 s ⁻¹)	R = -0.066; p = 0.195	R = -0.027; p = 0.599	R = 0.051; p = 0.322	R = 0.038; p = 0.451	R = 0.054; p = 0.286	R = -0.045; p = 0.378	R = -0.011; p = 0.836	R = -0.021; p = 0.682	R = -0.063; p = 0.216	R = 0.008; p = 0.875	R = 0.090; p = 0.078	R = 0.106; p = 0.039	R = -0.056; p = 0.274	R = -0.045; p = 0.381	R = 0.046; p = 0.374	R = -0.020; p = 0.690
Wingate fatigue resistance	R = -0.066; p = 0.197	R = 0.042; p = 0.410	R = -0.083; p = 0.101	R = 0.021; p = 0.677	R = 0.017; p = 0.742	R = 0.040; p = 0.429	R = -0.026; p = 0.606	R = 0.024; p = 0.632	R = 0.038; p = 0.458	R = -0.054; p = 0.287	R = 0.027; p = 0.592	R = 0.202; p < 0.001	R = 0.077; p = 0.136	R = -0.038; p = 0.459	R = 0.006; p = 0.901	R = 0.027; p = 0.596

(Continued)

Table 3. (Continued)

	ACE (rs4341)	MYLK1 (rs2700352)	MYLK1 H2IP (rs28497577)	MYLK1 (rs820336)	MYLK2 (rs6060965)	MYLK2 (rs4911532)	MYLK2 (rs6119729)	MYLK3 (rs36471)	MYLK3VI180L (rs28407821)	MSTNK153R (rs1805086)	ACVR2B (rs3792527)	ACVR2B (rs7372545)	ACTN3R577X (rs1815739)	ADRA2A (rs553668)	ADRA2B (rs4066772)
Pull-ups (no.)	R = 0.041; p = 0.423	R = -0.036; p = 0.487	R = -0.073; p = 0.154	R = 0.009; p = 0.864	R = 0.043; p = 0.378	R = -0.052; p = 0.315	R = 0.078; p = 0.130	R = 0.049; p = 0.340	R = -0.031; p = 0.539	R = 0.002; p = 0.970	R = -0.018; p = 0.729	R = -0.014; p = 0.791	R = -0.086; p = 0.094	R = -0.088; p = 0.092	R = -0.034; p = 0.502
VO ₂ max (ml·kg ⁻¹ ·min ⁻¹)	R = -0.089; p = 0.097	R = 0.004; p = 0.938	R = -0.066; p = 0.216	R = 0.054; p = 0.311	R = -0.035; p = 0.513	R = 0.055; p = 0.308	R = -0.016; p = 0.760	R = 0.016; p = 0.766	R = -0.006; p = 0.904	R = -0.216; p < 0.001	R = -0.069; p = 0.201	R = -0.005; p = 0.932	R = 0.040; p = 0.460	R = 0.057; p = 0.294	R = 0.031; p = 0.562

If there is more than 1 row for a phenotype, the second row indicates which polymorphism is most explanatory in the darkest grey, with the contribution of additional polymorphisms explaining the phenotype to a lesser and lesser, but still significant, extent by lighter grey fields (showing increasing R²_{adj} and P values).
VO₂max, maximal oxygen uptake.

<https://doi.org/10.1371/journal.pone.0275179.t003>

Table 4. Adjusted determination coefficients (R^2_{adj}) of height, age, group and polymorphisms on the muscularity and fitness.

	Height	Age	Group	ACVR1B	ACVR2B	MYLK1	MYLK1 H21P	ADRA2B
				(rs2854464)	(rs379527)	(rs2700352)	(rs28497577)	(rs4066772)
Thigh girth	<u>0.096</u> ***	+0.006***→0.111	+0.009***→0.105	+0.020***→ 0.131				
Upper arm girth	<u>0.057</u> ***	+0.038***→0.095	+0.017***→ 0.146	+0.036***→0.129				
Handgrip strength	<u>0.014</u> ***							
Vertical jump power ($W \cdot kg^{-1}$)			0.080***	+0.015***→ 0.095				
Isokinetic knee extension torque/thigh girth	<u>0.248</u> ***		+0.053***→ 0.301					
Shuttle agility run (s)					<u>0.018</u> **		+0.013**→0.031	+0.008**→ 0.039
Wingate fatigue resistance		+0.014***→ 0.096	<u>0.049</u> ***		+0.033**→0.082			
VO ₂ max ($ml \cdot min^{-1} \cdot kg^{-1}$)	+0.009***→0.102	<u>0.093</u> ***	+0.009***→ 0.120	+0.009***→0.111				
Knee extension peak torque	+0.038***→0.188		<u>0.150</u> ***	+0.008***→0.196		+0.008***→ 0.204		

Note. Only the significant determination coefficients

(*** @ $P < 0.001$

** @ $P < 0.01$) are presented. The coefficient of prime determinant of the phenotype is underscored, and the final estimation of the explained variance is highlighted in **bold**.

VO₂max, maximal oxygen uptake.

<https://doi.org/10.1371/journal.pone.0275179.t004>

cardiac patients [46], and the A allele (rs2854464) has been reported to be associated with sprint/power performance, depending on the ethnicity of the athletes [47]. Although it has been reported that the genetic polymorphisms of the myostatin signalling pathway are more strongly associated with muscle strength than muscle mass [16, 48], in our cohort of young men the ACVR1B genotype was associated more with muscle size than strength (Table 3). While a positive association between ACVR1B A allele and muscle strength has been found in untrained men [18, 19], we found that the G allele was positively associated with muscle fitness phenotypes in our cohort of young individuals, and was also more prevalent among power and team sports athletes than in endurance athletes and non-athletes. This suggests that contradictory findings between the studies may occur also because of the different training status of subjects.

In another case-control study there was a trend towards lower frequency of ACVR1B A allele in Brazilian athletes, while Caucasian Power athletes had a higher frequency of the A allele compared to controls [47]. One may argue that these conflicting findings are due to different gene interactions in the two ethnicities, or just limitations of sample size. However, a

more likely explanation for the discrepancy is the dissimilar representation of the Power sport disciplines and training background between the two cohorts as in our Caucasian cohort we found that the A allele of the *ACVR1B* gene was underrepresented in Team sports athletes compared to Controls, and the A allele was also less frequent in both Team and Power compared to Endurance athletes. In this context, it is important to note that downregulated myostatin pathway activity is counterproductive for the endurance capacity [49, 50].

Caution should be exercised as to whether the inhibition of MSTN signaling benefits, as occurs in the *ACVR1B* G allele, muscle function since, for instance, the downregulation of activin 1 receptor (*Alk4*) expression not only decreased skeletal muscle mass to an even lower level in mdx mice (model of muscular dystrophy) but also induced marked muscle atrophy in normal animals [51]. In addition, lower myostatin signalling activity has recently been reported in patients with inflammatory myopathies as compared with healthy subjects, and there was no clear association between the activity of the myostatin signalling pathway and the functional state of the patients [52].

As with the ACE I/D polymorphism, studies on the effects of *ACTN3* R577X polymorphism on muscle characteristics have yielded so far inconsistent results [53], but the general impression is that the R-allele harbours benefits for sprinting performance and muscle power [54]. Also, the muscle damage response to intense exercise is not consistently associated with the *ACTN3* genotype [37, 55, 56]. Here we detected only a minor contribution of *ACTN3* R577X SNP to muscular fitness (adding not more than 1% of the explanation of the variance in two of the muscle fitness tests out of eight implemented) and any association to sprinting or jumping performance, which confirms the limited role of this genotype on strength and power in the healthy young population of European decent. This is in line with our previous observation that even though 2.4% of the variance in 40 m sprint run time has been explained by R577X polymorphism in Greek schoolboys, this was not the case in schoolgirls [38].

It has been suggested that polymorphisms of the skeletal muscle isoform of myosin light chain kinase (*MYLK2*) associates with exertional muscle damage [29] and may thus modulate the training response. It appears, however, that Clarkson et al. 2005 [29] did not study the skeletal muscle isoforms of *MYLK* and this may explain the discrepancy from the current study where we found no associations to muscular phenotypes with variants of *MYLK2*, while there were associations with genetic variants of *MYLK1* (smooth muscle isoform) and *MYLK3* (myocardial isoform). The *MYLK1* rs28497577 GT genotype has recently been shown to associate with slower recovery after knee injury in soccer players [57]. Intriguingly, polymorphisms of *MYLK1* might modulate the pro-inflammatory state of the bowel and nutrient absorption [58], which are important for the anabolic response in resistance training. It was surprising to see that polymorphism of cardiac myosin light chain kinase isoform (*MYLK3*) associates with muscle strength. This finding is new and needs confirmation by other studies. However, *MYLK3* plays an important role in sarcomerogenesis in the heart [59] and its detection in skeletal muscle [60] suggests a similar role in skeletal muscle sarcomere assembly.

Study limitations

One limitation is that we have pre-selected the set of candidate genes rather than screened for polymorphisms across the genome to pick up most of genetic markers associated with the muscular fitness of young healthy subjects. As in most other genotype-phenotype association studies, especially those of large cohorts of athletes, we have limited our study population to men and therefore direct extrapolation of the findings to women may not be applicable, as some other studies on functional SNPs on muscular fitness phenotypes have shown sexually dimorphic outcome [38, 61].

Study participants included into each of the Endurance, Power and Team sports category were from different sports and performance level. In addition, heterogeneity within the groups of athletes was high due to some of them being trained largely for their upper body while most of the athletes largely trained the leg or all body muscles. This disproportionally enhanced training of the upper body occurred particularly in the Power athletes where participants included not only sprinters but also body builders that clearly require different adaptations to succeed in their sport. However, they all trained their muscles with mostly high to maximal intensity short bursts of exercise, thus must develop overlapping adaptations which are somewhat contrasting to those of the endurance athletes.

Finally, while the study population used in the current study is smaller than that used in some association-seeking studies, it is not of unusual sample size of many other similar studies [62, 63]. In fact, the sample size was adequate to detect significant, and hence physiologically meaningful, associations, and one may query the physiological significance of any additional association if it can only be detected with investigating very large populations.

Conclusions

Genetic polymorphisms in myostatin (*MSTNK153R*), the myostatin receptors Activins 1B (*ACVR1B*) and 2B (*ACVR2B*), myosin light chain kinases (*MYLK*) isoforms 1 and 3, and α -2-adrenergic receptor, and actinin (*ACTN3R577X*) were found to explain 1–4% of the variation in muscle fitness in young healthy male Caucasians, either separately or when combined. *ACVR1B* and *ACVR2B* polymorphisms associated independently to both muscle size and functional measures of muscular fitness, and the allele frequency distribution of *ACVR1B* rs2854464 differed between groups, with power and team athletes having higher frequency of the G allele. Variation in *MYLK* 1 and 3 genes contributes to some of the phenotypic variation of muscle size and power. Even though the contribution of these genetic polymorphisms to muscular fitness was quite large (up to 4.1%) given that only a limited number of gene polymorphisms were analysed, this also indicates that many more polymorphisms contribute to the 50% heredity of muscle mass and strength as has been recently shown by another group of researchers [64].

Author Contributions

Conceptualization: Tomas Venckunas, Hans Degens.

Data curation: Tomas Venckunas.

Formal analysis: Tomas Venckunas, Hans Degens.

Investigation: Tomas Venckunas.

Methodology: Tomas Venckunas, Hans Degens.

Resources: Tomas Venckunas.

Validation: Tomas Venckunas.

Writing – original draft: Tomas Venckunas, Hans Degens.

Writing – review & editing: Tomas Venckunas, Hans Degens.

References

1. Artero EG, Ruiz JR, Ortega FB, España-Romero V, Vicente-Rodríguez G, Molnar D, et al. Muscular and cardiorespiratory fitness are independently associated with metabolic risk in adolescents: the

- HELENA study. *Pediatr Diabetes*. 2011 Dec; 12(8):704–12. <https://doi.org/10.1111/j.1399-5448.2011.00769.x> PMID: 21470352
2. Schousboe K, Visscher PM, Erbas B, Kyvik KO, Hopper JL, Henriksen JE, et al. Twin study of genetic and environmental influences on adult body size, shape, and composition. *Int J Obes Relat Metab Disord*. 2004 Jan; 28(1):39–48. <https://doi.org/10.1038/sj.ijo.0802524> PMID: 14610529
 3. Pescatello LS, Devaney JM, Hubal MJ, Thompson PD, Hoffman EP. Highlights from the functional single nucleotide polymorphisms associated with human muscle size and strength or FAMuSS study. *Biomed Res Int*. 2013; 2013:643575. <https://doi.org/10.1155/2013/643575> PMID: 24455711
 4. Fagard R, Bielen E, Amery A. Heritability of aerobic power and anaerobic energy generation during exercise. *J Appl Physiol* (1985). 1991 Jan; 70(1):357–62. <https://doi.org/10.1152/jappl.1991.70.1.357> PMID: 2010392
 5. Thompson PD, Moyna N, Seip R, Price T, Clarkson P, Angelopoulos T, et al. Functional polymorphisms associated with human muscle size and strength. *Med Sci Sports Exerc*. 2004 Jul; 36(7):1132–9. <https://doi.org/10.1249/01.mss.0000132274.26612.23> PMID: 15235316
 6. Bray MS, Hagberg JM, Pérusse L, Rankinen T, Roth SM, Wolfarth B, et al. The human gene map for performance and health-related fitness phenotypes: the 2006–2007 update. *Med Sci Sports Exerc*. 2009 Jan; 41(1):35–73. <https://doi.org/10.1249/mss.0b013e3181844179> PMID: 19123262
 7. Santiago C, Ruiz JR, Muniesa CA, González-Freire M, Gómez-Gallego F, Lucia A. Does the polygenic profile determine the potential for becoming a world-class athlete? Insights from the sport of rowing. *Scand J Med Sci Sports*. 2010 Feb; 20(1):e188–194. <https://doi.org/10.1111/j.1600-0838.2009.00943.x> PMID: 19422651
 8. Sarzynski MA, Loos RJF, Lucia A, Pérusse L, Roth SM, Wolfarth B, et al. Advances in Exercise, Fitness, and Performance Genomics in 2015. *Med Sci Sports Exerc*. 2016 Oct; 48(10):1906–16. <https://doi.org/10.1249/MSS.0000000000000982> PMID: 27183119
 9. Broos S, Van Leemputte M, Deldicque L, Thomis MA. History-dependent force, angular velocity and muscular endurance in ACTN3 genotypes. *Eur J Appl Physiol*. 2015 Aug; 115(8):1637–43. <https://doi.org/10.1007/s00421-015-3144-6> PMID: 25761730
 10. Erskine RM, Williams AG, Jones DA, Stewart CE, Degens H. The individual and combined influence of ACE and ACTN3 genotypes on muscle phenotypes before and after strength training. *Scand J Med Sci Sports*. 2014 Aug; 24(4):642–8. <https://doi.org/10.1111/sms.12055> PMID: 23384112
 11. Williams AG, Day SH, Folland JP, Gohlke P, Dhamrait S, Montgomery HE. Circulating angiotensin converting enzyme activity is correlated with muscle strength. *Med Sci Sports Exerc*. 2005 Jun; 37(6):944–8. PMID: 15947718
 12. Nazarov IB, Woods DR, Montgomery HE, Shneider OV, Kazakov VI, Tomilin NV, et al. The angiotensin converting enzyme I/D polymorphism in Russian athletes. *Eur J Hum Genet*. 2001 Oct; 9(10):797–801. <https://doi.org/10.1038/sj.ejhg.5200711> PMID: 11781693
 13. Schuelke M, Wagner KR, Stolz LE, Hübner C, Riebel T, Kömen W, et al. Myostatin mutation associated with gross muscle hypertrophy in a child. *N Engl J Med*. 2004 Jun 24; 350(26):2682–8. <https://doi.org/10.1056/NEJMoa040933> PMID: 15215484
 14. Saunders MA, Good JM, Lawrence EC, Ferrell RE, Li WH, Nachman MW. Human adaptive evolution at Myostatin (GDF8), a regulator of muscle growth. *Am J Hum Genet*. 2006 Dec; 79(6):1089–97. <https://doi.org/10.1086/509707> PMID: 17186467
 15. Amthor H, Macharia R, Navarrete R, Schuelke M, Brown SC, Otto A, et al. Lack of myostatin results in excessive muscle growth but impaired force generation. *Proc Natl Acad Sci U S A*. 2007 Feb 6; 104(6):1835–40. <https://doi.org/10.1073/pnas.0604893104> PMID: 17267614
 16. Huygens W, Thomis MA, Peeters MW, Aerssens J, Vlietinck R, Beunen GP. Quantitative trait loci for human muscle strength: linkage analysis of myostatin pathway genes. *Physiol Genomics*. 2005 Aug 11; 22(3):390–7. <https://doi.org/10.1152/physiolgenomics.00010.2005> PMID: 15914581
 17. Walsh S, Metter EJ, Ferrucci L, Roth SM. Activin-type II receptor B (ACVR2B) and follistatin haplotype associations with muscle mass and strength in humans. *J Appl Physiol* (1985). 2007 Jun; 102(6):2142–8.
 18. Windelinckx A, De Mars G, Huygens W, Peeters MW, Vincent B, Wijmenga C, et al. Comprehensive fine mapping of chr12q12-14 and follow-up replication identify activin receptor 1B (ACVR1B) as a muscle strength gene. *Eur J Hum Genet*. 2011 Feb; 19(2):208–15. <https://doi.org/10.1038/ejhg.2010.173> PMID: 21063444
 19. He L, Van Roie E, Bogaerts A, Morse CI, Delecluse C, Verschueren S, et al. Genetic predisposition score predicts the increases of knee strength and muscle mass after one-year exercise in healthy elderly. *Exp Gerontol*. 2018 Oct 1; 111:17–26. <https://doi.org/10.1016/j.exger.2018.06.030> PMID: 29991458

20. Klimentidis YC, Bea JW, Thompson P, Klimecki WT, Hu C, Wu G, et al. Genetic Variant in ACVR2B Is Associated with Lean Mass. *Med Sci Sports Exerc.* 2016 Jul; 48(7):1270–5. <https://doi.org/10.1249/MSS.0000000000000889> PMID: 26848890
21. Kostek MA, Angelopoulos TJ, Clarkson PM, Gordon PM, Moyna NM, Visich PS, et al. Myostatin and follistatin polymorphisms interact with muscle phenotypes and ethnicity. *Med Sci Sports Exerc.* 2009 May; 41(5):1063–71. <https://doi.org/10.1249/MSS.0b013e3181930337> PMID: 19346981
22. Seibert MJ, Xue QL, Fried LP, Walston JD. Polymorphic variation in the human myostatin (GDF-8) gene and association with strength measures in the Women's Health and Aging Study II cohort. *J Am Geriatr Soc.* 2001 Aug; 49(8):1093–6. <https://doi.org/10.1046/j.1532-5415.2001.49214.x> PMID: 11555072
23. Santiago C, Ruiz JR, Rodríguez-Romo G, Fiuza-Luces C, Yvert T, Gonzalez-Freire M, et al. The K153R polymorphism in the myostatin gene and muscle power phenotypes in young, non-athletic men. *PLoS One.* 2011 Jan 20; 6(1):e16323. <https://doi.org/10.1371/journal.pone.0016323> PMID: 21283721
24. Kim J, Park K, Lee J. Myostatin A55T Genotype is Associated with Strength Recovery Following Exercise-Induced Muscle Damage. *Int J Environ Res Public Health.* 2020 Jul 7; 17(13):E4900. <https://doi.org/10.3390/ijerph17134900> PMID: 32645967
25. Li X, Wang SJ, Tan SC, Chew PL, Liu L, Wang L, et al. The A55T and K153R polymorphisms of MSTN gene are associated with the strength training-induced muscle hypertrophy among Han Chinese men. *J Sports Sci.* 2014; 32(9):883–91. <https://doi.org/10.1080/02640414.2013.865252> PMID: 24479661
26. Urano T, Inoue S. Recent genetic discoveries in osteoporosis, sarcopenia and obesity. *Endocr J.* 2015; 62(6):475–84. <https://doi.org/10.1507/endocrj.EJ15-0154> PMID: 25866211
27. Zillikens MC, Demissie S, Hsu YH, Yerges-Armstrong LM, Chou WC, Stolk L, et al. Large meta-analysis of genome-wide association studies identifies five loci for lean body mass. *Nat Commun.* 2017 Jul 19; 8(1):80. <https://doi.org/10.1038/s41467-017-00031-7> PMID: 28724990
28. Hernandez Cordero AI, Gonzales NM, Parker CC, Sokolof G, Vandenberg DJ, Cheng R, et al. Genome-wide Associations Reveal Human-Mouse Genetic Convergence and Modifiers of Myogenesis, CPNE1 and STC2. *Am J Hum Genet.* 2019 Dec 5; 105(6):1222–36. <https://doi.org/10.1016/j.ajhg.2019.10.014> PMID: 31761296
29. Clarkson PM, Devaney JM, Gordish-Dressman H, Thompson PD, Hubal MJ, Urso M, et al. ACTN3 genotype is associated with increases in muscle strength in response to resistance training in women. *J Appl Physiol (1985).* 2005 Jul; 99(1):154–63. <https://doi.org/10.1152/jappphysiol.01139.2004> PMID: 15718405
30. Eynon N, Alves AJ, Meckel Y, Yamin C, Ayalon M, Sagiv M, et al. Is the interaction between HIF1A P582S and ACTN3 R577X determinant for power/sprint performance? *Metabolism.* 2010 Jun; 59(6):861–5. <https://doi.org/10.1016/j.metabol.2009.10.003> PMID: 20005538
31. Charlier R, Caspers M, Knaeps S, Mertens E, Lambrechts D, Lefevre J, et al. Limited potential of genetic predisposition scores to predict muscle mass and strength performance in Flemish Caucasians between 19 and 73 years of age. *Physiol Genomics.* 2017 Mar 1; 49(3):160–6. <https://doi.org/10.1152/physiolgenomics.00085.2016> PMID: 28039429
32. Karaliute R, Raugaliene R, Venckunas T. Relationship between left ventricular structure and post-exercise blood pressure in endurance athletes. *Acta Cardiol.* 2011 Jun; 66(3):359–63. <https://doi.org/10.1080/ac.66.3.2114136> PMID: 21744706
33. Malinauskas R, Dumciene A, Mamkus G, Venckunas T. Personality traits and exercise capacity in male athletes and non-athletes. *Percept Mot Skills.* 2014 Feb; 118(1):145–61. <https://doi.org/10.2466/29.25.PMS.118k13w1> PMID: 24724519
34. Venckunas T, Emeljanovas A, Mieziene B, Volbekiene V. Secular trends in physical fitness and body size in Lithuanian children and adolescents between 1992 and 2012. *J Epidemiol Community Health.* 2017 Feb; 71(2):181–7. <https://doi.org/10.1136/jech-2016-207307> PMID: 27485051
35. Degens H, Stasiulis A, Skurvydas A, Statkeviciene B, Venckunas T. Physiological comparison between non-athletes, endurance, power and team athletes. *Eur J Appl Physiol.* 2019 Jun; 119(6):1377–86. <https://doi.org/10.1007/s00421-019-04128-3> PMID: 30919126
36. Christou M, Smilios I, Sotiropoulos K, Volaklis K, Piliandis T, Tokmakidis SP. Effects of resistance training on the physical capacities of adolescent soccer players. *J Strength Cond Res.* 2006 Nov; 20(4):783–91. <https://doi.org/10.1519/R-17254.1> PMID: 17194231
37. Venckunas T, Skurvydas A, Brazaitis M, Kamandulis S, Snieckus A, Moran CN. Human alpha-actinin-3 genotype association with exercise-induced muscle damage and the repeated-bout effect. *Appl Physiol Nutr Metab.* 2012 Dec; 37(6):1038–46. <https://doi.org/10.1139/h2012-087> PMID: 22891846
38. Moran CN, Yang N, Bailey MES, Tsiokanos A, Jamurtas A, MacArthur DG, et al. Association analysis of the ACTN3 R577X polymorphism and complex quantitative body composition and performance

- phenotypes in adolescent Greeks. *Eur J Hum Genet*. 2007 Jan; 15(1):88–93. <https://doi.org/10.1038/sj.ejhg.5201724> PMID: 17033684
39. Weyerstraß J, Stewart K, Wesselius A, Zeegers M. Nine genetic polymorphisms associated with power athlete status—A Meta-Analysis. *J Sci Med Sport*. 2018 Feb; 21(2):213–20. <https://doi.org/10.1016/j.jsams.2017.06.012> PMID: 28666769
 40. Murtagh CF, Brownlee TE, Rienzi E, Roquero S, Moreno S, Huertas G, et al. The genetic profile of elite youth soccer players and its association with power and speed depends on maturity status. *PLoS One*. 2020; 15(6):e0234458. <https://doi.org/10.1371/journal.pone.0234458> PMID: 32569264
 41. Ruiz JR, Arteta D, Buxens A, Artieda M, Gómez-Gallego F, Santiago C, et al. Can we identify a power-oriented polygenic profile? *J Appl Physiol* (1985). 2010 Mar; 108(3):561–6. <https://doi.org/10.1152/jappphysiol.01242.2009> PMID: 20044471
 42. Thomis MAI, Huygens W, Heuninckx S, Chagnon M, Maes HHM, Claessens AL, et al. Exploration of myostatin polymorphisms and the angiotensin-converting enzyme insertion/deletion genotype in responses of human muscle to strength training. *Eur J Appl Physiol*. 2004 Jul; 92(3):267–74. <https://doi.org/10.1007/s00421-004-1093-6> PMID: 15083369
 43. Ferrell RE, Conte V, Lawrence EC, Roth SM, Hagberg JM, Hurley BF. Frequent sequence variation in the human myostatin (GDF8) gene as a marker for analysis of muscle-related phenotypes. *Genomics*. 1999 Dec 1; 62(2):203–7. <https://doi.org/10.1006/geno.1999.5984> PMID: 10610713
 44. Fernández-Santander A, Valveny N, Harich N, Kandil M, Luna F, Martín MÁ, et al. Polymorphisms influencing muscle phenotypes in North-African and Spanish populations. *Ann Hum Biol*. 2012 Mar; 39(2):166–9. <https://doi.org/10.3109/03014460.2012.657243> PMID: 22324844
 45. Muniesa CA, González-Freire M, Santiago C, Lao JI, Buxens A, Rubio JC, et al. World-class performance in lightweight rowing: is it genetically influenced? A comparison with cyclists, runners and non-athletes. *Br J Sports Med*. 2010 Sep; 44(12):898–901. <https://doi.org/10.1136/bjism.2008.051680> PMID: 18801770
 46. Thomaes T, Thomis M, Onkelinx S, Goetschalckx K, Fagard R, Lambrechts D, et al. Genetic predisposition scores associate with muscular strength, size, and trainability. *Med Sci Sports Exerc*. 2013 Aug; 45(8):1451–9. <https://doi.org/10.1249/MSS.0b013e31828983f7> PMID: 23439425
 47. Voisin S, Guilherme JPFL, Yan X, Pushkarev VP, Cieszczyk P, Massidda M, et al. ACVR1B rs2854464 Is Associated with Sprint/Power Athletic Status in a Large Cohort of Europeans but Not Brazilians. *PLoS One*. 2016; 11(6):e0156316. <https://doi.org/10.1371/journal.pone.0156316> PMID: 27253421
 48. Huygens W, Thomis MA, Peeters MW, Aerssens J, Janssen R, Vlietinck RF, et al. Linkage of myostatin pathway genes with knee strength in humans. *Physiol Genomics*. 2004 May 19; 17(3):264–70. <https://doi.org/10.1152/physiolgenomics.00224.2003> PMID: 15026560
 49. Mouisel E, Relizani K, Mille-Hamard L, Denis R, Hourd C, Agbulut O, et al. Myostatin is a key mediator between energy metabolism and endurance capacity of skeletal muscle. *Am J Physiol Regul Integr Comp Physiol*. 2014 Aug 15; 307(4):R444–454. <https://doi.org/10.1152/ajpregu.00377.2013> PMID: 24965795
 50. Relizani K, Mouisel E, Giannesini B, Hourd C, Patel K, Morales Gonzalez S, et al. Blockade of ActRIIB signaling triggers muscle fatigability and metabolic myopathy. *Mol Ther*. 2014 Aug; 22(8):1423–33. <https://doi.org/10.1038/mt.2014.90> PMID: 24861054
 51. Pasteuning-Vuhman S, Boertje-van der Meulen JW, van Putten M, Overzier M, Ten Dijke P, Kielbasa SM, et al. New function of the myostatin/activin type I receptor (ALK4) as a mediator of muscle atrophy and muscle regeneration. *FASEB J*. 2017 Jan; 31(1):238–55. <https://doi.org/10.1096/fj.201600675R> PMID: 27733450
 52. Vernerová L, Horváthová V, Kropáčková T, Vokurková M, Klein M, Tomčík M, et al. Alterations in activin A-myostatin-follistatin system associate with disease activity in inflammatory myopathies. *Rheumatology (Oxford)*. 2020 Sep 1; 59(9):2491–501. <https://doi.org/10.1093/rheumatology/kez651> PMID: 31990347
 53. Gineviciene V, Jakaitiene A, Aksenov MO, Aksenova AV, Druzhevskaya AM, Astratenkova IV, et al. Association analysis of ACE, ACTN3 and PPARGC1A gene polymorphisms in two cohorts of European strength and power athletes. *Biol Sport*. 2016 Sep; 33(3):199–206. <https://doi.org/10.5604/20831862.1201051> PMID: 27601773
 54. Houweling PJ, Papadimitriou ID, Seto JT, Pérez LM, Coso JD, North KN, et al. Is evolutionary loss our gain? The role of ACTN3 p.Arg577Ter (R577X) genotype in athletic performance, ageing, and disease. *Hum Mutat*. 2018 Dec; 39(12):1774–87. <https://doi.org/10.1002/humu.23663> PMID: 30281865
 55. Vincent B, Windelinckx A, Nielens H, Ramaekers M, Van Leemputte M, Hespel P, et al. Protective role of alpha-actinin-3 in the response to an acute eccentric exercise bout. *J Appl Physiol* (1985). 2010 Aug; 109(2):564–73.

56. Coelho DB, Pimenta EM, Rosse IC, Veneroso C, Pussieldi GDA, Becker LK, et al. Alpha-Actinin-3 R577X Polymorphism Influences Muscle Damage and Hormonal Responses After a Soccer Game. *J Strength Cond Res.* 2019 Oct; 33(10):2655–64. <https://doi.org/10.1519/JSC.0000000000002575> PMID: 29781940
57. Hall ECR, Baumert P, Larruskain J, Gil SM, Lekue JA, Rienzi E, et al. The genetic association with injury risk in male academy soccer players depends on maturity status. *Scand J Med Sci Sports.* 2022 Feb; 32(2):338–50. <https://doi.org/10.1111/sms.14077> PMID: 34633711
58. Graham WV, He W, Marchiando AM, Zha J, Singh G, Li HS, et al. Intracellular MLCK1 diversion reverses barrier loss to restore mucosal homeostasis. *Nat Med.* 2019 Apr; 25(4):690–700. <https://doi.org/10.1038/s41591-019-0393-7> PMID: 30936544
59. Seguchi O, Takashima S, Yamazaki S, Asakura M, Asano Y, Shintani Y, et al. A cardiac myosin light chain kinase regulates sarcomere assembly in the vertebrate heart. *J Clin Invest.* 2007 Oct; 117(10):2812–24. <https://doi.org/10.1172/JCI30804> PMID: 17885681
60. Liu R, Han M, Liu X, Yu K, Bai X, Dong Y. Genome-Wide Identification and Characterization of Long Non-Coding RNAs in Longissimus dorsi Skeletal Muscle of Shandong Black Cattle and Luxi Cattle. *Front Genet.* 2022; 13:849399. <https://doi.org/10.3389/fgene.2022.849399> PMID: 35651943
61. Sprouse C, Gordish-Dressman H, Orkunoglu-Suer EF, Lipof JS, Moeckel-Cole S, Patel RR, et al. SLC30A8 nonsynonymous variant is associated with recovery following exercise and skeletal muscle size and strength. *Diabetes.* 2014 Jan; 63(1):363–8. <https://doi.org/10.2337/db13-1150> PMID: 24101675
62. Khanal P, He L, Stebbings G, Onambele-Pearson GL, Degens H, Williams A, et al. Prevalence and association of single nucleotide polymorphisms with sarcopenia in older women depends on definition. *Sci Rep.* 2020 Feb 19; 10(1):2913. <https://doi.org/10.1038/s41598-020-59722-9> PMID: 32076017
63. Khanal P, Morse CI, He L, Herbert AJ, Onambélé-Pearson GL, Degens H, et al. Polygenic Models Partially Predict Muscle Size and Strength but Not Low Muscle Mass in Older Women. *Genes (Basel).* 2022 May 30; 13(6):982.
64. Moreland E, Borisov OV, Semenova EA, Larin AK, Andryushchenko ON, Andryushchenko LB, et al. Polygenic Profile of Elite Strength Athletes. *J Strength Cond Res.* 2020 Dec 3. <https://doi.org/10.1519/JSC.0000000000003901> PMID: 33278272