

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

---

4-28-2022

## Agricultural intensification can help protect the Amazon Forest and reduce global warming / Protecting the Amazon forest and reducing global warming via agricultural intensification

Fabio R. Marin

*Luiz de Queiroz College of Agriculture (ESALQ), University of São Paulo*

Alencar J. Zanon

*Universidade Federal de Santa Maria*

Juan P. Monzon

*University of Nebraska-Lincoln*

José F. Andrade

*University of Nebraska-Lincoln, jandrade@unl.edu*

Evandro H.F.M. Silva

*Luiz de Queiroz College of Agriculture (ESALQ), University of São Paulo*

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Botany Commons](#), [Horticulture Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

---

Marin, Fabio R.; Zanon, Alencar J.; Monzon, Juan P.; Andrade, José F.; Silva, Evandro H.F.M.; Richter, Gean L.; Antolin, Luis A.S.; Ribeiro, Bruna S.M.R.; Ribas, Giovanna G.; Battisti, Rafael; Heinemann, Alexandre B.; and Grassini, Patricio, "Agricultural intensification can help protect the Amazon Forest and reduce global warming / Protecting the Amazon forest and reducing global warming via agricultural intensification" (2022). *Agronomy & Horticulture -- Faculty Publications*. 1584.  
<https://digitalcommons.unl.edu/agronomyfacpub/1584>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Fabio R. Marin, Alencar J. Zanon, Juan P. Monzon, José F. Andrade, Evandro H.F.M. Silva, Gean L. Richter, Luis A.S. Antolin, Bruna S.M.R. Ribeiro, Giovanna G. Ribas, Rafael Battisti, Alexandre B. Heinemann, and Patricio Grassini

# Agricultural intensification can help protect the Amazon Forest and reduce global warming

**Fabio Marin**

Luiz de Queiroz College of Agriculture (ESALQ), University of São Paulo

**Alencar Zanon**

Universidade Federal de Santa Maria

**Juan Pablo Monzon**

University of Nebraska-Lincoln <https://orcid.org/0000-0001-6992-1842>

**Jose Andrade**

University of Nebraska-Lincoln <https://orcid.org/0000-0002-8635-7320>

**Evandro Silva**

Luiz de Queiroz College of Agriculture (ESALQ), University of São Paulo

**Luis Antolin**

Luiz de Queiroz College of Agriculture (ESALQ), University of São Paulo

**Bruna San Martin**

Universidade Federal de Santa Maria

**Giovana Ribas**

Universidade Federal de Santa Maria

**Rafael Battisti**

Federal University of Goiás

**Alexandre Heinemann**

EMBRAPA Arroz e Feijão <https://orcid.org/0000-0002-7037-488X>

**Patricio Grassini** (✉ [pgrassini2@unl.edu](mailto:pgrassini2@unl.edu))

[pgrassini2@unl.edu](mailto:pgrassini2@unl.edu) <https://orcid.org/0000-0002-7501-842X>

---

## Article

### Keywords:

**Posted Date:** April 28th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1581982/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

**Version of Record:** A version of this preprint was published at Nature Sustainability on October 10th, 2022. See the published version at <https://doi.org/10.1038/s41893-022-00968-8>.

# Abstract

The Amazon basin includes 550 M ha covered with rainforests, with 60% of this area being in Brazil. Conversion of rainforest for soybean production raises concerns about the degree to which Brazil can reconcile production and environmental goals. Here we investigated the degree to which intensification could help Brazil produce more soybean without further encroachment of the Amazon Forest. Our analysis shows that continuation of current trends in soybean yield and area would lead to conversion of additional 5.7 M ha of forests and savannas during the next 15 years, with an associated 2550 Mt of CO<sub>2</sub>eq released into the atmosphere. In contrast, acceleration of yield improvement, coupled with expansion of soybean area only in areas currently used for livestock production, would allow Brazil to achieve similar economic benefits without deforestation and with substantially lower global climate warming.

# Full Text

The COVID-19 pandemic, together with the war in Ukraine, brought two signals that can have a massive impact on developing countries that rely on commodity crops as a main source of income. One is a sharp increase in crop commodity prices, which have risen *ca.* 75% compared with pre-pandemic levels<sup>1</sup>. The other signal is a strong desire of national governments to quickly recover from the negative economic impact by making use of their comparative advantages<sup>2</sup>. These signals are of critical importance for developing countries with vast tracts of land suitable for farming that are currently covered with fragile ecosystems such as rainforests and savannas, because they can trigger massive land conversion in a relatively short period of time, leading to biodiversity loss and global warming<sup>3-9</sup>.

Brazil hosts one of the largest pools of biodiversity in the world, with 516 M ha of forests and savannas (MAPBIOMAS Project – Collection 5.0)<sup>10</sup>. Of special relevance are the vast areas of rainforests located within the Amazon basin, summing to 330 M ha. At the same time, Brazil is the main soybean exporting country, accounting for *ca.* 40% of global exports (FAOSTAT, 2017–2019)<sup>11</sup>. Soybean production has driven massive deforestation during the late 1990s and early 2000s<sup>12,13</sup>. Fortunately, Brazil has made tangible progress in subsequent years to reduce deforestation rates *via* moratoriums and incentive programs funded by foreign countries<sup>14,15</sup>. At question is whether these measures alone will be sufficient to prevent conversion of fragile ecosystems in a context of high grain prices and with governments seeking economic growth *via* increased agricultural output.

Here we investigated the degree to which agricultural intensification, that is, increasing the productivity of existing cropland, could serve as a means of enabling Brazil to simultaneously reconcile production and environmental goals. To evaluate the potential of achieving both outcomes, we combined crop modeling and spatial analysis to investigate different scenarios of intensification and land-use change and associated impact on production, land conversion, and climate change. We discuss the resultant

implications for policy makers and priorities in agricultural research and development (AR&D) programs to foster agricultural intensification and protection of fragile ecosystems.

## Recent patterns of soybean area expansion and productivity gains

Soybean area in Brazil has expanded at 1.4 M ha per year during the 2007–2019 period, with most of this expansion occurring in four regions: Pampa, Atlantic Forest, Cerrado, and Amazonia (Fig. 1a; **Extended Data Figs. 1 and 2; Supplementary Table S1; Supplementary Material Section 1**). The first two regions experienced a massive process of land conversion for agriculture many decades ago and only a small portion of the native vegetation now remains. In contrast, large tracks of pristine forest and savanna remain in Cerrado and especially Amazonia. Unfortunately, one third of the annual land converted for soybean production in Brazil is now occurring in Amazonia, with half of the soybean expansion in this region occurring at expense of tropical rainforest (**Extended Data Fig. 2**). Soybean expansion into the Amazonia and Cerrado has been driven by availability of suitable soils for crop production and favorable weather, which allow farmers to achieve high and stable soybean yields and to cultivate an additional maize crop (hereafter referred to as ‘second-crop maize’) in the same cropping season (**Extended Data Fig. 1**). Yield improvement has been comparably slower in Cerrado and Amazonia compared to that in other regions (Fig. 1b), which has led to soybean production increasing mostly from cropland expansion (Fig. 1c).

## Available opportunity for increasing crop yields

Yield potential is the yield of a well-adapted cultivar when grown without nutrient limitations and in absence of yield-reducing factors such as weeds, insect pests, and pathogens (**Supplementary Material Section 2**). Here we estimated yield potential for soybean and second-crop maize in Brazil using well-validated crop simulation models and detailed weather, soil, and agronomic data (Fig. 2, **Extended Data Fig. 3, and Supplementary Tables 2 and 3**). Our assessment revealed an average yield potential of 5.5 Mg ha<sup>-1</sup> for soybean in Brazil, but with higher and more stable yield potential in Cerrado and Amazonia than in Atlantic Forest and Pampa. At question is how much room exists in each region to increase soybean production on existing cropland via intensification. Reaching yield potential in farmer fields is difficult, as it requires copious amounts of inputs and a high degree of sophistication to eliminate yield-reducing factors, leading to reduced profit and a large negative environmental impact. In contrast, attaining 80% of the simulated yield potential (hereafter referred to as ‘attainable yield’) is considered a reasonable level of yield-gap closure for a population of farmers with adequate access to inputs, market, and extension services (**Supplementary Material Section 2**). Here, we estimated the exploitable yield gap as the difference between the attainable yield and average yield and found that the exploitable yield gap for soybean in Brazil increases northwards, from 10% and 25% in Pampa and Atlantic Forest, respectively, to 35% in Amazonia and Cerrado (Fig. 2, **Supplementary Table 2**). In the case of second-crop maize, the exploitable yield gap represents 32–44% of the attainable yield across regions (**Supplementary Table 3**).

# Production and land use change scenarios

We first explored a business-as-usual (BAU) scenario in which historical yield improvement and land-use change patterns were projected to persist during the next 15 years (Figs. 3–4, **Supplementary Table 4**). In the BAU scenario, soybean area expands from 36 to 59 M ha while current yield increases from 3.1 to 3.6 Mg ha<sup>-1</sup>, leading to a national soybean production of 212 MMT by 2035. However, this production increase in the BAU scenario would occur at expense of converting 5.7 M ha of forests and savanna into soybean cultivation that would be accompanied by an associated 2550 Mt CO<sub>2</sub>eq released to the atmosphere due to land conversion.

As an alternative pathway, we evaluated an intensification (INT) scenario in which no further expansion of soybean area is allowed in any of the regions, but which would allow further adoption of the second-crop maize occurs on existing soybean areas where climate makes double cropping possible (Figs. 3 and 4; **Supplementary Table 4**). This scenario assumes a massive investment on AR&D so that the exploitable yield gap is narrowed by half in Cerrado and Amazonia. Such a level of yield gap closure would require rates of yield improvement that would be two to three times larger than the historical rates in those two regions but similar to those in Pampa and Atlantic Forest (Fig. 1). In the case of Pampa and Atlantic Forest, current rates of yield improvement are already high, and we assumed yield gain rates in these two regions to remain the same, thus allowing full closure of the exploitable yield gap in Brazil by 2035. In the INT scenario, soybean average yield reaches 3.9 Mg ha<sup>-1</sup> by 2035, while second-crop maize average yield attains 8.6 Mg ha<sup>-1</sup>. However, the yield-gap closure on existing cropland in the INT scenario would not be sufficient to reach a production output comparable to that in the BAU scenario, and results in 34% lower soybean and second-crop maize production. Hence, while effective to avoid deforestation and reduce global warming potential (GWP), the INT scenario incurs into a huge opportunity cost in terms of economic output (US\$ 380 billion over the 15-year study period), leading to a soybean production that falls short of that projected by the Brazilian Government<sup>17</sup> (Figs. 3 and 4).

We further assessed another pathway to reduce the negative environmental impact of land conversion for soybean production while minimizing the negative economic impact. Our third scenario (INT + TE) was identical to the INT, except that further soybean area expansion would be allowed into low-carbon (C) ecosystems such as pastures and grasslands in all regions but Amazonia (Figs. 3 and 4; **Supplementary Table 4**). This scenario would require intensification of the pasture-based livestock systems (*ca.* 10% increase in stocking rate), so that *ca.* 11 M ha of pastureland and grassland would be freed up for soybean production (**see Methods**). Expansion at expense of other food crop areas would not be allowed in this scenario to avoid indirect land-use change. Notably, the INT + TE scenario achieves a reasonable balance between increasing agricultural output while reducing the negative environmental impact. Soybean production increases to 185 MMT by 2035, which is only 12% lower than that in the BAU scenario, achieving an aggregated gross income from soybean and second-maize crop that represents 92% of that in the BAU scenario. Because there is no deforestation, total GWP decreases by 32% in the INT + TE *versus* BAU scenario, achieving a GWP intensity (*i.e.*, GWP per unit of gross income) that is

comparable to that in the INT scenario. Overall, the GWP reduction due to land saving is equivalent to the total fossil fuel GHG emissions at national level over two years and a half.

## Discussion

Brazil has made remarkable progress on fostering agricultural production during the past 50 years, becoming a major soybean, maize, and beef exporting country. However, much of the increase in agricultural output has occurred proportionately more from cropland expansion than cropland productivity (Fig. 1). Here we showed that nearly one third of the recent soybean expansion in Brazil has occurred in the Amazonia (Fig. 2), which is consistent with changes in C balance (from being a net CO<sub>2</sub> sink to a source) and record deforestation rates reported elsewhere<sup>7,8</sup>. Our study assessed the aggregated economic and environmental impact of different trajectories in crop intensification and land-use change at national level, including assessing the potential for intensification in the second-maize crop as well as allowing soybean expansion into pasture-based livestock area (Figs. 3 and 4). On the one hand, continuation of current trends in soybean area expansion would drive massive encroachment of the Amazon rainforest, biodiversity loss, and increased global warming. On the other hand, agricultural intensification without further soybean area expansion would lead to a substantial negative economic impact due to lower national crop production. However, we believe that there is an alternative pathway that would balance out environmental and economic goals, and we show here that Brazil can reverse current patterns of soybean expansion *via* a dual intensification of crop and livestock systems, coupled with land-use planning, without incurring significant trade-offs in crop production and economic profit.

Fostering an intensification in Brazilian agriculture will require strategic investments on AR&D programs in both crop and livestock sectors. Our detailed yield-gap analysis identifies areas with largest yield gaps, providing a roadmap to orient AR&D programs. For example, our study shows that yield gaps are larger in the relatively new soybean areas across Cerrado and Amazonia than in the traditional soybean area in southern Brazil (Fig. 2). Given favorable soil and weather endowments and adequate farmer access to markets, inputs, and extension services, we are optimistic about reaching the desired level of crop intensification in these new regions within a relatively short timeframe *via* a targeted investment on AR&D programs. In relation to the specific interventions needed to close the current yield gap, including judicious choice of sowing date and cultivars, application of nutrient fertilizer, better crop protection, and improved soil and water management, and use of cover crops<sup>18-21</sup>. Intensification would also benefit from a cropping-system perspective, rather than one focusing on individual crops, to optimize the productivity of the whole crop sequence. For example, using early-maturing soybean cultivars may reduce soybean yield but would allow earlier maize sowing and higher yield, leading to higher annual productivity and profit for the whole cropping system. In relation to pasture-based livestock systems, previous studies have shown that ample room exists for intensification considering that current productivity represents only 20–30% of the potential<sup>22-28</sup>. Hence, our assumption of a modest 10% increase in stocking rate leading to land sparing for soybean production is reasonable and feasible to achieve within a relatively short period.



In the current context of high grain prices and food supply disruptions, we believe there is a critical need for major crop producing countries to re-assess their potential to produce more on existing cropland. Our national assessment for Brazil moves beyond previous efforts to quantify yield gaps at local level<sup>29,30</sup>, showing that intensification can help achieve a reasonable balance between crop production and protection of fragile ecosystems. We are aware of other approaches to protect natural ecosystems. For example, previous studies in Brazil have shown that moratoriums, certification, and incentive programs can help protect fragile ecosystems from conversion<sup>15,31</sup>. However, recent examples for Brazil and other countries showed that these programs fall short in protecting forests in countries that depend heavily on crop commodity exports, especially when the socio-economic context is favorable for converting natural ecosystems to agricultural production (*e.g.*, high crop prices, poor enforcement of land-use policy)<sup>7,32</sup>. Intensification can complement these other approaches to protect fragile ecosystems, providing a means to reconcile economic and environmental goals. To be effective, however, intensification would require proper policy and enforcement to ensure that land savings derived from crop yield improvement led to land sparing for nature. The main message still remains: without an emphasis on intensifying crop production within the existing agricultural area, it would be difficult to protect the last bastions of forests and biodiversity in the planet.

## Methods

### Study regions and recent trends in land-use change

Our analysis focuses on four biomes (referred to as ‘regions’ in the rest of the text), accounting for nearly all soybean area in Brazil: Pampa, Atlantic Forest, Cerrado, and Amazonia (**Supplementary Material, Section 1**). Soybean production is negligible in Pantanal and Caatinga so these two regions were excluded from our analysis. We focused on soybean-based systems in Brazil, either those that include one crop per year (single soybean), or those including a second-crop maize. In the latter system, soybean is sown in Sept-Oct and maize is sown right after soybean harvest in late Jan-Feb. Single soybean is common in Pampa, where drier climate does not allow double cropping. In contrast, higher precipitation allows double cropping in Amazonia, Cerrado, and most of the Atlantic Forest (**Supplementary Material, Section 2**).

Recent trends in yield, area, and production for soybean and second-crop maize were derived from official statistics for the 2007–2019 period<sup>16</sup>. We fitted linear models to derive the annual rate of yield improvement and harvested area for soybean and second-crop maize, separately for each region (**Fig. 1 and Extended Data Fig. 1**). Land use change arising from soybean expansion was estimated using data from the MAPBIOMAS project (v5.0)<sup>10</sup> (**Supplementary Table 1**). Our estimation of land-use change accounted for the time lag between land conversion and beginning of soybean production, which can include transitional stages such as cultivation of upland rice or short-term pasture-based livestock systems<sup>33</sup>. To do so, we looked at the new land brought into soybean production during the 2008–2019

period, and we analyzed how much of this land was under a different land use type (forest, savanna, grassland, pasture, other crops) around the year 2000 (**Extended Data Fig. 2**).

## Estimation of yield potential and yield gaps

We used results on yield potential for Brazil generated by the authors through the Global Yield Gap Atlas project (GYGA)<sup>34</sup> using well-validated process-based crop models and best available sources of weather, soil, and management data. Briefly, we selected a total of 32 sites to portray the distribution of the soybean harvested area within the country, following protocols that ensure representativeness and a reasonable coverage of national crop area<sup>35</sup>. The 32 sites collectively accounted for half of soybean harvested area in Brazil. In turn, these sites were located within agro-climatic zones accounting for 86% national soybean production and accounted for 72–92% of soybean area in each region. Following protocols that give preference to measured data at a high level of spatial and temporal resolution<sup>36</sup>, databases on weather, soil, management, and crop yields were collected for soybean for each site, and also for second-crop maize at those sites where double cropping is practiced (**Supplementary Tables 2–3; Supplementary Material Section 3**).

Yield potential was simulated for widespread cultivars in each region using CROPGRO soybean model embedded in DSSAT v 4.5<sup>37</sup> and Hybrid-Maize model<sup>38</sup>. We first evaluated CROPGRO and Hybrid Maize models on the ability to reproduce measured phenology and yields across 40 well-managed experiments located across the four regions. Models showed satisfactory performance at reproducing measured values (**Extended Fig. 3**). Subsequently, we simulated soybean yield potential for the dominant agricultural soils at each site (usually two to three), as determined from the soil maps generated by the Radambrasil project<sup>39</sup>. Simulations were based on long-term (1999–2018) measured daily weather data retrieved from Brazilian Institute of Meteorology (INMET)<sup>40</sup>. We also simulated yield potential for second-crop maize crop for those sites where double cropping is practiced. To estimate average yield potential for each site, simulated values for each soil types was weighted by soil area fraction at each site. In all cases, simulations assumed no limitations to crop growth due to nutrient deficiencies and incidence of biotic stresses such as weeds, insect pests, and pathogens. Results were up-scaled from site to region and then to country following van Bussel et al.<sup>35</sup>. Briefly, average yield potential for each region was estimated by averaging simulated yields across the sites located within each region, weighing sites according to their share of the soybean area within each region. A similar approach was followed to upscale yield potential from region to national level. Details on crop modeling, data sources, and upscaling is provided in **Supplementary Material Section 3**.

Average farmer yield was calculated separately for soybean and second-crop maize based on the average yield reported over the 2012–2017 period for the municipalities that overlap with each site, weighing municipalities based on their share of the soybean or maize area within each site<sup>16</sup>. Including more years before 2012 would have led to a biased estimate of average actual yield due to the technological yield trend in Brazil. Average farmer yields were estimated at region and country levels following the same upscaling approach as for yield potential. Finally, the exploitable yield gap was

calculated as the difference between attainable yield and average farmer yield. The attainable yield was calculated as 80% of the simulated yield potential, which is considered a reasonable yield for farmers with adequate access to inputs, markets, and technical information (**Supplementary Material Section 2**).

## **Assessment of different scenarios of intensification and land-use change**

We explored three scenarios with different soybean and maize yields and area by year 2035 and assessed their outcomes in terms of production, land use change, and GWP (**Supplementary Table 4**). A 15-year future timespan is long enough to facilitate the implementation of long-term policies, investments, and technologies devoted to close the exploitable yield gap and to implement land-use policies, but it is short enough to minimize long-term effects from climate change on crop yields and cropping systems. In the business-as-usual (BAU) scenario, historical (2007–2019) trends of soybean and second-crop maize area and yield (**Extended Data Fig. 1**) remain unchanged in all regions between the baseline year (2019) and final year (2035). Likewise, soybean area expands following the same pattern of land-use change observed during 2008–2019 (**Extended Data Fig. 2**).

To explore the available opportunity for increasing production on existing production area, we considered an intensification (INT) scenario in which there is no physical expansion of cropland while full closure of the exploitable yield gap occurs in the regions where current yield gaps are small (Pampa and Atlantic Forest) and 50% closure of the exploitable yield gap takes place in regions where current yield gap is large (Amazonia and Cerrado) (**Supplementary Table 4**). These rates are comparable to historical yield gains in Pampa and Atlantic Forest. In contrast, a scenario of full yield closure in Amazonia and Cerrado would have been unrealistic as it would have required rates of yield improvement that are three-to-four times higher than historical rates and much higher than those in Pampa and Atlantic Forest and well beyond those reported for main soybean producing countries. In the case of second-maize crop, we assumed full closure of the exploitable yield gap by 2035 because historical rates of yield improvement are adequate to reach that yield level. In the case of second-crop maize area, we projected the proportion of double cropping to increase from current 47% (Amazonia), 39% (Cerrado), and 31% (Atlantic Forest) to 100%, 70%, and 50%, respectively, as determined based on the degree of water limitation in each region (**Supplementary Material Section 4**).

Finally, we explored a third scenario of intensification plus target area expansion (INT + TE), in which identical yield gain rates and adoption of double cropping equivalent to those in the INT scenario were assumed, but with physical expansion of the soybean-maize system allowed in low-C ecosystems. In this scenario, soybean expansion is limited to 10% of existing pastures and grasslands in Pampa, Atlantic Forest, and Cerrado (total of 11.4 M ha) as a result of a parallel intensification in the pasture-based livestock sector that frees up land for soybean production. Such intensification would require a modest 10% increase in stocking rate, which is a reasonable target within a relatively short timeframe, as reported in previous studies<sup>25,26</sup>. Another assumption is that the yield potential of pasture and grasslands converted for soybean production is similar to that in existing soybean areas in each region. Cropland

expansion into grassland and pastures was allowed in all regions, except for Amazonia to prevent 'leaking' effects and the impact of road development on land clearing<sup>41,42</sup>. Similarly, conversion of area cultivated with food crops for soybean production is not allowed to avoid the negative impact of indirect land use change<sup>43</sup>.

## Estimation of global warming potential and gross income

We estimated greenhouse gas emissions (GHG), including carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrous oxides (N<sub>2</sub>O) associated with land conversion (GHG<sub>LU</sub>) and crop production (GHG<sub>PROD</sub>) for the baseline year (2019) and for the three scenarios by year 2035 (BAU, INT, INT + TE). The GHG<sub>LU</sub> includes emissions associated with changes in C stocks from aboveground and belowground biomass when land is converted for soybean production (GHG<sub>BIO</sub>), and also GHG emissions derived from associated changes in soil organic C (GHG<sub>SOC</sub>). For each land use type, annual GHG<sub>BIO</sub> was estimated based on the change in C stocks between the land use type that was converted for production (**Supplementary Table 5**) and, depending on the scenario and region, the average C stocks of the new cropping system<sup>44-46</sup>:

$$GHG_{CON} = \sum (TDM_i - TDM_{crop}) * Ai \text{ Eq. [1]}$$

where *i* is the land cover type, *TDM* is the total dry matter (t C ha<sup>-1</sup>) in land cover type *i* and in cropland (*crop*), and *Ai* is the annual area converted from land use type *i* for soybean cultivation (**Supplementary Table 4**). C stocks for single soybean and soybean-second maize systems were assumed at 2 and 5 t C ha<sup>-1</sup>, respectively<sup>44-46</sup>. Changes in soil organic C (SOC) stocks was estimated following the IPCC 2019 guidelines<sup>45</sup> and the SOC estimated for each region<sup>47,48</sup>:

$$GHG_{SOC} = \sum (SOC_{REF_i} * F_{LU} * F_{MG} * F_I) * Ai \text{ Eq. [2]}$$

where *i* is the land cover type, *SOC<sub>REF</sub>* is the soil organic C stock for mineral soils in the upper 30 cm for the reference condition (t C ha<sup>-1</sup>)<sup>47</sup> in land cover type *i* (**Supplementary Table 5**), *F<sub>LU</sub>* is the stock change factor for SOC land-use systems for a particular land-use, *F<sub>MG</sub>* is the stock change factor for SOC for management regime, *F<sub>I</sub>* is the stock change factor for SOC for the input of organic amendments, and *Ai* is the annual area converted from land use type *i* (**Supplementary Table 4**). We used a *F<sub>LU</sub>* = 0.84 that corresponds to area converted to annual crops in tropical wet regions, *F<sub>MG</sub>* = 1.04 that corresponds to reduced tillage in tropical wet regions, and *F<sub>I</sub>* = 1.0 representative for annual cropping with cereals where all crop residues are returned to the field. We assumed that GHG<sub>BIO</sub> and GHG<sub>SOC</sub> occurred during the first year after land conversion and were expressed as CO<sub>2</sub> equivalents by multiplying changes in C stocks by 3.67.

Annual GHG emissions derived from soybean and second-crop maize production (GHG<sub>PROD</sub>) were calculated for each scenario and included those derived from manufacturing, packaging, and transportation of agricultural inputs, fossil fuel use for field operations, and soil N<sub>2</sub>O emissions derived

from application of nitrogen (N) fertilizer, and domestic grain transportation. For the baseline year (2019), annual GHG from N, phosphorous (P), and potassium (K) fertilizers and other inputs (lime, pesticides, and fuel) was calculated based on current average input rates for soybean and second-crop maize in each region as derived from the crop management data collected for each region (**Supplementary Table 6; Supplementary Material Section 3.4**). To calculate GHG emissions associated with manufacturing, packaging, and transportation of N, phosphorous (P), and potassium (K) fertilizers and lime, we used specific updated emissions factors for South America<sup>49</sup>, selecting those fertilizer sources that are most commonly used for soybean and second-crop maize production: urea (N), monoammonium phosphate (P), and potassium chloride (K). Our calculations also included the extra lime application that is needed to correct soil acidity converted areas. Emission factors associated with seed production, pesticides and diesel were derived from Lal (2014)<sup>50</sup>. Soil N<sub>2</sub>O emissions derived from N fertilizer application were calculated assuming a N<sub>2</sub>O emission factor of 1.6% of the applied N fertilizer applied based on the IPCC emission factor for mineral soils in tropical regions<sup>45</sup>. Emissions derived from domestic grain transportation for each region were estimated using the GHG per ton of grain as reported by previous studies for each region<sup>51</sup>. We assumed that inputs other than nutrient fertilizer will not change relative to the baseline in the BAU scenario. In the case of the INT and INT-TE scenarios, applied inputs were calculated based on those reported for current high-yield fields where the yield gap is small. In the case of fertilizer nutrient rates, we estimated them for the three scenarios following a nutrient-balance approach that depends upon the projected yield for each scenario (**Supplementary Table 6; Supplementary Material Section 3.4**)

The GHG<sub>PROD</sub> in the baseline year (2019) and for the three scenarios in 2035 (BAU, INT, and INT-TE) was estimated for each region by multiplying the emissions per unit of area by the annual soybean harvested area, summing them up to estimate GHG emissions at national level. Overall 100-y GWP was estimated as the sum of GHG<sub>LUC</sub> and GHG<sub>PROD</sub>, both expressed as CO<sub>2</sub> equivalents (CO<sub>2</sub>eq) to account for the higher warming potential of CH<sub>4</sub> and N<sub>2</sub>O, which are 25 and 298 times the intensity of CO<sub>2</sub> on per mass basis, respectively. The gross income was estimated for each scenario by multiplying annual crop production by the average price for soybean and maize grain during the past ten years (453 and 184 US\$ t<sup>-1</sup> for soybean and maize, respectively<sup>1</sup>). Finally, to combine the environmental and economic impact into one metric, we calculated the GWP intensity as the ratio between GWP and gross income.

## Uncertainties and limitations

Our study is subject to several uncertainties. For example, our yield gap estimates could be biased due to inaccuracies in the underpinning weather, soil, and management databases. To the extent it was possible, we relied on the best sources of data that exists, given preference to measured data at the finest level of spatial resolution. Similarly, our estimates of GWP can be biased because of uncertainty in emission factors used to compute GHGs from land conversion and crop production. Again, we relied on most updated emission factors published for tropical regions and dominant management practices as determined *via* experts in Brazil. Our scenario assessment may also have limitations. For example,

changes in yield potential because of potential expansion of irrigated area in Brazil and/or genetic improvement per se were not considered in our assessment. Similarly, the adverse impact of climate change on yield potential is not accounted for. We note, however, that the impact of these factors is likely to be negligible considering the short time of our assessment (16 years) and, hence, are not expected to influence the overall conclusions from our study. Another limitation is to assume the productivity of new land converted for soybean production to have same yield potential as the land currently under production, which, indeed, is not the case in many instances<sup>52</sup>. If the new land has lower yield potential, it will further emphasize the need to intensify production on existing cropland so that the overall conclusion of our study will not change.

### Data availability

Data on yield potential and yield gaps that support the findings of this study are publicly available via the Global Yield Gap Atlas website ([www.yieldgap.org](http://www.yieldgap.org)). Data source files associated with the figures shown in the article are also provided.

## References

1. Commodity Markets. *The World Bank* <https://www.worldbank.org/en/research/commodity-markets>.
2. Morris, M. *et al.* *Future Foodscapes: Re-imagining Agriculture in Latin America and the Caribbean*. vol. 1 (2020).
3. Bonan, G. B. Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science* (80-). **320**, 1444–1449 (2008).
4. Johnson, C. N. *et al.* Biodiversity losses and conservation responses in the Anthropocene. *Science* (80-). **356**, 270–275 (2017).
5. Seymour, F. & Harris, N. L. Reducing tropical deforestation. *Science* (80-). **365**, 756–757 (2019).
6. Instituto Nacional de Pesquisas Espaciais. <http://inpe.br/>.
7. Silva Junior, C. H. L. *et al.* The Brazilian Amazon deforestation rate in 2020 is the greatest of the decade. *Nat. Ecol. Evol.* **5**, 144–145 (2021).
8. Gatti, L. V. *et al.* Amazonia as a carbon source linked to deforestation and climate change. *Nature* **595**, 388–393 (2021).
9. Boulton, C. A., Lenton, T. M. & Boers, N. Pronounced loss of Amazon rainforest resilience since the early 2000s. *Nat. Clim. Chang.* **12**, 271–278 (2022).
10. MapBiomas Project- Collection [5.0] of the Annual Series of Land Use and Land Cover Maps of Brazil. <https://mapbiomas.org/en>.
11. FAOSTAT. Food and Agriculture Organization. FAOSTAT Statistical Database. <http://www.fao.org/faostat/en/#data/QC>.
12. DeFries, R. S., Rudel, T., Uriarte, M. & Hansen, M. Deforestation driven by urban population growth and agricultural trade in the twenty-first century. *Nat. Geosci.* **3**, 178–181 (2010).

13. Zalles, V. *et al.* Near doubling of Brazil's intensive row crop area since 2000. *Proc. Natl. Acad. Sci.* **116**, 428–435 (2019).
14. Kastens, J. H., Brown, J. C., Coutinho, A. C., Bishop, C. R. & Esquerdo, J. C. D. M. Soy moratorium impacts on soybean and deforestation dynamics in Mato Grosso, Brazil. *PLoS One* **12**, e0176168 (2017).
15. Heilmayr, R., Rausch, L. L., Munger, J. & Gibbs, H. K. Brazil's Amazon Soy Moratorium reduced deforestation. *Nat. Food* **1**, 801–810 (2020).
16. IBGE. Instituto Brasileiro de Geografia e Estatística. Agricultural Production at County Level: Produção Agrícola Municipal. .
17. CONAB, Companhia Nacional de Abastecimento. [www.conab.gov.br](http://www.conab.gov.br).
18. Zanon, A. J., Streck, N. A. & Grassini, P. Climate and management factors influence Soybean yield potential in a subtropical environment. *Agron. J.* **108**, 1447–1454 (2016).
19. Grassini, P. *et al.* Soybean. in *In Crop Physiology Case Histories for Major Crops* 289–319 (Academic Press, 2021).
20. Tagliapietra, E. L. *et al.* Biophysical and management factors causing yield gap in soybean in the subtropics of Brazil. *Agron. J.* **113**, 1882–1894 (2021).
21. Battisti, R. *et al.* Rules for grown soybean-maize cropping system in Midwestern Brazil: Food production and economic profits. *Agric. Syst.* **182**, 102850 (2020).
22. Martha, G. B., Alves, E. & Contini, E. Land-saving approaches and beef production growth in Brazil. *Agric. Syst.* **110**, 173–177 (2012).
23. Cohn, A. S. *et al.* Cattle ranching intensification in Brazil can reduce global greenhouse gas emissions by sparing land from deforestation. *Proc. Natl. Acad. Sci.* **111**, 7236–7241 (2014).
24. de Oliveira Silva, R. *et al.* Sustainable intensification of Brazilian livestock production through optimized pasture restoration. *Agric. Syst.* **153**, 201–211 (2017).
25. de Oliveira Silva, R., Barioni, L. G., Queiroz Pellegrino, G. & Moran, D. The role of agricultural intensification in Brazil's Nationally Determined Contribution on emissions mitigation. *Agric. Syst.* **161**, 102–112 (2018).
26. Arantes, A. E., Couto, V. R. de M., Sano, E. E. & Ferreira, L. G. Livestock intensification potential in Brazil based on agricultural census and satellite data analysis. *Pesqui. Agropecuária Bras.* **53**, 1053–1060 (2018).
27. Gil, J. D. B. *et al.* Tradeoffs in the quest for climate smart agricultural intensification in Mato Grosso, Brazil. *Environ. Res. Lett.* **13**, 064025 (2018).
28. Strassburg, B. B. N. *et al.* When enough should be enough: Improving the use of current agricultural lands could meet production demands and spare natural habitats in Brazil. *Glob. Environ. Chang.* **28**, 84–97 (2014).
29. Sentelhas, P. C. *et al.* The soybean yield gap in Brazil – magnitude, causes and possible solutions for sustainable production. *J. Agric. Sci.* **153**, 1394–1411 (2015).

30. de Souza Nória Júnior, R. & Sentelhas, P. C. Yield gap of the double-crop system of main-season soybean with off-season maize in Brazil. *Crop Pasture Sci.* **71**, 445 (2020).
31. Gibbs, H. K. *et al.* Brazil's Soy Moratorium. *Science* (80-). **347**, 377–378 (2015).
32. Carlson, K. M. *et al.* Effect of oil palm sustainability certification on deforestation and fire in Indonesia. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 121–126 (2018).
33. Nemecek, T. *et al.* World Food LCA Database: Methodological guidelines for the life cycle inventory of agricultural products. Version 3.5. 88 (2019).
34. Global Yield Gap and Water Productivity Atlas. <https://www.yieldgap.org>.
35. van Bussel, L. G. J. *et al.* From field to atlas: Upscaling of location-specific yield gap estimates. *F. Crop. Res.* **177**, 98–108 (2015).
36. Grassini, P. *et al.* How good is good enough? Data requirements for reliable crop yield simulations and yield-gap analysis. *F. Crop. Res.* **177**, 49–63 (2015).
37. Jones, J. W. *et al.* *The DSSAT cropping system model*. *European Journal of Agronomy* vol. 18 (2003).
38. Yang, H. Hybrid-maize—a maize simulation model that combines two crop modeling approaches. *F. Crop. Res.* **87**, 131–154 (2004).
39. Cooper, M., Mendes, L. M. S., Silva, W. L. C. & Sparovek, G. A National Soil Profile Database for Brazil Available to International Scientists. *Soil Sci. Soc. Am. J.* **69**, 649–652 (2005).
40. Brazilian Institute of Meteorology (INMET). <https://www.gov.br/agricultura/pt-br/assuntos/inmet>.
41. Byerlee, D., Stevenson, J. & Villoria, N. Does intensification slow crop land expansion or encourage deforestation? *Glob. Food Sec.* **3**, 92–98 (2014).
42. Laurance, W. F., Goosem, M. & Laurance, S. G. W. Impacts of roads and linear clearings on tropical forests. *Trends Ecol. Evol.* **24**, 659–669 (2009).
43. Searchinger, T. *et al.* Use of U.S. Croplands for Biofuels Increases Greenhouse Gases Through Emissions from Land-Use Change. *Science* (80-). **319**, 1238–1240 (2008).
44. Bonini, I. *et al.* Collapse of ecosystem carbon stocks due to forest conversion to soybean plantations at the Amazon-Cerrado transition. *For. Ecol. Manage.* **414**, 64–73 (2018).
45. IPCC. 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4: Agriculture, Forestry and Other Land Use: Mapping Tables. *Texts Wait. Hist.* 1–48 (2019).
46. Dionizio, E. A., Pimenta, F. M., Lima, L. B. & Costa, M. H. Carbon stocks and dynamics of different land uses on the Cerrado agricultural frontier. *PLoS One* **15**, e0241637 (2020).
47. Bernoux, M., da Conceição Santana Carvalho, M., Volkoff, B. & Cerri, C. C. Brazil's Soil Carbon Stocks. *Soil Sci. Soc. Am. J.* **66**, 888–896 (2002).
48. Brazil. Ministry of Science, T. and I. S. of P. and P. & Change., of R. and D. G. C. of G. C. *Third National Communication of Brazil to the United Nations Framework Convention on Climate Change – Volume III/ Ministry of Science, Technology and Innovation. Brasília: Ministério da Ciência, Tecnologia e Inovação.* (2016).



49. Brentrup, F., Lammel, J., Stephani, T. & Christensen, B. Updated carbon footprint values for mineral fertilizer from different world regions. *11th Int. Conf. Life Cycle Assess. Food 2018 (LCA Food) conjunction with 6th LCA AgriFood Asia 7th Int. Conf. Green Sustain. Innov. "Global food challenges Towar. Sustain. Consum 2018*, 19–22 (2018).
50. Lal, R. Carbon emission from farm operations. *Environ. Int.* **30**, 981–990 (2004).
51. Escobar, N. *et al.* Spatially-explicit footprints of agricultural commodities: Mapping carbon emissions embodied in Brazil's soy exports. *Glob. Environ. Chang.* **62**, 102067 (2020).
52. Andrade, J. F. *et al.* Impact of urbanization trends on production of key staple crops. *Ambio* (2021) doi:10.1007/s13280-021-01674-z.
53. Fehr, W. R., Caviness, C. E., Burmood, D. T. & Pennington, J. S. Stage of development descriptions for soybean, *Glycine max* (L.). Merrill. *Crop Sci.* 929–931 (1971).
54. Ritchie, J.. & Hanway, J.. How a corn plant develops. Iowa State Univ. Sci. Technol. Coop. Ext. Serv. Ames, IOWA. Spec. Rep. 48. (1982).
55. Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. & Kent, J. Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858 (2000).
56. Buainain, A., Favareto, A., Contini, E., Chaves, F. T. & Henz, G. P. *Desafios para a agricultura nos biomas brasileiros*. *Journal of Chemical Information and Modeling* vol. 53 (2020).
57. *Agricultura tropical: quatro décadas de inovações tecnológicas, institucionais e políticas*. (Embrapa Informação Tecnológica, 2008).
58. Castro, A. A. J. F., Martins, F. R., Tamashiro, J. Y. & Shepherd, G. J. How Rich is the Flora of Brazilian Cerrados? *Ann. Missouri Bot. Gard.* **86**, 192 (1999).
59. Bridgewater, S., Ratter, J. A. & Felipe Ribeiro, J. Biogeographic patterns, -diversity and dominance in the cerrado biome of Brazil. *Biodivers. Conserv.* **13**, 2295–2317 (2004).
60. Dirzo, R. & Raven, P. H. Global State of Biodiversity and Loss. *Annu. Rev. Environ. Resour.* **28**, 137–167 (2003).
61. Mantovani, W. A degradação dos biomas brasileiros. in *Patrimônio ambiental brasileiro* (ed. Ribeiro, W. C.) 367–439 (Editora Universidade de São Paulo, São Paulo., 2003).
62. Oliveira-Filho, A. T. & Fontes, M. A. L. Patterns of Floristic Differentiation among Atlantic Forests in Southeastern Brazil and the Influence of Climate<sup>1</sup>. *Biotropica* **32**, 793–810 (2000).
63. Overbeck, G. E. *et al.* Brazil's neglected biome: The South Brazilian Campos. *Perspect. Plant Ecol. Evol. Syst.* **9**, 101–116 (2007).
64. Evans, L. T. *Crop evolution, adaptation and yield*. (Cambridge University Press, 1993).
65. van Ittersum, M. K. *et al.* Yield gap analysis with local to global relevance—A review. *F. Crop. Res.* **143**, 4–17 (2013).
66. Lobell, D. B., Cassman, K. G. & Field, C. B. Crop yield gaps: Their importance, magnitudes, and causes. *Annu. Rev. Environ. Resour.* **34**, 179–204 (2009).

67. Cassman, K. G., Dobermann, A., Walters, D. T. & Yang, H. Meeting Cereal Demand While Protecting Natural Resources and Improving Environmental Quality. *Annu. Rev. Environ. Resour.* **28**, 315–358 (2003).
68. Institute, I. F. P. R. Global Spatially-Disaggregated Crop Production Statistics Data for 2010 Version 2.0. doi:10.7910/DVN/PRFF8V.
69. van Wart, J. *et al.* Use of agro-climatic zones to upscale simulated crop yield potential. *F. Crop. Res.* **143**, 44–55 (2013).
70. Allen, R. G., Pereira, L. S., Raes, D. & Smith, M. FAO Irrigation and Drainage Paper No. 56 - Crop Evapotranspiration. (1998).
71. Aramburu Merlos, F. *et al.* Potential for crop production increase in Argentina through closure of existing yield gaps. *F. Crop. Res.* **184**, 145–154 (2015).
72. Monteiro, L. A., Sentelhas, P. C. & Pedra, G. U. Assessment of NASA/POWER satellite-based weather system for Brazilian conditions and its impact on sugarcane yield simulation. *Int. J. Climatol.* **38**, 1571–1581 (2018).
73. NASA LaRC POWER Project. <https://power.larc.nasa.gov/>.
74. Pivetta, L. A., Castoldi, G., dos Santos, G. P. & Rosolem, C. A. Crescimento e atividade de raízes de soja em função do sistema de produção. *Pesqui. Agropecu. Bras.* **46**, 1547–1554 (2011).
75. Battisti, R. *et al.* Assessment of soybean yield with altered water-related genetic improvement traits under climate change in Southern Brazil. *Eur. J. Agron.* **83**, 1–14 (2017).
76. Franchini, J. C., Junior, A. A. B., Debiasi, H. & Nepomuceno, A. L. Root growth of soybean cultivars under different water availability conditions. *Semin. Agrar.* **38**, 715–724 (2017).
77. Tomasella, J., Hodnett, M. G. & Rossato, L. Pedotransfer Functions for the Estimation of Soil Water Retention in Brazilian Soils. *Soil Sci. Soc. Am. J.* **64**, 327–338 (2000).
78. Reichardt, K. *et al.* Critical analysis of the field determination of soil hydraulic conductivity functions using the flux-gradient approach. *Soil Tillage Res.* **48**, 81–89 (1998).
79. Tomasella, J. & Hodnett, M. Pedotransfer functions for tropical soils. in *Development of Pedotransfer Functions in Soil Hydrology* vol. 30 415–429 (Elsevier, 2004).
80. Ribeiro, B. & Zanon, A. J. *Ecofisiologia do Milho: visando altas produtividades*.
81. Roquetti, D. Potencial de produção de grãos brasileiros via fertilização e impactos nas emissões co<sub>2</sub> eq. [http://anda.org.br/wp-content/uploads/2018/10/Dissertacao\\_David\\_Roquetti\\_Filho.pdf](http://anda.org.br/wp-content/uploads/2018/10/Dissertacao_David_Roquetti_Filho.pdf) (2014).
82. Embrapa Milho e Sorgo. <https://www.embrapa.br/milho-e-sorgo/biblioteca/acervo>.
83. Embrapa Solos. <https://www.embrapa.br/solos/publicacoes>.
84. Embrapa Soja. <https://www.embrapa.br/soja/publicacoes>.
85. IPNI Estimates of Nutrient Uptake and Removal. <http://www.ipni.net/article/IPNI-3296>.
86. Zhang, X. *et al.* Managing nitrogen for sustainable development. *Nature* **528**, 51–59 (2015).

## Declarations

## **Acknowledgments**

This project was funded by the International Plant Nutrition Institute (grant INS-19/0007 to P.G.), the Research Foundation of the State of São Paulo (FAPESP 433 grants 2017/20925-0, 2018/06396-7, 2021/00720-0 to F.R.M.), the Brazilian Research Council (CNPq 130972/2019-3, 425174/2018-2 and 300916/2018-3 to F.R.M.), the Research Foundation of the State of Rio Grande do Sul (FAPERGS - 17/2551-0000775-1 to A.J.Z.), and the Global Engagement Office at the Institute of Agriculture and Natural Resources at University of Nebraska-Lincoln (UNL) through the FAPESP-UNL SPRINT Program (2017/50445-0 to P.G.). The first author (F.M.) received financial support from the Fulbright program to support a 6-month stay at UNL.


## **Author Contribution**

F.R.M., P.G., and A.J.Z. conceived the project. E.H.F.M.S., G.R., L.A.S.A., B.S.M. collected the data and ran the model simulations with input from J.P.M., P.G., A.B.H., and R.B. P.G., J.P.M., and J.F.A. analyzed the data. P.G., J.F.A., J.P.M., F.R.M., and A.J.Z. wrote the manuscript with input from all authors.

## **Competing interests**

The authors declare no competing interests.

## **Figures**



**Figure 1**

*Trends in soybean area and yield in main producing areas in Brazil. (a. b) Trends in soybean area and average yield. Also shown in (b) is the annual rate of yield improvement for each region during 2007-2019 and 95% confidence interval. (c) Contribution of cropland expansion and yield improvement to soybean production increase in each region. Each region is shown with a different color. Size of the pie chart in each region is proportional to the share of national soybean production. Inset shows soybean area distribution. Data were retrieved from IBGE (2021)<sup>16</sup>.*




Figure 2

*Attainable yield and yield gaps across soybean producing areas in Brazil. Pie charts show the actual yield (green) and exploitable yield gap (yellow) as percentage of the attainable yield. Size of the pie chart is proportional to the attainable yield. Inset shows current average yield for each region: Pampa (PAM), Atlantic Forest (ATL), Cerrado (CER), and Amazonia (AMA).*





Figure 3

*Changes in soybean production with different yield and land-use change scenarios in Brazil. Three scenarios are shown: business as usual (BAU), intensification (INT), intensification plus target area expansion (INT+TE). Also shown is the projected soybean production by the Brazilian government for year 2029 (CONAB, 2014).*



**Figure 4**

*Land conversion, global warming potential (GWP), and gross income associated with different scenarios of intensification and land-use change in soybean-based systems in Brazil. Three scenarios are shown: business as usual (BAU), intensification (INT), intensification plus target area expansion (INT+TE). (a) Accumulated area of forest, savanna, grasslands plus pastures, and other land uses converted for soybean production. (b) Accumulated GWP disaggregated by source: land-use change (LUC), crop*

production, and domestic grain transportation. **(c)** Annual GWP and gross income in 2019 and for the three scenarios by year 2035. Arrows show the trajectory between 2019 and 2035 for each scenario while values show GWP intensity, expressed as GWP per unit of income (kg CO<sub>2</sub>-eq per US\$).

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFile.docx](#)