



Review

Russeting of Fruits: Etiology and Management

Andreas Winkler , Thomas Athoo and Moritz Knoche *

Institute of Horticultural Production Systems, Leibniz University Hannover, Herrenhäuser Straße 2, 30419 Hannover, Germany; andreas.winkler@obst.uni-hannover.de (A.W.); t.athoo@obst.uni-hannover.de (T.A.)
* Correspondence: moritz.knoche@obst.uni-hannover.de; Tel.: +49-511-762-9020

Abstract: The skin of a fruit protects the vulnerable, nutrient-rich flesh and seed(s) within from the hostile environment. It is also responsible for the fruit's appearance. In many fruitcrop species, russeting compromises fruit appearance and thus commercial value. Here, we review the literature on fruit russeting, focusing on the factors and mechanisms that induce it and on the management and breeding strategies that may reduce it. Compared with a primary fruit skin, which is usually distinctively colored and shiny, a secondary fruit skin is reddish-brown, dull and slightly rough to the touch (i.e., russeted). This secondary skin (periderm) comprises phellem cells with suberized cell walls, a phellogen and a phelloderm. Russeted (secondary) fruit skins have similar mechanical properties to non-russeted (primary) ones but are more plastic. However, russeted fruit skins are more permeable to water vapor, so russeted fruits suffer higher postharvest water loss, reduced shine, increased shrivel and reduced packed weight (most fruit is sold per kg). Orchard factors that induce russeting include expansion-growth-induced strain, surface wetness, mechanical damage, freezing temperatures, some pests and diseases and some agrochemicals. All these probably act via an increased incidence of cuticular microcracking as a result of local concentrations of mechanical stress. Microcracking impairs the cuticle's barrier properties. Potential triggers of russeting (the development of a periderm), consequent on cuticular microcracking, include locally high concentrations of O₂, lower concentrations of CO₂ and more negative water potentials. Horticulturists sometimes spray gibberellins, cytokinins or boron to reduce russeting. Bagging fruit (to exclude surface moisture) is also reportedly effective. From a breeding perspective, genotypes having small and more uniform-sized epidermal cells are judged less likely to be susceptible to russeting.

Keywords: disorder; periderm; repair mechanism



Citation: Winkler, A.; Athoo, T.; Knoche, M. Russeting of Fruits: Etiology and Management. *Horticulturae* **2022**, *8*, 231. <https://doi.org/10.3390/horticulturae8030231>

Academic Editors: Isabel Lara Ayala and Brian Farneti

Received: 28 January 2022

Accepted: 4 March 2022

Published: 8 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The skin of a fruit lies at the interface between the vulnerable, nutrient-rich fleshy tissues and seed(s) inside and the surrounding 'hostile' environment outside. The fruit skin is exposed to a broad range of abiotic and biotic challenges, thus serving as a critical barrier protecting the fruit tissues against (a) uncontrolled water loss/uptake [1], (b) uncontrolled exchanges of respiratory gasses (O₂, CO₂) and the hormone ethylene (C₂H₄) [2], (c) UV-radiation [2,3] and (d) invasion by pathogens [4,5]. In modern horticulture, some of these functions require opposing properties, such as protection against cell content leakage, while at the same time permitting penetration of foliar-applied nutrients, growth regulators or other agrochemicals [6]. To fulfill these functions, the fruit skin must remain intact throughout the period of fruit growth and development. This review deals with the development of a secondary surface (a periderm) on the skins of commercial fruit types that usually retain their shiny, distinctively-colored, primary surfaces through to harvest and consumption.

A plant organ's primary surface comprises a complex of materials. On the outside, there is a polymeric cuticle that overlies a cellular structure usually consisting of a single epidermal cell layer, which itself overlies one to several layers of hypodermal cells. In most fruit crops, the epidermis and hypodermis are responsible for the skin's mechanical

properties, while the cuticle is responsible for the skin's barrier properties [7,8]. It is also the primary fruit skin that determines the appearance and attractiveness of the fruit to end consumers, as well as to seed-dispersing animals. After all, the wild types of most commercial fruitcrop species evolved to attract animals as their agents of seed dispersal. Thus, the epicuticular waxes on the cuticle are responsible for the skin's gloss, and the pigments in the cuticle and subtending cell layers for the skin's distinctive color [9,10].

However, the skins of a significant number of commercial fruit types are partially or wholly covered by areas of a secondary surface. Horticulturists refer to this as russetting [11,12]. The proportion of the surface of a mature fruit that is primary vs. secondary is genetically determined. Thus, in apple, some cultivars rarely exhibit areas of russetting (Royal Gala); in some, russetting is a cultivar characteristic (Cox's Orange Pippin); while in others, the whole fruit surface is usually russeted (Egremont Russet). Russetting is seen as a market defect (market value is reduced) only in the first case or in the second if the russeted area is excessive.

A secondary surface forms when the primary surface fails. Failure may occur for various reasons. Potential reasons include the normal internal processes of ontogeny (e.g., growth strain) or external factors such as mechanical or chemical damage or harsh environmental conditions such as freezing [7]. A periderm forms to (partially) restore the impaired barrier properties of the damaged primary fruit skin. The proportion of the fruit-skin affected by russetting ranges from small patches in particular regions of the fruit surface to a uniform layer that covers the entire fruit.

From a horticultural point of view, the dull, reddish-brown appearance of russetting is usually unattractive to the consumer. Russetting is therefore considered to be a fruit surface disorder in many fruitcrop species and in all 'smooth-skinned' cultivars. In many russet-susceptible cultivars, russetting is readily accepted as being 'normal'. Thus, the entire fruit skin is russeted in most kiwifruit cultivars/species. Similarly, russetting is considered normal and acceptable in the 'Reinette' apple cultivars and in pear cultivars, such as Bosc, Conference and Gold La France, the latter being a russeted sport of the non-russeted cultivar La France [13]. Similarly, in Asian pear, smooth-skinned cultivars, as well as entirely russeted cultivars, are known. Melons form a notable exception. Here, the 'netting' pattern of russetting of the fruit skin is seen as a positive indicator of fruit quality and, so, is considered highly desirable [14].

Russetting is not a new phenomenon. The first research publications date back nearly two centuries [12,15–22]. Most of the literature on russetting relates to pome fruit, and especially to apple. Fewer studies relate to pear, kiwifruit, mango, tomato, bell pepper and others (Table 1). Occasionally, russetting has been reported for plums and grapes. Sweet and sour cherries, peaches, apricots and most 'berryfruit' crops, including currants, blueberries, raspberries, blackberries and strawberries, are essentially free of russet. The published information on russetting is scattered and often not conclusive. The objective of this paper is to review the literature on the practical aspects of russetting, including its occurrence, triggers, mechanical bases and management strategies adopted to reduce russetting under orchard conditions by cultivation and breeding. For a comprehensive review of the biochemistry and molecular biology of russet formation, the reader is referred to the excellent recent reviews by Macnee et al. [23] and Wang et al. [24].

2. Occurrence and Symptoms of Russet

Russetting occurs in a large number of fruitcrop species (Table 1). Often, russet-susceptible and non-susceptible cultivars are known within a species. In apple, some highly russet-susceptible cultivars are identified by the cultivar name. Examples include Red Russet, Golden Russet, Roxbury Russet [22] or Egremont Russet [25], as well as the Reinette-type cultivars [26].

Table 1. Occurrence, symptoms, causes and management of russeting. Results are compiled from literature sources.

| Cultivar | Symptoms | Causes | Management |
|-------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Apple | Russet as rough and brown skin [17,28], often in stem [29] and calyx cavities [30], some cultivars with entire surface russeted [31], high susceptibility during early fruit development [12,29,32–34] | Moisture [35–38] or high humidity [17,35], damage by pesticides, growth regulators, surfactants and other substances [19,39–61], frost [27,34], fungi [62–66], viruses [67–69], insects [70,71] | Spray application of growth regulators [30,32,33,56,58,72–81], CaCl ₂ [82], prohexadione-calcium [72], organic/mineral bio-stimulators [83], chlorogenic acid [84], coatings [82], insecticides to prevent insect damage [71], shading nets [85], rain shelters [48], bagging [17,48,82,86] |
| Pear | Russet as dull-brownish skin patches, more in calyx and cheek than in neck [87], some cultivars completely russeted [23], high susceptibility during early development [87] | Surface moisture [88–90], high humidity [90], growth stress [87], fungicides, thinners and growth regulators [89–93], insects [94–96], bacteria [97], fungi [98,99] | Bagging [100–104], spray application of GA ₄₊₇ [105], mancozeb + sulfur [105] or kaolin ± mancozeb [106] |
| Citrus | Russet as rough texture, brownish-black, greyish discoloration [107]. | (Rust) mites, thrips and other sucking insects [107–109], mechanical damage by wind, hail, contact with branches [110,111] | Zineb against citrus mites [108,109,112] |
| Prune | On immature fruit: longitudinal stripes at stylar end [113], mature: rough, brown, dried surface [113] | Copper spray [113], mechanical damage by wind, abrasion by leaves, shoots, adjacent fruits [113,114], exposure to surface wetness or free water, high humidity [113,114], scab [115] | Captafol, ziram for scab control [113] |
| Loquat | Deep brown stripes, approx. 1 mm wide [116,117] | Growth stress [116,117], microclimate (high temperature) [116], very high light intensities [116] | Shading using nets to decrease growth rate during cell division phase [116,118] |
| Tomato | Russet as rough corky discolored surface [119], also referred to as ‘shoulder check’ [119] or ‘cuticle cracking’ [120] | Rust mites [71,121], growth stress [120,122,123], surface moisture [119] | Non-susceptible cultivars [123], moderate thinning [123], spray application of Ca+B [119,124] |
| Melon | Rind netting common in some cultivars [14], russet as dry, white to brownish ridges [14] | Growth stress [125,126], surface moisture [125], wounding [14] | Rind netting desirable, countermeasures not needed |
| Grape berry | Brown patches of russet [127] | Surfactants [128], fungicides [129], insects [71,130], surface moisture [131] | Spray applications of GA ₃ , GA ₃ + CPPU [132], insecticides [71], Ca [133] |
| Mango | Rough brownish irregular patches of russet [134], beginning at lenticels [134] | Surface moisture, cold nights [134] | Bagging [135] |
| Pomegranate | Corky surface [136] | High humidity [136,137], heat waves [138], temperature fluctuation during maturation [136], pomegranate mite [139] | Spray applications of GA ₃ , CPPU [137,140], acetylsalicylic acid [136], sulfur dust against pomegranate mites [139] |

The symptoms of russet are similar between different fruitcrop species. A rough, reddish-brown and corky appearance is characteristic of a russeted surface (Table 1 and Figure 1). The region of the fruit surface affected by russet can differ. In apples, the spatial distribution of russetting differs depending on the cause. Russet induced by growth strain or by exposure to high humidity or dew occurs in large, uniform patches and may cover the entire fruit surface. Russetting limited to the stem cavity is more likely the result of long wetness durations and high growth strains. Russetting in response to mechanical wounding (e.g., scratching or abrasion from contact with a neighboring fruit or leaf or stem) is typically well defined spatially, being strictly limited to the region of direct physical contact. Russetting caused by spray chemicals occurs in regions of the fruit surface where spray droplets collect and later concentrate excessively during drying. Small fruitlets that come into contact with spray solutions during the particularly russet-susceptible early stages of fruit development may be entirely russeted [27]. A net-like pattern of russetting on apple is characteristic of infection with powdery mildew. Russetting caused by the feeding of pests (thrips, stink bugs, mites, etc.) is limited to the site of the puncture wound and the immediately surrounding cells. Forms of russetting caused by frost are typically in rings. These are induced by freezing temperatures when only part of the flower or fruitlet is damaged (Table 1).




Figure 1. Russeting in different fruit crops. (A) Apple; (B) pear; (C) citrus; (D) plum; (E) pomegranate. Images: (A,C) Andreas Winkler, (B,D,E) Martin Brüggewirth.

3. Some Fruit Skin Disorders Not Related to Periderm Formation

There are some fruit skin disorders that can be confused with russetting. These include skin spots and scarf skin in apple and maturity bronzing (sometimes also called maturity stain) in banana (Table 2). These disorders can bear a visual similarity to russetting. However, they differ from russetting in that a periderm does not develop.

In skin spots, cuticular microcracks are causal. These form due to moisture exposure during late-stage fruit development [36]. In this stage, the apple fruit skin is no longer able to form a periderm [36,38]. Here, the impaired barrier properties of the skin

are restored to some extent by the deposition of lignin in the cell walls immediately underlying a microcrack. This process hydraulically isolates the portion of the fruit skin underlying a microcrack. The characteristic spot-like appearance is caused by the resulting cell death [141].

Apples with scarf skin symptoms look as if they have a thin and very 'soft' periderm. However, periderm formation is not involved in scarf skin. Instead, scarf skin is thought to result from the formation of subepidermal air spaces [142]. The cause of this is not yet known. Surface moisture may be involved since bagging during early fruit development (when russet susceptibility is particularly high) reduces scarf skin [143].

Maturity bronzing in banana is also connected to fractures in the cuticle, which propagate into the epidermis [144]. Maturity bronzing occurs primarily in the tropical wet season when temperatures and humidities are especially high, and the sky is overcast [145]. These conditions result in high rates of growth strain, which may be causal in maturity bronzing [144].

Table 2. Fruit surface disorders that bear some similarity with russet, but where no periderm is involved. Data are compiled from literature sources.

| Disorder | Crop Affected | Symptoms | Causes | Management |
|-------------------------------------|---------------|----------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| Skin spots | Apple | Irregular patches of small, round and brown spots, develops in CA-storage, promoted by 1-MCP [141] | Moisture-induced microcracks late in the season [36,141] | Reducing surface wetness duration, for susceptible batches, no storage or cool-storage only [36,141] |
| Scarf skin | Apple | Whitish lines or stripes [146], whitish or opalescent sheen [147], due to formation of subepidermal air spaces [142] | Unknown | - |
| Maturity bronzing or maturity stain | Banana | Pre-harvest necrosis of the skin, bronze coloration [144] | Growth stress [148], water stress [144] | Bagging [149], reducing the number of leaves [150] |

4. Anatomy of Russeted Fruit Skin

In botanical terms, a russeted fruit skin represents a periderm consisting of phellem, phellogen and phelloderm [7,23]. The phellem is the outermost layer of this composite, the phelloderm the innermost. The phellogen is the interfacing sheet-like meristematic layer. The phellogen is formed in the hypodermal cell layer by dedifferentiation of hypodermal cells [16,151]. Periclinal cell division in the phellogen generates stacks of phellem cells where each cell of a stack originates from the division of a single underlying mother cell of the phellogen [7].

Phellem cells have suberized cell walls. When the stacks of phellem cells reach the surface, they come into contact with the atmosphere. Here, the suberized cell walls turn brown. It is the suberin that is responsible for the dull and reddish/brown color of a russeted fruit surface [152]. Due to the lipophilic character of suberin, suberized cell walls present a significant barrier to water loss [153].

From the above, it is evident that during the early stages of periderm formation, the periderm may still be covered by a cuticle, epidermal cells and some hypodermal cells. The periderm reaches the surface as growth proceeds and as the residues of the primary fruit skin (now hydraulically isolated and desiccated) tear and are sloughed off.

5. Physiology of Russeted Fruit Skin

The physiological properties of the fruit skin change with russetting. For the fruit of a particular apple cultivar, the water vapor permeance of a russeted area of skin is higher than that of a non-russeted area [11,134]. Furthermore, a non-russeted area of the primary surface of a russet-susceptible apple cultivar has a higher water vapor permeance than a

non-russeted area of a non-russet-susceptible cultivar [38]. This latter finding is likely due to a higher incidence of microcracking of the cuticles of the russet-susceptible cultivars. Compared to non-russet-susceptible cultivars, the higher water vapor permeance results in greater water loss during storage, and thus, a higher mass loss and (possibly) more shrivel. In this way, russeted fruit have reduced cool-storage potential and shorter supermarket shelf-lives compared to non-russeted fruit. We are unaware of studies that measure the fruit skin permeances to O₂, CO₂ or ethylene of russeted fruit.

The mechanical properties of fruit skins differ slightly between russeted and non-russeted fruit surfaces. The maximum stress and maximum strain that the fruit skin can withstand without failure are of similar magnitude for non-russeted and russeted skins [18]. Enzymatically isolated periderms of apple and pear are more plastic than isolated cuticles as indexed by a higher strain at maximum stress and a lower modulus of elasticity [18]. The higher plasticity renders the periderm a very suitable 'repair patch' for an overly-strained fruit surface. It allows the periderm to cope with ongoing area expansion during growth without excessive increases in stress build up [18,154].

6. Factors in Russet Formation

Russetting has been related to a number of factors. Growth strains are considered causal in russet formation in apple [18], pear [87], loquat [116,117], tomato [120,122,123] and melon [125,126]. During growth, the skin of developing fruit is subject to considerable tangential strain [7], arising from the increase in fruit volume and hence in fruit surface (area strain). Support for the idea that excessive growth strain lies behind the formation of russet comes from the following observations. First, susceptibility to russet is highest during early fruit development [12,29,32–34,155]. During early development, the relative surface area growth rate is at a maximum, resulting in maximum rates of strain [156]. The relative surface area growth rate equals the increase in surface area per unit time (cm² d⁻¹) divided by the surface area (cm²) at that time. Relative surface area growth rate, thus, has the units d⁻¹. Second, the calyx and cheek regions of pear are more russeted than the neck [87]. Both these regions have higher relative surface area growth rates than the neck [87]. Third, the stem cavity of apple fruit is often russeted. Here, stress concentration is at maximum due to the small radius of curvature of the fruit surface [154].

Extended periods of exposure of fruit surfaces to moisture, either as liquid water or as high water-vapor concentration (high relative humidity), has been identified as causal in russetting. Typical examples include russet in apple, pear, prune, tomato, melon, grape and mango (Table 1). Surface moisture is particularly critical during the early stages of fruit development when susceptibility to russetting is high [38]. The following observations support a role for moisture in russetting: First, the development of fruit under cool, rainy and high-humidity conditions stimulates russet formation in apple [156,157] and pear [88–90]. Second, experimental exposure of fruit surfaces to water, by immersion [73], by mounting a test tube filled with water on the fruit surface [38] or by overhead sprinkling [36], results in enhanced russetting. Indeed, these techniques are often used experimentally to induce russetting [37,158].

Mechanical damage of the fruit surface is also a trigger for russetting. Mechanical damage may be caused by a combination of wind and contact of fruit with a neighboring branch, shoot, leaf or fruit. Hail also damages fruit skin and causes russetting [110,111,113,114].

Pests and diseases may cause russetting in several fruitcrops. Examples of such pests include the citrus rust mite [107–109] and the tomato rust mite [71,121]. Similarly, fungi, such as powdery mildew in apple or epiphytic yeast species in apple and pear, have been reported to be causal in russetting (Table 1).

Exposure of fruit to freezing temperatures may result in formation of russet. Characteristic shapes of russet due to frost in apple are "periderm tongues" that run from the stem cavity downwards to the equatorial plane along one side of the fruit or rings of russetting that completely surround the fruit [27,34]. Why these characteristic shapes arise is unknown.

Application of agrochemicals may increase, not affect or decrease russetting. Compounds known to induce russetting include lime sulfur, copper hydroxide and thinners such as ammonium thiosulfate or ethephon (Table 3). Surfactants such as Tween 20 or Citowett that are often used in agrochemical formulations are reported to induce russetting in some fruitcrops (Table 3). An important factor would seem to be the developmental stage at the time of agrochemical application. Applications made during periods of high susceptibility to russet (e.g., during early fruit development) are more likely to induce russet. Meanwhile, the same chemical compounds may have no effect on russet formation when applied at a later stage when susceptibility is lower. In addition, environmental conditions, such as high temperatures, that favor the rapid uptake of agrochemicals are more likely to induce russetting. Rapid uptake may result in overloading of the contacted cells and thus a phytotoxic reaction. This occurs particularly in regions of the fruit surface where spray droplets collect; the droplets coalesce, and highly-concentrated chemical deposits form as the droplets dry. Then, when the critical concentration is exceeded, the cells collapse.

Reduced incidence of russetting has been found following the application of fungicides, such as mancozeb. This effect is accounted for by a reduction in the population of fungal species that induce russet.

Table 3. Effect of fungicides, surfactants and foliar fertilizers on russet. Results are compiled from literature sources.

| Chemical | Category | Crop | Cultivar | Time of Application | Effect on Russet | Reference |
|-----------------------------------------------------------------------------------------------------------------------|--------------------------------|---------------|----------------------------|----------------------------------------------------------|------------------------|---------------|
| Di-l-p-methene (2.5%) B (300 mg L ⁻¹) | Antitranspirant | Apple | Golden Delicious | 4, 13, 21 and 27 DAFB | Increased | [48] |
| B (300 mg L ⁻¹) + Ca (2 g L ⁻¹) | Foliar fertilizer | Tomato | Mountain Spring | Weekly | Decreased | [119] |
| Dithane (4 kg ha ⁻¹) Packhard (0.5% Ca) | Foliar fertilizer | Apple | Golden Delicious Spur | Flowering, PF and FS | Increased | [39] |
| Zn (100 g ha ⁻¹) Captafol (1.8 g a.i. L ⁻¹) | Foliar fertilizer Fungicide | Apple Plum | Elstar French | Green and pink stage and at bloom beginning 60–90% FB | Increased Decreased | [40] [113] |
| Chlorothalonil (3.37 kg ha ⁻¹) Kocide (Copper hydroxide) (0.32 g L ⁻¹) | Fungicide | Grape berry | Concord | 10 DAFB | Increased | [129] |
| Kocide (Copper hydroxide) (1.5 g L ⁻¹) | Fungicide | Apple | Braeburn | Weekly starting at pink tip stage | Increased | [41] |
| Kocide (Copper hydroxide) (16 or 63 g L ⁻¹) | Fungicide | Apple | Golden Delicious | 3 to 9 weeks after FB | Increased | [42] |
| Kocide (Copper hydroxide) (16 or 63 g L ⁻¹) | Fungicide | Apple | Granny Smith | Pink bud, FB and PF | Increased | [43] |
| Copper hydroxide (50%) (2.5 kg ha ⁻¹) + amino acids (10%) (2 L ha ⁻¹) | Fungicide | Pear | Conference | PF and 1 week after PF | Increased | [89] |
| Copper hydroxide (0.3 g L ⁻¹) | Fungicide | Pear | Beurré Bosc | PF, 7, 14 and 21 DAPF | Increased | [91] |
| Copper hydroxide | Fungicide | Pear | Bosc | PF | Increased | [90] |
| Copper hydroxide Copper oxychloride | Fungicide | Apple | Idared | FB | Increased | [44] |
| Copper oxychloride (4 g L ⁻¹) | Fungicide | Apple | Red Fuji | Green tip stage | Increased | [19] |
| Lime sulfur (6 g L ⁻¹) | Fungicide | Apple | Fuji More | 90% FB | Increased | [45] |
| Lime sulfur (2%) | Fungicide | Apple | Honeycrisp | Fruitlet stage | Increased | [46] |
| Lime sulfur (2%) + winter oil Fish emulsion (3%) + 2% fish oil | Fungicide | Apple | Gala | 20% and 80% FB | Increased | [47] |
| Fish emulsion (3%) + Tween 20 (0.125%) Mancozeb (2 g L ⁻¹) + Sulfur (2 g L ⁻¹) | Fungicide | Pear | Packham's Triumph | 80% FB | Decreased | [105] |
| Wettable sulfur (17 kg ha ⁻¹) | Fungicide | Apple | Golden Delicious Spur | FB, PF and FS | Decreased | [39] |
| Ziram (2.4 g a.i. L ⁻¹) Diazinon (0.08%) | Fungicide Insecticide | Plum Apple | French Golden Delicious | 60–90% FB 18 DAFB | Decreased Increased | [113] [48] |
| Rape oil (10 or 30 g L ⁻¹), Sunflower oil (30 g L ⁻¹), Soya oil (30 g L ⁻¹) | Oil | Apple | Golden Delicious | FB | Increased | [49] |
| Superior oil (0.5%) Citowett (>1%) | Oil Surfactant | Apple | Golden Delicious | 18 DAFB FB, PF and 10 weeks after FB | Increased Increased | [48] [50] |
| Tween 20 (≥1%) Ortho X-77 (1.0%) | Surfactant | Apple | Suntan | 3 weeks after FB | Increased | [51] |

Table 3. Cont.

| Chemical | Category | Crop | Cultivar | Time of Application | Effect on Russet | Reference |
|-----------------------------------------------------------------------------------------------|------------|-------|------------------------------|---------------------------------------|------------------|-----------|
| Polysorbate 20 (0.5%), Polysorbate 60 (0.5%), Polysorbate 80 (0.5%), Lecithin (0.5%) | Surfactant | Apple | Golden Delicious | At 12.5 + 18 or 18 + 20 mm diameter | Increased | [52] |
| Polysorbate 20 (0.5%), Polysorbate 60 (0.5%) | Surfactant | Apple | Fuji | At 12.5 + 18 mm diameter | Increased | [52] |
| Potassium soap (500 mg L ⁻¹) | Surfactant | Apple | Golden Delicious Smoothie | FB and 2 DAFB | Increased | [53] |
| Ammonium thiosulphate (4%) | Thinner | Apple | Golden Delicious | 20% FB | Increased | [55] |
| Ammonium thiosulphate (1.2%) | Thinner | Pear | Conference | 20% or 50% FB | Increased | [92] |
| Endothal (0.8–1.2 mL L ⁻¹) + CyLex (150 mg L ⁻¹) | Thinner | Apple | Oregon Spur Red Delicious | 80% FB | Increased | [54] |
| Apasil (silicon dioxide) (2.5%) | Other | Apple | Golden Delicious | 1 to 4 applications between 4–25 DAFB | Decreased | [48] |
| PEG 20000 (2.5%) | Other | Apple | Golden Delicious | 4, 13, 21 and 27 DAFB | Increased | [48] |

PEG = polyethylene glycol; FB: full bloom; PF: petal fall; FS: fruit set; DAFB: days after full bloom; DAPF: days after petal fall; Dithane: ethylene-bis dithy-ocarbamate manganese 62%, Mn 16%, Zn 2%; Packard: 8% Ca, 6% carboxylic acids, 0.5% B, pH < 3.0.

7. The Mechanism of Russeting—A Central Role for Cuticular Microcracks

Microscopic cracks in the cuticle, so-called microcracks, play a key role in russet formation [12,159]. Microcracks are invisible, or barely visible, to the naked eye. They are limited to the thickness of the cuticle and do not propagate deeper into the underlying cell layers [160]. Importantly, the formation of microcracks provides a unifying explanation for a diverse list of factors found to trigger russeting.

7.1. Temporal and Spatial Heterogeneity

High growth strains represent the critical factor for microcracking of the cuticle. The skin of a developing fruit is subject to ongoing tangential strain as the fruit volume and, hence, the fruit surface area increases during growth [7]. In the epidermal and hypodermal cell layers, the increase in skin surface area is accommodated by a combination of cell division (more cells) and cell extension (larger cells). Furthermore, some epidermal cells change their shape from ‘portrait’ to ‘landscape’ (in anticlinal view) as they increase in periclinal area and decrease in anticlinal height, but without significant change in (anticlinal) perimeter [16,151,161,162]. The change in cell shape implies that areas of previously anticlinal cell walls de-bond and change their orientation to form part of the expanding periclinal cell wall [162]. Such a re-orientation of cell wall material will focus the associated cuticular strain on the narrow region immediately above the anticlinal cell walls. Because the cuticle is a non-living polymer, it cannot divide but instead is dragged along (stretched) as the underlying surface expands. The strain concentration above the anticlinal cell wall (see just above) makes the cuticle particularly vulnerable to microcracking in this region. This explains the characteristic pattern of microcracks above the anticlinal cell walls as seen in a number of fruit crops, including in apple [162,163]. It also explains why fruits of many species are particularly susceptible to microcracking and russet formation during early-stage development [73]. In early-stage fruit development, the relative surface area growth rate is maximal.

Whether the microcracks propagate more deeply to traverse the entire cuticle or instead remain shallow and limited to the outer (older) volume of the cuticle depends on the relativity between the rate of deposition of new cuticular material (on the inside, adjacent to the cell wall) and the rate of fruit area growth. As mimicked in a uniaxial tensile test of a portion of fruit skin, a high surface area growth rate, in the absence of an appropriately high cuticle deposition, causes the cuticle to thin and thus fail. This occurs before the cellular components fail [164]. Correspondingly, a high rate of cuticle deposition in the absence of an appropriate surface area occurs and results in an increase in cuticle thickness. In apple fruit skin, the rate of cutin and wax deposition usually exceeds that required to match the increase in fruit surface area. Hence, cuticle thickness increases during development [165].

As previously noted, the deposition of cutin occurs on the inner surface of the cuticle (i.e., adjacent to the cell wall) [166]. Thus, the outer cuticle layers are older and, thus, have a longer history of being stretched and are more strained than the younger, inner layers [167]. This results in a radial gradient in strain across the cuticle. The gradient also accounts for the occurrence of shallow microcracks in the outer layers of the apple fruit cuticle that do not extend through to the inner layers [12,168,169]. Because cuticular microcracks differ in depth, the extent of impairment of the cuticle’s barrier properties differ. These factors explain why shallow microcracks occur on fruit surfaces without triggering periderm formation, whereas deep ones do trigger it.

Temporal and spatial heterogeneity in fruit expansion during growth is another factor in strain concentration and thus microcracking of the cuticle. The heterogeneity may be due to irregular and variable cell sizes in the epidermis [151,161,170]. Moreover, structures in the epidermis may vary cuticle stiffness—structures such as stomata [171], lenticels [172] and trichomes. Thus, cuticular microcracks may be associated with trichomes and lenticels [173]. Furthermore, cellular heterogeneity may also arise from damage caused by browsing pests,

diseases, agrochemical phytotoxicity or freezing injury. Again, periods of high rates of surface area growth result in high susceptibility to microcracking.

Moisture induces microcracking and subsequent russetting—occurring either as liquid-phase water on the fruit surface or as high concentrations of vapor-phase water close by (high humidity). While these trigger effects are well documented for a number of fruitcrop species, including apple, grape and sweet cherry [27,37,62,158], the mechanistic bases for these effects are not known. A possible explanation for moisture-induced microcracking is a higher state of hydration of the cuticle. Cuticular hydration decreases its modulus of elasticity, stiffness and fracture force, whereas its fracture strain increases [8,73]. All these changes increase the likelihood of cuticular microcracking. Other possible explanations include a weakening of cell-to-cell adhesion due to the swelling of cell walls [174].

7.2. Trigger and Signal Transmission

The question remains, how does cuticular microcracking trigger periderm formation? Microcracking occurs in the cuticle, but periderm formation occurs in the hypodermis, several cell layers below. This implies that some signals are transmitted across several cell layers that connect the two processes.

We know that microcracks impair the barrier properties of the cuticle and that this seems to trigger periderm formation. We hypothesize that these two are related, with the reduction in barrier properties somehow triggering the initiation of the periderm. What support is there for this hypothesis? First, when periderm formation is induced experimentally in apple fruit by exposing the fruit surface to moisture, the periderm begins to form only after the surface moisture is removed [37,158]. Apparently, although surface moisture has induced the cuticular microcracking, the periderm formation has been induced by the re-exposure of the (now) microcracked cuticle to the atmosphere. This conclusion is based on histological evidence [37] and gene expression analysis [158]. Second, in another experiment with apple, the formation of a wound periderm was markedly delayed when the periderm-inducing wound was sealed by silicone rubber (Chen, unpublished data). Both these experimental results indicate that the trigger is related to the impaired barrier function. Potential candidate triggers are (1) a decrease in the tissue water potential (more negative) as a result of an increase in transpiration through the microcrack and/or (2) an increase in internal O₂ concentration and/or a decrease in internal CO₂ concentration [37,158]. Based on the literature, an increase in the internal O₂ concentration is the more likely trigger. Thus, in kiwifruit, O₂ is essential for wound-induced suberization [175]; in grape, the O₂ concentrations just below the cuticle is lower than in the ambient atmosphere and decreases with increasing distance from the surface [176]; in apple, similar results have been reported [177] and, in potato, periderm and suberin formation are inhibited by a low O₂ concentration and a high CO₂ concentration [178,179].

8. Management

Various approaches have been investigated to reduce or eliminate russetting: (1) Spray applications of gibberellins and other plant growth regulators (PGRs), (2) applications of foliar fertilizers and other compounds, (3) the exclusion of moisture using bagging and (4) selective breeding.

8.1. Application of PGRs

The gibberellins A₃ (GA₃) and A₄₊₇ (GA₄₊₇) are used to improve peel finish and reduce russet in russet-susceptible cultivars of apple, pear, grape and pomegranate (Table 1). Typically, four sprays of 10 mg L⁻¹ gibberellic acid (GA) at 10 d intervals starting from petal fall are applied. Russet is reduced significantly (Table 4). The modes of action of GA in decreasing russet formation are several-fold. First, GA results in more uniform and smaller epidermal cells [30]. Skins comprising smaller epidermal cells are likely to be mechanically stiffer. Furthermore, the structural support of the cuticle provided by smaller cells is more uniform. This decreases stress concentrations, a critical factor in microcracking. Second,

GA decreases moisture-induced microcracking in russet-susceptible 'Golden Delicious' apple [73]. Applications of GA have no effect on cuticle mass, wax content or mechanical strength of the isolated apple fruit cuticle [73].

Often, GA is combined with the cytokinin benzyladenine (BA). In this combination, BA is thought to offset certain adverse effects that GA may have on flowering [58,80]. Further, GA₄₊₇ plus BA (known commercially as 'Promalin') increases fruit size and alters fruit shape. The length to width ratio of the fruit increases, particularly in the calyx region, with the result that fruit have more extended calyx lobes [80,180]. If BA is applied alone, it increases russeting [58,93]. The reason for this negative effect is unknown. The combination GA₄₊₇ plus BA decreases russet only to the same extent as GA₄₊₇ (Table 4).

In grapes, GA₃ plus the cytokinin *N*-(2-chloro-4-pyridyl)-*N'*-phenylurea (CPPU) reduces russeting, but GA₃ alone has little effect on russeting [181]. It is thought that CPPU stimulates cell division with the result that fruit have larger numbers of smaller cells [182,183]. Whether these effects also apply for the epidermis and whether microcracking of the cuticle is decreased, as observed in apple, is unknown. We suggest that such an effect would not be unlikely, and it would also account for reduced russeting following CPPU application.

8.2. Foliar Sprays of Fertilizers and Other Compounds

Insufficient supplies of boron (B) cause a number of fruit disorders, including russeting [184]. In mango, sprays of B plus Ca result in thicker cell walls and smaller intercellular spaces. As a consequence, cells are more densely packed, thereby providing greater mechanical stiffness and thus better support for the cuticle [185]. The potential roles of B in russeting also include effects on cell wall synthesis, lignification and cell wall structure, for example, by cross-linking cell wall constituents, such as pectins [186]. It is thought that B also helps maintain cell wall extensibility. In B-deficient plants, cell walls become less elastic and more rigid [184]. This causes cell walls to crack more easily and/or cells to separate from one another under tension along their middle lamellae. A separation of epidermal and/or hypodermal cells weakens the cellular support substrate for the cuticle and is therefore likely to increase cuticular microcracking. There were no effects on russeting following applications of B in pomegranate [137]. However, B applied alone or in combination with Ca did reduce russeting in tomato [119,124]. Several studies have reported decreased microcracking of fruit following applications of B, with or without Ca [187–190]. Since the initial steps in fruit cracking (macrocracking) and russeting would seem to be the same, in that both processes first require cuticular microcracking [191], it would not seem unlikely that applications of B will also decrease microcracking and russeting.

A small number of studies have reported on the effects of 'exotic' compounds on russeting. Thus, chlorogenic acid applied during early development reduced russet formation in 'Golden Delicious' apples. The authors suggest inhibition of lignin synthesis is the underlying mechanism [84]. In other studies, calmodulin and various fruit coatings have been applied, and these are reported to reduce russeting [82]. While the mode of action of calmodulin in inhibiting russeting is unknown, fruit coatings are likely to cover and thus help seal cuticular microcracks and thereby may help restore the impaired barrier functions of a microcracked cuticle. Unfortunately, direct evidence for the effect is lacking. For such an effect, the permeance of the 'exotic' coating to O₂, CO₂ and ethylene should be similar to that of an intact cuticle. Ideally, the coating should be waterproof if it is to be rain-fast. Lastly, the stomatal conductance of the leaves must not be compromised by these exotic coatings, or photosynthesis will be adversely affected—note that it is commercially impracticable to apply these coatings to the fruit without also applying them to the leaves.

Table 4. Effect of the plant growth regulators (PGR) benzyladenine (BA), 4-(2,2-Dimethylhydrazin-1-yl)-4-oxobutanoic acid (daminozide), ethephon, gibberellin A₄₊₇ (GA₄₊₇), gibberellic acid (GA₃) acid (GA), *N*-(2-chloro-4-pyridyl)-*N'*-phenylurea (CPPU), (2RS, 3RS)-1-(4-chlorophenyl)-4,4-dimethyl-2-(1H,2,4-triazol-1-yl) pentan-3-ol (Paclobutrazol) on russetting. Results are compiled from literature sources.

| PGR | Crop | Cultivar | Concentration (mg L ⁻¹) | Time and Frequency of Application | Effect on Russet | Reference |
|---------------------------------------------------------|-------------|-----------------------------------------|-------------------------------------|-------------------------------------------------------------|------------------|-----------|
| BA | Apple | Golden Delicious, Jonathan | 50 or 150 | 80% PF and once after 7–10 d | Increased | [56] |
| BA | Apple | Elstar | 200 or 300 | At 10–12 mm fruit diameter | Increased | [57] |
| BA | Apple | Golden Delicious | 50 | 0, 4, 12, 26, 42, 57 DAFB | Increased | [58] |
| BA | Pear | Bartlett | 150 | PF and 10 mm stage | Increased | [93] |
| Daminozide | Apple | Golden Delicious | 2000 | 3 DAFB | Increased | [48] |
| Ethephon | Apple | Fuji | 400 | FB | Increased | [59] |
| GA ₄ , GA ₇ and GA ₄₊₇ | Apple | Golden Delicious, Karmijn de Sonnaville | 10 | 4 applications in 10 d intervals beginning at PF | Decreased | [32] |
| GA ₄₊₇ | Apple | Golden Delicious | 200 | 4 applications in 7 d intervals beginning at PF | Decreased | [75] |
| GA ₄₊₇ | Apple | Golden Delicious | 62.5, 125 or 250 | 8–15 DAFB | Decreased | [76] |
| GA ₄₊₇ | Apple | Golden Delicious | 25 | 0, 4, 12, 26, 42, 57 DAFB | Decreased | [58] |
| GA ₄₊₇ | Apple | Golden Delicious | 10 | 4 applications in 10 d intervals beginning at PF | Decreased | [73] |
| GA ₄₊₇ | Apple | Golden Delicious | 10 | 4 applications in 10 d intervals beginning at PF | Decreased | [77] |
| GA ₄₊₇ | Apple | Golden Delicious | 10 | PF | Decreased | [33] |
| GA ₄₊₇ | Apple | Golden Delicious | 15 or 30 | 5 applications in 7 to 10 d intervals beginning at PF | Decreased | [30] |
| GA ₄₊₇ | Apple | Golden Delicious | 10 | 5 applications in 7 d intervals beginning at FB | Decreased | [78] |
| GA ₄₊₇ | Apple | Golden Delicious | 20 | 3 applications in 10 d intervals beginning at PF | Decreased | [72] |
| GA ₄₊₇ | Apple | Golden Delicious, Jonathan | 25–200 | 80% PF and once after 7–10 d | Decreased | [56] |
| GA ₄₊₇ | Apple | Golden Delicious | 5, 10 | 4 applications in 7 d intervals beginning at PF | Decreased | [79] |
| GA ₄₊₇ | Pear | Packham's Triumph | 5, 10, 20 | 80% PF and 0 to 3 additional sprays at 10 or 15 d intervals | Decreased | [105] |
| GA ₄₊₇ | Apple | Golden Delicious | 6, 12, 24, 50 | Beginning of bloom and 3 additional sprays at 7 d intervals | Decreased | [80] |
| GA ₄₊₇ + BA | Apple | Golden Delicious | 6, 12, 24, 50 | Beginning of bloom and 3 additional sprays at 7 d intervals | Decreased | [80] |
| GA ₄₊₇ + BA | Apple | Scarlet Spur II | 1, 2.5 or 5 | 3 to 4 applications in 10 d intervals beginning at PF | Decreased | [81] |
| GA ₃ | Apple | Karmijn de Sonnaville | 10 | 4 applications in 10 d intervals beginning at PF | Decreased | [32] |
| GA ₃ | Apple | Golden Delicious | 100 or 200 | PF | Decreased | [33] |
| GA ₃ | Apple | Golden Delicious | 100 | PF | Increased | [60] |
| GA ₃ | Pomegranate | G-137 | 50 | Mid May–Mid June | Decreased | [137] |
| CPPU | Apple | Scarlet Spur II | 2, 5 or 10 | 3 to 4 applications in 10 d intervals beginning at PF | Decreased | [81] |
| CPPU | Apple | Golden Delicious | 20 | PF | Increased | [60] |
| CPPU | Pomegranate | Kandhari | 5 or 10 | Mid May | Decreased | [140] |
| CPPU | Pomegranate | G-137 | 5 | Mid May–Mid June | Decreased | [137] |

Table 4. Cont.

| PGR | Crop | Cultivar | Concentration (mg L ⁻¹) | Time and Frequency of Application | Effect on Russet | Reference |
|------------------------------------------------|--------|------------------------------|---------------------------------------------------|-----------------------------------|------------------|-----------|
| CPPU+GA ₃ | Grapes | Shine Muscat | 10 CPPU + 25 GA ₃ | FB | Decreased | [132] |
| CPPU+GA ₃ , CPPU+GA ₄ | Apple | Golden Delicious | 20 CPPU + 100 GA ₃ /GA ₄ | PF | Increased | [60] |
| Paclobutrazol | Apple | Suntan | 120 or 240 | 3 weeks after FB | Increased | [51] |
| Paclobutrazol | Apple | Smoothie Golden Delicious | 250 | Between early bloom and PF | Increased | [61] |

FB: full bloom; PF: petal fall; DAFB: days after full bloom.

Where russetting is induced primarily by insect pests or fungi, spray applications of suitable agrochemicals will likely be successful in decreasing russetting. Examples reported include applications of zineb for citrus mites [108,109,112] or captafol or ziram for scab in prune [113]. However, the right dose and timing must be chosen, or the product may itself cause russetting.

8.3. Bagging

Fruit bagging is reported to be a successful countermeasure to inhibit russetting in several fruitcrop species (Table 5). Bagging prevents russetting by keeping the fruit surface dry. However, selecting a suitable material for the bag is critical as the bag material must prevent contact of the fruit surface with liquid water and, at the same time, avoid an elevated humidity in the microclimate of the enclosed fruit. A high-humidity environment inside the bag severely increases russetting [17], probably by increasing cuticular microcracking [73].

Furthermore, the bagged fruit must not overheat [192]. The spectral properties of the bagging material affect the amount and wavelengths of light reaching the fruit surface [193]. In those fruitcrop species and cultivars with colored skins, and where light absorption by the bag impairs pre-harvest fruit coloring, the bag is removed shortly before harvest to induce coloring. With this, there is an increased risk of sunburn, so removal of the bag must be done cautiously, possibly stepwise—for example, by using multi-layer bags [192]. Other benefits of pre-harvest bagging include a decreased incidence of sunburn [194,195], pest infestation and hail damage [196]. However, bagging fruit is laborious, so it requires a high-value product, a high-end market and/or a low labor cost for it to be economic.

Table 5. Effect of bagging fruit on russet and color. Data are compiled from literature sources.

| Type of Bag | Crop | Cultivar | Time of Bagging | Effect on Russet | Effect on Color | Reference |
|-------------------------------------------------|-------|----------------------------------|--------------------|------------------|---------------------|-----------|
| Polythene bag (Kordite freeze bags) | Apple | Golden Delicious, Rome Beauty | 18 mm diameter | Increased | Greener groundcolor | [17] |
| Polythene bag with aluminum paper | Pear | Packham's Triumph | Fruit set | Increased | Not determined | [105] |
| Microperforated polypropylene bags | Pear | Doyenne du Comice | 30 DAFB | Decreased | Not determined | [100] |
| Nylon (polyamide) | Apple | Golden Delicious, Rome Beauty | 18 mm diameter | Increased | Decreased red color | [17] |
| Kraft paper bags | Apple | Golden Delicious, Rome Beauty | 18 mm diameter | Decreased | Decreased red color | [17] |
| Kraft paper bags | Mango | Apple | 70 DAFB | Reduced | Decreased red color | [135] |
| Kraft paper bags | Apple | Golden Delicious | 5 DAFB | Reduced | Not determined | [48] |
| White, yellow and discoloration bags | Apple | Gamhong | 20, 30 and 40 DAFB | Reduced | No change | [82] |
| Light impermeable double layer paper bags | Apple | Golden Delicious | 20 DAFB | No russet | Not determined | [86] |
| Paper bags (single layer) | Pear | Cuiguan | 35 DAFB | Decreased | More yellow | [101] |
| Paper bags (white, single layer) | Pear | Cuiguan | 20 DAFB | No russet | No change | [102] |
| Paper bags (yellow-white, double layer) | Pear | Cuiguan | 40 DAFB | No russet | No change | [102] |
| Papers bags (single layer + double layers) | Pear | Cuiguan | 28 DAFB | Decreased | Lighter color | [103] |
| Paper bags (double layer) | Pear | Cuiguan | 20 + 45 DAFB | No russet | Greener groundcolor | [101] |
| Paper bag (double layered with attached filter) | Pear | Niitaka | 30–40 DAFB | Decreased | Lighter color | [104] |
| Paper bag (triple layer) | Pear | Concorde | After June drop | Increased | Lighter color | [197] |

DAFB = days after full bloom.

8.4. Breeding

In the long term, a breeding approach to control russet will likely be the most successful since russet susceptibility is a genetically controlled trait [198–201]. For a review on the molecular biology of russet formation, the reader is referred to the recent reviews by Macnee et al. [23] and Wang et al. [24].

In apple, the anatomies of the skins of russet-susceptible and russet-non-susceptible cultivars have been compared. The cellular layers of the skin differ [151,161]. The russet-susceptible cultivars have larger cells and more variable cell sizes in both the epidermis and hypodermis [161,170]. These result in higher stiffness and lower strain at fracture during early fruit development when russet susceptibility is highest [161]. When subjected to a tangential growth strain, skin cells of irregular size and shape result in greater stress concentrations and increased likelihood of failure. Comparisons of russet-susceptible and russet-non-susceptible cultivars reveal no consistent differences in cuticular properties—such as mass per unit area of the cuticular membrane, the dewaxed cuticular membrane or wax content [31]. Furthermore, there were no significant differences relating to russet susceptibility in cuticular strain or cuticular mechanical properties, as determined in uniaxial tensile tests (i.e., maximum force, strain at maximum force or stiffness of the cuticular membrane) [18]. Genotypes meeting the following criteria are likely to exhibit low susceptibility to russetting: (1) A long period of skin cell division, so the increase in fruit surface area is substantially accounted for by increases in the numbers of cells, rather than by increased cell expansion (which is often associated with changes in epidermal cell aspect ratio). (2) Smaller epidermal and hypodermal cells are also of more uniform size. These are better able to sustain high tensile forces and offer less stress concentration and lower chances of failure. (3) Lack of stress concentration at stomata, lenticels and trichomes. Susceptibility to failure at these sites may be checked by monitoring formation of microcracks following moisture exposure of the fruit surface. Incubating fruit in the fluorescent tracer acridine orange permits localized penetration through microcracks. When viewed using a fluorescence microscope, microcracks are easily identified by the fluorescing ‘halo’ surrounding sites of preferential uptake.

9. Conclusions

The locally impaired barrier properties of the cuticle due to a microcrack and, probably, increased O₂ diffusion seem to be the primary trigger for periderm formation. Microcracking is likely the integrator of a range of factors that induce russetting. These factors include growth stress, surface moisture and high humidity, but also pests and diseases, mechanical wounds and freezing temperatures. Significant progress has been made in our understanding of molecular biology and of the physiology of russetting.

The classical concepts of reducing russetting by spray applications of gibberellins, with or without cytokinins, or of B and/or of Ca have a sound mechanistic basis and are reported to be effective in a range of fruitcrop species. The identification of impaired barrier properties of the cuticle as the trigger causing periderm formation now provides promising options for russet management that merit further research. These include applications of ‘exotic’ coatings during critical phases of fruit development, especially when relative surface area growth rates are high. In addition, the prevention of radial extension (i.e., deepening) of microcracks by stimulating the rate of cuticle deposition is not an unrealistic strategy.

Recently, evidence has been presented that feeding oleic acid to the apple fruit surface results in significant incorporation of oleic acid into the cutin fraction [202]. If this treatment could be upscaled in the field to generate gravimetrically detectable increases in cuticle thickness following spray application of a suitable precursor, the increased cutin deposition could hinder cuticle microcracks from propagating so as to fully traverse the cuticle. When applied during phases of high rates of relative surface area growth, the formation of traversing microcracks and, hence, of russet may be prevented or reduced. Several of these aspects merit further study.

Author Contributions: A.W. and T.A. compiled the tables; A.W., T.A. and M.K. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: The study was funded by a grant (KN 402/21-1 from the Deutsche Forschungsgemeinschaft) to M.K. and a stipend from the German Academic Exchange Service (DAAD) to T.A. The publication of this article was funded by the Open Access fund of Leibniz Universität Hannover.

Acknowledgments: We thank Alexander Lang for helpful comments on an earlier version of this manuscript, Martin Brüggewirth for permission to use images and the DFG and the DAAD for financial support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Becker, M.; Kerstiens, G.; Schönherr, J. Water permeability of plant cuticles: Permeance, diffusion and partition coefficients. *Trees* **1986**, *1*, 54–60. [[CrossRef](#)]
2. Yeats, T.H.; Rose, J.K. The formation and function of plant cuticles. *Plant Physiol.* **2013**, *163*, 5–20. [[CrossRef](#)] [[PubMed](#)]
3. Krauss, P.; Markstadter, C.; Riederer, M. Attenuation of UV radiation by plant cuticles from woody species. *Plant Cell Environ.* **1997**, *20*, 1079–1085. [[CrossRef](#)]
4. Serrano, M.; Coluccia, F.; Torres, M.; L'Haridon, F.; Metraux, J.P. The cuticle and plant defense to pathogens. *Front. Plant Sci.* **2014**, *5*, 274. [[CrossRef](#)]
5. Reina-Pinto, J.J.; Yephremov, A. Surface lipids and plant defenses. *Plant Physiol. Biochem.* **2009**, *47*, 540–549. [[CrossRef](#)] [[PubMed](#)]
6. Bukovac, M.J.; Petracek, P.D. Characterizing pesticide and surfactant penetration with isolated plant cuticles. *Pestic. Sci.* **1993**, *37*, 179–194. [[CrossRef](#)]
7. Knoche, M.; Lang, A. Ongoing growth challenges fruit skin integrity. *Crit. Rev. Plant Sci.* **2017**, *36*, 190–215. [[CrossRef](#)]
8. Khanal, B.P.; Knoche, M. Mechanical properties of cuticles and their primary determinants. *J. Exp. Bot.* **2017**, *68*, 5351–5367. [[CrossRef](#)]
9. Piringer, A.A.; Heinze, P.H. Effect of light on the formation of a pigment in the tomato fruit cuticle. *Plant Physiol.* **1954**, *29*, 467–472. [[CrossRef](#)]
10. Lancaster, J.E. Regulation of skin color in apples. *Crit. Rev. Plant Sci.* **1992**, *10*, 487–502. [[CrossRef](#)]
11. Khanal, B.P.; Ikigu, G.M.; Knoche, M. Russetting partially restores apple skin permeability to water vapour. *Planta* **2019**, *249*, 849–860. [[CrossRef](#)] [[PubMed](#)]
12. Faust, M.; Shear, C.B. Russetting of apples, an interpretive review. *HortScience* **1972**, *7*, 233–235.
13. Charoenchongsuk, N.; Matsumoto, D.; Itai, A.; Murayama, H. Ripening characteristics and pigment changes in russeted pear fruit in response to ethylene and 1-MCP. *Horticulturae* **2018**, *4*, 22. [[CrossRef](#)]
14. Gerchikov, N.; Keren-Keiserman, A.; Perl-Treves, R.; Ginzberg, I. Wounding of melon fruits as a model system to study rind netting. *Sci. Hortic.* **2008**, *117*, 115–122. [[CrossRef](#)]
15. Sorauer, P.; Lindau, G.; Reh, L. *Manual of Plant Diseases*, 3rd ed.; The Record Press: Wilkes-Barré, PA, USA, 1922.
16. Bell, H.P. The origin of russetting in the Golden Russet apple. *Can. J. Res.* **1937**, *15*, 560–566. [[CrossRef](#)]
17. Tukey, L.D. Observations on the russetting of apples growing in plastic bags. *Proc. Am. Soc. Hortic. Sci.* **1959**, *74*, 30–39.
18. Khanal, B.P.; Grimm, E.; Knoche, M. Russetting in apple and pear: A plastic periderm replaces a stiff cuticle. *AoB Plants* **2013**, *5*, pls048. [[CrossRef](#)]
19. Jones, K.M.; Bound, S.A.; Oakford, M.J.; Wilson, D. A strategy for reducing russet in Red Fuji apples while maintaining control of black spot (*Venturia inaequalis*). *Aust. J. Exp. Agric.* **1994**, *34*, 127–130. [[CrossRef](#)]
20. Gossard, H.A. Commercial apple orcharding in Ohio. *Ohio Agric. Exp. Stn.* **1911**, *12*, 3–15.
21. Hall, F.H.; Stewart, F.C.; Blodgett, F.H. Fruit diseases found along the Hudson. *N. Y. Agric. Exp. Stn. Bull.* **1899**, *167*, 1–6.
22. Cole, S.W. *The American Fruit Book; Containing Directions for Raising, Propagating, and Managing Fruit Trees, Shrubs, and Plants; with a Description of the Best Varieties of Fruit, Including New and Valuable Kinds*; A.O. Moore Agricultural Book Publisher: New York, NY, USA, 1858.
23. Macnee, N.C.; Rebstock, R.; Hallett, I.C.; Schaffer, R.J.; Bulley, S.M. A review of current knowledge about the formation of native peridermal exocarp in fruit. *Funct. Plant. Biol.* **2020**, *47*, 1019–1031. [[CrossRef](#)] [[PubMed](#)]
24. Wang, Y.; Dai, M.; Cai, D.; Zhang, S.J.; Shi, Z. A review for the molecular research of russet/semi-russet of sand pear exocarp and their genetic characters. *Sci. Hortic.* **2016**, *210*, 138–142. [[CrossRef](#)]
25. Duggan, J.B. The training of apple trees. *Sci. Hortic.* **1957**, *13*, 62–73.
26. Biedenfeld, F. *Handbuch Aller Bekanntest Obstsorten, Band 2*; Frommann: Jena, Germany, 1854.
27. Thalheimer, M. About the russetting of apples in 2018. *Laimburg J.* **2019**, *1*, 1–4.
28. Legay, S.; Guerriero, G.; Andre, C.; Guignard, C.; Cocco, E.; Charton, S.; Boutry, M.; Rowland, O.; Hausman, J.F. MdMyb93 is a regulator of suberin deposition in russeted apple fruit skins. *New Phytol.* **2016**, *212*, 977–991. [[CrossRef](#)]
29. Skene, D.S. The development of russet, rough russet and cracks on the fruit of the apple Cox's Orange Pippin during the course of the season. *J. Hortic. Sci.* **1982**, *57*, 165–174. [[CrossRef](#)]

30. Curry, E. Increase in epidermal planar cell density accompanies decreased russetting of 'Golden Delicious' apples treated with Gibberellin A₄₊₇. *HortScience* **2012**, *47*, 232–237. [[CrossRef](#)]
31. Khanal, B.P.; Shrestha, R.; Hückstädt, L.; Knoche, M. Russetting in apple seems unrelated to the mechanical properties of the cuticle at maturity. *HortScience* **2013**, *48*, 1135–1138. [[CrossRef](#)]
32. Wertheim, S.J. Fruit russetting in apple as affected by various gibberellins. *J. Hortic. Sci.* **1982**, *57*, 283–288. [[CrossRef](#)]
33. Taylor, B.K. Effects of gibberellin sprays on fruit russet and tree performance of Golden Delicious apple. *J. Hortic. Sci.* **1978**, *53*, 167–169. [[CrossRef](#)]
34. Simons, R.K.; Chu, M.C. Periderm morphology of mature Golden Delicious apple with special reference to russetting. *Sci. Hortic.* **1978**, *8*, 333–340. [[CrossRef](#)]
35. Knoche, M.; Grimm, E. Surface moisture induces microcracks in the cuticle of 'Golden Delicious' apple. *HortScience* **2008**, *43*, 1929–1931. [[CrossRef](#)]
36. Winkler, A.; Grimm, E.; Knoche, M.; Lindstaedt, J.; Köpcke, D. Late-season surface water induces skin spot in apple. *HortScience* **2014**, *49*, 1324–1327. [[CrossRef](#)]
37. Chen, Y.H.; Straube, J.; Khanal, B.P.; Knoche, M.; Debener, T. Russetting in apple is initiated after exposure to moisture ends-I. Histological evidence. *Plants* **2020**, *9*, 1293. [[CrossRef](#)]
38. Khanal, B.P.; Imoro, Y.; Chen, Y.H.; Straube, J.; Knoche, M. Surface moisture increases microcracking and water vapour permeance of apple fruit skin. *Plant. Biol.* **2021**, *23*, 74–82. [[CrossRef](#)]
39. Sánchez, E.; Soto, J.M.; Uvalle, J.X.; Hernández, A.P.; Ruiz, J.M.; Romero, L. Chemical treatments in "Golden Delicious Spur" fruits in relation to russetting and nutritional status. *J. Plant. Nutr.* **2001**, *24*, 191–202. [[CrossRef](#)]
40. Wójcik, P.; Filipczak, J.; Wójcik, M. Effects of prebloom sprays of tryptophan and zinc on calcium nutrition, yielding and fruit quality of 'Elstar' apple trees. *Sci. Hortic.* **2019**, *246*, 212–216. [[CrossRef](#)]
41. Palmer, J.W.; Davies, S.B.; Shaw, P.W.; Wünsche, J.N. Growth and fruit quality of 'Braeburn' apple (*Malus domestica*) trees as influenced by fungicide programmes suitable for organic production. *N. Z. J. Crop Hortic. Sci.* **2003**, *31*, 169–177. [[CrossRef](#)]
42. Brown, G.S.; Kitchener, A.E.; Barnes, S. Calcium hydroxide sprays for the control of black spot on apples—Treatment effects on fruit quality. *Acta Hortic.* **1998**, *513*, 47–52. [[CrossRef](#)]
43. Teviotdale, B.L.; Viveros, M. Fruit russetting and tree toxicity symptoms associated with copper treatments of Granny Smith apple trees (*Malus sylvestris* Mill.). *Acta Hortic.* **1999**, *489*, 565–571. [[CrossRef](#)]
44. Momol, M.T.; Norelli, J.L.; Aldwinckle, H.S. Evaluation of biological control agents, systemic acquired resistance inducers and bactericides for the control of fire blight on apple blossom. *Acta Hortic.* **1999**, *489*, 553–557. [[CrossRef](#)]
45. Marchioretto, L.D.; De Rossi, A.; do Amaral, L.O.; Ribeiro, A.M.A.D. Efficacy and mode of action of blossom thinners on 'Fuji More' apple trees. *Sci. Hortic.* **2019**, *246*, 634–642. [[CrossRef](#)]
46. Peck, G.M.; DeLong, C.N.; Combs, L.D.; Yoder, K.S. Managing apple crop load and diseases with bloom thinning applications in an organically managed 'Honeycrisp'/'MM.111' orchard. *HortScience* **2017**, *52*, 377–381. [[CrossRef](#)]
47. Bound, S.A. Alternate thinning chemicals for apples. *Acta Hortic.* **2010**, *884*, 229–236. [[CrossRef](#)]
48. Creasy, L.L.; Swartz, H.J. Agents influencing russet on Golden Delicious apple fruits. *J. Am. Soc. Hortic. Sci.* **1981**, *106*, 203–206.
49. Stopar, M. Vegetable oil emulsions, NaCl, CH₃COOH and CaS_x as organically acceptable apple blossom thinning compounds. *Eur. J. Hortic. Sci.* **2008**, *73*, 55–61.
50. Noga, G.J.; Bukovac, M.J. Impact of surfactants on fruit quality of 'Schattenmorelle' sour cherries and 'Golden Delicious' apples. *Acta Hortic.* **1986**, *179*, 771–778. [[CrossRef](#)]
51. Richardson, P.J.; Webster, A.D.; Quinlan, J.D. The effect of paclobutrazol sprays with or without the addition of surfactants on the shoot growth, yield and fruit quality of the apple cultivars Cox and Suntan. *J. Hortic. Sci.* **1986**, *61*, 439–446. [[CrossRef](#)]
52. Stopar, M.; Hladnik, J. Polysorbates 20, 60 and 80 are apple thinning agents. *Acta Hortic.* **2020**, *1295*, 57–62. [[CrossRef](#)]
53. Alegre, S.; Alins, G. The flower thinning effect of different compounds on organic 'Golden Smoothie (R)' apple trees. *Acta Hortic.* **2007**, *737*, 67–69. [[CrossRef](#)]
54. Bound, S.A. The influence of endothal and 6-benzyladenine on crop load and fruit quality of red 'Delicious' apple. *J. Hortic. Sci. Biotechnol.* **2001**, *76*, 691–699. [[CrossRef](#)]
55. Bound, S.A.; Jones, K.M. Ammonium thiosulphate as a blossom thinner of 'Delicious' apple, 'Winter Cole' pear and 'Hunter' apricot. *Aust. J. Exp. Agric.* **2004**, *44*, 931–937. [[CrossRef](#)]
56. Taylor, B.K. Reduction of apple skin russetting by gibberellin A₄₊₇. *J. Hortic. Sci.* **1975**, *50*, 169–172. [[CrossRef](#)]
57. Maas, F. Thinning 'Elstar' apple with benzyladenine. *Acta Hortic.* **2006**, *727*, 415–421. [[CrossRef](#)]
58. McLaughlin, J.M.; Greene, D.W. Effects of BA, GA₄₊₇, and daminozide on fruit set, fruit quality, vegetative growth, flower initiation, and flower quality of 'Golden Delicious' apple. *J. Am. Soc. Hortic. Sci.* **1984**, *109*, 34–39.
59. Jones, K.M.; Koen, T.B.; Bound, S.A.; Oakford, M.J. Some reservations in thinning 'Fuji' apples with naphthalene acetic acid (NAA) and ethephon. *N. Z. J. Crop Hortic. Sci.* **1991**, *19*, 225–228. [[CrossRef](#)]
60. Bangerth, F.; Schröder, M. Strong synergistic effects of gibberellins with the synthetic cytokinin N-(2-Chloro-4-Pyridyl)-N-Phenylurea on parthenocarpic fruit set and some other fruit characteristics of apple. *Plant. Growth Regul.* **1994**, *15*, 293–302. [[CrossRef](#)]
61. El-Khoreiby, A.M.; Unrath, C.R.; Lehman, L.J. Paclobutrazol spray timing influences apple tree growth. *HortScience* **1990**, *25*, 310–312. [[CrossRef](#)]

62. Daines, R.; Weber, D.J.; Bunderson, E.D.; Roper, T. Effect of early sprays on control of powdery mildew fruit russet on apples. *Plant. Dis.* **1984**, *68*, 326–328. [[CrossRef](#)]
63. Heidenreich, M.C.M.; Corral-Garcia, M.R.; Momol, E.A.; Burr, T.J. Russet of apple fruit caused by *Aureobasidium pullulans* and *Rhodotorula glutinis*. *Plant Dis.* **1997**, *81*, 337–342. [[CrossRef](#)]
64. Gildemacher, P.; Heijne, B.; Silvestri, M.; Houbraken, J.; Hoekstra, E.; Theelen, B.; Boekhout, T. Interactions between yeasts, fungicides and apple fruit russetting. *FEMS Yeast Res.* **2006**, *6*, 1149–1156. [[CrossRef](#)] [[PubMed](#)]
65. Goffinet, M.C.; Burr, T.J.; Heidenreich, M.C.; Welsel, M.J. Developmental anatomy of russet of ‘McIntosh’ apple fruit induced by the fungus *Aureobasidium pullulans*. *HortScience* **2006**, *41*, 983. [[CrossRef](#)]
66. Gildemacher, R.; Heijne, B.; Houbraken, J.; Vromans, T.; Hoekstra, S.; Boekhout, T. Can phyllosphere yeasts explain the effect of scab fungicides on russetting of Elstar apples? *Eur. J. Plant. Pathol.* **2004**, *110*, 929–937. [[CrossRef](#)]
67. Li, C.J.; Yaegashi, H.; Kishigami, R.; Kawakubo, A.; Yamagishi, N.; Ito, T.; Yoshikawa, N. Apple russet ring and apple green crinkle diseases: Fulfillment of Koch’s postulates by virome analysis, amplification of full-length cDNA of viral genomes, in vitro transcription of infectious viral RNAs, and reproduction of symptoms on fruits of apple trees inoculated with viral RNAs. *Front. Microbiol.* **2020**, *11*, 1627. [[CrossRef](#)] [[PubMed](#)]
68. Wood, G.A. Russet ring and some associated virus disorders of apple (*Malus sylvestris* (L.) Mill) in New England. *N. Z. J. Agric. Res.* **1972**, *15*, 405–412. [[CrossRef](#)]
69. Welsh, M.F.; May, J. Virus etiology of foliar vein-flecking or ring pattern and fruit russetting or blotch on apple. *Can. J. Plant. Sci.* **1967**, *47*, 703–708. [[CrossRef](#)]
70. Easterbrook, M.A.; Fuller, M.M. Russetting of apples caused by apple rust mite *Aculus schlechtendali* (Acarina, Eriophyidae). *Ann. Appl. Biol.* **1986**, *109*, 1–9. [[CrossRef](#)]
71. Duso, C.; Castagnoli, M.; Simoni, S.; Angeli, G. The impact of eriophyids on crops: Recent issues on *Aculus schlechtendali*, *Calepitrimerus vitis* and *Aculops lycopersici*. *Exp. Appl. Acarol.* **2010**, *51*, 151–168. [[CrossRef](#)]
72. McArtney, S.; Obermiller, J.D.; Green, A. Prohexadione-Ca reduces russet and does not negate the efficacy of GA₄₊₇ sprays for russet control on ‘Golden Delicious’ apples. *HortScience* **2007**, *42*, 550–554. [[CrossRef](#)]
73. Knoche, M.; Khanal, B.P.; Stopar, M. Russetting and microcracking of ‘Golden Delicious’ apple fruit concomitantly decline due to gibberellin A₄₊₇ application. *J. Am. Soc. Hortic. Sci.* **2011**, *136*, 159–164. [[CrossRef](#)]
74. Fogelman, E.; Redel, G.; Doron, I.; Naor, A.; Ben-Yashar, E.; Ginzberg, I. Control of apple russetting in a warm and dry climate. *J. Hortic. Sci. Biotechnol.* **2009**, *84*, 279–284. [[CrossRef](#)]
75. Eccher, T. Russetting of Golden Delicious apples as related to endogenous and exogenous gibberellins. *Acta Hortic.* **1978**, *80*, 381–386. [[CrossRef](#)]
76. Eccher, T.; Boffelli, G. Effects of dose and time of application of GA₄₊₇ on russetting, fruit set and shape of ‘Golden Delicious’ apples. *Sci. Hortic.* **1981**, *14*, 307–314. [[CrossRef](#)]
77. Wertheim, S.J. Chemical thinning of Golden Delicious apple with NAA and/or carbaryl in combination with a spreader and the anti-russetting agent GA₄₊₇. *Acta Hortic.* **1986**, *179*, 659–666. [[CrossRef](#)]
78. Steenkamp, J.; Vanzyl, H.J.; Westraad, I. A preliminary evaluation of various chemical substances for the control of calyx-end russetting in Golden Delicious apples. *J. Hortic. Sci.* **1984**, *59*, 501–505. [[CrossRef](#)]
79. Bubán, T.; Rátz, M.; Oláh, L. Improved fruit shape and less russetting of apples by using gibberellins. *Acta Hortic.* **1993**, *329*, 137–139. [[CrossRef](#)]
80. Eccher, T. Control of russetting of Golden Delicious apples by growth regulator treatments. *Acta Hortic.* **1983**, *137*, 375–382. [[CrossRef](#)]
81. Sharma, S.; Sharma, N.; Sharma, D.P.; Chauhan, N. Effects of GA₄₊₇ + BA and CPPU on russetting and fruit quality in apple (*Malus × domestica*). *Indian J. Agric. Sci.* **2020**, *90*, 74–77.
82. Moon, Y.J.; Nam, K.W.; Kang, I.K.; Moon, B.W. Effects of tree-spray of calcium agent, coating agent, GA₄₊₇ + BA and paper bagging on russet prevention and quality of ‘Gamhong’ apple fruits. *Korean J. Hortic. Sci. Technol.* **2016**, *34*, 528–536. [[CrossRef](#)]
83. Basak, A.; Bielicki, P. Effect of novel organic/mineral biostimulators on fruit quality parameters in apple. *Acta Hortic.* **2010**, *873*, 295–302. [[CrossRef](#)]
84. Wang, L.; Li, J.; Gao, J.; Feng, X.; Shi, Z.; Gao, F.; Xu, X.; Yang, L. Inhibitory effect of chlorogenic acid on fruit russetting in ‘Golden Delicious’ apple. *Sci. Hortic.* **2014**, *178*, 14–22. [[CrossRef](#)]
85. Dayioglu, A.; Hepaksoy, S. Effects of shading nets on sunburn and quality of ‘Granny Smith’ apple fruits. *Acta Hortic.* **2016**, *1139*, 523–528. [[CrossRef](#)]
86. Yuan, G.; Bian, S.; Han, X.; He, S.; Liu, K.; Zhang, C.; Cong, P. An integrated transcriptome and proteome analysis reveals new insights into russetting of bagging and non-bagging “Golden Delicious” apple. *Int. J. Mol. Sci.* **2019**, *20*, 4462. [[CrossRef](#)] [[PubMed](#)]
87. Scharwies, J.D.; Grimm, E.; Knoche, M. Russetting and relative growth rate are positively related in ‘Conference’ and ‘Condo’ pear. *HortScience* **2014**, *49*, 746–749. [[CrossRef](#)]
88. Shi, C.H.; Qi, B.X.; Wang, X.Q.; Shen, L.Y.; Luo, J.; Zhang, Y.X. Proteomic analysis of the key mechanism of exocarp russet pigmentation of semi-russet pear under rainwater condition. *Sci. Hortic.* **2019**, *254*, 178–186. [[CrossRef](#)]
89. Asin, L.; Torres, E.; Vilardell, P. Orchard cooling with overtree microsprinkler irrigation to increase fruit russet on ‘Conference’ pear. *Acta Hortic.* **2011**, *909*, 557–564. [[CrossRef](#)]

90. Sugar, D.; Villardel, P.; Asin, L. Relationship of weather factors to russet incidence in 'Comice' and 'Bosc' pear fruit. *Acta Hortic.* **2015**, *1094*, 533–538. [[CrossRef](#)]
91. Sugar, D.; Basile, S.R. Russet induction in 'Beurre Bosc' and 'Taylor's Gold' pears. *Acta Hortic.* **2008**, *800*, 257–261. [[CrossRef](#)]
92. Maas, F.M.; Kanne, H.J.; van der Steeg, P.A.H. Chemical thinning of 'Conference' pears. *Acta Hortic.* **2010**, *884*, 293–304. [[CrossRef](#)]
93. Greene, D.W. Influence of abscisic acid and benzyladenine on fruit set and fruit quality of 'Bartlett' pears. *HortScience* **2012**, *47*, 1607–1611. [[CrossRef](#)]
94. Civolani, S. The past and present of pear protection against the pear psylla, *Cacopsylla pyri* L. In *Insecticides—Pest Engineering*; Perveen, F., Ed.; IntechOpen: London, UK, 2012.
95. Sanchez, J.A.; Carrasco-Ortiz, A.; López-Gallego, E.; Ramírez-Soria, M.J.; La Spina, M. Ants reduce fruit damage caused by psyllids in Mediterranean pear orchards. *Pest. Manag. Sci.* **2020**, *77*, 1886–1892. [[CrossRef](#)]
96. Westigard, P.H. Pest status of insects and mites on pear in Southern Oregon. *J. Econ. Entomol.* **1973**, *66*, 227–232. [[CrossRef](#)]
97. Lindow, S.E.; Desurmont, C.; Elkins, R.; McGourty, G.; Clark, E.; Brandl, M.T. Occurrence of indole-3-acetic acid-producing bacteria on pear trees and their association with fruit russet. *Phytopathology* **1998**, *88*, 1149–1157. [[CrossRef](#)]
98. Serdani, M.; Spotts, R.A.; Calabro, J.M.; Postman, J.D. Powdery mildew resistance in *Pyrus* germplasm. *Acta Hortic.* **2005**, *671*, 609–613. [[CrossRef](#)]
99. Spotts, R.A.; Cervantes, L.A. Involvement of *Aureobasidium pullulans* and *Rhodotorula glutinis* in russet of d'Anjou pear fruit. *Plant. Dis.* **2002**, *86*, 625–628. [[CrossRef](#)] [[PubMed](#)]
100. Amarante, C.; Banks, N.H.; Max, S. Preharvest bagging improves packout and fruit quality of pears (*Pyrus communis*). *N. Z. J. Crop Hortic. Sci.* **2002**, *30*, 93–98. [[CrossRef](#)]
101. Lin, J.; Wang, Z.H.; Li, X.G.; Chang, Y.H. Effects of bagging twice and room temperature storage on quality of 'Cuiguan' pear fruit. *Acta Hortic.* **2012**, *934*, 837–840. [[CrossRef](#)]
102. Zhang, J.; Zhang, Y.F.; Zhang, P.F.; Bian, Y.H.; Liu, Z.Y.; Zhang, C.; Liu, X.; Wang, C.L. An integrated metabolic and transcriptomic analysis reveals the mechanism through which fruit bagging alleviates exocarp semi-russetting in pear fruit. *Tree Physiol.* **2021**, *41*, 1306–1318. [[CrossRef](#)]
103. Lin, J.; Chang, Y.H.; Yan, Z.M.; Li, X.G. Effects of bagging on the quality of pear fruit and pesticide residues. *Acta Hortic.* **2008**, *772*, 315–318. [[CrossRef](#)]
104. Seo, H.H.; Lee, J.Y.; Jung, H.W. Fruit appearance improvement by using filter-attached paper bags in 'Niitaka' Pears. *Hortic. Environ. Biotechnol.* **2010**, *51*, 73–77.
105. Yuri, J.A.; Castelli, R. Pear russet control with gibberellins and other products, in cv. Packham's Triumph. *Acta Hortic.* **1998**, *475*, 303–310. [[CrossRef](#)]
106. Sugar, D.; Powers, K.A.; Basile, S.R. Mancozeb and kaolin applications can reduce russet of 'Comice' pear. *HortTechnology* **2005**, *15*, 272–275. [[CrossRef](#)]
107. McCoy, C.W. Damage and control of eriophyoid mites in crops: Styolar feeding injury and control of eriophyoid mites in citrus. In *Eriophyoid Mites: Their Biology, Natural Enemies and Control*; Lindquist, E.E., Sabelis, M.W., Bruin, J., Eds.; Elsevier: Amsterdam, The Netherlands, 1996; Volume 6, pp. 481–490.
108. Fisher, F.E. Control of citrus fruit russet in Florida with zineb. *Phytopathology* **1957**, *47*, 433–437.
109. Johnson, R.B.; King, J.R.; McBride, J.J. Zineb controls citrus rust mite. *Proc. Fla. State Hortic. Soc.* **1957**, *70*, 38–48.
110. Smoot, J.J.; Houck, L.G.; Johnson, H.B. *Market. Diseases of Citrus and Other Subtropical Fruits*; U.S. Department of Agriculture, Agricultural Research Service: Washington, DC, USA, 1971.
111. Winston, J.R. *Tear-Stain of Citrus Fruits*; U.S. Department of Agriculture: Washington, DC, USA, 1921; Volume 924.
112. Johnson, R.B. The effect of copper compounds on control of citrus rust mite with zineb. *J. Econ. Entomol.* **1960**, *53*, 395–397. [[CrossRef](#)]
113. Michailides, T.J. Russetting and russet scab of prune, an environmentally induced fruit disorder: Symptomatology, induction, and control. *Plant Dis.* **1991**, *75*, 1114–1123. [[CrossRef](#)]
114. Michailides, T.J.; Ogawa, J.M. Control and induction of russet scab and wind bruise damage ("wind scab") of French prunes. In *Prune Research Reports*; California Prune Board: San Francisco, CA, USA, 1988.
115. Corbin, J.B.; Lider, J.V.; Roberts, K.O. Controlling prune russet scab. *Calif. Agric.* **1968**, *22*, 6–7.
116. Avidan, B.; Klein, I. Physiological disorders in loquat (*Eriobotrya japonica* Lindl.). I. Russetting. *Adv. Hortic. Sci.* **1998**, *12*, 190–195.
117. Wang, L.; Wang, H.C.; Hu, Y.L.; Huang, X.M. Loquat fruit physiological disorders: Creasing and russetting. *Acta Hortic.* **2007**, *750*, 269–273. [[CrossRef](#)]
118. Barone, F.; Farina, V.; Lo Bianco, R. Growth, yield and fruit quality of 'Peluche' loquat under windbreak nets. *Acta Hortic.* **2011**, *887*, 155–159. [[CrossRef](#)]
119. Huang, J.S.; Snapp, S.S. The effect of boron, calcium, and surface moisture on shoulder check, a quality defect in fresh-market tomato. *J. Am. Soc. Hortic. Sci.* **2004**, *129*, 599–607. [[CrossRef](#)]
120. Bakker, J.C. Russetting (cuticle cracking) in glasshouse tomatoes in relation to fruit growth. *J. Hortic. Sci.* **1988**, *63*, 459–463.
121. Kamau, A.W.; Mueke, J.M.; Khaemba, B.M. Resistance of tomato varieties to the tomato russet mite, *Aculops lycopersici* (Masse) (Acarina, Eriophyidae). *Insect Sci. Appl.* **1992**, *13*, 351–356. [[CrossRef](#)]
122. Ehret, D.L.; Hill, B.D.; Raworth, D.A.; Estergaard, B. Artificial neural network modelling to predict cuticle cracking in greenhouse peppers and tomatoes. *Comput. Electron. Agric.* **2008**, *61*, 108–116. [[CrossRef](#)]

123. Demers, D.A.; Dorais, M.; Papadopoulos, A.P. Yield and russetting of greenhouse tomato as influenced by leaf-to-fruit ratio and relative humidity. *HortScience* **2007**, *42*, 503–507. [[CrossRef](#)]
124. Jobin-Lawler, F.; Simard, K.; Gosselin, A.; Papadopoulos, A.P.; Dorais, M. The influence of solar radiation and boron-calcium fruit application on cuticle cracking of a winter tomato crop grown under supplemental lighting. *Acta Hort.* **2002**, *580*, 235–239. [[CrossRef](#)]
125. Meissner, F. Die Korkbildung der Früchte von *Aesculus*- und *Cucumis*-Arten. *Osterr. Bot. Z.* **1952**, *99*, 606–624. [[CrossRef](#)]
126. Keren-Keiserman, A.; Tanami, Z.; Shoseyov, O.; Ginzberg, I. Differing rind characteristics of developing fruits of smooth and netted melons (*Cucumis melo*). *J. Hortic. Sci. Biotechnol.* **2015**, *79*, 107–113. [[CrossRef](#)]
127. Rose, D.H.; Bratley, C.O.; Pentzer, W.T. *Market. Diseases of Fruits and Vegetables: Grapes and Other Small Fruits*; U.S. Department of Agriculture: Washington, DC, USA, 1939.
128. Sholberg, P.L.; Boule, J. Palmolive (R) detergent controls apple, cherry, and grape powdery mildew. *Can. J. Plant. Sci.* **2009**, *89*, 1139–1147. [[CrossRef](#)]
129. Goffinet, M.C.; Pearson, R.C. Anatomy of russetting induced in concord grape berries by the fungicide chlorothalonil. *Am. J. Enol. Vitic.* **1991**, *42*, 281–289.
130. Araya, J.E.; Merino, C.; Santibanez, F.; Sazo, L. Ring spots by feeding of *Frankliniella occidentalis* (Thysanoptera: Thripidae) on white table grapes. *Rev. Colomb. Entomol.* **2014**, *40*, 1–6.
131. De Villiers, F.J. Physiological studies of the grape. *Union S. Afr. Dep. Agric. Sci.* **1926**, *45*, 38–48.
132. Xu, Y.S.; Hou, X.D.; Feng, J.; Khalil-Ur-Rehman, M.; Tao, J.M. Transcriptome sequencing analyses reveals mechanisms of eliminated russet by applying GA₃ and CPPU on ‘Shine Muscat’ grape. *Sci. Hortic.* **2019**, *250*, 94–103. [[CrossRef](#)]
133. Huang, Y.; Wang, P.; Wang, X.; Wang, J.; Liu, F. Effects of calcium on the formation of berry russet and phenolic compounds in ‘Shine Muscat’ grape. *IOP Conf. Ser. Earth Environ. Sci.* **2021**, *792*, 012038. [[CrossRef](#)]
134. Athoo, T.O.; Winkler, A.; Knoche, M. Russetting in ‘Apple’ mango: Triggers and mechanisms. *Plants* **2020**, *9*, 898. [[CrossRef](#)]
135. Mathooko, F.M.; Kahangi, E.M.; Runkuab, J.M.; Onyangob, C.A.; Owinob, W.O. Preharvest mango (*Mangifera indica* L. ‘Apple’) fruit bagging controls lenticel discolouration and improves postharvest quality. *Acta Hort.* **2011**, *906*, 55–62. [[CrossRef](#)]
136. Drogoudi, P.; Pantelidis, G.E.; Vekiari, S.A. Physiological disorders and fruit quality attributes in pomegranate: Effects of meteorological parameters, canopy position and acetylsalicylic acid foliar sprays. *Front. Plant Sci.* **2021**, *12*, 645547. [[CrossRef](#)]
137. Sharma, N.; Belsare, C. Effect of plant bio-regulators and nutrients on fruit cracking and quality in pomegranate (*Punica granatum* L.) ‘G-137’ in Himachal Pradesh. *Acta Hort.* **2011**, *890*, 347–352. [[CrossRef](#)]
138. Joshi, M.; Schmilovitch, Z.; Ginzberg, I. Pomegranate fruit growth and skin characteristics in hot and dry climate. *Front. Plant. Sci.* **2021**, *12*, 725479. [[CrossRef](#)]
139. Ebeling, W.; Pence, R.J. New pomegranate mite: Russetting and cracking of peel characterize injury responsible for much culling. *Calif. Agric.* **1949**, *3*, 11–14.
140. Sahu, P.; Sharma, N. Fruit cracking and quality of pomegranate (*Punica granatum* L.) cv. Kandhari as influenced by CPPU and boron. *J. Pharmacogn. Phytochem.* **2019**, *8*, 2644–2648. [[CrossRef](#)]
141. Grimm, E.; Khanal, B.P.; Winkler, A.; Knoche, M.; Köpcke, D. Structural and physiological changes associated with the skin spot disorder in apple. *Postharvest Biol. Technol.* **2012**, *64*, 111–118. [[CrossRef](#)]
142. Byers, M.A. A scarf skin-like disorder of apples. *HortScience* **1977**, *12*, 226–227. [[CrossRef](#)]
143. Ferree, D.C.; Ellis, M.A.; Bishop, B.L. Scarf skin on ‘Rome Beauty’: Time of origin and influence of fungicides and GA₄₊₇. *J. Am. Soc. Hortic. Sci.* **1984**, *109*, 422–427.
144. Williams, M.H.; Vesk, M.; Mullins, M.G. Development of the banana fruit and occurrence of the maturity bronzing disorder. *Ann. Bot.* **1990**, *65*, 9–19. [[CrossRef](#)]
145. Campbell, S.J.; Williams, W.T. Factors associated with maturity bronzing of banana fruit. *Aust. J. Exp. Agric.* **1976**, *16*, 428–432. [[CrossRef](#)]
146. Beach, S.A.; Booth, N.O.; Taylor, O.M. *The Apples of New York*; J.B. Lyon: Albany, GA, USA, 1905; Volume 1.
147. Weber, R.W.S.; Zabel, D. White haze and scarf skin, two little-known cosmetic defects of apples in northern Germany. *Eur. J. Hortic. Sci.* **2011**, *76*, 45–50.
148. Williams, M.H.; Vesk, M.; Mullins, M.G. Characteristics of the surface of banana peel in cultivars susceptible and resistant to maturity bronzing. *Can. J. Bot.* **1989**, *67*, 2154–2160. [[CrossRef](#)]
149. Daniells, J.W.; Lisle, A.T.; Ofarrell, P.J. Effect of bunch-covering methods on maturity bronzing, yield, and fruit quality of bananas in North Queensland. *Aust. J. Exp. Agric.* **1992**, *32*, 121–125. [[CrossRef](#)]
150. Daniells, J.W.; Lisle, A.T.; Bryde, N.J. Effect of bunch trimming and leaf removal at flowering on maturity bronzing, yield, and other aspects of fruit quality of bananas in North Queensland. *Aust. J. Exp. Agric.* **1994**, *34*, 259–265. [[CrossRef](#)]
151. Meyer, A. A study of the skin structure of Golden Delicious apples. *Proc. Am. Soc. Hortic. Sci.* **1944**, *45*, 105–110.
152. Schreiber, L.; Franke, R.; Hartmann, K. Wax and suberin development of native and wound periderm of potato (*Solanum tuberosum* L.) and its relation to peridermal transpiration. *Planta* **2005**, *220*, 520–530. [[CrossRef](#)] [[PubMed](#)]
153. Franke, R.; Schreiber, L. Suberin—A biopolyester forming apoplastic plant interfaces. *Curr. Opin. Plant. Biol.* **2007**, *10*, 252–259. [[CrossRef](#)] [[PubMed](#)]
154. Considine, J.; Brown, K. Physical aspects of fruit growth: Theoretical analysis of distribution of surface growth forces in fruit in relation to cracking and splitting. *Plant Physiol.* **1981**, *68*, 371–376. [[CrossRef](#)] [[PubMed](#)]

155. Eccher, T.; Hajnajari, H. Fluctuations of endogenous gibberellin A₄ and A₇ content in apple fruits with different sensitivity to russet. *Acta Hort.* **2006**, *727*, 537–543. [[CrossRef](#)]
156. Creasy, L.L. The correlation of weather parameters with russet of Golden Delicious apples under orchard conditions. *J. Am. Soc. Hort. Sci.* **1980**, *105*, 735–738.
157. Barcelo-Vidal, C.; Bonany, J.; Martin-Fernandez, J.A.; Carbo, J. Modelling of weather parameters to predict russet on ‘Golden Delicious’ apple. *J. Hort. Sci. Biotechnol.* **2013**, *88*, 624–630. [[CrossRef](#)]
158. Straube, J.; Chen, Y.H.; Khanal, B.P.; Shumbusho, A.; Zeisler-Diehl, V.; Suresh, K.; Schreiber, L.; Knoche, M.; Debener, T. Russetting in apple is initiated after exposure to moisture ends: Molecular and biochemical evidence. *Plants* **2021**, *10*, 65. [[CrossRef](#)]
159. Faust, M.; Shear, C.B. Fine structure of the fruit surface of three apple cultivars. *J. Am. Soc. Hort. Sci.* **1972**, *97*, 351–355.
160. Peschel, S.; Knoche, M. Characterization of microcracks in the cuticle of developing sweet cherry fruit. *J. Am. Soc. Hort. Sci.* **2005**, *130*, 487–495. [[CrossRef](#)]
161. Khanal, B.P.; Le, T.L.; Si, Y.; Knoche, M. Russet susceptibility in apple is associated with skin cells that are larger, more variable in size, and of reduced fracture strain. *Plants* **2020**, *9*, 1118. [[CrossRef](#)] [[PubMed](#)]
162. Maguire, K.M. *Factors Affecting Mass Loss of Apples*; Massey University: Palmerston North, New Zealand, 1998.
163. Knoche, M.; Khanal, B.P.; Brüggewirth, M.; Thapa, S. Patterns of microcracking in apple fruit skin reflect those of the cuticular ridges and of the epidermal cell walls. *Planta* **2018**, *248*, 293–306. [[CrossRef](#)] [[PubMed](#)]
164. Khanal, B.P.; Knoche, M. Mechanical properties of apple skin are determined by epidermis and hypodermis. *J. Am. Soc. Hort. Sci.* **2014**, *139*, 139–147. [[CrossRef](#)]
165. Lai, X.; Khanal, B.P.; Knoche, M. Mismatch between cuticle deposition and area expansion in fruit skins allows potentially catastrophic buildup of elastic strain. *Planta* **2016**, *244*, 1145–1156. [[CrossRef](#)] [[PubMed](#)]
166. Si, Y.; Khanal, B.P.; Schlüter, O.K.; Knoche, M. Direct evidence for a radial gradient in age of the apple fruit cuticle. *Front. Plant Sci.* **2021**, *12*, 730837. [[CrossRef](#)] [[PubMed](#)]
167. Khanal, B.P.; Knoche, M.; Bußler, S.; Schlüter, O. Evidence for a radial strain gradient in apple fruit cuticles. *Planta* **2014**, *240*, 891–897. [[CrossRef](#)]
168. Konarska, A. The structure of the fruit peel in two varieties of *Malus domestica* Borkh. (Rosaceae) before and after storage. *Protoplasma* **2013**, *250*, 701–714. [[CrossRef](#)]
169. Roy, S.; Conway, W.S.; Watada, A.E.; Sams, C.E.; Erbe, E.F.; Wergin, W.P. Heat treatment affects epicuticular wax structure and postharvest calcium uptake in ‘Golden Delicious’ apples. *HortScience* **1994**, *29*, 1056–1058. [[CrossRef](#)]
170. Eccher, T. Influenza di alcuni fitormoni sulla rugginosità della “Golden Delicious”. *Riv. Ortoflorofruttic. Ital.* **1975**, *59*, 246–261.
171. Knoche, M.; Peschel, S. Deposition and strain of the cuticle of developing European plum fruit. *J. Am. Soc. Hort. Sci.* **2007**, *132*, 597–602. [[CrossRef](#)]
172. Athoo, T.O.; Khanal, B.P.; Knoche, M. Low cuticle deposition rate in ‘Apple’ mango increases elastic strain, weakens the cuticle and increases russet. *PLoS ONE* **2021**, *16*, e0258521. [[CrossRef](#)] [[PubMed](#)]
173. Konarska, A. Morphological, histological and ultrastructural changes in fruit epidermis of apple *Malus domestica* cv. Ligol (Rosaceae) at fruit set, maturity and storage. *Acta Biol. Crac. Ser. Bot.* **2014**, *56*, 35–48. [[CrossRef](#)]
174. Brüggewirth, M.; Knoche, M. Cell wall swelling, fracture mode, and the mechanical properties of cherry fruit skins are closely related. *Planta* **2017**, *245*, 765–777. [[CrossRef](#)] [[PubMed](#)]
175. Wei, X.; Mao, L.; Han, X.; Lu, W.; Xie, D.; Ren, X.; Zhao, Y. High oxygen facilitates wound induction of suberin polyphenolics in kiwifruit. *J. Sci. Food Agric.* **2018**, *98*, 2223–2230. [[CrossRef](#)]
176. Xiao, Z.; Rogiers, S.Y.; Sadras, V.O.; Tyerman, S.D. Hypoxia in grape berries: The role of seed respiration and lenticels on the berry pedicel and the possible link to cell death. *J. Exp. Bot.* **2018**, *69*, 2071–2083. [[CrossRef](#)]
177. Ho, Q.T.; Verboven, P.; Verlinden, B.E.; Schenk, A.; Delele, M.A.; Rolletschek, H.; Vercammen, J.; Nicolai, B.M. Genotype effects on internal gas gradients in apple fruit. *J. Exp. Bot.* **2010**, *61*, 2745–2755. [[CrossRef](#)]
178. Wigginton, M.J. Effects of temperature, oxygen tension and relative humidity on the wound-healing process in the potato tuber. *Potato Res.* **1974**, *17*, 200–214. [[CrossRef](#)]
179. Lipton, W.J. Some effects of low-oxygen atmospheres on potato tubers. *Am. Potato J.* **1967**, *44*, 292–298. [[CrossRef](#)]
180. Jindal, K.K.; Pal, S.; Chauhan, P.S.; Mankotia, M.S. Effect of promalin and mixtadol on fruit growth, yield efficiency and quality of ‘Starking Delicious’ apple. *Acta Hort.* **2004**, *636*, 533–536. [[CrossRef](#)]
181. Hou, X.; Wei, L.; Xu, Y.; Khalil-Ur-Rehman, M.; Feng, J.; Zeng, J.; Tao, J. Study on russet-related enzymatic activity and gene expression in ‘Shine Muscat’ grape treated with GA₃ and CPPU. *J. Plant. Interact.* **2018**, *13*, 195–202. [[CrossRef](#)]
182. Flaishman, M.A.; Shargal, A.; Shlizerman, L.; Stern, R.A.; Lev-Yadun, S.; Grafi, G. The synthetic cytokinins CPPU and TDZ prolong the phase of cell division in developing pear (*Pyrus communis* L.) fruit. *Acta Hort.* **2005**, *671*, 151–157. [[CrossRef](#)]
183. Kano, Y. Effects of CPPU treatment on fruit and rind development of watermelons (*Citrullus lanatus* Matsum. et Nakai). *J. Hort. Sci. Biotechnol.* **2015**, *75*, 651–654. [[CrossRef](#)]
184. Ganie, M.A.; Akhter, F.; Bhat, M.A.; Malik, A.R.; Junaid, J.M.; Shah, M.A.; Bhat, A.H.; Bhat, T.A. Boron—A critical nutrient element for plant growth and productivity with reference to temperate fruits. *Curr. Sci.* **2013**, *104*, 76–85.
185. Muengkaew, R.; Whangchai, K.; Chaiprasart, P. Application of calcium—boron improve fruit quality, cell characteristics, and effective softening enzyme activity after harvest in mango fruit (*Mangifera indica* L.). *Hortic. Environ. Biotechnol.* **2018**, *59*, 537–546. [[CrossRef](#)]

186. Broadley, M.; Brown, P.; Cakmak, I.; Rengel, Z.; Zhao, F. Function of Nutrients: Micronutrients. In *Marschner's Mineral Nutrition of Higher Plants*, 3rd ed.; Marschner, P., Ed.; Academic Press: San Diego, CA, USA, 2012; pp. 191–248.
187. Ferri, V.C.; Rombaldi, C.V.; Silva, J.A.; Pegoraro, C.; Nora, L.; Antunes, P.L.; Girardi, C.L.; Tibola, C.S. Boron and calcium sprayed on 'Fuyu' persimmon tree prevent skin cracks, groove and browning of fruit during cold storage. *Cienc. Rural.* **2008**, *38*, 2146–2150. [[CrossRef](#)]
188. Ghanbarpour, E.; Rezaei, M.; Lawson, S. Reduction of cracking in pomegranate fruit after foliar application of humic acid, calcium-boron and kaolin during water stress. *Erwerbs Obstbau* **2019**, *61*, 29–37. [[CrossRef](#)]
189. Singh, A.; Shukla, A.K.; Meghwal, P.R. Fruit cracking in pomegranate: Extent, cause, and management—A review. *Int. J. Fruit Sci.* **2020**, *20*, S1234–S1253. [[CrossRef](#)]
190. Kavvadias, V.; Daggas, T.; Paschalidis, C.; Vavoulidou, E.; Theocharopoulos, S. Effect of boron application on yield, quality, and nutritional status of peach cultivar Andross. *Commun. Soil Sci. Plant. Anal.* **2012**, *43*, 134–148. [[CrossRef](#)]
191. Schumann, C.; Winkler, A.; Brüggewirth, M.; Köpcke, K.; Knoche, M. Crack initiation and propagation in sweet cherry skin: A simple chain reaction causes the crack to 'run'. *PLoS ONE* **2019**, *14*, e0219794. [[CrossRef](#)]
192. Racsko, J.; Schrader, L.E. Sunburn of apple fruit: Historical background, recent advances and future perspectives. *Crit. Rev. Plant Sci.* **2012**, *31*, 455–504. [[CrossRef](#)]
193. Chonhenchob, V.; Kamhangwong, D.; Krueenate, J.; Khongrat, K.; Tangchantra, N.; Wichai, U.; Singh, S.P. Preharvest bagging with wavelength-selective materials enhances development and quality of mango (*Mangifera indica* L.) cv. Nam Dok Mai #4. *J. Sci. Food Agric.* **2011**, *91*, 664–671. [[CrossRef](#)] [[PubMed](#)]
194. Abdel Gawad-Nehad, M.A.; EL-Gioushy, S.F.; Baiea, M.H.M. Impact of different bagging types on preventing sunburn injury and quality improvement of Keitt mango fruits. *Middle East J. Agric. Res.* **2017**, *6*, 484–494.
195. Sarkomi, F.H.; Moradinezhad, F.; Khayat, M. Pre-harvest bagging influences sunburn, cracking and quality of pomegranate fruits. *J. Hortic. Postharvest Res.* **2019**, *2*, 131–142. [[CrossRef](#)]
196. Buthelezi, N.M.D.; Mafeo, T.P.; Mathaba, N. Preharvest bagging as an alternative technique for enhancing fruit quality: A review. *HortTechnology* **2021**, *31*, 4–13. [[CrossRef](#)]
197. Hudina, M.; Stampar, F. Bagging of 'Concorde' pears (*Pyrus communis* L.) influences fruit quality. *Acta Hortic.* **2011**, *909*, 625–630. [[CrossRef](#)]
198. Macnee, N.; Hilario, E.; Tahir, J.; Currie, A.; Warren, B.; Rebstock, R.; Hallett, I.C.; Chagne, D.; Schaffer, R.J.; Bulley, S.M. Peridermal fruit skin formation in *Actinidia* sp. (kiwifruit) is associated with genetic *loci* controlling russetting and cuticle formation. *BMC Plant Biol.* **2021**, *21*, 334. [[CrossRef](#)]
199. Petit, J.; Bres, C.; Mauxion, J.P.; Bakan, B.; Rothan, C. Breeding for cuticle-associated traits in crop species: Traits, targets, and strategies. *J. Exp. Bot* **2017**, *68*, 5369–5387. [[CrossRef](#)]
200. Falginella, L.; Cipriani, G.; Monte, C.; Gregori, R.; Testolin, R.; Velasco, R.; Troggio, M.; Tartarini, S. A major QTL controlling apple skin russetting maps on the linkage group 12 of 'Renetta Grigia di Torriana'. *BMC Plant Biol.* **2015**, *15*, 150. [[CrossRef](#)]
201. Legay, S.; Guerriero, G.; Deleruelle, A.; Lateur, M.; Evers, D.; André, C.M.; Hausman, J.F. Apple russetting as seen through the RNA-seq lens: Strong alterations in the exocarp cell wall. *Plant Mol. Biol.* **2015**, *88*, 21–40. [[CrossRef](#)]
202. Si, Y.; Khanal, B.P.; Sauheiß, L.; Knoche, M. Cutin synthesis in developing, field-grown apple fruit examined by external feeding of labelled precursors. *Plants* **2021**, *10*, 497. [[CrossRef](#)]