

12-1-2022

Cognitive composites for genetic frontotemporal dementia: GENFI-Cog

Jackie M. Poos
Erasmus MC

Katrina M. Moore
UCL Queen Square Institute of Neurology

Jennifer Nicholas
London School of Hygiene & Tropical Medicine

Lucy L. Russell
UCL Queen Square Institute of Neurology

Georgia Peakman
UCL Queen Square Institute of Neurology

See next page for additional authors

Follow this and additional works at: https://ir.lib.uwo.ca/neurosci_inst_pubs

Citation of this paper:

Poos, Jackie M.; Moore, Katrina M.; Nicholas, Jennifer; Russell, Lucy L.; Peakman, Georgia; Convery, Rhian S.; Jiskoot, Lize C.; van der Ende, Emma; van den Berg, Esther; Papma, Janne M.; Seelaar, Harro; Pijnenburg, Yolande A.L.; Moreno, Fermin; Sanchez-Valle, Raquel; Borroni, Barbara; Laforce, Robert; Masellis, Mario; Tartaglia, Carmela; Graff, Caroline; Galimberti, Daniela; Rowe, James B.; Finger, Elizabeth; Synofzik, Matthias; Vandenberghe, Rik; de Mendonça, Alexandre; Tiraboschi, Pietro; Santana, Isabel; Ducharme, Simon; Butler, Chris; and Gerhard, Alexander, "Cognitive composites for genetic frontotemporal dementia: GENFI-Cog" (2022). *Neuroscience Institute Publications*. 198.
https://ir.lib.uwo.ca/neurosci_inst_pubs/198

Authors


Jackie M. Poos, Katrina M. Moore, Jennifer Nicholas, Lucy L. Russell, Georgia Peakman, Rhian S. Convery, Lize C. Jiskoot, Emma van der Ende, Esther van den Berg, Janne M. Papma, Harro Seelaar, Yolande A.L. Pijnenburg, Fermin Moreno, Raquel Sanchez-Valle, Barbara Borroni, Robert Laforce, Mario Masellis, Carmela Tartaglia, Caroline Graff, Daniela Galimberti, James B. Rowe, Elizabeth Finger, Matthis Synofzik, Rik Vandenberghe, Alexandre de Mendonça, Pietro Tiraboschi, Isabel Santana, Simon Ducharme, Chris Butler, and Alexander Gerhard

RESEARCH

Open Access



Cognitive composites for genetic frontotemporal dementia: GENFI-Cog

Jackie M. Poos^{1,2}, Katrina M. Moore², Jennifer Nicholas³, Lucy L. Russell², Georgia Peakman², Rhian S. Convery², Lize C. Jiskoot^{1,2}, Emma van der Ende¹, Esther van den Berg¹, Janne M. Papma¹, Harro Seelaar¹, Yolande A. L. Pijnenburg⁴, Fermin Moreno⁵, Raquel Sanchez-Valle⁶, Barbara Borroni⁷, Robert Laforce⁸, Mario Masellis⁹, Carmela Tartaglia¹⁰, Caroline Graff¹¹, Daniela Galimberti^{12,13}, James B. Rowe¹⁴, Elizabeth Finger¹⁵, Matthis Synofzik^{16,17}, Rik Vandenberghe¹⁸, Alexandre de Mendonça¹⁹, Pietro Tiraboschi²⁰, Isabel Santana²¹, Simon Ducharme²², Chris Butler²³, Alexander Gerhard²⁴, Johannes Levin^{25,26,27}, Adrian Danek²⁵, Markus Otto²⁸, Isabel Le Ber^{29,30,31}, Florence Pasquier^{32,33,34}, John C. van Swieten¹, Jonathan D. Rohrer^{2*}  and on behalf of the Genetic FTD Initiative (GENFI)

Abstract

Background: Clinical endpoints for upcoming therapeutic trials in frontotemporal dementia (FTD) are increasingly urgent. Cognitive composite scores are often used as endpoints but are lacking in genetic FTD. We aimed to create cognitive composite scores for genetic frontotemporal dementia (FTD) as well as recommendations for recruitment and duration in clinical trial design.

Methods: A standardized neuropsychological test battery covering six cognitive domains was completed by 69 *C9orf72*, 41 *GRN*, and 28 *MAPT* mutation carriers with CDR[®] plus NACC-FTLD ≥ 0.5 and 275 controls. Logistic regression was used to identify the combination of tests that distinguished best between each mutation carrier group and controls. The composite scores were calculated from the weighted averages of test scores in the models based on the regression coefficients. Sample size estimates were calculated for individual cognitive tests and composites in a theoretical trial aimed at preventing progression from a prodromal stage (CDR[®] plus NACC-FTLD 0.5) to a fully symptomatic stage (CDR[®] plus NACC-FTLD ≥ 1). Time-to-event analysis was performed to determine how quickly mutation carriers progressed from CDR[®] plus NACC-FTLD = 0.5 to ≥ 1 (and therefore how long a trial would need to be).

Results: The results from the logistic regression analyses resulted in different composite scores for each mutation carrier group (i.e. *C9orf72*, *GRN*, and *MAPT*). The estimated sample size to detect a treatment effect was lower for composite scores than for most individual tests. A Kaplan-Meier curve showed that after 3 years, ~ 50% of individuals had converted from CDR[®] plus NACC-FTLD 0.5 to ≥ 1 , which means that the estimated effect size needs to be halved in sample size calculations as only half of the mutation carriers would be expected to progress from CDR[®] plus NACC-FTLD 0.5 to ≥ 1 without treatment over that time period.

Discussion: We created gene-specific cognitive composite scores for *C9orf72*, *GRN*, and *MAPT* mutation carriers, which resulted in substantially lower estimated sample sizes to detect a treatment effect than the individual cognitive

*Correspondence: j.rohrer@ucl.ac.uk

² Dementia Research Centre, Department of Neurodegenerative Disease, National Hospital for Neurology and Neurosurgery, UCL Institute of Neurology, 8-11 Queen Square, Box 16, London WC1N 3BG, UK
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

tests. The GENFI-Cog composites have potential as cognitive endpoints for upcoming clinical trials. The results from this study provide recommendations for estimating sample size and trial duration.

Keywords: Frontotemporal dementia, Cognition, Neuropsychology, Composite score, Language, Attention, Executive function, Memory, Social cognition

Background

Frontotemporal dementia (FTD) encompasses a heterogeneous group of early-onset neurodegenerative disorders caused by prominent frontal and/or temporal lobe degeneration with a wide range of overlapping clinical features [1]. The two main phenotypes are behavioural variant FTD (bvFTD), with prominent behavioural changes and executive dysfunction [2], and primary progressive aphasia (PPA), with impairment in language comprehension and/or production [3]. FTD is a highly heritable disease, with 20–30% of cases having an autosomal dominant pattern of inheritance [4]. The most common causes of genetic FTD are mutations in the microtubule-associated protein tau (*MAPT*), progranulin (*GRN*), and chromosome 9 open reading frame 72 (*C9orf72*) genes [4].

Clinical trials testing disease-modifying treatments for FTD are now underway, and clinical endpoints to monitor treatment response are therefore urgently needed. It is believed that interventions may have the most profound effect if initiated in the earliest stages of the disease; however, a major challenge facing these clinical trials is the lack of outcome measures that are sensitive enough to track the effects of treatment in the early stages of the disease [5–7].

Traditional outcomes such as progression to clinical diagnosis or cognitive measures developed for other forms of dementia such as Alzheimer's disease (AD) might not be well-suited to serve as endpoints for early-stage FTD treatment trials because of the large sample size and long trial duration that would be required to measure possible treatment effects or due to the psychometric properties of the tests themselves [8–10]. Sensitive outcome measures in patients with clinically diagnosed AD, such as the Alzheimer's Disease Assessment Scale Cognitive Subscale (ADAS-Cog), might not be sensitive to decline in patients with FTD [10, 11]. Multiple genetic FTD cohort studies have investigated a wide range of cognitive instruments and found gene-specific cognitive impairment and/or decline in language, executive function, social cognition, attention/processing speed, and memory, in symptomatic and presymptomatic stages [12–27]. However, due to the subtlety of cognitive decline in the early stages of the disease, using individual tests as outcome measures might not be sensitive enough to detect a treatment effect. Furthermore, an

individual cognitive test is limited to measuring only one specific symptom, and due to the heterogeneity of clinical features between FTD patients, tests from multiple cognitive domains would need to be included. A selection of the most sensitive tests for each genetic group would enable shortening of the neuropsychological test battery thereby significantly minimizing time and other resource costs compared to using a broad range of individual cognitive tests [28].

Composite scores are often used in clinical trials to reduce the number of variables used as outcome measures [8]. A composite score is any measure which combines the results of multiple cognitive and clinical assessments into a single summary score [29]. As a result, it provides a measure of multiple domains but can serve as a single primary endpoint in clinical trials [8]. Such composites have been developed for several neurodegenerative disorders, such as AD (e.g. the ADAS-Cog [11]), Parkinson's disease (PD) (e.g. the Unified Parkinson's Disease Rating Scale (UPDRS) [30]), and Huntington's disease (HD) (e.g. the Unified Huntington's Disease Rating Scale (UHDRS) [29]) but are, as of yet, lacking in FTD.

Therefore, the aim of this study was to create gene-specific cognitive composite scores for *MAPT*, *GRN*, and *C9orf72* mutation carriers in the early symptomatic stage by empirically determining the combination of neuropsychological tests most sensitive to differentiate mutation carriers from non-carriers. Data was collected within the Genetic FTD Initiative (GENFI), an international genetic FTD cohort study aimed at developing novel markers of disease onset and progression [14]. To evaluate their performance, we compared the sample size requirements between each of the proposed composites and individual cognitive tests for a theoretical trial aimed at preventing progression from a prodromal stage ($CDR^{\text{®}}$ plus NACC-FTLD [31] = 0.5) to a fully symptomatic stage ($CDR^{\text{®}}$ plus NACC-FTLD ≥ 1). Lastly, we performed time-to-event analyses to determine how many people progressed from a $CDR^{\text{®}}$ plus NACC-FTLD 0.5 to ≥ 1 , to provide recommendations on the duration of such clinical trials.

Methods

Participants

Data was included from the fifth GENFI data freeze in which participants from confirmed genetic FTD families were recruited between 30 January 2012 and 31

May 2019 in 24 centres across Europe and Canada. A total of 69 *C9orf72*, 41 *GRN*, and 28 *MAPT* mutation carriers with a CDR[®] plus NACC FTLD ≥ 0.5 and 275 mutation-negative controls (i.e. family members who tested negative for the mutation) were included in this study. Of the mutation carrier group, 41 *C9orf72*, 17 *GRN*, and 16 *MAPT* mutation carriers fulfilled the diagnostic criteria for bvFTD [2] (*C9orf72* = 36, *GRN* = 11, *MAPT* = 16), PPA [3] (*GRN* = 6), or FTD with amyotrophic lateral sclerosis (FTD-ALS) [32] (*C9orf72* = 5). Participant characteristics are summarized in Table 1,

and the number of participants included in each of the statistical analysis steps can be found in Fig. S1.

Procedure

All participants completed a comprehensive neuropsychological test battery covering six cognitive domains: language (modified Camel and Cactus Test [33]; Boston Naming Test (BNT, short 30 item version) [34]; category fluency (animals) [35]), attention/processing speed and executive function (WMS-R Digit span [34]; Trail Making Test (TMT) [36]; WAIS-R Digit Symbol test [34]; D-KEFS Colour-Word Interference Test (CWIT) [37];

Table 1 Participant characteristics and neuropsychological test results

	<i>C9orf72</i>	<i>GRN</i>	<i>MAPT</i>	Controls
Number of participants	69	41	28	275
Sex, f:m	30:39	20:21	14:14	160:115
Age	55 (12.0)	53.0 (11.4)	51.1 (12.6)	45.8 (12.7)
Education	13.7 (3.1)	14.0 (3.5)	14.3 (3.4)	14.6 (3.4)
MMSE	27.1 (3.2)	26.6 (7.0)	27.5 (3.0)	29.3 (2.1)
CDR[®] plus NACC FTLD sob	5.9 (5.5)	3.4 (4.8)	4.8 (5.0)	0.2 (0.6)
Language				
Camel and Cactus Test	- 1.81 (2.81)	- 0.57 (1.36)	- 2.10 (3.08)	-
Boston Naming Test	- 1.77 (3.32)	- 0.68 (1.62)	- 2.63 (3.16)	-
Category fluency	- 1.20 (1.05)	- 0.54 (1.04)	- 0.84 (1.14)	-
Attention and mental processing speed				
Digit span forward	- 0.39 (1.19)	- 0.08 (1.26)	0.13 (1.23)	-
Trail Making Test – part A	- 1.37 (2.17)	- 0.69 (1.63)	- 0.72 (1.54)	-
Digit symbol	- 1.18 (1.30)	- 0.62 (1.23)	- 0.67 (1.31)	-
D-KEFS CWIT – colour naming	- 2.85 (3.58)	- 0.52 (1.85)	- 1.30 (2.17)	-
D-KEFS CWIT – word naming	- 1.86 (3.11)	- 0.02 (1.46)	- 0.54 (1.47)	-
Executive function				
Digit span backward	- 0.53 (1.23)	- 0.49 (1.23)	- 0.19 (0.98)	-
Trail Making Test – part B	- 2.44 (2.95)	- 1.81 (3.06)	- 1.37 (2.58)	-
D-KEFS CWIT – ink naming	- 3.46 (3.91)	- 1.13 (2.21)	- 1.16 (2.54)	-
Phonemic fluency	- 1.18 (1.18)	- 0.08 (1.33)	- 0.64 (1.28)	-
Visuoconstruction				
Benson figure copy	- 0.90 (1.90)	- 0.06 (1.16)	- 0.46 (1.39)	-
Memory				
Benson figure recall	- 0.72 (1.57)	- 0.75 (1.46)	- 1.27 (1.91)	-
FCSRT free recall	- 1.68 (1.36)	- 0.72 (1.49)	- 1.71 (1.80)	-
FCSRT total recall	- 2.20 (3.56)	- 1.42 (3.05)	- 2.86 (3.62)	-
FCSRT delayed free recall	- 1.59 (1.59)	- 0.97 (1.58)	- 1.72 (2.04)	-
FCSRT delayed total recall	- 2.10 (3.81)	- 1.13 (3.09)	- 2.82 (4.02)	-
Social cognition				
Facial Emotion Recognition Test	- 1.67 (1.87)	- 1.00 (1.47)	- 1.04 (1.59)	-

Values are mean Z-scores (raw score – mean score controls/standard deviation of controls) corrected for age, years of education, and sex, with standard deviation in parentheses unless otherwise specified. For the FCSRT and letter fluency, an additional correction was made for language as stimuli differed between languages
 Abbreviations: *C9orf72* chromosome 9 open reading frame 72, *GRN* progranulin, *MAPT* microtubule-associated protein tau, *MMSE* Mini-Mental State Examination, *CDR[®] plus NACC FTLD sob* Clinical Dementia Rating scale plus National Alzheimer's Coordinating Center Frontotemporal Lobar Degeneration sum of boxes, *D-KEFS CWIT* Delis-Kaplan Executive Function System Colour-Word Interference Test, *FCSRT* Free and Cued Selective Reminding Test

phonemic fluency [35]), verbal and visuospatial memory (Free and Cued Selective Reminding Test (FCSRT) [20]; Benson figure recall), social cognition (Facial Emotion Recognition test [38]), and visuoconstruction (Benson figure copy). The Mini-Mental State Examination (MMSE) [39] was administered to measure global cognitive functioning, and clinical status was determined by means of a structured clinical interview, including the CDR[®] plus NACC FTLD [31].

Statistical methods

Statistical analyses were performed using Stata version 14 and R version 3.6.2. We compared the continuous demographic data between the mutation carrier groups with Kruskal-Wallis and post hoc Mann-Whitney tests. A chi-square test was used to compare sex between the groups.

All neuropsychological data were converted to Z-scores corrected for age, education, and sex compared to the control group collected within GENFI (i.e. mutation-negative participants). The FCSRT and letter fluency scores were also corrected for language as the test stimuli differed by language across the different GENFI sites. The control data available in each language can be found in Additional file 1: Table S1. Z-scores for tests with reaction times (i.e. TMT and D-KEFS CWIT) were inverted so that lower Z-scores indicated worse performance on all tests. A detailed description of how the corrected Z-scores were calculated can be found in Additional file 1.

Creating the composite scores

Least absolute shrinkage and selection operator (LASSO) [40] logistic regression models with 10-fold cross-validation were used to identify the combination of neuropsychological tests that discriminated best between each mutation carrier group and controls. Participants with missing data were excluded from this analysis. A separate model was fitted for each genetic group with carrier status as the outcome and the neuropsychological tests as the predictors. A detailed description of the statistical methods can be found in Additional file 1. The glmnet package in R was used to fit the LASSO models and carry out the cross-validation.

From the resulting model, two different cognitive composite scores were calculated: (1) an average of the scores for all cognitive tests that were selected in the model and (2) a weighted average of the scores for all cognitive tests that were selected in the model, using the regression coefficients to determine the weights.

Sample size calculation

For each outcome, the sample size was calculated for a hypothetical two-arm study with 1:1 randomization to

placebo versus active drug with 80% power to detect a treatment effect at a 5% significance level [41]. The focus of future studies is likely to be on treating people with very early symptomatic disease, and so, we focused on calculating the sample sizes for a trial of prodromal mutation carriers (i.e. CDR[®] plus NACC FTLD = 0.5) where the therapeutic drug had an effect on the progression to being fully symptomatic (i.e. CDR[®] plus NACC FTLD = 1). We therefore calculated sample sizes for a 10%, 20%, and 40% effect size where a 100% treatment effect would be the difference in the mean between the CDR[®] plus NACC FTLD 0.5 and 1 groups. Choosing the effect size in this way assumes that the hypothetical treatment will prevent a given proportion of the decline in cognitive scores seen between these two groups. For example, a 20% treatment effect assumes that the untreated group will experience the change seen between CDR[®] plus NACC FTLD 0.5 and 1 groups, but the treated group will only experience 80% of this change (i.e. 20% less). See Additional file 1 for more details on the sample size calculations and the parameters used (Additional file 1: Table S2) [41].

Time-to-event analysis

To provide recommendations on the timeline for the hypothesized trial, we present Kaplan-Meier curves showing the cumulative proportion of participants who progressed from a CDR[®] plus NACC FTLD 0.5 to ≥ 1 within the GENFI cohort over time. In this analysis, the censoring date was the date of conversion or the date of the last follow-up. As this is an ongoing prospective cohort study, not all mutation carriers completed all study visits which resulted in missing data. There were 62 mutation carriers (19 *C9orf72*, 27 *GRN*, and 16 *MAPT*) that had a CDR[®] plus NACC FTLD of 0.5 and one or multiple follow-up visits and were included in the time-to-event analysis (Additional file 1: Table S4 and Fig. 1). A log rank test was performed to compare the rate of progression between the genetic groups.

Results

Demographics

Participant characteristics for all mutation carriers are summarized in Table 1. Overall, the number of males to females differed between the groups ($p = 0.020$). *C9orf72*, *GRN*, and *MAPT* mutation carriers were older and had lower MMSE and higher CDR[®] plus NACC FTLD sum of boxes scores than controls (all $p < 0.010$). In addition, *C9orf72* mutation carriers had higher CDR[®] plus NACC FTLD sum of boxes scores than *GRN* mutation carriers ($p = 0.007$). There were no differences between the groups in years of education ($p = 0.290$). The characteristics of participants




Fig. 1 Kaplan-Meier estimates of mutation carriers that converted from CDR[®] plus NACC FTLD 0.5 to ≥ 1 . The number of mutation carriers included, and the number that progressed or were lost to follow-up are reported in Additional file 1: Table S4. *C9orf72*, chromosome 9 open reading frame 72; *GRN*, progranulin; *MAPT*, microtubule-associated protein tau; CDR plus NACC FTLD, Clinical Dementia Rating scale plus National Alzheimer's Coordinating Center Frontotemporal Lobar Degeneration

when individually stratified by CDR[®] plus NACC FTLD global score (i.e. in 0.5, 1, 2, and 3 groups) can be found in Additional file 1: Table S3.

Logistic regression analyses

The results from the logistic regression model can be seen in Table 2. A combination of category fluency, D-KEFS CWIT – colour, word and ink naming, TMT – part B, the Benson figure copy, FCSRT free recall, and the Facial Emotion Recognition Test was most sensitive to discriminate *C9orf72* repeat expansion carriers from controls. For *GRN* mutation carriers, a combination of the Camel and Cactus Test, TMT – part B, D-KEFS CWIT – ink naming, Benson figure recall, FCSRT total and delayed free recall, and the Facial Emotion Recognition Test was most sensitive. In *MAPT* mutation carriers, a combination of the Camel and Cactus Test, BNT, D-KEFS CWIT – colour naming, Benson figure recall, FCSRT free, total and delayed free recall, and the Facial Emotion Recognition Test was most sensitive to differentiate from controls. For each mutation carrier group, the average and weighted composite scores were calculated, including the tests with a negative coefficient in Table 2. A summary of the included tests that

were included in each GENFI-Cog per gene group can be seen in Fig. 2.

Sample size calculation

Sample size estimates can be observed in Table 3. In *C9orf72* repeat expansion carriers, both the average and weighted composite scores resulted in lower sample sizes than most individual cognitive tests. The only test that resulted in a lower sample size than the composite score was the D-KEFS CWIT – ink naming, with the digit symbol test also resulting in a lower sample size than the average but not the weighted composite score. In *GRN* mutation carriers, again both composite scores resulted in lower sample sizes than for most individual cognitive tests except the TMT – part B. The TMT – part A also resulted in a lower sample size than the weighted composite, but not the average composite. In addition, the D-KEFS CWIT – ink naming resulted in a sample size of less than 100, albeit not lower than the composites. In *MAPT* mutation carriers, both composites resulted in estimated sample sizes smaller than 130 with an effect size of 0.1, but the TMT – part A, digit symbol test, and D-KEFS CWIT – colour and ink naming resulted in even lower sample sizes ($n < 100$). In *C9orf72* and *MAPT* mutation carriers, the weighted composite score resulted in a lower estimated sample size than the

Table 2 Regression coefficients and corresponding weights

	<i>C9orf72</i>		<i>GRN</i>		<i>MAPT</i>	
	Coef.	Weight	Coef.	Weight	Coef.	Weight
Language						
Camel and Cactus Test			-0.004	0.003	-0.04	0.04
Boston Naming Test					-0.39	0.40
Category fluency	-0.13	0.09				
Attention and mental processing speed						
Digit span forward						
Trail Making Test – part A						
Digit symbol						
D-KEFS CWIT – colour naming	-0.06	0.04			-0.09	0.09
D-KEFS CWIT – word naming	-0.04	0.03	0.09 ^a			
Executive function						
Digit span backward						
Trail Making Test – part B	-0.07	0.05	-0.28	0.23		
D-KEFS CWIT – ink naming	-0.29	0.20	-0.24	0.20		
Phonemic fluency			0.24 ^a			
Visuoconstruction						
Benson figure copy	-0.09	0.06				
Memory						
Benson figure recall			-0.06	0.05	-0.01	0.01
FCSRT free recall	-0.50	0.35			-0.06	0.06
FCSRT total recall			-0.05	0.04	-0.30	0.31
FCSRT delayed free recall			-0.16	0.13	-0.01	0.01
FCSRT delayed total recall						
Social cognition						
Facial Emotion Recognition Test	-0.26	0.18	-0.42	0.35	-0.08	0.08

Data are presented as coefficients and weights. Coefficient gives the change in log odds of being a mutation carrier for each Z-score increase in the score on the cognitive test. Weight gives the weighting used when calculating the weighted cognitive composite score

Abbreviations: *C9orf72* chromosome 9 open reading frame 72, *GRN* progranulin, *MAPT* microtubule-associated protein tau, *D-KEFS CWIT* Delis-Kaplan Executive Function System Colour-Word Interference Test, *FCSRT* Free and Cued Selective Reminding Test

^a Positive coefficients indicate better performance in mutation carriers compared to controls and were not included in the composite score




Table 3 Sample size per arm for a hypothetical clinical trial using different cognitive outcome measures

Outcome measures	<i>C9orf72</i>			<i>GRN</i>			<i>MAPT</i>		
	ES 10%	ES 20%	ES 40%	ES 10%	ES 20%	ES 40%	ES 10%	ES 20%	ES 40%
Cognitive composite scores									
Average composite	306	76	19	27	7	2	124	31	8
Weighted composite	214	53	13	53	13	3	90	23	6
Language									
Camel and Cactus Test	4946	1237	309	292	73	18	357	89	22
Boston Naming Test	1109	277	69	213	53	13	223	56	14
Category fluency	1584	396	99	781	195	49	400	100	25
Attention and mental processing speed									
Digit span forward	13,0210	32,553	8138	2677	669	167	17,773	4443	1111
Trail Making Test – part A	2272	568	142	45	11	3	69	17	4
Digit symbol	254	64	16	925	231	58	80	20	5
D-KEFS CWIT – colour naming	866	216	54	502	126	31	66	17	4
D-KEFS CWIT – word naming	19,224	4806	1202	3310	828	207	150	37	9
Executive functioning									
Digit span backward	1724	431	108	840	210	52	26,218	6555	1639
Trail Making Test – part B	1275	319	80	25	6	2	81	20	5
D-KEFS CWIT – ink naming	61	15	4	70	17	4	26	7	2
Phonemic fluency	558	139	35	2229	557	139	161	40	10
Visuoconstruction									
Benson figure copy	5911	1478	369	2119	530	132	6,282,036	1,570,509	392,627
Memory									
Benson figure recall	1044	261	65	657	164	41	7611	1903	476
FCSRT free recall	1302	326	81	294	74	18	521	130	33
FCSRT total recall	1020	255	64	477	119	30	524	131	33
FCSRT delayed free recall	606	152	38	767	192	48	261	65	16
FCSRT delayed total recall	358	89	22	193	48	12	681	170	43
Social cognition									
Facial Emotion Recognition Test	7570	1892	473	7805	1951	488	147	37	9

The sample size per arm was estimated as $n = (1 - \rho^2)(2\sigma^2)/\delta^2 f(\alpha, \beta)$, where ρ is the correlation between baseline and follow-up measures of the outcome, σ is the standard deviation of the outcome in the CDR[®] plus NACC-FTLD 0.5 group, δ is the treatment effect (effect size multiplied by the difference in mean between CDR[®] plus NACC-FTLD 0.5 and 1 group), α is the significance level (0.05), and $1 - \beta$ is the power to detect a treatment effect (80%)

Abbreviations: *C9orf72* chromosome 9 open reading frame 72, *GRN* progranulin, *MAPT* microtubule-associated protein tau, *ES* effect size as a proportion of the difference between the outcome in the CDR[®] plus NACC-FTLD 0.5 group and the outcome in the CDR[®] plus NACC-FTLD 1 group, *D-KEFS CWIT* Delis-Kaplan Executive Function System Colour-Word Interference Test, *FCSRT* Free and Cued Selective Reminding Test

average composite, whereas in *GRN* mutation carriers the average composite resulted in a lower sample size. For *GRN* (all $n < 60$) and *MAPT* (all $n < 125$) mutation carriers, lower sample sizes would be necessary to detect a treatment effect than for *C9orf72* repeat expansion carriers (all $n \leq 306$).

Time-to-event analysis

Kaplan-Meier curves can be seen in Fig. 1, and details on the sample included in the time-to-event analysis are reported in Additional file 1: Table S4. For *C9orf72* repeat expansion carriers, the probability of converting to a CDR[®] plus NACC FTLD of ≥ 1 increases from 6% after 2

years (SE = 0.06, 95% CI 0.01–0.39) to 53% after 3 years (SE = 0.33, 95% CI 0.12–0.99). In *GRN* mutation carriers, the probability of converting to a CDR[®] plus NACC FTLD of ≥ 1 increased from 4% after 1 year (SE = 0.04, 95% CI 0.01–0.24) to 43% after 3 years (SE = 0.14, 95% CI 0.22–0.72). In *MAPT* mutation carriers, the probability of converting to a global score of ≥ 1 increased from 10% after 1 year (SE = 0.10, 95% CI 0.01–0.49) to 42% during the second year (SE = 0.20, 95% CI 0.14–0.85). The Kaplan-Meier curve for *MAPT* mutation increased to 100% after 3 years in Fig. 1 because only one mutation carrier had follow-up up to this point and this individual progressed to a CDR[®] plus NACC FTLD of ≥ 1 . There

was no significant difference between the progression rates of different genetic groups ($\chi^2(2) = 1.18, p = 0.55$). In the total group of mutation carriers, the probability of converting to a CDR[®] plus NACC FTLD of ≥ 1 was 21% after 2 years (SE = 0.03, 95% CI 0.11–0.40) and 52% after 3 years (SE = 0.16, 95% CI 0.26–0.83). This means that for a 3-year trial where drug treatment is assumed to have a 20% effect (i.e. only 80% of the treated group will experience the change seen between CDR[®] plus NACC FTLD 0.5 and 1 groups), the sample size corresponding to a 10% effect in Table 3 needs to be included in order to demonstrate a treatment effect, because only ~ 50% of mutation carriers would be expected to progress from CDR[®] plus NACC FTLD 0.5 to 1 without treatment (i.e. effect size needs to be divided by 2).

Discussion

We have empirically developed gene-specific cognitive composite scores in *MAPT*, *GRN*, and *C9orf72* mutation carriers (GENFI-Cog) and demonstrated that they provide feasible sample sizes for clinical trials to evaluate the effect of treatment on clinical progression from the prodromal to the fully symptomatic stage. Time-to-event analyses revealed that roughly 50% of the patients with a CDR[®] plus NACC FTLD of 0.5 progress to 1 or higher after a period of 3 years. The results from this study show that GENFI-Cog has potential as a cognitive endpoint in upcoming clinical trials and provide important guidelines on sample size recruitment and clinical trial duration.

The GENFI-Cog composites can be regarded as attractive clinical outcome measures because they produce substantially lower sample size estimates than most individual neuropsychological tests. Depending on the effect size (40% to 10%), sample size estimates ranged between 13 and 214 for *C9orf72*, 3 and 53 for *GRN*, and 6 and 90 for *MAPT* per study arm for the weighted GENFI-Cog. A practical problem in trial design for FTD spectrum disorders is recruiting enough patients to test candidate therapeutics as FTD is much less common than AD, with an estimated prevalence of 15/100,000 and approximately 10–20% of cases being caused by mutations in *C9orf72*, *GRN*, and *MAPT* genes [4, 7, 42]. It is therefore unlikely that a trial would be able to include many hundreds of patients per study arm, which our results showed would be necessary for most individual neuropsychological tests. There were some individual neuropsychological tests that required reasonable sample sizes similar to that of GENFI-Cog, e.g. TMT and D-KEFS CWIT. These tests are typically included in clinical trials such as the current AL001 study of *GRN*-related FTD [7]. Yet, due to the heterogeneity in cognitive symptoms between patients even with the same genetic mutation, individually examining each cognitive test might not provide a sensitive and

clinically meaningful primary outcome measure. Using GENFI-Cog will allow a single cognitive outcome to be used when analysing treatment effect, although validation in other large cohorts is warranted.

The CDR[®] plus NACC FTLD is currently often used as an inclusion criterion for clinical trials as well as for tracking disease progression. The results showed that roughly 50% of the patients with a CDR[®] plus NACC FTLD 0.5 progress to 1 or higher after a period of 3 years. This indicates that for trials with a duration of 3 years, around 50% of patients with CDR[®] plus NACC FTLD of 0.5 on entry to the trial would be expected to progress to CDR[®] plus NACC FTLD of 1 in the absence of effective disease-modifying treatment. This means that if a treatment is expected to have a 20% effect, the sample size corresponding to a 10% effect needs to be included per study arm to be able to demonstrate a treatment effect, because only half of the mutation carriers would be expected to progress from CDR[®] plus NACC FTLD 0.5 to 1 without treatment. This is important to consider when planning trial duration and recruitment with the currently available clinical measures.

The optimal gene-specific cognitive composite score incorporated tests from different cognitive domains. For *GRN* mutation carriers, tests for executive function and social cognition contributed the most to the composite score, with the addition of tests for memory and language. In *MAPT* mutation carriers, there was a strong focus on semantic and episodic memory tests in the composite score with the addition of tests for attention and mental processing speed. A combination of tests from all cognitive domains was most sensitive in *C9orf72* mutation carriers, with the strongest contribution from tests within the domains of executive function, social cognition, and memory. These results complement recent studies showing cognitive decline in the early stages of FTD with widespread cognitive impairment covering multiple domains in *C9orf72* [22, 43], dysexecutive functioning as the key feature in *GRN* [13, 22] and a specific impairment in episodic and semantic memory in *MAPT*-associated FTD [13, 20, 22]. Impairment of social cognition appears to be a key feature in all three genetic groups [38], which was probably due to the high number of bvFTD cases in the sample. Neuroimaging studies have indeed shown that the neurodegenerative process in *C9orf72* mutation carriers typically is reflected by widespread degeneration in frontal, temporal, and cerebellar and subcortical structures [43], whereas focal atrophy of the anteromedial temporal lobe, an area important for memory and semantic functioning, is often seen in *MAPT*-associated FTD [44]. In *GRN* mutation carriers, the typical pattern of degeneration includes the inferior frontal regions as well as the cingulate cortex, areas known to be critical

in executive function [44]. Thus, although the GENFI-Cog was empirically derived, the selected tests are clinically meaningful and in line with a theoretically driven approach where the composite would be constructed a priori from cognitive tests that are known to decline in the early stages of each genetic group.

This is to our knowledge the first study that has created cognitive composites for genetic forms of FTD by selecting the most sensitive combinations of cognitive variables based on systematic comparisons with controls. A major strength of this study is the use of a large cohort of genetic FTD mutation carriers allowing gene-specific analyses, but also the use of a matched control group of mutation-negative family members. Another strength is the use of LASSO with cross-validation to avoid overfitting bias to ensure that results have generalizability [41].

Limitations

There are some limitations to the present study, however. The results from the logistic regression analysis revealed two neuropsychological tests in *GRN* mutation carriers with a positive coefficient, indicating better performance compared to controls, and were excluded from the composite scores. Development of GENFI-Cog was constrained by the neuropsychological test battery that is used in the GENFI cohort [14], which made validation in an independent sample not possible and limited the generalizability of the findings. Validation in other cohorts (such as ALLFTD [45] or DINAD) is therefore recommended. Although the LASSO model with 10-fold cross-validation included an internal cross-validation step to select the penalization term for the selection of the cognitive tests, the findings were not externally validated in an independent sample thereby limiting the generalizability of GENFI-Cog. Future collaborations within the FTD Prevention Initiative (FPI) could be a starting point to cross-validate our findings. The sample size estimates serve as a guide on the sensitivity and power of GENFI-Cog compared to individual cognitive tests and should be interpreted with caution as they were calculated from the cross-sectional difference between a small number of patients with CDR[®] plus NACC FTLD 0.5 and 1, assuming that the difference between these groups is representative of the change over time that would be seen in longitudinal scores in a clinical trial as patients progress from a score of 0.5 to 1, i.e. prodromal to fully symptomatic. Future research using longitudinal data and larger sample sizes is necessary to examine the validity of this assumption and to examine if the cognitive composites presented in the current study are similar to those derived using longitudinal change in scores. Importantly, it is essential for future clinical trials of FTD to also include other biomarkers such as neuroimaging,

neurofilament light chain, or other fluid protein levels as endpoints. As such, it would be interesting to include such biomarkers in addition to GENFI-Cog within a future longitudinal multimodal analysis. Lastly, as GENFI is a prospective cohort study with ongoing recruitment, not all participants completed the same number of visits contributing to low sample sizes at later visits in the time-to-event analysis. The time-to-event analysis was performed to provide insight on the possible duration required for a clinical trial, but validation with larger sample sizes where all participants have completed the same number of visits is warranted.

Conclusions

In summary, we examined the cognitive data from the GENFI cohort and conducted a search for the combination of cognitive assessments most sensitive to differentiate *MAPT*, *GRN*, and *C9orf72* mutation carriers from non-carriers. As a result, we created three gene-specific cognitive composite scores, GENFI-Cog, that were sensitive to track progression on the clinical progression of the CDR[®] plus NACC FTLD 0.5 to 1 stage as it resulted in smaller sample sizes than most individual neuropsychological tests. To conclude, GENFI-Cog has the potential to be a primary cognitive outcome measure in upcoming clinical trials for *C9orf72*, *GRN*, and *MAPT* mutation carriers.

Abbreviations

AD: Alzheimer's disease; ADAS-Cog: Alzheimer's Disease Assessment Scale Cognitive Subscale; ALLFTD: ARTFL-LEFFTDS Longitudinal Frontotemporal Lobar Degeneration Study; BNT: Boston Naming Test; bvFTD: Behavioural variant frontotemporal dementia; C9orf72: Chromosome 9 open reading frame 72; CDR plus NACC-FTLD: Clinical Dementia Rating scale plus National Alzheimer's Coordinating Center – Frontotemporal Lobar Degeneration; CWIT: Colour Word Interference Test; DINAD: Dominantly inherited non-Alzheimer dementias; D-KEFS: Delis-Kaplan Execution Function System; FCSRT: Free and Cued Selective Reminding Test; FTD: Frontotemporal dementia; FTD-ALS: Frontotemporal dementia with amyotrophic lateral sclerosis; GENFI: Genetic FTD Initiative; GRN: Progranulin; HD: Huntington's disease; LASSO: Least absolute shrinkage and selection operator; MAPT: Microtubule-associated protein tau; MMSE: Mini-Mental State Examination; PD: Parkinson's disease; PPA: Primary progressive aphasia; TMT: Trail Making Test; UPDRS: Unified Parkinson's Disease Rating Scale; UHDRS: Unified Huntington's Disease Rating Scale; WAIS-R: Wechsler Adult Intelligence Scale - Revised; WMS-R: Wechsler Memory Scale - Revised.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13195-022-00958-0>.

Additional file 1: Table S1. Number of control data available in each language per cognitive test. **Table S2.** Parameters included in the sample size calculations. **Table S3.** Participants characteristics and neuropsychological test results per CDR[®] plus NACC FTLD global score. **Table S4.** Number of mutation carriers that progressed on the CDR[®] plus NACC FTLD. **Figure S1.** STROBE flowchart.

Acknowledgements

We thank the research participants and their families for their contribution to the study. Several authors of this publication are members of the European Reference Network for Rare Neurological Diseases - Project ID No 739510. Group authorship for the Genetic FTD Initiative:

Arabella Bouzigues MSc¹, Martin N. Rossor MD FRCP¹, Nick C. Fox MD FRCP¹, Jason D. Warren PhD FRACP¹, Martina Bocchetta PhD¹, Imogen J. Swift MSc¹, Rachelle Shafei MRCP¹, Carolin Heller BSc¹, Emily Todd MSc¹, David Cash PhD¹, Ione Woollacott PhD¹, Henrik Zetterberg¹, Annabel Nelson BSc¹, Rita Guerreiro PhD², Jose Bras PhD², David L. Thomas PhD³, Simon Mead PhD⁴, Lieke Meeter MD⁵, Jessica Panman MSc⁵, Rick van Minkelen PhD⁵, Myriam Barandiaran PhD^{7,8}, Begoña Indakoetxea MD^{7,8}, Alazne Gabilondo MD⁸, Mikel Tainta MD⁸, Ana Gorostidi PhD⁸, Miren Zulaica BSc⁸, Alina Díez MSc⁸, Jorge Villanua MD PhD⁹, Sergi Borrego-Ecija MD¹⁰, Jaume Olives MSc¹⁰, Albert Lladó PhD¹⁰, Mircea Balasa PhD¹⁰, Anna Antonell PhD¹⁰, Nuria Bargallo PhD¹¹, Enrico Premi MD¹², Stefano Gazzina MD¹³, Roberto Gasparotti MD¹⁴, Silvana Archetti MBiolSci¹⁵, Sandra Black MD¹⁶, Sara Mitchell MD¹⁶, Ekaterina Rogaeva PhD¹⁷, Morris Freedman MD¹⁸, Ron Keren MD¹⁹, David Tang-Wai MD²⁰, Hakan Thonberg MD²¹, Linn Ojterstedt MD^{21,22}, Christin Andersson PhD²³, Vesna Jelic MD²⁴, Andrea Arighi MD^{25,26}, Chiara Fenoglio PhD^{25,26}, Elio Scarpini MD^{25,26}, Giorgio Fumagalli MD^{25,26}, Thomas Cope MRCP²⁷, Carolyn Timberlake BSc²⁷, Timothy Rittman MRCP²⁷, Christen Shoesmith MD²⁸, Robart Bartha PhD^{29,30}, Rosa Rademakers PhD³¹, Carlo Wilke MD^{32,33}, Hans-Otto Karnath MD³⁴, Benjamin Bender MD³⁵, Rose Bruffaerts MD PhD³⁶, Philip Vandamme MD PhD³⁷, Mathieu Vandenbulcke MD PhD^{38,39}, Catarina B. Ferreira MSc⁴⁰, Gabriel Miltenberger PhD⁴¹, Carolina Maruta MPsych PhD⁴², Ana Verdelho MD PhD⁴³, Sónia Afonso BSc⁴⁴, Ricardo Taipa MD PhD⁴⁵, Paola Caropponi MD PhD⁴⁶, Giuseppe Di Fede MD PhD⁴⁶, Giorgio Giaccone MD⁴⁶, Sara Prioni PsyD⁴⁶, Veronica Redaelli MD⁴⁶, Giacomina Rossi MSc⁴⁶, Diana Duro NPsych⁴⁷, Maria Rosario Almeida PhD⁴⁷, Miguel Castelo-Branco MD PhD⁴⁷, Maria João Leitão BSc⁴⁸, Miguel Tabuas-Pereira MD⁴⁹, Beatriz Santiago MD⁴⁹, Serge Gauthier MD⁵⁰, Pedro Rosa-Neto MD PhD⁵¹, Michele Veldsman PhD⁵², Paul Thompson PhD⁵³, Tobias Langheinrich MD⁵³, Catharina PRIX MD⁵⁴, Tobias Hoegen MD⁵⁴, Elisabeth Wlasich Mag. rer. nat.⁵⁴, Sandra Loosli MD⁵⁴, Sonja Schonecker MD⁵⁴, Sarah Anderl-Straub Dr.hum.biol Dipl.Psych⁵⁵, Jolina Lombardi⁵⁵, Nuria Bargallo MD PhD⁵⁶, Alberto Benussi MD⁵⁷, Valentina Cantoni⁵⁷, Maxime Bertoux PhD^{58,59}, Anne Bertrand MD PhD⁶⁰, Alexis Brice MD PhD⁶⁰, Agnès Camuzat⁶⁰, Olivier Colliot PhD⁶⁰, Sabrina Sayah⁶⁰, Aurélie Funkiewiez^{60,61}, Daisy Rinaldi^{60,61}, Gemma Lombardi⁶¹, Benedetta Nacmias⁶¹, Dario Saracino^{60,61,62}, Valentina Bessi⁶³, Camilla Ferrari⁶³, Marta Cañada⁶⁴, Vincent Deramecourt⁶⁵, Gregory Kuchcinski⁶⁵, Thibaud Lebouvier⁶⁵, Sebastien Ourselin⁶⁶, Cristina Polito⁶⁷, and Adeline Rollin⁶⁸

¹Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK; ²Center for Neurodegenerative Science, Van Andel Institute, Grand Rapids, Michigan, MI 49503, USA; ³Division of Neuroscience and Experimental Psychology, Wolfson Molecular Imaging Centre, University of Manchester, Manchester, UK; ⁴MRC Prion Unit, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK; ⁵Department of Neurology, Erasmus Medical Center, Rotterdam, The Netherlands; ⁶Department of Clinical Genetics, Erasmus Medical Center, Rotterdam, The Netherlands; ⁷Cognitive Disorders Unit, Department of Neurology, Donostia University Hospital, San Sebastian, Gipuzkoa, Spain; ⁸Neuroscience Area, Biodonostia Health Research Institute, San Sebastian, Gipuzkoa, Spain; ⁹OSATEK, University of Donostia, San Sebastian, Gipuzkoa, Spain; ¹⁰Alzheimer's Disease and Other Cognitive Disorders Unit, Neurology Service, Hospital Clinic, Barcelona, Spain; ¹¹Imaging Diagnostic Center, Hospital Clinic, Barcelona, Spain; ¹²Stroke Unit, ASST Brescia Hospital, Brescia, Italy; ¹³Neurology, ASST Brescia Hospital, Brescia, Italy; ¹⁴Neuroradiology Unit, University of Brescia, Brescia, Italy; ¹⁵Biotechnology Laboratory, Department of Diagnostics, ASST Brescia Hospital, Brescia, Italy; ¹⁶Sunnybrook Health Sciences Centre, Sunnybrook Research Institute, University of Toronto, Toronto, Canada; ¹⁷Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto, Canada; ¹⁸Baycrest Health Sciences, Rotman Research Institute, University of Toronto, Toronto, Canada; ¹⁹The University Health Network, Toronto Rehabilitation Institute, Toronto, Canada; ²⁰The University Health Network, Krembil Research Institute, Toronto, Canada; ²¹Center for Alzheimer Research, Division of Neurogeriatrics, Department of Neurobiology, Care Sciences and Society, Bioclinicum, Karolinska Institutet, Solna, Sweden; ²²Unit for Hereditary Dementias, Theme Aging, Karolinska University Hospital, Solna, Sweden; ²³Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden; ²⁴Division of Clinical Geriatrics, Karolinska Institutet, Stockholm, Sweden; ²⁵Fondazione IRCCS Ca' Granda

Ospedale Maggiore Policlinico, Neurodegenerative Diseases Unit, Milan, Italy; ²⁶University of Milan, Centro Dino Ferrari, Milan, Italy; ²⁷Department of Clinical Neuroscience, University of Cambridge, Cambridge, UK; ²⁸Department of Clinical Neurological Sciences, University of Western Ontario, London, Ontario, Canada; ²⁹Department of Medical Biophysics, The University of Western Ontario, London, Ontario, Canada; ³⁰Centre for Functional and Metabolic Mapping, Roberts Research Institute, The University of Western Ontario, London, Ontario, Canada; ³¹Department of Neurosciences, Mayo Clinic, Jacksonville, FL, USA; ³²Department of Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research and Center of Neurology, University of Tübingen, Tübingen, Germany; ³³Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany; ³⁴Division of Neuropsychology, Hertie-Institute for Clinical Brain Research and Center of Neurology, University of Tübingen, Tübingen, Germany; ³⁵Department of Diagnostic and Interventional Neuroradiology, University of Tübingen, Tübingen, Germany; ³⁶Laboratory for Cognitive Neurology, Department of Neurosciences, KU Leuven, Leuven, Belgium; ³⁷Neurology Service, University Hospitals Leuven, Belgium; Laboratory for Neurobiology, VIB-KU Leuven Centre for Brain Research, Leuven, Belgium; ³⁸Geriatric Psychiatry Service, University Hospitals Leuven, Leuven, Belgium; ³⁹Neuropsychiatry, Department of Neurosciences, KU Leuven, Leuven, Belgium; ⁴⁰Laboratory of Neurosciences, Institute of Molecular Medicine, Faculty of Medicine, University of Lisbon, Lisbon, Portugal; ⁴¹Faculty of Medicine, University of Lisbon, Lisbon, Portugal; ⁴²Laboratory of Language Research, Centro de Estudos Egas Moniz, Faculty of Medicine, University of Lisbon, Lisbon, Portugal; ⁴³Department of Neurosciences and Mental Health, Centro Hospitalar Lisboa Norte - Hospital de Santa Maria & Faculty of Medicine, University of Lisbon, Lisbon, Portugal; ⁴⁴Instituto Ciências Nucleares Aplicadas a Saude, Universidade de Coimbra, Coimbra, Portugal; ⁴⁵Neuropathology Unit and Department of Neurology, Centro Hospitalar do Porto - Hospital de Santo António, Oporto, Portugal; ⁴⁶Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy; ⁴⁷Faculty of Medicine, University of Coimbra, Coimbra, Portugal; ⁴⁸Centre of Neurosciences and Cell Biology, Universidade de Coimbra, Coimbra, Portugal; ⁴⁹Neurology Department, Centro Hospitalar e Universitario de Coimbra, Coimbra, Portugal; ⁵⁰Alzheimer Disease Research Unit, McGill Centre for Studies in Aging, Department of Neurology & Neurosurgery, McGill University, Montreal, Québec, Canada; ⁵¹Translational Neuroimaging Laboratory, McGill Centre for Studies in Aging, McGill University, Montreal, Québec, Canada; ⁵²Nuffield Department of Clinical Neurosciences, Medical Sciences Division, University of Oxford, Oxford, UK; ⁵³Division of Neuroscience and Experimental Psychology, Wolfson Molecular Imaging Centre, University of Manchester, Manchester, UK; ⁵⁴Neurologische Klinik, Ludwig-Maximilians-Universität München, Munich, Germany; ⁵⁵Department of Neurology, University of Ulm, Ulm, Germany; ⁵⁶Imaging Diagnostic Center, Hospital Clinic, Barcelona, Spain; ⁵⁷Centre for Neurodegenerative Disorders, Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy; ⁵⁸Inserm 1172, Lille, France; ⁵⁹CHU, CNR-MAJ, Labex Distalz, LiCEND, Lille, France; ⁶⁰Sorbonne Université, Paris Brain Institute – Institut du Cerveau – ICM, Inserm U1127, CNRS UMR 7225, AP-HP - Hôpital Pitié-Salpêtrière, Paris, France; ⁶¹Centre de référence des démences rares ou précoces, IM2A, Département de Neurologie, AP-HP - Hôpital Pitié-Salpêtrière, Paris, France; ⁶²Inria, Aramis project-team, F-75013, Paris, France ⁶³Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Florence, Italy; ⁶⁴CITA Alzheimer, San Sebastian, Gipuzkoa, Spain; ⁶⁵University of Lille, Lille, France; ⁶⁶School of Biomedical Engineering & Imaging Sciences, King's College London, London, UK; ⁶⁷Department of Biomedical, Experimental and Clinical Sciences "Mario Serio", Nuclear Medicine Unit, University of Florence, Florence, Italy; ⁶⁸CHU, CNR-MAJ, Labex Distalz, LiCEND, Lille, France

Authors' contributions

JMP, KMM, JN, and JDR contributed to the conception and design of the work and the analysis of the data. JMP drafted the original work. All authors contributed to the acquisition and interpretation of the data and revised the work. All authors read and approved the final manuscript.

Funding

The Dementia Research Centre is supported by Alzheimer's Research UK, Alzheimer's Society, Brain Research UK, and The Wolfson Foundation. This work was supported by the NIHR UCL/H Biomedical Research Centre, the Leonard Wolfson Experimental Neurology Centre (LWENC) Clinical Research Facility, and the UK Dementia Research Institute, which receives its funding from UK DRI Ltd., funded by the UK Medical Research Council, Alzheimer's Society

and Alzheimer's Research UK. JDR is supported by an MRC Clinician Scientist Fellowship (MR/M008525/1) and has received funding from the NIHR Rare Disease Translational Research Collaboration (BRC149/NS/MH). This work was also supported by the MRC UK GENFI grant (MR/M023664/1), the Bluefield Project, the JPND GENFI-PROX grant (2019-02248), the Dioraphte Foundation [grant numbers 09-02-00], the Association for Frontotemporal Dementias Research Grant 2009, The Netherlands Organization for Scientific Research (NWO) (grant HCM1 056-13-018), ZonMw Memorabel (Deltaplan Dementie, (project numbers 733 050 103 and 733 050 813), and JPND PreFrontAls Consortium (project number 733051042). JM Poos is supported by a fellowship award from Alzheimer Nederland (WE.15-2019.02). This work was conducted using the MRC Dementias Platform UK (MR/L023784/1 and MR/009076/1).

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request

Declarations

Ethics approval and consent to participate

All GENFI sites had local ethical approval for the study, and all participants gave written informed consent.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Neurology, Erasmus MC University Medical Center, Rotterdam, The Netherlands. ²Dementia Research Centre, Department of Neurodegenerative Disease, National Hospital for Neurology and Neurosurgery, UCL Institute of Neurology, 8-11 Queen Square, Box 16, London WC1N 3BG, UK. ³Department of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK. ⁴Department of Neurology, Alzheimer Center, Amsterdam University Medical Center, Amsterdam Neuroscience, Amsterdam, The Netherlands. ⁵Cognitive Disorders Unit, Department of Neurology, Donostia University Hospital, San Sebastian, Gipuzkoa, Spain. ⁶Alzheimer's disease and Other Cognitive Disorders Unit, Neurology Service, Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi I Sunyer, University of Barcelona, Barcelona, Spain. ⁷Centre for Neurodegenerative Disorders, Neurology Unit, Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy. ⁸Clinique Interdisciplinaire de Mémoire, Département des Sciences Neurologiques, Université Laval, Québec, Canada. ⁹Sunnybrook Health Sciences Centre, Sunnybrook Research Institute, University of Toronto, Toronto, Canada. ¹⁰Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto, Canada. ¹¹Department of Geriatric Medicine, Karolinska University Hospital-Huddinge, Stockholm, Sweden. ¹²University of Milan, Centro Dino Ferrari, Milan, Italy. ¹³Neurodegenerative Diseases Unit, Fondazione IRCCS Ca'Granda, Ospedale Policlinico, Milan, Italy. ¹⁴Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK. ¹⁵Department of Clinical Neurological Sciences, University of Western Ontario, London, Ontario, Canada. ¹⁶Department of Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research and Center of Neurology, University of Tübingen, Tübingen, Germany. ¹⁷German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany. ¹⁸Laboratory for Cognitive Neurology, Department of Neurosciences, KU Leuven, Leuven, Belgium. ¹⁹Faculty of Medicine, University of Lisbon, Lisbon, Portugal. ²⁰Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan, Italy. ²¹Faculty of Medicine, University of Coimbra, Coimbra, Portugal. ²²Department of Psychiatry, McGill University Health Centre, McGill University, Montreal, Québec, Canada. ²³Department of Clinical Neurology, University of Oxford, Oxford, UK. ²⁴Faculty of Medical and Human Sciences, Institute of Brain, Behaviour and Mental Health, University of Manchester, Manchester, UK. ²⁵Department of Neurology, Ludwig-Maximilians-University, Munich, Germany. ²⁶German Center for Neurodegenerative Diseases (DZNE), Munich, Germany. ²⁷Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ²⁸Department of Neurology, University of Ulm, Ulm, Germany. ²⁹Paris Brain Institute – Institut du Cerveau – ICM, Inserm U1127, CNRS UMR 7225,

AP-HP - Hôpital Pitié-Salpêtrière, Sorbonne Université, Paris, France. ³⁰Centre de référence des démences rares ou précoces, IM2A, Département de Neurologie, AP-HP - Hôpital Pitié-Salpêtrière, Paris, France. ³¹Département de Neurologie, AP-HP - Hôpital Pitié-Salpêtrière, Paris, France. ³²University of Lille, Lille, France. ³³Inserm 1172, Lille, France. ³⁴CHU, CNR-MAJ, Labex Distalz, LICEND, Lille, France.

Received: 7 September 2021 Accepted: 28 December 2021

Published online: 19 January 2022

References

- Seelaar H, Rohrer JD, Pijnenburg YA, Fox NC, van Swieten JC. Clinical, genetic and pathological heterogeneity of frontotemporal dementia: a review. *J Neurol Neurosurg Psychiatry*. 2011;82(5):476–86.
- Rascovsky K, Hodges JR, Knopman D, Mendez MF, Kramer JH, Neuhaus J, et al. Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain*. 2011;134(Pt 9):2456–77.
- Gorno-Tempini ML, Hillis AE, Weintraub S, Kertesz A, Mendez M, Cappa SF, et al. Classification of primary progressive aphasia and its variants. *Neurology*. 2011;76(11):1006–14.
- Lashley T, Rohrer JD, Mead S, Revesz T. An update on clinical, genetic and pathological aspects of frontotemporal lobar degenerations. *Neuro-pathol Appl Neurobiol*. 2015;41(7):858–81.
- Desmarais P, Rohrer JD, Nguyen QD, Herrmann N, Stuss DT, Lang AE, et al. Therapeutic trial design for frontotemporal dementia and related disorders. *J Neurol Neurosurg Psychiatry*. 2019;90(4):412–23.
- Tsai RM, Boxer AL. Therapy and clinical trials in frontotemporal dementia: past, present, and future. *J Neurochem*. 2016;138:211–21.
- Panza F, Lozupone M, Seripa D, Daniele A, Watling M, Giannelli G, et al. Development of disease-modifying drugs for frontotemporal dementia spectrum disorders. *Nat Rev Neurol*. 2020;16(4):213–28.
- Langbaum JB, Hendrix SB, Ayutyanont N, Chen K, Fleisher AS, Shah RC, et al. An empirically derived composite cognitive test score with improved power to track and evaluate treatments for preclinical Alzheimer's disease. *Alzheimers Dement*. 2014;10(6):666–74.
- Silverberg NB, Ryan LM, Carrillo MC, Sperling R, Petersen RC, Posner HB, et al. Assessment of cognition in early dementia. *Alzheimers Dement*. 2011;7(3):e60–76.
- Cano SJ, Posner HB, Moline ML, Hurt SW, Swartz J, Hsu T, et al. The ADAS-cog in Alzheimer's disease clinical trials: psychometric evaluation of the sum and its parts. *J Neurol Neurosurg Psychiatry*. 2010;81(12):1363–8.
- Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. *Am J Psychiatry*. 1984;141(11):1356–64.
- Jiskoot LC, Dopfer EG, Heijer T, Timman R, van Minkelen R, van Swieten JC, et al. Presymptomatic cognitive decline in familial frontotemporal dementia: a longitudinal study. *Neurology*. 2016;87(4):384–91.
- Jiskoot LC, Panman JL, Meeter LH, Dopfer EGP, Donker Kaat L, Franzen S, et al. Longitudinal multimodal MRI as prognostic and diagnostic biomarker in presymptomatic familial frontotemporal dementia. *Brain*. 2018;142(1):193–208.
- Rohrer JD, Nicholas JM, Cash DM, van Swieten J, Dopfer E, Jiskoot L, et al. Presymptomatic cognitive and neuroanatomical changes in genetic frontotemporal dementia in the Genetic Frontotemporal dementia Initiative (GENFI) study: a cross-sectional analysis. *Lancet Neurol*. 2015;14(3):253–62.
- Barandíaran M, Estanga A, Moreno F, Indakoetxea B, Alzuale A, Balluerka N, et al. Neuropsychological features of asymptomatic c. 709-1G> A progranulin mutation carriers. *J Int Neuropsychol Soc*. 2012;18(6):1086–90.
- Bertrand A, Wen J, Rinaldi D, Houot M, Sayah S, Camuzat A, et al. Early cognitive, structural, and microstructural changes in presymptomatic C9orf72 carriers younger than 40 years. *JAMA Neurol*. 2018;75(2):236–45.
- Floeter MK, Traynor BJ, Farren J, Braun LE, Tierney M, Wiggs EA, et al. Disease progression in C9orf72 mutation carriers. *Neurology*. 2017;89(3):234–41.
- Hallam BJ, Jacova C, Hsiung G-YR, Wittenberg D, Sengdy P, Bouchard-Kerr P, et al. Early neuropsychological characteristics of progranulin mutation carriers. *Journal of the International Neuropsychological Society*. 2014;20(7):694.

19. Staffaroni AM, Bajorek L, Casaletto KB, Cobigo Y, Goh SYM, Wolf A, et al. Assessment of executive function declines in presymptomatic and mildly symptomatic familial frontotemporal dementia: NIH-EXAMINER as a potential clinical trial endpoint. *Alzheimers Dement*. 2020;16(1):11–21.
20. Poos JM, Russell LL, Peakman G, Bocchetta M, Greaves CV, Jiskoot LC, et al. Impairment of episodic memory in genetic frontotemporal dementia: a GENFI study. *Alzheimer's Dement Diagn Assess Dis Monit*. 2021;13(1):e12185.
21. Barandiaran M, Moreno F, de Arriba M, Indakoetxea B, Boda I, Gabilondo A, et al. Longitudinal neuropsychological study of presymptomatic c. 709-1G> A progranulin mutation carriers. *J Int Neuropsychol Soc*. 2019;25(1):39–47.
22. Poos JM, Jiskoot LC, Leijdesdorff SMJ, Seelaar H, Panman JL, van der Ende EL, et al. Cognitive profiles discriminate between genetic variants of behavioral frontotemporal dementia. *J Neurol*. 2020;267(6):1603–12.
23. Le Ber I, Camuzat A, Hannequin D, Pasquier F, Guedj E, Rovelet-Lecrux A, et al. Phenotypic variability in progranulin mutation carriers: a clinical, neuropsychological, imaging and genetic study. *Brain*. 2008;131(3):732–46.
24. Cheran G, Wu L, Lee S, Manoochehri M, Cines S, Fallon E, et al. Cognitive indicators of preclinical behavioral variant frontotemporal dementia in MAPT carriers. *J Int Neuropsychol Soc*. 2019;25(2):184–94.
25. Geschwind DH, Robidoux J, Alarcón M, Miller BL, Wilhelmsen KC, Cummings JL, et al. Dementia and neurodevelopmental predisposition: cognitive dysfunction in presymptomatic subjects precedes dementia by decades in frontotemporal dementia. *Ann Neurol*. 2001;50(6):741–6.
26. Pappa JM, Jiskoot LC, Panman JL, Dopfer EG, Den Heijer T, Kaat LD, et al. Cognition and gray and white matter characteristics of presymptomatic C9orf72 repeat expansion. *Neurology*. 2017;89(12):1256–64.
27. Jiskoot LC, Panman JL, van Asseldonk L, Franzen S, Meeter LHH, Kaat LD, et al. Longitudinal cognitive biomarkers predicting symptom onset in presymptomatic frontotemporal dementia. *J Neurol*. 2018;265(6):1381–92.
28. Tsou E, Ernhoff SJ, Goode CA, Dorsman KA, Kanjanapong S, Lindbergh CA, et al. BHA-CS: a novel cognitive composite for Alzheimer's disease and related disorders. *Alzheimer's Dement Diagn Assess Dis Monit*. 2020;12(1):e12042.
29. Jones R, Stout JC, Labuschagne I, Say M, Justo D, Coleman A, et al. The potential of composite cognitive scores for tracking progression in Huntington's disease. *J Huntington's Dis*. 2014;3(2):197–207.
30. Fahn S. Unified Parkinson's disease rating scale. Recent development in Parkinson's disease; 1987.
31. Miyagawa T, Brushaber D, Syrjanen J, Kremers W, Fields J, Forsberg LK, et al. Utility of the global CDR® plus NACC FTD rating and development of scoring rules: data from the ARTFL/LEFFTDS Consortium. *Alzheimers Dement*. 2020;16(1):106–17.
32. Brooks BR, Miller RG, Swash M, Munsat TL. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Later Scleros Motor Neuron Disord*. 2000;1(5):293–9.
33. Moore K, Convery R, Bocchetta M, Neason M, Cash DM, Greaves C, et al. A modified Camel and Cactus Test detects presymptomatic semantic impairment in genetic frontotemporal dementia within the GENFI cohort. *Appl Neuropsychol Adult*. 2020:1–8. Ahead of print
34. Morris JC, Weintraub S, Chui HC, Cummings J, DeCarli C, Ferris S, et al. The Uniform Data Set (UDS): clinical and cognitive variables and descriptive data from Alzheimer disease centers. *Alzheimer Dis Assoc Disord*. 2006;20(4):210–6.
35. Tombaugh TN, Kozak J, Rees L. Normative data stratified by age and education for two measures of verbal fluency: FAS and animal naming. *Arch Clin Neuropsychol*. 1999;14(2):167–77.
36. Corrigan JD, Hinkley NS. Relationships between parts A and B of the Trail Making Test. *J Clin Psychol*. 1987;43(4):402–9.
37. Delis DC, Kaplan E, Kramer J, den Buysch HO, Noens ILJ, Berckelaer-Onnes IA. D-KEFS: Delis-Kaplan executive function system: color-word interference test: handleiding. Amsterdam: Pearson; 2008.
38. Russell LL, Greaves CV, Bocchetta M, Nicholas J, Convery RS, Moore K, et al. Social cognition impairment in genetic frontotemporal dementia within the GENFI cohort. *Cortex*. 2020.
39. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189–98.
40. Kukreja SL, Löfberg J, Brenner MJ. A least absolute shrinkage and selection operator (LASSO) for nonlinear system identification. *IFAC Proc*. 2006;39(1):814–9.
41. Friedman LM, Furberg C, DeMets DL. Fundamentals of clinical trials. 5th ed. New York: Springer; 2015. Print
42. Coyle-Gilchrist ITS, Dick KM, Patterson K, Rodríguez PV, Wehmann E, Wilcox A, et al. Prevalence, characteristics, and survival of frontotemporal lobar degeneration syndromes. *Neurology*. 2016;86(18):1736–43.
43. Mahoney CJ, Downey LE, Ridgway GR, Beck J, Clegg S, Blair M, et al. Longitudinal neuroimaging and neuropsychological profiles of frontotemporal dementia with C9ORF72 expansions. *Alzheimers Res Ther*. 2012;4(5):41.
44. Rohrer JD, Warren JD. Phenotypic signatures of genetic frontotemporal dementia. *Curr Opin Neurol*. 2011;24(6):542–9.
45. Boeve B, Bove J, Brannely P, Brushaber D, Coppola G, Dever R, et al. The longitudinal evaluation of familial frontotemporal dementia subjects protocol: framework and methodology. *Alzheimers Dement*. 2019.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

