

Interactions between climate and COVID-19

James D Ford, Carol Zavaleta-Cortijo, Triphini Ainembabazi, Cecilia Anza-Ramirez, Ingrid Arotoma-Rojas, Joana Bezerra, Victoria Chicmana-Zapata, Eranga K Galappaththi, Martha Hangula, Christopher Kazaana, Shuaib Lwasa, Didacus Namanya, Nosipho Nkwinti, Richard Nuwagira, Samuel Okware, Maria Osipova, Kerrie Pickering, Chandni Singh, Lea Berrang-Ford, Keith Hyams, J Jaime Miranda, Angus Naylor, Mark New, Bianca van Bavel, in collaboration with the COVID Observatory



In this Personal View, we explain the ways that climatic risks affect the transmission, perception, response, and lived experience of COVID-19. First, temperature, wind, and humidity influence the transmission of COVID-19 in ways not fully understood, although non-climatic factors appear more important than climatic factors in explaining disease transmission. Second, climatic extremes coinciding with COVID-19 have affected disease exposure, increased susceptibility of people to COVID-19, compromised emergency responses, and reduced health system resilience to multiple stresses. Third, long-term climate change and prepandemic vulnerabilities have increased COVID-19 risk for some populations (eg, marginalised communities). The ways climate and COVID-19 interact vary considerably between and within populations and regions, and are affected by dynamic and complex interactions with underlying socioeconomic, political, demographic, and cultural conditions. These conditions can lead to vulnerability, resilience, transformation, or collapse of health systems, communities, and livelihoods throughout varying timescales. It is important that COVID-19 response and recovery measures consider climatic risks, particularly in locations that are susceptible to climate extremes, through integrated planning that includes public health, disaster preparedness, emergency management, sustainable development, and humanitarian response.

Introduction

First reported to WHO in 2019 as a cluster of cases of pneumonia of unknown cause, COVID-19 (SARS-CoV-2) rapidly spread around the globe via highly connected global travel networks. It has now reached the most remote areas of the world, including Indigenous Amazonian communities and the polar regions,^{1,2} with 599 071 265 confirmed cumulative cases and 646 702 3 cumulative deaths worldwide reported as of Aug 30, 2022.³ COVID-19 is an unprecedented challenge to humanity, with widespread implications for health, economic development, and wellbeing that weaken progress made in achieving the UN Sustainable Development Goals.⁴

The COVID-19 pandemic is exacerbating existing inequalities, with marginalised and vulnerable groups experiencing higher numbers of infections and deaths, increased poverty, and growing deficits in education.⁴⁻⁷ The reasons for this exacerbation of inequalities vary by context, but include the necessity of social contact for work, high population density, variable health-care provision and availability, and minimal or absent social protection. Unemployment as a result of COVID-19 has tended to affect individuals who work in the informal sector or in insecure jobs the most, as their livelihoods are at greatest risk of being made redundant due to government attempts to control the spread of the SARS-CoV-2 virus (eg, through lockdowns, and limits to transportation and mobility).⁸ These individuals in precarious employment are more likely to be part of marginalised communities.⁸ Diverse policies have been implemented to control the COVID-19 pandemic, including national and regional lockdowns, physical distancing measures, and compulsory face mask wearing in public spaces. Vaccines began to be administered to the public in 2020 in some, mostly high-income, countries.

These policies have had varying degrees of success, but have also created new challenges to health, livelihoods, and wellbeing. At various stages of the pandemic, governments have been forced to develop and adjust policies to restrict the effects of COVID-19 response measures (eg, supplement lost wages or campaign to get people to return to restaurants). However, efforts to control other diseases (eg, tuberculosis, malaria, and dengue fever) have been affected by these policies, introducing difficulties for conditions that require chronic care.⁹

In a highly interconnected world, COVID-19 intersects with diverse regional and global challenges, such as inequality, conflict, nationalism, increasing commodity prices, and environmental change. Although the dynamics of the interactions between COVID-19 and these challenges are poorly understood, with little research taking a holistic look at how COVID-19 intersects with these diverse challenges, their co-occurrence can create effects that have consequences in areas other than the health domain and can weaken the resilience of multiple sectors of social-ecological systems.^{4,10-12} The potential for compound risks or complex health emergencies could be counteracted by changes that have enhanced health and livelihoods, including technological developments, food sovereignty, improved communications technology, community empowerment, and lessons from previous pandemics.

The emergence of COVID-19 coincided with one of the hottest years on record globally (2020). 2020 and 2021 were characterised by notable climatic extremes, creating substantial interest in the intersection of the COVID-19 pandemic with climate risks. Research to date has investigated lessons learned from COVID-19 for climate policy,^{10,13-15} disaster preparedness,^{16,17} and support for climate action,^{18,19} compared the politics of responding to COVID-19 with the politics of responding to climate

Lancet Planet Health 2022;
6: e825-33

Priestley International Centre for Climate, University of Leeds, Leeds, UK (Prof J D Ford PhD, I Arotoma-Rojas MA, Prof L Berrang-Ford PhD, B van Bavel PhD); Intercultural Citizenship and Indigenous Health Unit (C Zavaleta-Cortijo PhD), School of Public Health and Administration (V Chicmana-Zapata BS), Cayetano Heredia University, Lima, Peru; School of Public Health and Social Policy, University of Victoria, Victoria, BC, Canada (A Naylor PhD); Department of Geography, Geo-Informatics, and Climatic Sciences, Makerere University, Kampala, Uganda (T Ainembabazi MA, Prof S Lwasa PhD); Center of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia, Lima, Peru (C Anza-Ramirez MSc, Prof J J Miranda PhD); Community Engagement, Rhodes University, Makhanda, South Africa (J Bezerra PhD, N Nkwinti BA); Department of Geography, Virginia Tech, Blacksburg, VA, USA (E K Galappaththi PhD); Department of Livestock Production, Agribusiness, and Economics, University of Namibia, Oshakati, Namibia (M Hangula MSc); Makerere University Business School, Makerere University, Kampala, Uganda (C Kazaana MA); Ministry of Health (D Namanya MA), Uganda National Health Research Organisation (R Nuwagira BA, S Okware PhD), Entebbe, Uganda; Arctic State Institute of Culture and Arts, North-Eastern Federal University, Yakutsk, Russia (M Osipova PhD); Sustainability Research Centre, University of the Sunshine Coast, Buderim, QLD, Australia (K Pickering PhD); School of Environment and Sustainability, Indian Institute for Human Settlements, Bangalore, India (C Singh PhD); Department of Politics and

International Studies,
University of Warwick,
Coventry, UK
(Prof K Hyams PhD);
Environmental and
Geographical Science,
University of Cape Town,
Cape Town, South Africa
(Prof M New PhD)

Correspondence to:
Prof J D Ford, Priestley
International Centre for Climate,
University of Leeds,
Leeds LS2 9JT, UK
j.ford2@leeds.ac.uk
For the UN Sustainable
Development Goals see [https://
sdgs.un.org/](https://sdgs.un.org/)

Panel: Summary of case studies to show how COVID-19 and climate interact

- The Sri Lanka case study shows the interactions between air quality and deaths, examining the climatic risks and responses of Coastal-Vedda communities to COVID-19 (appendix pp 2–4)
- The India case study reports the risks of cyclones and COVID-19 and how they are affecting the livelihoods of people and jeopardising physical distancing (appendix pp 4–5)
- The Russia case study focuses on wildfires that have affected remote Indigenous communities during the COVID-19 pandemic (appendix p 6)
- The Fiji case study reports on how COVID-19 restrictions delayed aid responding to tropical cyclones in a population with a high prevalence of non-communicable diseases (appendix pp 7–8)
- The Uganda case study shows how flooding and landslides before and during the COVID-19 pandemic lockdowns increased the risk to health, livelihoods, and wellbeing for Indigenous populations (appendix pp 9–10)
- The South Africa case study reports the long-term effects of climate change on food and water security, increasing vulnerability to water and food insecurity, and weakening community resilience to COVID-19 (appendix pp 11–13)
- The Peruvian case study shows how poorly prepared the health system in Peru was to respond to the COVID-19 pandemic, especially in the Amazon region, and how endemic climate-sensitive diseases (eg, dengue fever) increased the difficulty of identifying COVID-19 cases (appendix pp 14–15)

change;²⁰ examined implications of the pandemic for greenhouse gas emissions^{21,22} and the environment as a whole;^{23,24} and focused on how to improve the sustainability of recovery packages to fulfil the UN Build Back Better campaign goals.^{25–27} Studies have also focused on the associations between weather and COVID-19 transmission and, to a lesser extent, the ways climatic factors affect lived experience of and response to COVID-19.^{28–35} These studies have improved understanding of how environmental change and the pandemic interact, but typically only focus on particular regions, sectors, or geographical scales, and rarely focus on the interactions between multiple risks.^{9,12,36,37}

In this Personal View, we explain the ways climatic risks can affect the transmission, perception, response, and lived experience of COVID-19. We also provide case studies that show ongoing work with Indigenous people and other marginalised communities as part of the COVID Observatories, an international research programme working in multiple nations of varying incomes to document lived experiences of, effects of, and policy responses to COVID-19 in the context of climate risks (panel; appendix pp 2–15). This Personal View is influenced

by concepts from health geography,³⁸ sustainability science,³⁹ and planetary health.⁴⁰ These concepts bring attention to the complex interactions between natural and human systems that affect the emergence, spread, and effects of disease in the context of environmental risks. Research in these areas emphasises the importance of political, economic, social, and biomedical factors operating across local to global scales, and views health as encompassing not only the presence or absence of illness but also physical, mental, and sociocultural wellbeing.⁴¹

Climatic risks influencing the transmission of COVID-19

COVID-19 transmission mostly occurs through close contact between an infected person and an uninfected person through exposure to aerosols or droplets carrying the infectious SARS-CoV-2 virus. Contact with contaminated surfaces or objects can also transmit COVID-19, but the risk of infection is much lower.⁴² This implies that the presence, survival, and infectivity of a virus in the air and on object surfaces could be affected by environmental conditions, such as temperature, humidity, precipitation, air quality, and type of surface.^{29,43,44} As the SARS-CoV-2 virus has been isolated from stool and urine samples, other methods of transmission and the association with environmental factors, including water and food sources, are also being investigated.⁴⁵

Studies examining the links between climatic factors and COVID-19 have drawn inconclusive results.^{29,33,43,46–49} In Brazil, for example, some research has shown that COVID-19 infection rates are inversely correlated with temperature,⁵⁰ whereas other studies have documented that high mean temperatures were associated with increased COVID-19 transmission.^{51,52} There is little understanding about the effects of humidity, wind, and solar radiation on COVID-19 transmission. Whether climatic factors affect SARS-CoV-2 itself or influence transmission by changing the behaviours, and therefore exposures, of people remains unclear. Some of the differences in results across studies reflect the wide range of statistical and modelling approaches used and the temporal scope of data analysed, and suggest that non-climatic factors are typically more important than climatic factors in disease transmission⁵³ as climatic factors alone are not large enough to control the COVID-19 pandemic.^{28,33–35} As the pandemic has progressed and COVID-19 has become global, the robustness of early studies suggesting strong links between climate and COVID-19 have been questioned.^{33,34,47,54,55} Reviewing these studies, Gutiérrez-Hernández and García³⁴ argue that evidence of climate influences on COVID-19 is not robust enough to be considered in public health policies.

There is negligible evidence that climate change will directly or substantively affect COVID-19 transmission in the long term. There is, however, evidence of links between air pollution and COVID-19 transmission. Multiple studies show strong correlations between

See Online for appendix

particulate matter (PM) from fossil fuel combustion (eg, PM_{2.5}) and high rates of COVID-19 infection and mortality (appendix pp 2–4),^{30,56–60} although the evidence remains merely associational.

Coinciding risks of climatic extremes and COVID-19

Globally, the beginning of the COVID-19 pandemic intersected with one of the hottest years on record (2020). 2020 and 2021 also had several extreme climatic events, including wildfires (eg, in Siberia, North America, and Türkiye); hurricanes and tropical storms (eg, in Fiji and the Bay of Bengal); floods (eg, in southern China, Pakistan, and Germany); droughts (eg, in west USA); and heatwaves (eg, in Canada). Although not all of these events are directly attributable to climate change, trends in the intensity and frequency of some climate and weather extremes have been detected. The World Meteorological Organization and Intergovernmental Panel on Climate Change note a clear effect of anthropogenic climate change on many of these extreme events.^{61,62} The people and communities with the highest exposure to climatic extremes have also experienced more infections of and deaths from COVID-19,^{63–66} creating a double burden to affected individuals. Evidence of the interaction between climatic extremes and COVID-19 has only just begun to emerge, with several discernible potential links (eg, direct effects on livelihoods or indirect effects on maternal health and nutrition). Different types of inequalities are increasingly known to interact to exacerbate vulnerability to a range of different health, social, and environmental hazards; COVID-19 has revealed how ubiquitous these interactions are.

Modified disease exposure

Extreme weather events result in mass displacement (eg, evacuation, migration, and mass sheltering), thereby introducing diseases into new regions, clustering survivors in temporary accommodation with little or no ventilation, increasing amounts of social contact, and making face masks difficult to use (eg, during temperature extremes).^{30,31,67} In these situations, physical distancing is more challenging than in typical conditions, with guidelines for separation insufficient in enclosed spaces, such as shelters, and challenging to practise during evacuations (appendix pp 4–5).^{64,68} Regular handwashing can also be difficult in these situations, in which hygiene facilities might be disrupted due to breakage or contamination of water and sanitation systems.^{69,70} The COVID-19 pandemic could also amplify pre-existing inequalities in access to safe drinking water.⁷¹ For example, in Namibia, movement restrictions and lockdowns hindered the ability of individuals to earn daily wages to pay for drinking and washing water, leading to an inability to observe all the COVID-19 protocols. Heavy rainfall events might potentially increase the risks of faecal-oral transmission of COVID-19 by causing sewage system

overflows; the virus has been detected in human faeces and urine where it can remain viable for days.^{72,73} Although research suggests COVID-19 transmission is largely airborne,⁷⁴ handwashing remains one of the most recommended preventive measures.

Increased susceptibility to COVID-19 through wildfire smoke inhalation

Previous exposure to smoke from wildfires has been linked to increased morbidity from influenza,⁷⁵ leading to concern about increased risk of COVID-19 infection associated with wildfires.^{76–78} Wildfire smoke contains small particles (PM_{2.5}) that penetrate the lungs, causing irritation and inflammation and conceivably increasing the risk of COVID-19 infection. Individuals most at risk of enhanced susceptibility to respiratory infection due to wildfire smoke exposure are children and infants with small lung size and individuals with pre-existing lung or heart conditions.^{78,79} Several large wildfires have coincided with the COVID-19 pandemic—notably in Siberia, California (USA), British Columbia (Canada), Türkiye, and the Brazilian Amazon—with studies documenting trends of increasing wildfire activity in some regions compared with previous years, driven partly by climate change and other anthropogenic factors, such as land management practices (appendix p 6).^{77,80,81} Individuals with COVID-19-related pulmonary or cardiac impairment are at high risk of illness and death from wildfire-related smoke (and heatwaves).³⁰ Smoke from the 2020 California wildfire season also impacted people's lives during the pandemic by necessitating further restrictions on outdoor activities and exacerbating social isolation; with potential mental health implications.⁷⁶

Compromised emergency response and reduced health system resilience

Climate-related disasters can challenge COVID-19 responses from governments and public health systems by disrupting supply chains, restricting access to humanitarian aid, and reducing the number of mobilisable health-care staff and resources (appendix pp 7–10).^{30,82,83} Access to health-care services is an important factor in assessing the capacity of a health system to respond to risks and effects of climate-related disasters.^{84,85} Extreme climatic events reduce patients' ability to seek treatment and access care for COVID-19. Climate hazards also cause problems in the delivery of public services (eg, disruption of power supplies and emergency services).^{30,82} These events can also make contact tracing difficult as people move and intermingle (eg, in response to flooding).⁷³

Actions taken to control COVID-19 could also increase the vulnerability of individuals and health-care services to climatic risks. Orders to stay at home and concerns about COVID-19 might result in people failing to heed evacuation orders, leading to preventable injury and death.^{31,86} COVID-19 has also created challenges for managing extreme climatic events. For example, efforts to

control wildfires in California were restricted by high heat, low humidity, high winds, and no people living in prison who are available to work (an integral part of fire control in the state) due to high infection rates within the prison population.⁷⁶ Measures for responding to climatic extremes can be incompatible with efforts to control COVID-19, leading to confusion, contradictory messaging, and uncertainty about whether to follow COVID-19 control efforts or not. These examples show how multiple risks can all affect health systems, leading to negative outcomes for people and locations with low capacities to respond to COVID-19.

Long-term climate change, community vulnerability, and increased COVID-19 risk

The effects of climate change have been observed globally in diverse regions,^{87,88} with implications for community vulnerability to diverse threats, including COVID-19. In a modelling study, Ozkan and colleagues⁸⁹ showed that countries with high climatic risk also had a higher COVID-19 mortality rate than countries with low or no climatic risk. The apparent clustering of climate and COVID-19 vulnerability in particular regions and communities shows the potential for direct or indirect effects of the interaction between COVID-19 and climate, and that similar factors often drive both climate risk and COVID-19 risk, with climate risk also leading to specific conditions that result in increased COVID-19 risk.

Direct effects of climate change include damaging or undermining health and emergency response infrastructure. For example, in Peru, the 2017 El Niño Costero event severely damaged 934 health posts (primary care health centres) via flooding. Few of these posts had been rebuilt when COVID-19 happened, compromising pandemic planning and management.³⁷ Severe and long-lasting drought in some regions has negatively affected food systems, leading to malnourishment, poverty, increased sensitivity of individuals to COVID-19, and restricting access to clean water, which is essential for controlling the transmission of COVID-19 (appendix pp 11–13).^{11,12,24,31} Heavy rainfall in south Namibia aggravated food insecurity by depleting household food stocks and increasing reliance on markets, which are costly and were difficult to reach due to flooding.

Indirect effects of climate change include long-term climate change interacting with social, economic, political, and cultural factors; undermining adaptive capacity to manage COVID-19; and increasing sensitivity of individuals and health-care systems to COVID-19. For example, Indigenous populations in South America are among some of the communities most affected by the COVID-19 pandemic.⁹⁰ Climate change has exacerbated pre-existing problems from structural inequalities, marginalisation, land dispossession, and government policy and has increased vulnerability to a diversity of threats (appendix pp 14–15).^{91–94} In the Peruvian Amazon, changes to precipitation regimes, seasonality, hydrology,

and temperatures have negatively affected Indigenous food systems by disrupting community fisheries, reducing access to and availability of highly valued forest animals, and reducing opportunities to develop and acquire traditional knowledge.⁹⁵ As some Indigenous communities in the Peruvian Amazon are currently encountering different forms of malnutrition,⁹⁵ climate-related food shortages risk affecting both the availability and the quality of food sources (eg, fresh vegetables and fish), suggesting that nutritional consequences will be observed in future generations.⁹⁶ Among reindeer herders in Siberia, Stammner and Ivanova³² report how the cumulative effects of frequent climate-induced disasters (eg, large numbers of reindeer dying because of pasture icing events or an anthrax outbreak) have resulted in an increase in conspiracy theories, in which disasters such as COVID-19 are viewed as being caused by malevolent human forces that are beyond the control of herders. In this context, lockdowns imposed by regional or federal governments are seen as an attack on nomadic life, with implications for acceptance of control measures.

Discussion

In this Personal View, we explain the ways climatic risks might affect the transmission, perception, response, and lived experience of COVID-19, focusing on long-term climate change and sporadic extremes during the pandemic. The COVID-19 pandemic is ongoing, and therefore some interactions between climate and COVID-19 are probably not discernible at the time of writing and will emerge in the future. Understanding the links between the COVID-19 pandemic and climate change is essential to advance the field of planetary health, particularly in terms of the effects across daily life activities and professions. The potential interactions between COVID-19 and climate described in this Personal View offer a portent of how society might be affected by future compounding climatic risks as the COVID-19 pandemic continues to develop alongside an increase in severe and frequent climatic extremes.

In this Personal View we propose a conceptual model of how climate and COVID-19 risks interact (figure). The model places COVID-19 risk in specific locations and populations at the centre of analysis, emphasising both direct and indirect effects of COVID-19 on these populations. The model shows the complex societal, environmental, and health factors affecting how climatic risks and COVID-19 interact at different scales. These interactions could lead to vulnerability, resilience, transformation, or collapse of health systems, communities, and livelihoods throughout varying timescales, which changes depending on the phase of the COVID-19 pandemic a society is in. The model also shows important drivers and considerations for understanding the interaction between climate risks and COVID-19 using the published literature and research done as part of the COVID Observatories project.

Climate and other interacting factors affecting COVID-19 risk

Relationships between human society, the natural environment, and disease are affected by multiple, synchronous stressors.³⁹ Stressors come from rapidly developing threats, such as COVID-19 emergence and policy response, climate extremes, ecosystemic stress, changes to livelihood, food prices, protests, and market access. These threats, known as fast variables, interact with and are affected by long-term changes in society, human–environment interactions, and the natural environment. So-called slow variables include globalisation, conflict, development, colonisation, demographics, and environmental change. The interaction of fast and slow variables affects how climate and COVID-19 interact in specific locations, in which feedbacks, thresholds, and non-linear responses can result in rapid and immediate or slow and transitional outcomes. The complexities of these interactions are poorly understood and in some instances could lead to resilience to multiple interacting stressors (appendix pp 2–5, 11–13). In remote Indigenous communities in the Arctic, for example, a high amount of vigilance, travel restrictions, and rapid vaccinations have managed to keep infection rates low, despite the presence of underlying COVID-19 risk factors in Indigenous populations and climatic warming rates that are 3 times faster than the global average.² In Amazonia, some Indigenous communities have enforced strict measures regarding entry to their communities, showing self-determination and implementation of medical Indigenous strategies to control COVID-19.^{92,97,98} For the Asháninka people in Peru, for example, the inability to travel to markets was an opportunity for people to revalue traditional knowledge of food sovereignty. Cities in South Africa have seen a rise in home gardens since the beginning of the COVID-19 pandemic to improve food security.⁹⁹

Interaction between climate and COVID-19

The ways that climate and COVID-19 interact vary between and within populations and regions, intersecting with gender, ethnicity, culture, livelihoods, income, and access to health care to create differential risks. Evidence indicates that proactive responses to climatic risks and COVID-19 can moderate risks (eg, health, livelihoods, and wellbeing), emphasising the importance of planning and implementation of early and ongoing action, communication, flexible funding mechanisms, capacity building, and monitoring and evaluation of policies and responses.^{37,100} Japan and South Korea had COVID-19 resurgences during the summer of 2020 after a category four typhoon, but were able to control the spread.¹⁰¹ Responses to COVID-19 might only require minor alterations to existing hazard preparedness plans,³⁷ although substantial challenges, including an absence of tools for managing compound risks, unprecedented uncertainty, data scarcity, competing policy options,




Figure: Conceptual model of the complex interactions between COVID-19 and climate risk (A) The multiple environmental, socioeconomic, and health factors affecting COVID-19 risk. (B) Different phases of the COVID-19 pandemic (eg, increasing and decreasing numbers of cases) across time and geographical scales.

marginalisation, and inequality, have been identified as major constraints.^{24,102,103} The effectiveness of a health system depends on its ability to help vulnerable populations; consider the intersections of gender, ethnicity, rights, and other influential factors; and consider the influence of these factors on differential health outcomes.⁸⁵ Integrated services and collaboration across sectors will be crucial in establishing effective and equitable health system prevention, detection, and responses to COVID-19.^{84,85,104}

The characteristics of individuals and their interaction with the pathological agent (SARS-Cov-2) modulate the severity of the effects of climate risk on COVID-19. Some of these characteristics have also been reported to be associated with adverse outcomes as a result of climatic extreme events (eg, fast variables),¹⁰⁵ suggesting that people with these characteristics would be more vulnerable to risks of COVID-19 and climate than people without these characteristics. For example, older adults and people living with particular non-communicable diseases are more likely to die during heatwaves.^{106–108} The risk of developing severe forms of COVID-19 infection are higher in people with diabetes.¹⁰⁹ These two groups might therefore be vulnerable to compounding health risks from extreme heat and COVID-19 outbreaks. A high BMI is also associated with increased risk of severe COVID-19 disease.¹¹⁰ The prevalence of overweight in many low-income and middle-income countries is increasing as food systems and food environments change (eg, slow variables), meaning that these populations could become more susceptible to severe health impacts from outbreaks of infectious diseases, such as COVID-19 (appendix p 7).⁸³

COVID-19 measures, climate change, and increased climate variability

Policy responses to COVID-19 have prioritised controlling the spread of the SARS-CoV-2 virus.¹¹¹ While these public health measures were often necessary tools to confront the COVID-19 pandemic they have also increased poverty, reduced market access, and restricted mobility in many locations, and might consequently have increased vulnerability to climatic extremes and long-term climate change. As the COVID-19 pandemic continues to affect communities globally, it is important that planning and response considers these implications, as well as the complications from ongoing climate extremes.¹⁰ Connecting research, practitioner, and policy-making communities within public health, disaster preparedness, emergency management, humanitarian response, and development planning sectors is imperative for informing more integrated responses across sectors and geographic scales. For example, responses to challenges affecting health, livelihoods, and wellbeing have had success in reducing disruption caused by COVID-19 and climate extremes (appendix pp 16–17).

The COVID-19 pandemic has shown the importance of challenging underlying inequalities and injustices that increase vulnerability to health emergencies and climate change. It also highlights deficits in existing efforts to adapt to climate change. For example, few climate adaptation actions integrate pandemic preparedness or have contingency plans for major or rare events, which might affect their implementation and effectiveness.¹¹² To fulfil the UN Build Back Better campaign goals, a comprehensive analysis of the multiple climatic risks that influence transmission, lived

experiences, and responses to COVID-19 is required; as is an analysis of the underlying socioeconomic, political, and cultural factors that create pandemic vulnerability or provide opportunities to protect human health. Political incentives for focusing on short-term projects and results and nationalism motivate inadequate responses to both climate and pandemic risk. New political institutions and interventions that increase the focus on long-term projects and results and global cooperation will help with both climate change and future pandemic risk.

Contributors

JDF and CZ-C conceptualised the piece. JDF, CZ-C, DN, CS, LB-F, KH, JJM, and MN acquired funding. JDF, CZ-C, TA, CA-R, IA-R, JB, VC-Z, EKG, MH, CK, SL, DN, NN, RN, SO, MO, KP, and CS did investigations for the case studies. JDF, CZ-C, and LBF conducted the method. Project administration was done by CA-R. CZ-C, CA-R, EKG, KP, and CS created the figures. The original draft was written by JDF, CZ-C, LB-F, TA, CA-R, IA-R, JB, VC-Z, EKG, MH, CK, SL, DN, MN, NN, RN, SO, MO, KP, and CS. Reviewing and editing was done by JDF, CZ-C, TA, CA-R, IA-R, JB, VC-Z, EKG, MH, CK, SL, DN, NN, RN, SO, MO, KP, CS, LB-F, KH, JJM, AN, MN, and BvB.

Declaration of interests

The COVID-19 Observatory (project reference number: EP/V043102/1) is funded by a collective fund award from the UK Research and Innovation Global Challenges Research Fund and Newton Fund; the Foreign, Commonwealth & Development Office; and International Programme—Russia (Arctic Voices, 2021 006). CZ-C was supported by the National Institute for Health Research (NIHR) through the Official Development Assistance Funding of the UK and Wellcome (218743_Z_19_Z) under the NIHR-Wellcome Partnership for Global Health Research. The views expressed are those of the authors and not necessarily those of Wellcome, the NIHR, or the Department of Health and Social Care. All other authors declare no competing interests.

Acknowledgments

The COVID-19 Observatory would like to acknowledge the Indigenous Peoples and representatives from Uganda (Batwa), Namibia (Ovambo), Ghana (Dagaaba), South Africa (Xhosa), Peru (Asháninka, Shawi), Bolivia (Guarani, Amazónicos, Quechuas, and Aymaras), Sri Lanka (coastal-Vedda), India (Irula, Badaga, Toda, Kota, and Kurumba), Fiji (iTaukei), Russia (Even), Australia (Gubbi Gubbi), and Canada (Inuit), without whose collaboration this project would not be possible. In Peru, we would like to thank Ketty Marcelo Lopez and the Indigenous organisation she belongs to, Organización de Mujeres Indígenas Amazónicas Ashaninkas de la Selva Central (OMIAASEC). We are also grateful for the support in the visualisation of this Personal View prepared by Carlee Wright.

References

- Zavaleta-Cortijo C, Ford JD, Arotoma-Rojas I, et al. Climate change and COVID-19: reinforcing Indigenous food systems. *Lancet Planet Health* 2020; 4: e381–82.
- Petrov AN, Welford M, Golosov N, et al. The “second wave” of the COVID-19 pandemic in the Arctic: regional and temporal dynamics. *Int J Circumpolar Health* 2021; 80: 1925446.
- WHO. WHO coronavirus (COVID-19) dashboard. 2022. <https://covid19.who.int/> (accessed June 14, 2022).
- The Lancet COVID-19 Commissioners, Task Force Chairs, and Commission Secretariat. *Lancet* COVID-19 Commission Statement on the occasion of the 75th session of the UN General Assembly. *Lancet* 2020; 396: 1102–24.
- Krieger N. Enough: COVID-19, structural racism, police brutality, plutocracy, climate change and time for health justice, democratic governance, and an equitable, sustainable future. *Am J Public Health* 2020; 110: 1620–23.
- Douglas M, Katikireddi SV, Taulbut M, McKee M, McCartney G. Mitigating the wider health effects of COVID-19 pandemic response. *BMJ* 2020; 369: m1557.

- 7 Nicola M, Alsafi Z, Sohrabi C, et al. The socio-economic implications of the coronavirus pandemic (COVID-19): a review. *Int J Surg* 2020; **78**: 185–93.
- 8 Blustein DL, Duffy R, Ferreira JA, Cohen-Scali V, Cinamon RG, Allan BA. Unemployment in the time of COVID-19: a research agenda. *J Vocat Behav* 2020; **119**: 103436.
- 9 Cousins T, Pentecost M, Alvergne A, et al. The changing climates of global health. *BMJ Glob Health* 2021; **6**: e005442.
- 10 Ringsmuth AK, Otto IM, van den Hurk B, et al. Lessons from COVID-19 for managing transboundary climate risks and building resilience. *Clim Risk Manage* 2022; **35**: 100395.
- 11 Mishra A, Bruno E, Zilberman D. Compound natural and human disasters: managing drought and COVID-19 to sustain global agriculture and food sectors. *Sci Total Environ* 2021; **754**: 142210.
- 12 Cazzolla Gatti R, Menéndez LP, Laciny A, et al. Diversity lost: COVID-19 as a phenomenon of the total environment. *Sci Total Environ* 2021; **756**: 144014.
- 13 Botzen W, Duijndam S, van Beukering P. Lessons for climate policy from behavioral biases towards COVID-19 and climate change risks. *World Dev* 2021; **137**: 105214.
- 14 Klenert D, Funke F, Mattauich L, O'Callaghan B. Five lessons from COVID-19 for advancing climate change mitigation. *Environ Resour Econ* 2020; **76**: 751–78.
- 15 Manzanedo RD, Manning P. COVID-19: lessons for the climate change emergency. *Sci Total Environ* 2020; **742**: 140563.
- 16 Sheehan MC, Fox MA. Early warnings: the lessons of COVID-19 for public health climate preparedness. *Int J Health Serv* 2020; **50**: 264–70.
- 17 Pelling M, Chow WTL, Chu E, et al. A climate resilience research renewal agenda: learning lessons from the COVID-19 pandemic for urban climate resilience. *Clim Dev* 2021; published online Aug 22. <https://doi.org/10.1080/17565529.2021.1956411>.
- 18 Dupont C, Oberthur S, von Homeyer I. The COVID-19 crisis: a critical juncture for EU climate policy development? *J Eur Integr* 2020; **42**: 1095–110.
- 19 Howarth C, Bryant P, Corner A, et al. Building a social mandate for climate action: lessons from COVID-19. *Environ Resour Econ* 2020; **76**: 1107–15.
- 20 van der Ven H, Sun YX. Varieties of crises: comparing the politics of COVID-19 and climate change. *Glob Environ Polit* 2021; **21**: 13–22.
- 21 Forster PM, Forster HI, Evans MJ, et al. Erratum: publisher correction: current and future global climate impacts resulting from COVID-19. *Nat Clim Chang* 2020; **10**: 971.
- 22 Zerefos CS, Solomos S, Kapsomenakis J, et al. Lessons learned and questions raised during and post-COVID-19 anthropopause period in relation to the environment and climate. *Environ Dev Sustain* 2021; **23**: 10623–45.
- 23 Praveena SM, Aris AZ. The impacts of COVID-19 on the environmental sustainability: a perspective from the southeast Asian region. *Environ Sci Pollut Res Int* 2021; **28**: 63829–36.
- 24 Sarkar P, Debnath N, Reang D. Coupled human-environment system amid COVID-19 crisis: a conceptual model to understand the nexus. *Sci Total Environ* 2021; **753**: 141757.
- 25 Engler D, Groh ED, Gutsche G, Ziegler A. Acceptance of climate-oriented policy measures under the COVID-19 crisis: an empirical analysis for Germany. *Climate Policy* 2021; **10**: 1281–97.
- 26 Rosenbloom D, Markard J. A COVID-19 recovery for climate. *Science* 2020; **368**: 447.
- 27 UNA-UK. Sustainable Development Goals: building back better. <https://una.org.uk/sustainable-development-goals-building-back-better> (accessed Sept 1, 2022).
- 28 Pan J, Yao Y, Liu Z, et al. Warmer weather unlikely to reduce the COVID-19 transmission: an ecological study in 202 locations in 8 countries. *Sci Total Environ* 2021; **753**: 142272.
- 29 Kubota Y, Shiono T, Kusumoto B, Fujinuma J. Multiple drivers of the COVID-19 spread: the roles of climate, international mobility, and region-specific conditions. *PLoS One* 2020; **15**: e0239385.
- 30 Salas RN, Shultz JM, Solomon CG. The climate crisis and COVID-19—a major threat to the pandemic response. *N Engl J Med* 2020; **383**: e70.
- 31 Phillips CA, Caldas A, Cleetus R, et al. Compound climate risks in the COVID-19 pandemic. *Nat Clim Chang* 2020; **10**: 586–88.
- 32 Stammeler FM, Ivanova A. From spirits to conspiracy? Nomadic perceptions of climate change, pandemics and disease. *Anthropol Today* 2020; **36**: 8–12.
- 33 Briz-Redon A, Serrano-Aroca A. The effect of climate on the spread of the COVID-19 pandemic: a review of findings, and statistical and modelling techniques. *Prog Phys Geogr* 2020; **44**: 591–604.
- 34 Gutiérrez-Hernández O, García LV. On the usefulness of the bioclimatic correlative models of SARS-CoV-2. *Environ Res* 2021; **195**: 110818.
- 35 Sera F, Armstrong B, Abbott S, et al. A cross-sectional analysis of meteorological factors and SARS-CoV-2 transmission in 409 cities across 26 countries. *Nat Commun* 2021; **12**: 5968.
- 36 Shakil MH, Munim ZH, Tasnia M, Sarowar S. COVID-19 and the environment: a critical review and research agenda. *Sci Total Environ* 2020; **745**: 141022.
- 37 de la Poterie AT, Clatworthy Y, Easton-Calabria E, de Perez EC, Lux S, van Aalst M. Managing multiple hazards: lessons from anticipatory humanitarian action for climate disasters during COVID-19. *Clim Dev* 2021; **14**: 374–88.
- 38 Rosenberg M. Health geography II: 'dividing' health geography. *Prog Hum Geogr* 2016; **40**: 546–54.
- 39 Naylor A, Ford J, Pearce T, Van Alstine J. Conceptualizing climate vulnerability in complex adaptive systems. *One Earth* 2020; **2**: 444–54.
- 40 Whitmee S, Haines A, Beyrer C, et al. Safeguarding human health in the Anthropocene epoch: report of The Rockefeller Foundation-Lancet Commission on planetary health. *Lancet* 2015; **386**: 1973–2028.
- 41 Ford JD. Indigenous health and climate change. *Am J Public Health* 2012; **102**: 1260–66.
- 42 Centers for Disease Control and Prevention. Science Brief: SARS-CoV-2 and surface (fomite) transmission for indoor community environments. Atlanta, GA: Centers for Disease Control and Prevention, 2021.
- 43 Sasikumar K, Nath D, Nath R, Chen W. Impact of extreme hot climate on COVID-19 outbreak in India. *Geohealth* 2020; **4**: GH000305.
- 44 Kumar S, Singh R, Kumari N, et al. Current understanding of the influence of environmental factors on SARS-CoV-2 transmission, persistence, and infectivity. *Environ Sci Pollut Res Int* 2021; **28**: 6267–88.
- 45 Han J, Zhang X, He S, Jia P. Can the coronavirus disease be transmitted from food? A review of evidence, risks, policies and knowledge gaps. *Environ Chem Lett* 2021; **19**: 5–16.
- 46 Zaitchik BF, Sweijd N, Shumake-Guillemot J, et al. A framework for research linking weather, climate and COVID-19. *Nat Commun* 2020; **11**: 5730.
- 47 Carlson CJ, Gomez ACR, Bansal S, Ryan SJ. Misconceptions about weather and seasonality must not misguide COVID-19 response. *Nat Commun* 2020; **11**: 4312.
- 48 Fan JL, Da YB, Zeng B, et al. How do weather and climate change impact the COVID-19 pandemic? Evidence from the Chinese mainland. *Environ Res Lett* 2021; **16**: 014026.
- 49 To T, Zhang K, Maguire B, et al. Correlation of ambient temperature and COVID-19 incidence in Canada. *Sci Total Environ* 2021; **750**: 141484.
- 50 Pequeno P, Mendel B, Rosa C, et al. Air transportation, population density and temperature predict the spread of COVID-19 in Brazil. *PeerJ* 2020; **8**: e9322.
- 51 Prata DN, Rodrigues W, Bermejo PH. Temperature significantly changes COVID-19 transmission in (sub)tropical cities of Brazil. *Sci Total Environ* 2020; **729**: 138862.
- 52 Auler AC, Cássaro FAM, da Silva VO, Pires LF. Evidence that high temperatures and intermediate relative humidity might favor the spread of COVID-19 in tropical climate: a case study for the most affected Brazilian cities. *Sci Total Environ* 2020; **729**: 139090.
- 53 Kerr GH, Badr HS, Gardner LM, Perez-Saez J, Zaitchik BF. Associations between meteorology and COVID-19 in early studies: inconsistencies, uncertainties, and recommendations. *One Health* 2021; **12**: 100225.
- 54 Malanson GP. COVID-19 and physical geography—yellow card. *Prog Phys Geogr* 2020; **44**: 447–48.
- 55 Gutiérrez-Hernández O, García LV. Multiplicity eludes peer review: the case of COVID-19 research. *Int J Environ Res Public Health* 2021; **18**: 9304.

- 56 Ali N, Islam F. The effects of air pollution on COVID-19 infection and mortality—a review on recent evidence. *Front Public Health* 2020; **8**: 580057.
- 57 Srivastava A. COVID-19 and air pollution and meteorology—an intricate relationship: a review. *Chemosphere* 2021; **263**: 128297.
- 58 Petroni M, Hill D, Younes L, et al. Hazardous air pollutant exposure as a contributing factor to COVID-19 mortality in the United States. *Environ Res Lett* 2020; **15**: 0940a9.
- 59 Wu X, Nethery RC, Sabath MB, Braun D, Dominici F. Air pollution and COVID-19 mortality in the United States: strengths and limitations of an ecological regression analysis. *Sci Adv* 2020; **6**: eabd4049.
- 60 Sarkodie SA, Owusu PA. Global effect of city-to-city air pollution, health conditions, climatic and socio-economic factors on COVID-19 pandemic. *Sci Total Environ* 2021; **778**: 146394.
- 61 World Meteorological Organization. United in science report: climate change has not stopped for COVID19. 2020. <https://public.wmo.int/en/media/press-release/united-science-report-climate-change-has-not-stopped-covid19> (accessed Sept 1, 2022).
- 62 Intergovernmental Panel on Climate Change. Climate change 2021: the physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press, 2021.
- 63 Rahman MM, Bodrud-Doza M, Shammi M, Md Towfiqul Islam AR, Moniruzzaman Khan AS. COVID-19 pandemic, dengue epidemic, and climate change vulnerability in Bangladesh: scenario assessment for strategic management and policy implications. *Environ Res* 2021; **192**: 110303.
- 64 Simonovic SP, Kundzewicz ZW, Wright N. Floods and the COVID-19 pandemic—a new double hazard problem. *WIREs Water* 2021; **8**: e1509.
- 65 Flores-Ramírez R, Berumen-Rodríguez AA, Martínez-Castillo MA, Alcántara-Quintana LE, Díaz-Barriga F, Díaz de León-Martínez L. A review of environmental risks and vulnerability factors of indigenous populations from Latin America and the Caribbean in the face of the COVID-19. *Glob Public Health* 2021; **16**: 975–99.
- 66 Power T, Wilson D, Best O, et al. COVID-19 and Indigenous Peoples: an imperative for action. *J Clin Nurs* 2020; **29**: 2737–41.
- 67 Li Y, Tokura H, Guo YP, et al. Effects of wearing N95 and surgical facemasks on heart rate, thermal stress and subjective sensations. *Int Arch Occup Environ Health* 2005; **78**: 501–09.
- 68 Somsen GA, van Rijn C, Kooij S, Bem RA, Bonn D. Small droplet aerosols in poorly ventilated spaces and SARS-CoV-2 transmission. *Lancet Respir Med* 2020; **8**: 658–59.
- 69 Armitage R, Nellums LB. Water, climate change, and COVID-19: prioritising those in water-stressed settings. *Lancet Planet Health* 2020; **4**: e175.
- 70 Ebrahim SH, Rahman NMM, Imtiaz R, et al. Forward planning for disaster-related mass gatherings amid COVID-19. *Lancet Planet Health* 2020; **4**: e379–80.
- 71 Hernández-Vasquez A, Rojas-Roque C, Marques Sales D, et al. Inequalities in access to safe drinking water in Peruvian households according to city size: an analysis from 2008 to 2018. *Int J Equity Health* 2021; **20**: 133.
- 72 Han J, He S. Urban flooding events pose risks of virus spread during the novel coronavirus (COVID-19) pandemic. *Sci Total Environ* 2021; **755**: 142491.
- 73 Guo Y, Wu Y, Wen B, et al. Floods in China, COVID-19, and climate change. *Lancet Planet Health* 2020; **4**: e443–44.
- 74 Greenhalgh T, Jimenez JL, Prather KA, Tufekci Z, Fisman D, Schooley R. Ten scientific reasons in support of airborne transmission of SARS-CoV-2. *Lancet* 2021; **397**: 1603–05.
- 75 Landguth EL, Holden ZA, Graham J, et al. The delayed effect of wildfire season particulate matter on subsequent influenza season in a mountain west region of the USA. *Environ Int* 2020; **139**: 105668.
- 76 Kizer KW. Extreme wildfires—a growing population health and planetary problem. *JAMA* 2020; **324**: 1605–06.
- 77 Xu R, Yu P, Abramson MJ, et al. Wildfires, global climate change, and human health. *N Engl J Med* 2020; **383**: 2173–81.
- 78 Henderson SB. The COVID-19 pandemic and wildfire smoke: potentially concomitant disasters. *Am J Public Health* 2020; **110**: 1140–42.
- 79 Centers for Disease Control and Prevention. Wildfire smoke and COVID-19. 2022. https://www.cdc.gov/disasters/covid-19/wildfire_smoke_covid-19.html (accessed Sept 1, 2022).
- 80 Williams AP, Abatzoglou JT, Gershunov A, et al. Observed impacts of anthropogenic climate change on wildfire in California. *Earths Futur* 2019; **7**: 892–910.
- 81 Schoennagel T, Balch JK, Brenkert-Smith H, et al. Adapt to more wildfire in western North American forests as climate changes. *Proc Natl Acad Sci USA* 2017; **114**: 4582–90.
- 82 Frausto-Martinez O, Aguilar-Becerra CD, Colin-Olivares O, et al. COVID-19, storms, and floods: impacts of Tropical Storm Cristobal in the western sector of the Yucatan Peninsula, Mexico. *Sustainability (Basel)* 2020; **12**: 9925.
- 83 Walton D, van Aalst M. Climate-related extreme weather events and COVID-19: a first look at number of people affected by intersecting disasters. <https://www.preventionweb.net/publication/climate-related-extreme-weather-events-and-covid-19-first-look-number-people-affected> (accessed Sept 1, 2022).
- 84 Kandel N, Chungong S, Omaar A, Xing J. Health security capacities in the context of COVID-19 outbreak: an analysis of International Health Regulations annual report data from 182 countries. *Lancet* 2020; **395**: 1047–53.
- 85 Haldane V, De Foo C, Abdalla SM, et al. Health systems resilience in managing the COVID-19 pandemic: lessons from 28 countries. *Nat Med* 2021; **27**: 964–80.
- 86 Shultz JM, Fugate C, Galea S. Cascading risks of COVID-19 resurgence during an active 2020 Atlantic hurricane season. *JAMA* 2020; **324**: 935–36.
- 87 Intergovernmental Panel on Climate Change. AR5 climate change 2014: impacts, adaptation, and vulnerability. 2014. <https://www.ipcc.ch/report/ar5/wg2/> (accessed Sept 1, 2022).
- 88 Intergovernmental Panel on Climate Change. Global warming of 1.5 °C. 2018. <https://www.ipcc.ch/sr15/> (accessed Sept 1, 2022).
- 89 Ozkan A, Ozkan G, Yalaman A, Yildiz Y. Climate risk, culture and the COVID-19 mortality: a cross-country analysis. *World Dev* 2021; **141**: 105412.
- 90 Ramírez JD, Sordillo EM, Gotuzzo E, et al. SARS-CoV-2 in the Amazon region: a harbinger of doom for Amerindians. *PLoS Negl Trop Dis* 2020; **14**: e0008686.
- 91 Rao N, Narain N, Chakraborty S, Bhanjdeo A, Pattnaik A. Destinations matter: social policy and migrant workers in the times of covid. *Eur J Dev Res* 2020; **32**: 1639–61.
- 92 Menton M, Milanez F, Souza JMD, Cruz FSM. The COVID-19 pandemic intensified resource conflicts and Indigenous resistance in Brazil. *World Dev* 2021; **138**: 138.
- 93 The Lancet. Self-determination and Indigenous health. *Lancet* 2020; **396**: 361.
- 94 Polidoro M, Mendonca FD, Meneghel SN, et al. Territories under siege: risks of the decimation of Indigenous and Quilombolas Peoples in the context of COVID-19 in south Brazil. *J Racial Ethn Health Disparities* 2020; **8**: 1119–29.
- 95 Zavaleta C, Berrang-Ford L, Ford J, et al. Multiple non-climatic drivers of food insecurity reinforce climate change maladaptation trajectories among Peruvian Indigenous Shawi in the Amazon. *PLoS One* 2018; **13**: e0205714.
- 96 Thiery W, Lange S, Rogelj J, et al. Intergenerational inequities in exposure to climate extremes. *Science* 2021; **374**: 158–60.
- 97 Richardson L, Crawford A. COVID-19 and the decolonization of Indigenous public health. *CMAJ* 2020; **192**: e1098–100.
- 98 Zavaleta C. COVID-19: review Indigenous peoples' data. *Nature* 2020; **580**: 185.
- 99 Climate and Development Knowledge Network. Cape Town, South Africa: urban farming on the rise to boost people's food security. https://cdkn.org/2020/12/feature-cape-town-south-africa-urban-farming-on-the-rise-to-boost-peoples-food-security/?loclang=en_gb (accessed Sept 13, 2021).
- 100 Sohrabizadeh S, Yousefian S, Bahramzadeh A, Vaziri MH. A systematic review of health sector responses to the coincidence of disasters and COVID-19. *BMC Public Health* 2021; **21**: 709.
- 101 Shultz JM, Kossin JP, Ali A, et al. Superimposed threats to population health from tropical cyclones in the prevaccine era of COVID-19. *Lancet Planet Health* 2020; **4**: e506–08.

- 102 Mallard A, Pesantes MA, Zavaleta-Cortijo C, Ward J. An urgent call to collect data related to COVID-19 and Indigenous populations globally. *BMJ Glob Health* 2021; **6**: e004655.
- 103 Anderson C, Leeson C, Valcourt A, Urajnik D. COVID-19 pandemic: implications for First Nations communities in Canada. *Univ Toronto Med J* 2021; **98**: 31–34.
- 104 Legido-Quigley H, Asgari N, Teo YY, et al. Are high-performing health systems resilient against the COVID-19 epidemic? *Lancet* 2020; **395**: 848–50.
- 105 Watts N, Amann M, Arnell N, et al. The 2020 report of The Lancet Countdown on health and climate change: responding to converging crises. *Lancet* 2021; **397**: 129–70.
- 106 Saleem SG, Ansari T, Ali AS, Fatima S, Rizvi MH, Samad MA. Risk factors for heat related deaths during the June 2015 heat wave in Karachi, Pakistan. *J Ayub Med Coll Abbottabad* 2017; **29**: 320–24.
- 107 Qiao Z, Guo Y, Yu W, Tong S. Assessment of short-and long-term mortality displacement in heat-related deaths in Brisbane, Australia, 1996–2004. *Environ Health Perspect* 2015; **123**: 766–72.
- 108 Centers for Disease Control and Prevention. People with certain medical conditions. 2022. https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fneed-extra-precautions%2Fgroups-at-higher-risk.html (accessed Aug 2, 2021).
- 109 Fadini GP, Morieri ML, Longato E, Avogaro A. Prevalence and impact of diabetes among people infected with SARS-CoV-2. *J Endocrinol Invest* 2020; **43**: 867–69.
- 110 Yang J, Hu J, Zhu C. Obesity aggravates COVID-19: a systematic review and meta-analysis. *J Med Virol* 2021; **93**: 257–61.
- 111 WHO. Responding to community spread of COVID-19: interim guidance. 2020. <https://apps.who.int/iris/handle/10665/331421> (accessed Sept 1, 2022).
- 112 Berrang-Ford L, Siders AR, Lesnikowski A, et al. A systematic global stocktake of evidence on human adaptation to climate change. *Nat Clim Chang* 2021; **11**: 989–1000.

Copyright © 2022 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.