

Kent Academic Repository

Full text document (pdf)

Citation for published version

Dolding-Smith, Jessica, Clement, A. and Mahoney, Patrick UNSPECIFIED In: UNSPECIFIED.

DOI

Link to record in KAR

<https://kar.kent.ac.uk/97685/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Analysing the occlusal wear of the hominins of Sima de los Huesos

JESSICA AM. DOLDING-SMITH^{1,2}, ANNA F. CLEMENT³, PATRICK MAHONEY², SIMON W. HILLSON³, JUAN-LUIS ARSUAGA⁴, JOSÉ MARÍA BERMÚDEZ DE CASTRO⁵ and MARÍA MARTINÓN-TORRES^{1,5}

¹Anthropology, UCL, ²School of Anthropology & Conservation, University of Kent, ³Institute of Archaeology, UCL, ⁴Centro de Investigación de la Evolución y Comportamiento Humanos, Universidad Complutense de Madrid, ⁵National Research Centre on Human Evolution, CENIEH

The occlusal surface of a tooth can provide evidence of past behaviour. In general terms, occlusal wear decreases through time, with industrialised modern humans having low occlusal wear because of the greater reliance upon processed foods. Neanderthals are thought to have exceptionally heavy occlusal wear, due to high mechanical loading from paramasticatory and masticatory behaviour.

Here, we compare occlusal wear of hominins from Sima de los Huesos (SH, n=19) to Neanderthals (n=21) and modern humans (Middle Palaeolithic, n=5; Upper/ Early Epi-Palaeolithic, n=26; Igloolik Inuit n=79; 19th-20th-century Madrileños individuals, CMH, n=14). The SH are thought to either be early Neanderthals or closely related to them, and share a number of cranial and dental traits with both Neanderthals (e.g. shovel-shaped incisors) and modern humans (e.g. frequent absence of the hypoconulid). Results deviate from the general and expected trend. The SH group had more wear on their upper I1 compared to Neanderthals ($p=0.022$), but not modern humans ($p>0.05$). We found that SH upper P3 is more worn than all other groups ($p<0.05$), except CMH ($p=0.201$), potentially pointing to masticatory behavioural differences. The mandibular dentition of the SH was significantly more worn compared to Neanderthals (C-M2, $p<0.05$), Inuit (I1-M2, $p<0.05$), and CMH (M2, $p=0.030$). It may be that SH were using their mandibular dentition less for paramasticatory activities or that the lower age range of the SH hominins used in the mandibular group meant that they generally had less wear.

Dirección General de Investigación of Spanish Ministerio de Economía y Competitividad (MINECO)/FEDER grant number: CGL2015-65387-C3-2-3-P and The Leakey Foundation (personal support of Dub Crook and Gordon Getty to author MM-T).