

Sleep architecture and obstructive sleep apnea in obese children with and without metabolic syndrome: a case control study

Shabnam Jalilolghadr¹ · Zohreh Yazdi² · Manoochehr Mahram¹ ·
Farkhondeh Babaei³ · Neda Esmailzadehha² · Hoormehr Nozari⁴ · Fatemeh Saffari²

Received: 4 July 2015 / Revised: 29 October 2015 / Accepted: 23 November 2015 / Published online: 28 December 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract

Purpose Obesity and biochemical parameters of metabolic disorders are both closely related to obstructive sleep apnea (OSA). The aim of this study was to compare sleep architecture and OSA in obese children with and without metabolic syndrome.

Methods Forty-two children with metabolic syndrome were selected as case group and 38 children without metabolic syndrome were matched for age, sex, and BMI as control group. The standardized Persian version of bedtime problems, excessive daytime sleepiness, awakenings during the night, regularity and duration of sleep, snoring (BEARS) and Children's Sleep Habits Questionnaires were completed, and polysomnography (PSG) was performed for all study subjects. Scoring was performed using the manual of American Academy of Sleep Medicine for children. Data were analyzed using chi-square test, *T* test, Mann–Whitney *U* test, and logistic regression analysis.

Results Non-rapid eye movement (NREM) sleep and N1 stage in the case group were significantly longer than the control group, while REM sleep was significantly shorter. Waking after sleep onset (WASO) was significantly different between two groups. Severe OSA was more frequent in the control group. Multivariate logistic regression analysis

showed that severe OSA (OR 21.478, 95 % CI 2.160–213.600; *P* = 0.009) and REM sleep (OR 0.856, 95 % CI 0.737–0.994; *P* = 0.041) had independent association with metabolic syndrome.

Conclusions Obese children with metabolic syndrome had increased WASO, N1 sleep stage, and severe OSA. But the results regarding sleep architecture are most likely a direct result of OSA severity. More longitudinal studies are needed to confirm the association of metabolic syndrome and OSA.

Keywords Metabolic syndrome X · Obesity · Obstructive sleep apnea · Sleep stages · Polysomnography

Introduction

Overweight and obesity are of the serious health problems of the twenty-first century. Prevalence of metabolic syndrome in overweight and obese children is higher than normal weight population [1]. Despite a lower prevalence of obesity in Asia compared with Europe, metabolic syndrome is progressed as a health problem in these countries [2]. Prevalence rates of metabolic syndrome in Turkey and Iran are unusually high, similar to the USA [2].

The risk of obstructive sleep apnea (OSA) is higher in obese children and adolescents. OSA occurs frequently and more severe in obese children compared with lean children [3]. In the study by Chay et al., 12.5 % of reportedly asymptomatic children were found to have OSA [4]. OSA is associated with active inflammatory mechanisms that are activated in obesity [5]. Obesity and OSA are associated with adverse cardiovascular, metabolic, and neuropsychological consequences [6]. Obesity is an important factor in the assessment of adverse metabolic outcomes in OSA [7].

✉ Fatemeh Saffari
drfa_saffari@yahoo.com

¹ Children Growth Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

² Metabolic Diseases Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

³ Qazvin University of Medical Sciences, Qazvin, Iran

⁴ Gaziantep Medical University, Gaziantep, Turkey