

2002

Ribbonfishes

John E. Olney


Follow this and additional works at: <https://scholarworks.wm.edu/vimsbooks>



Part of the [Aquaculture and Fisheries Commons](#)



THE LIVING MARINE RESOURCES OF THE
**WESTERN CENTRAL
ATLANTIC**



Volume 2 Bony fishes part 1 (Acipenseridae to Grammatidae)



AMERICAN
SOCIETY OF
ICHTHYOLOGISTS
AND
HERPETOLOGISTS



FOOD AND
AGRICULTURE
ORGANIZATION
OF THE
UNITED NATIONS

EUROPEAN
COMMISSION



FAO SPECIES IDENTIFICATION GUIDE FOR FISHERY PURPOSES
and
AMERICAN SOCIETY OF ICHTHYOLOGISTS AND HERPETOLOGISTS
SPECIAL PUBLICATION No. 5

THE LIVING MARINE RESOURCES OF THE WESTERN CENTRAL ATLANTIC

VOLUME 2

Bony fishes part 1 (Acipenseridae to Grammatidae)

edited by

Kent E. Carpenter

Department of Biological Sciences
Old Dominion University
Norfolk, Virginia, USA

with the support of the
American Society of Ichthyologists and Herpetologists
and the
European Commission

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS
Rome, 2002

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

ISBN 92-5-104825-8

All rights reserved. Reproduction and dissemination of material in this information product for educational or other non-commercial purposes are authorized without any prior written permission from the copyright holders provided the source is fully acknowledged. Reproduction of material in this information product for resale or other commercial purposes is prohibited without written permission of the copyright holders. Applications for such permission should be addressed to the Chief, Publishing Management Service, Information Division, FAO, Viale delle Terme di Caracalla, 00100 Rome, Italy or by e-mail to copyright@fao.org


© FAO 2002

TRACHIPTERIDAE

Ribbonfishes (dealfishes)

by J.E. Olney, Virginia Institute of Marine Science, USA

Diagnostic characters: Large-sized lampridiform fishes (to 2 m); body elongate, ribbon-like, compressed. In most species, body depth gradually decreasing from head to caudal peduncle. Upper jaw highly protrusible, maxilla broad; usually recurved, pointed teeth on jaws, vomer, and palatines; bones of head and jaws thin and fragile. Dorsal fin very long, extending along entire body length to tail; **anterior dorsal-fin elements consisting of 4 to 8 elongate, flexible spines that insert above eye**; total dorsal-fin elements 120 to 197; dorsal-fin rays bear strong lateral spinules that tend to interlock with adjacent soft rays and strengthen the fin. Anal fin absent. Caudal fin with 2 lobes; upper lobe sometimes upturned, conspicuous, and fan-like; total caudal-fin soft rays usually 13 to 18; usually 5 to 9 soft rays in lower fin lobe, some of which are elongate; usually 5 to 7 soft rays in the upper fin lobe, all of which are elongate in *Zu*. **Pelvic fins with 5 to 7 soft rays; often elongate in juveniles; sometimes lost at metamorphosis. Skin usually covered with bony, raised, bump-like tubercles. Scales absent, except for lateral-line scales that are tubular and bear sharp spines.** (Scalloped ribbonfish, *Zu cristatus*, with distinctive scalloped or wavy ventral margin, and possessing small deciduous scales). Total vertebrae, 62 to 102; thoracic vertebrae, 18 to 40. In trachipterids (and all lampridiforms), the anterior palatomaxillary ligament and palatine prong are absent; as a result, the maxilla is free to extend, along with premaxilla, well away from the ethmo-vomerine region during jaw protrusion. Other anatomical features of the trachipterids (and all lampridiforms): first dorsal-fin pterygiophore inserts anterior to first neural spine; elongate ascending processes of premaxilla and a large rostral cartilage insert into a front vault or cradle; mesethmoid posterior to lateral ethmoids. In trachipterids (and regalecids), the dorsal-, caudal-, and pelvic-fin rays bear spinules that project laterally; in trachipterids, the parapophyses of each thoracic vertebra are well developed, but ribs are lacking. **Colour:** head and body usually silver with oblique dusky bars or with dark spots; fins deep crimson-red.




Habitat, biology, and fisheries: Trachipterids are rare mesopelagic fishes that occur in all oceans. They consume pelagic crustaceans, small fishes, and squids. Eggs free-floating, large, and red. Very little is known of their habits and reproductive ecology. There is no fishery for the group.

Remarks: Trachipterids are distributed worldwide in tropical and temperate waters. There are approximately ten species in 3 genera (*Trachipterus*, *Zu*, and *Desmodema*), at least 3 of which are known from the area. There are a number of other nominal species whose validity is not widely recognized by all authors. In addition, there may be undescribed species in the area. The family is in need of revision.

Similar families occurring in the area

Regalecidae: also lacking anal fin, but with more dorsal-fin soft rays (260 to 412 versus 120 to 200), and attaining a far larger size.

All other lampridiform families possess an anal fin.



Regalecidae

Key to the species of Trachipteridae occurring in the area

- 1a. Caudal fin without 2 lobes and not sharply upturned; no long spines or bony tubercles along ventral edge of tail; dorsal fin with 120 to 124 elements *Desmodema polystictum*
- 1b. Caudal fin with 2 lobes, the upper lobe sharply upturned; ventral edge of tail bears long spiny plates or bony tubercles; dorsal fin usually with more than 124 elements → 2
- 2a. Posterior portion of lateral line runs along ventral edge of tail as a series of sharp spines that point in alternating directions; wavy or scalloped ventral body margin; dorsal fin with less than 150 elements *Zu cristatus*
- 2b. Posterior portion of lateral line runs well above the ventral edge of tail; lateral line spines project laterally, and do not point in alternating directions; wavy or scalloped ventral body margin; dorsal fin with more than 150 elements. *Trachipterus arcticus*

List of species occurring in the area

- Desmodema polystictum* (Ogilby, 1898). To about 100 cm. Mesopelagic in all oceans.
- Trachipterus arcticus* (Brünnich, 1788). To about 250 cm. Mesopelagic in all oceans.
- Zu cristatus* (Bonelli, 1819). To about 120 cm. Mesopelagic in all oceans.

References

Olney, J.E. 1984. Lampridiformes: development and relationships. In *Ontogeny and systematics of fishes*, edited by H.G. Moser, W.J. Richards, D.M. Cohen, M.P. Fahay, A.W. Kendall, Jr, and S.L. Richardson. American Society of Ichthyologists and Herpetologists, Publication 1, pp. 368-379.

Olney, J.E., G.D. Johnson, and C.C. Baldwin. 1993. Phylogeny of lampridiform fishes. *Bull. Mar. Sci.*, 52:137-169.

Palmer, G. 1961. The dealfishes (Trachipteridae) of the Mediterranean and north-east Atlantic. *Bull. Brit. Mus. (Nat. Hist.) Zool.*, 7(7):337-351.

Robins, C.R., G.C. Ray and J. Douglas. 1986. *A field guide to Atlantic coast fishes North America*. Boston, Houghton Mifflin Co., 354 p.