

ORIGINAL RESEARCH ARTICLE

Discussion on the value of 48h ambulatory electrocardiogram

Jining Zhou, Jintao Xiang*, Xin Lai, Fang Wang, Xiaochun Mu, Limei Hu, Lijuan Ye, Xiaoqing Li, Yuanxiu Chen

Department of Cardiology, Renmin Hospital of Wuhan University, Wuhan 430060, Hubei, China. E-mail: xjt0909@yeah.net

ABSTRACT

Objective: The statistical number of cases was 73 times. 66 patients entered the study, including 31 males and 35 females, aged from 17 to 86 years. Firstly, the accurate big data analysis was carried out on 66 cases of ambulatory ECG, and then the corresponding time that can be compared and analyzed before and after 24h was determined through the time (T) RR interval scatter diagram. The changes of 24h ECG scatter diagram were observed, and the total number of heart beats and the number of premature beats were counted by the scatter diagram technique, and then compared and analyzed according to the type of arrhythmia. **Results:** A frequent or occasional premature beats: 52 patients (59 times), 26 males and 26 females, aged (56.1 ± 15.4) years; the effective analysis duration was (22.40 ± 1.19) H. There was a significant correlation between the number of heart beats before and after treatment ($r=0.934$, $p=0.000$). b. Persistent arrhythmia: 4 cases of persistent atrial fibrillation (referred to as atrial fibrillation), 1 case of frequent atrial and ventricular concurrent arrhythmia, 2 males and 3 females, aged (68.4 ± 9.4) years; the effective analysis duration was (22.68 ± 0.74) H. The t-RR scatter plot and Lorenz RR scatter plot of these patients had self-similarity before and after 24h. There was no significant difference in the total heart beat number before and after 24h[(111796 ± 16439) vs (111262 ± 16421), $p=0.624$], and the total heart beat number before and after 24h was significantly correlated ($r=0.991$, $p=0.001$); the qualitative diagnosis of long interval in 3 patients with atrial fibrillation was consistent 24 hours before and after diagnosis. c. Paroxysmal arrhythmia: 9 cases, 3 cases of paroxysmal atrial flutter, 5 cases of paroxysmal atrial fibrillation, and 1 case of paroxysmal vertical separation of sinus node function. There were 3 males and 6 females. The age was (71.2 ± 12.7) years. The effective analysis duration was (22.67 ± 0.74) H. **Conclusion:** 24h ambulatory ECG can meet the requirements for patients with frequent or occasional premature beats and persistent arrhythmias, while 48h ambulatory ECG may be necessary for patients with paroxysmal arrhythmias.

Keywords: cardiology; dynamic ECG; big data analysis; ECG scatter diagram; arrhythmia; 48-hour ECG

1. Introduction

The earth, where human beings live, rotates once every 24 hours. When facing the sun, it is day.

When departing from the sun, it is night. It is the law of human life to work at sunrise and rest at sunset. The heart rhythm that is consistent with the change of human life law also shows regular changes. Therefore, 24h Holter detection can reflect the pe-

ARTICLE INFO

Received: April 16, 2021 | Accepted: May 26, 2021 | Available online: July 13, 2021

CITATION

Zhou J, Xiang J, Lai X, Wang F, Mu X, Hu L, Ye L, Li X, Chen Y. Discussion on the value of 48h ambulatory electrocardiogram. Cardiac and Cardiovascular Research 2021; 2(2): 8 pages.

COPYRIGHT

Copyright © 2021 by author(s) and Asia Pacific Academy of Science Pte. Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), permitting distribution and reproduction in any medium, provided the original work is cited.

riodic change of human heart rate or rhythm^[1]; the research shows that the occurrence of arrhythmia also shows a significant diurnal distribution^[2]. Therefore, the author speculates that understanding the change of a 24h overall heart rhythm of patients means that we can understand all the 24h overall heart rhythm; however, some arrhythmias may not be sporadic or frequent in a change cycle within 24h due to some sudden factors, and it may take longer monitoring to capture the sudden events. The author divided the 48h ambulatory electrocardiogram into two 24h, compared the rhythm of the two 24h monitoring, and discussed the detection and significance of the 48h ambulatory electrocardiogram.

2. Data and methods

2.1. Observation object

The people's Hospital of Wuhan University began in January, 2021. The outpatient doctors requested 48h or 72h dynamic ECG testing according to the needs of patients' conditions. The 48h or 72h dynamic ECG testing was completed by the Department of cardiac function of the hospital. The instrument was completed by the intelligent dynamic ECG analysis system produced by Hangzhou Baihui Medical Equipment Co., Ltd. The recording time could be set by computer software according to the needs at that time, when the patient has completed the corresponding recording time, import it into the computer through the corresponding software. The ECG doctor on duty will issue a report after the intelligent analysis system + manual analysis and correction, and archive the original recorded data for further analysis.

From January 4, 2021 to September 3, 2021, 59 patients (including 1 patient who performed 48h ambulatory ECG recording twice at different times, calculated as 2 patients) performed 48h ambulatory ECG recording, and 7 patients performed 72h ambulatory ECG recording. In order to compare the ECG records before and after 24h (48h in total), the author divided the 72h ambulatory ECG records into


three 24h periods (A, B, C). A and B, B and C were used to compare the changes of ECG records before and after 24h; that is, 7 patients were recorded 2 times per person, so the number of cases included in the analysis and statistics was 73 times; the number of cases was 66. 66 patients, 31 males and 35 females; the youngest is 17 years old and the oldest is 86 years old.

2.2. Analysis method

In order to analyze the 48h heart beat in detail, the author adopted modern ECG big data analysis methods: a. accuracy of 48h or 72h whole stroke: time (t) RR interval scatter plot and reverse technology combined with Lorenz RR scatter plot analysis technology can achieve accurate heart beat in general cases^[3-4], and Lorenz RR scatter plot + superimposed heart beat technology is required in some special cases^[5]; then refer to "histogram and histogram inversion technology" to accurately determine the heart rate of the minimum (fastest) and maximum (slowest) cardiac cycles. B. Determination and comparison of the available time course for analysis in the first 24 hours and the next 24 hours: for the patients who performed 48h or 72h ambulatory ECG detection, when they came to the hospital to remove the recording box after the second 24 hours or the third 24 hours, most of them failed to achieve the full 24h recording. In order to make the heart beats in the first 24 hours and the next 24 hours comparable, the author chose the time course (length) of ECG actually recorded in the next 24 hours as the standard available time course for analysis, Then, the t-RR scatter chart technology is used to take the same time course in the corresponding time frame in the first 24 hours as the time course available for analysis in the next 24 hours for statistical measurement. The detailed method is shown in Figure 1.

For paired atrial premature beats (abbreviated as atrial premature beats) and atrial tachycardia (abbreviated as atrial tachycardia), if it is difficult to measure from the Lorenz RR scatter diagram, the occurrence time can be read from the "atrial premature events" for 24h counting.

2.3. Statistical grouping



The previous figure is a t-RR scatter diagram, a: the duration of the whole bar is set to 48h, and the actual recording duration is 47h19min (frame timing duration); article B: the time course available for analysis (23h20min) recorded within the next 24 Hours, the starting time of which is 07-24, 09:16:04 (corresponding to the starting time of article C 07-23, 09:16:04, and the time course determined in the inner circle of the box is also 23h20min). Note that the starting time of article a is 07-23, 09:16:55 (the time indicated in a'), which lags behind 16:04 by 51s. It is speculated that 16:04 May be the power on time, and 16:55 Is the time recorded by ECG; article C: the available time for analysis recorded within the first 24 Hours, starting at 07-23, 09:16:04 (consistent with the starting time of article b); the figure below is Lorenz RR scatter diagram. A', B', C' respectively correspond to the figure generated by the rectangular box in figure a, B and C above; C'' is the "atrial premature" scattergram extracted from C'. The total number of heart beats can be calculated by framing the Lorenz RR scattergram. The total number of heart beats of a', B', C' and C'' are 139881, 69835, 67974 And 234 Respectively (atrial premature number includes all ectopic atrial beats, such as the number of atrial beats of paired atrial premature and atrial tachycardia). A', B', C' in the figure have obvious self-similarity, and the accompanying arrhythmia is atrial parallel arrhythmia

Figure 1. Available analysis time history and calculation method for 24h before and 24h after confirmation.

After accurate analysis and statistical measurement of all cases, the author found that for the convenience of scientific analysis, the following categories are more appropriate: ① patients with frequent or occasional premature beats are the first category; b. Persistent atrial fibrillation (AF) is the second type; c. Paroxysmal arrhythmias, such as paroxysmal atrial fibrillation and atrial flutter (referred to as atrial flutter) are classified into the third category. The diagnosis of types of atrial premature beats and ventricular premature beats (referred to as ventricular premature beats) is based on the graphic characteristics of big data in clinical ECG scatter^[6-7]. Longitudinal separation of sinus node

function and sinus escape reference^[8-9].

2.4. Statistical treatment

SPSS statistical software was used for statistical analysis. The data of measurement data are expressed in $X \pm s$, and paired t-test is adopted; if the distribution is skewed, it is expressed by median + quartile[m (P25, p75)], and rank sum test is used; the correlation was tested by paired samples. The difference was significant ($p < 0.05$).

3. Results

3.1. Type I: frequent or accidental prema-

ture beats

The recording time of 45 cases was 48h, and that of 7 cases was 72h (combined before and after 48h, 2 cases were calculated), a total of 52 patients (59 cases). Among them, there were 26 males and 26 females, the age was (56.1 ± 15.4) years old, and the duration was (2769.67 ± 496.60) min (mean 46.15h); the effective analysis duration was (1345.75 ± 71.27) min (mean 22.40h). All patients had premature beats, of which the least was 1 atrial premature beat and the most was 14410 ventricular premature beats. There were 37 cases of atrial concurrent rhythm and 2 cases of borderline concurrent rhythm; there were 1 case of reentrant atrial premature beats and 6 cases of atrial premature beats of undetermined nature (the number of atrial premature beats was too small to form a characteristic scatter diagram); there were 14 cases of ventricular concurrent rhythm, 5 cases of

reentrant ventricular tachycardia, 13 cases of undetermined ventricular tachycardia (the number of ventricular tachycardia was small and did not constitute a characteristic scatter plot), and 32 cases of both atrial and ventricular tachycardia.

The t-RR scatter plot and Lorenz RR scatter plot of this kind of patients before and after 24h have self-similarity (the shape of the formed graph is similar), as shown in **Figure 1**. There was no significant difference in the total number of heart beats, atrial premature beats, paired atrial premature beats, atrial tachycardia and ventricular premature beats between the first 24 hours and the second 24 hours in 59 patients ($P > 0.05$); see **Table 1**. There was a significant correlation between the total heart beat number before and after 24h ($r = 0.934$, $p = 0.000$).

Table 1. Comparison of total heart beats, atrial premature beats, atrial tachycardia array and ventricular premature beats recorded in 24h before and after 48h Holter recording (n=59)

Category	Total heart beats	Room breakfast / piece	Paired room breakfast / time	Atrial tachycardia / time	Room breakfast / piece
First 24h	92256±12151	26(5,101)	0(0,2)	0(0,1)	2(0,20)
After 24h	92638±11978	27(5,97)	0(0,1)	0(0,1)	1(0,10)
P	0.507	0.509	0.294	0.924	0.927

3.2. Type II: persistent arrhythmia

Among 4 patients with persistent atrial fibrillation, 1 patient with frequent atrial and ventricular fibrillation recorded a 48ht RR scatter plot like that of persistent atrial fibrillation (heart rate is abso-

lutely uneven), as shown in **Figure 2**, so 5 patients were included in the observation. There were 2 males and 3 females with an average age of (68.4 ± 9.4) years; the recording time course was (2800.40 ± 44.70) min (mean 46.67h), and the analysis time course was (1360.6 ± 44.66) min (mean 22.68h).




Figure 2. T-RR scatter plot of sinus rhythm with frequent atrial and ventricular concurrent rhythms.

The patient, a 79-year-old male, had sinus rhythm, atrial Union and ventricular union repeat-

edly and alternately in 48h, resulting in absolute irregularity of ventricular rate. Therefore, the scatter

pattern formed in the above graph is similar to that of atrial fibrillation; the following figure is an enlarged t-RR scatter diagram (the time length is relaxed, but the length of the vertical axis remains unchanged) of the 1H scatter diagram framed in the box in the previous figure. The “thin line” in the middle is a sinus rhythm point, forming a thin and straight “curve”, and its variability is reduced.

The t-RR scatter plot and Lorenz RR scatter plot of this kind of patients also have self-similarity. For example, the curve shape of t-RR scatter plot at 24h before and after Figure 2 remains unchanged. There was no significant difference in the total heart beats recorded in the five cases before and after 24 hours [(111796 ± 16439) vs (111262 ± 16421), $p=0.624$], and there was also a significant correlation between the total heart beats recorded in the five cases before and after 24 hours ($r=0.991$, $p=0.001$). There were 3 patients with persistent atrial fibrillation who had long intervals. The frequency of long intervals was the same in 2 patients before and after 24h (>2s were 7, 7 and 1, 1 respectively), while the number of long intervals in the first 24h was more than that in the next 24h (>2s, 124 times; >2.5s, 6 times vs >2s, 69 times; >2.5s, 1 time).

3.3. Category III: paroxysmal arrhythmia

Nine patients with paroxysmal arrhythmia, including 3 males and 6 females, aged (71.2 ± 12.7) years, were recorded for (2799.67 ± 45.09) min (mean 46.66h) and analyzed for (1360.00 ± 44.32) min (mean 22.67h). Among them, there were 3 cases of paroxysmal atrial flutter, 5 cases of paroxysmal atrial fibrillation, and 1 case of paroxysmal vertical separation of sinus node function. Basal heart rate: sinus rhythm combined with reentrant atrial tachycardia in 3 cases and atrial complication in 4 cases; one case had both reentrant atrial tachycardia and concurrent rhythmic atrial tachycardia.

The self-similarity of t-RR scatter plot and Lorenz RR scatter plot of this kind of patients disappeared 24 hours before and after treatment, as shown in Figure 3 and 4. There was no significant difference in the total number of heart beats in 9

cases before and after 48h [(100709 ± 23017) vs (99429 ± 17909), $p=0.823$]. There was a correlation between the total number of heart beats before and after 24h ($r=0.695$, $p=0.038$).

4. Discussion

4.1. Findings of this study


a. Through the 24h performance of common arrhythmia atrial premature and ventricular premature of 59 patients in this study, it is found that the morphological characteristics of big data scatter diagram have not changed and have significant “self similarity”, which is the “chaotic” performance of cardiac rhythm and reflects the “periodic phenomenon” of beating^[10], and it is also found in the 48h statistical data, There was no significant difference in total heart beat number, atrial premature beats, paired atrial premature beats, atrial tachycardia and the total number of ventricular premature beats before and after 24h, and there was a significant correlation between total heart beat number before and after 24h ($r=0.934$, $p=0.000$), indicating that for the detection of common arrhythmias (atrial premature beats and ventricular premature beats), 24h long-range ECG examination was enough to find their apparent characteristics, and 48h long-range ECG examination was not necessary. B. In this study, the ECG big data of patients with persistent arrhythmia (4 cases of atrial fibrillation, 1 case of frequent atrial and ventricular Union) were analyzed for 48 hours. It was also found that the morphology of the scatter plot before and after 24 hours was also self-similar, and there was no difference in the total number of heart beats before and after, and it was highly similar ($r=0.991$, $p=0.001$). It also showed that the long-range ECG examination for 24 hours in patients with this kind of arrhythmia was enough, and the examination for 48 hours was not necessary. C. Nine patients with paroxysmal arrhythmia were examined by dynamic electrocardiogram for 48 hours. It was found that the shape of the scatter diagram had changed and its self-similarity had disappeared, indicating that the nature of their heart rhythm had changed; according to the statistical data,

these 9 cases all had paroxysmal arrhythmias of different degrees, and each had its own nature and characteristics. However, the 24-hour overall rhythm characteristics could not show the nature and characteristics of their occurrence. Therefore, it is necessary to monitor the ECG of these patients for 48 hours.

4.2. Characteristics and long-term record of arrhythmia

Atrial premature and ventricular premature are the most common arrhythmias. In the 24-hour human activity cycle, the time length can provide the opportunity or opportunity for the occurrence of atrial premature and ventricular premature. Therefore, 24-hour ECG examination is enough to make qualitative or quantitative diagnosis for patients; for patients with persistent arrhythmia, there is no qual-


itative change in the nature of 24h and 48h duration and the number of heart beats, so only 24h ECG monitoring is required to detect problems, and 48h examination is not necessary; the occurrence and termination of paroxysmal arrhythmia may be greatly affected by sudden factors, and the length of 24h is not enough to reflect the characteristics of its occurrence, development and cessation, so it is necessary to monitor ECG for 48h or longer. So, before applying for the long-term ambulatory ECG examination, how do clinicians determine whether a patient needs 24h or 48h examination? This requires clinicians to ask the patient's symptoms and medical history in detail, observe the conventional 12 lead ECG, preliminarily determine the nature of arrhythmia, and then select the inspection duration of long-range ECG. Only in this way can we make rational use of resources and provide the best inspection means for patients.



The t-RR scatter gram (comparison of anterior and posterior segments) of 9 Patients before and after 24 Hours were significantly different, and the self-similarity disappeared. A female patient, 84 Years old, had

atrial flutter in the first 24 Hours, mainly passing down the ventricle at 2 : 1 And 4 : 1; in the next 24 Hours, the first 11 Hours were atrial flutter, and the last 10 Hours and 36 Minutes were sinus rhythm and atrial concurrent rhythm. B and a were the same patient, which were recorded after 9 Days of atrial flutter treatment. In the first 24 Hours, the first 15 Hours and 50 Minutes were sinus rhythm with occasional atrial fibrillation; the later stage was associated with frequent reentrant atrial tachycardia; in the next 24 Hours, frequent reentrant atrial premature beats continued to occur, and on this basis, short bursts of atrial flutter occurred 4 Times. The total duration was 10h42min, and then turned to sinus rhythm (during which a short atrial flutter occurred once, lasting for 9min). C patient was a 68 Year old male with sinus rhythm + borderline rhythm + borderline escape at the beginning of the first 24 Hours, lasting for 6h32min, followed by atrial tachycardia with different degrees of atrioventricular block; after that, the rate of atrial tachycardia was 2 : 1 Downward; after 24 Hours, the continuous constant atrial tachycardia was 2:1 Downward. D the patient was a 69-year-old male, who had frequent atrial tachycardia within the first 24 Hours, and the atrial tachycardia did not pass down; short paroxysmal atrial fibrillation occurred 6 Times, lasting from 23S to 1min12s; the whole course of 24 Hours was sinus rhythm. E patient was an 82-year-old male with sinus rhythm and frequent atrial fibrillation in the first 24 Hours; on the basis of atrial Union, paroxysmal atrial flutter / fibrillation occurred 24 Hours later, lasting for 1 Hour and 43 Minutes. F patient was a 74 Year old female with sinus rhythm and atrial fibrillation at 48 Hours; atrial flutter / fibrillation occurred at the junction of the first 24 Hours and the second 24 Hours, lasting for 4h26min. G patient was a 42 Year old male, whose basic rhythm was sinus rhythm with reentrant atrial tachycardia at 48 Hours. In the first 24 Hours, there were 17 Short bursts of atrial fibrillation, all lasting for about 1 Minute; atrial fibrillation with a long duration occurred 24 Hours later, lasting for a total of 5h9min. H female, 61 Years old, the basic rhythm of 48h was sinus rhythm with atrial Union, and atrial premature beats did not pass down; in the first 24 Hours, there were two periods of frequent atrial fibrillation, the duration of which was 10h11min and 3H respectively. In the later period, atrial fibrillation occurred on this basis (the duration of which was 1h22min); after 24 Hours, sinus rhythm combined with occasional atrial fibrillation. I a 77 Year old female patient whose basic rhythm was sinus rhythm with longitudinal separation of sinus node function and sinus escape; there were 6 Attacks in the first 24 Hours, and the duration was long. There were also 6 Attacks in the second 24 Hours, but the duration was short.

Figure 3. T-RR scatter diagram automatically drawn by computer in 48h ECG recording of patients with paroxysmal arrhythmia.



A, B, C and D are the 24h Lorenz RR scatter plots of the same patient, respectively. It can be seen that their self-similarity disappears, indicating that one of them has arrhythmia or has different degrees

Figure 4. Front and rear 24h Lorenz RR scatter plots of 2 Patients (B and I), selected from the t-RR scatter plot in Figure 3T

Conflict of interest

The authors declare no conflict of interest.

References

1. Zhang J, Chatham JC, Young ME. Circadian regulation of cardiac physiology: Rhythms that keep the heart beating. *Annurev Physiol* 2020; 28: 79.
2. Irwin JM, McCarthy EA, Wilkinson WE, et al. Circadian occurrence of symptomatic paroxysmal supraventricular tachycardia in untreated patients. *Circulation* 1988; 77(2): 298.
3. Xiang J. Timed RR-interval scatter plots and reverse technology. *Current Medical Science* 2020; 40(6): 1191
4. Xiang J. Lorenz RR scatter plot reverse technique and scatter plot block extraction technique. In: Xiang J, Jing Y. *Clinical electrocardiography*. Wuhan: Hubei Science and Technology Press 2016; p. 85–98.
5. Xiang Q, Yang Y, Liu W, et al. Scatter plot and ECG characteristics of a case of atypical second-degree type I atrioventricular block with atrioventricular reentrant atrial echo and reentrant tachycardia. *Chinese Journal of Cardiac Pacing and Electrophysiology* 2021; 35(4): 368
6. Xiang J. Lorenz RR scatter plot characteristics and electrophysiological significance of big data atrial parallel rhythm. In: Xiang J, Jing Y. *Clinical electrocardiography*. Wuhan: Hubei Science and Technology Press 2016; p. 143–148.
7. Xiang J. Lorenz RR scatter plot characteristics and electrophysiological significance of big data ventricular parallel rhythm. In: Xiang J, Jing Y. *Clinical electrocardiography*. Wuhan: Hubei Science and Technology Press 2016; p. 153–159
8. Xiang J, Li X, Chen Y. Separation of sinus node function revealed by ECG scatter gram. *Chinese Journal of Cardiac Pacing and Electrophysiology* 2013; 27(2): 101
9. Xiang J, Li X, Chen Y. A case of sinus rhythm complicated with sinus escape and premature beat was diagnosed by time RR interval scatter plot and reverse technique. *Chinese Journal of Cardiac Pacing and Electrophysiology* 2013; 27 (5): 457
10. Li F. Combination of chaos theory and ECG. *Chinese Journal of Cardiac Pacing and Electro-physiology* 2013; 17(6): 407.