

# Cardiac magnetic resonance imaging as screening for cardiac sarcoidosis or not?

Vasileios Kouranos<sup>1,3</sup>, Rakesh Sharma<sup>2,3</sup>

<sup>1</sup>Interstitial Lung Disease Unit, Royal Brompton Hospital, London, United Kingdom

<sup>2</sup>Department of Cardiology, Royal Brompton Hospital, London, United Kingdom

<sup>3</sup>Imperial College London, London, United Kingdom

## Related article

by Kyperska et al.

### Correspondence to:

Vasileios Kouranos, MD, PhD,  
Royal Brompton Hospital,  
Sydney Street, London SW3 6NP,  
United Kingdom,  
e-mail: v.kouranos@rbht.nhs.uk,  
vasileios.kouranos@nhs.net

Copyright by the Author(s), 2022

DOI: 10.33963/KPa2022.0229

### Received:

September 28, 2022

### Accepted:

September 28, 2022

### Early publication date:

September 30, 2022

In the last 2 decades, the wide use of cardiac magnetic resonance imaging (CMR) has revolutionized the diagnostic approach to patients with suspected cardiac sarcoidosis (CS). Several case series have shown that CMR alone was superior to the former (2007) Japanese Ministry of Health and Welfare (JMHW) guidelines on detecting myocardial involvement in patients with systemic sarcoidosis [1–8]. Furthermore, the identification of late gadolinium enhancement (LGE) on CMR; a marker for myocardial damage associated with cardiac sarcoidosis, was strongly associated with major adverse outcomes during follow-up [1–5]. As a result, CMR was considered a major diagnostic criterion in the Heart Rhythm Society expert consensus statement for diagnosis of CS in 2014 [9]. In that document, LGE on CMR in a pattern compatible with CS in patients with extra-cardiac sarcoidosis was consistent with at least probable cardiac involvement, when other causes were reasonably excluded. This was also acknowledged in the revised JMHW guidelines for diagnosis of CS [10].

While the role of CMR as a diagnostic tool has been widely accepted as the gold standard, its role as a screening tool in the general sarcoidosis population remains controversial. In the latest American Thoracic Society clinical practice guidelines in sarcoidosis, CMR was not recommended as part of the screening strategy [11]. Baseline evaluation of the general sarcoidosis population with cardiac symptoms and a 12-lead electrocardiogram (ECG) remain the recommended screening strategy [11]. Although CMR is expected to

detect a higher prevalence of myocardial damage in the sarcoidosis population, clinical implications for an asymptomatic patient without rhythm or morphological abnormalities remain unclear. On the other hand, CMR was strongly recommended as the first-choice imaging modality in patients with suspected CS patients with cardiac symptoms and/or ECG abnormalities.

In this issue of the *Kardiologia Polska (Polish Heart Journal)*, we read with interest the article regarding the role of CMR in the asymptomatic sarcoidosis population, which aims to shine light on the use of CMR in a subclinical setting [12]. In a cohort of 55 sarcoidosis patients with evidence of extra-cardiac disease, CMR managed to detect only 6% of cardiac involvement when used as a screening tool. None of the patients had cardiac symptoms, while all patients had no significant ECG abnormalities or morphological abnormalities on echocardiography or CMR at baseline. In addition, none of the patients was found to have elevated cardiac biomarkers such as troponin or BNP. Therefore, CS was an incidental finding in this population. No follow-up data were provided to evaluate the prognostic role of CS in this group of patients. A similar prevalence of CS (13%) was reported in a cohort of 61 Japanese sarcoidosis patients without any cardiac manifestations of CS [13]. In that study, the detection of LGE was not associated with any adverse events during follow-up. In a larger cohort of patients with extra-cardiac disease demonstrated by biopsy, CMR detected approximately 20% of CS in a subclinical

setting (lack of cardiac symptoms and/or ECG abnormalities) [3]. However, LGE on CMR was not associated with major adverse events during follow-up in the patients with subclinical disease in that study [3].

Current literature indicates that CMR has high sensitivity and specificity in identifying patients with CS and particularly those at higher risk of major arrhythmias during follow-up. CMR has clear superiority in evaluating cardiac morphology, in particular myocardial fibrosis, which has greater prognostic value than any other imaging modality in current use. In the study by Kouranos et al. [3], CMR was found to be superior to conventional tests such as 12-lead ECG, Holter monitoring, echocardiography, or a combination of those. What becomes crucial is identifying which suspected sarcoidosis patients should undergo CMR. Kyperska et al. [12] showed that it is unlikely to detect CS in the asymptomatic population without ECG or echocardiographic abnormalities, supporting the current guideline recommendations. However, we should acknowledge that the authors performed a comprehensive baseline assessment with all conventional tests such as 12-lead ECG, Holter monitoring, and echocardiography, as well as biomarker testing outside the guideline recommendations.

We support the measurement of serum biomarkers, such as NT-proBNP, and echocardiography for screening the sarcoidosis population. Modern echocardiographic techniques, such as speckle tracking, have been shown to be more sensitive in detecting myocardial involvement than conventional echocardiographic modalities. Such an approach in addition to the current strategy of assessment of cardiac symptoms and ECG abnormalities should be able to identify a higher number of patients suspected of CS. BNP and NT-proBNP are associated with both left and right ventricular strain, even at an early stage, and have been linked with CS [14]. Speckle tracking analysis is an echocardiographic technique that measures myocardial deformation and may be able to detect cardiac involvement of sarcoidosis earlier than conventional echocardiographic modalities [15]. In addition, regional wall motion abnormalities appear to be strongly associated with CS, which should be part of the routine echocardiographic assessment [15]. Finally, there was a weak association between elevated angiotensin converting enzyme levels and CS detection in the latest study presented in this issue of the *Kardiologia Polska* [12]. This would raise suspicion as to whether the clinical impression of disease activity should be included in the screening strategy, and it should indicate the performance of CMR as a screening test. Further studies are warranted to identify the optimal screening strategy for CS. The association of CS with sudden cardiac death and high morbidity and mortality requires early and accurate diagnosis.

### Article information

**Conflict of interest:** None declared.

**Funding:** None.

**Open access:** This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at [kardiologiapolska@ptkardio.pl](mailto:kardiologiapolska@ptkardio.pl).

### REFERENCES

- Patel MR, Cawley PJ, Heitner JF, et al. Detection of myocardial damage in patients with sarcoidosis. *Circulation*. 2009; 120(20): 1969–1977, doi: [10.1161/CIRCULATIONAHA.109.851352](https://doi.org/10.1161/CIRCULATIONAHA.109.851352), indexed in Pubmed: [19884472](https://pubmed.ncbi.nlm.nih.gov/19884472/).
- Greulich S, Deluigi CC, Gloekler S, et al. CMR imaging predicts death and other adverse events in suspected cardiac sarcoidosis. *JACC Cardiovasc Imaging*. 2013; 6(4): 501–511, doi: [10.1016/j.jcmg.2012.10.021](https://doi.org/10.1016/j.jcmg.2012.10.021), indexed in Pubmed: [23498675](https://pubmed.ncbi.nlm.nih.gov/23498675/).
- Kouranos V, Tzelepis GE, Rapti A, et al. Complementary role of CMR to conventional screening in the diagnosis and prognosis of cardiac sarcoidosis. *JACC Cardiovasc Imaging*. 2017; 10(12): 1437–1447, doi: [10.1016/j.jcmg.2016.11.019](https://doi.org/10.1016/j.jcmg.2016.11.019), indexed in Pubmed: [28330653](https://pubmed.ncbi.nlm.nih.gov/28330653/).
- Kandolin R, Lehtonen J, Airaksinen J, et al. Cardiac sarcoidosis: epidemiology, characteristics, and outcome over 25 years in a nationwide study. *Circulation*. 2015; 131(7): 624–632, doi: [10.1161/CIRCULATIONAHA.114.011522](https://doi.org/10.1161/CIRCULATIONAHA.114.011522), indexed in Pubmed: [25527698](https://pubmed.ncbi.nlm.nih.gov/25527698/).
- Ise T, Hasegawa T, Morita Y, et al. Extensive late gadolinium enhancement on cardiovascular magnetic resonance predicts adverse outcomes and lack of improvement in LV function after steroid therapy in cardiac sarcoidosis. *Heart*. 2014; 100(15): 1165–1172, doi: [10.1136/heartjnl-2013-305187](https://doi.org/10.1136/heartjnl-2013-305187), indexed in Pubmed: [24829369](https://pubmed.ncbi.nlm.nih.gov/24829369/).
- Mehta D, Lubitz SA, Frankel Z, et al. Cardiac involvement in patients with sarcoidosis: diagnostic and prognostic value of outpatient testing. *Chest*. 2008; 133(6): 1426–1435, doi: [10.1378/chest.07-2784](https://doi.org/10.1378/chest.07-2784), indexed in Pubmed: [18339784](https://pubmed.ncbi.nlm.nih.gov/18339784/).
- Smedema JP, Snoep G, van Kroonenburgh MPG, et al. Cardiac involvement in patients with pulmonary sarcoidosis assessed at two university medical centers in the Netherlands. *Chest*. 2005; 128(1): 30–35, doi: [10.1378/chest.128.1.30](https://doi.org/10.1378/chest.128.1.30), indexed in Pubmed: [16002912](https://pubmed.ncbi.nlm.nih.gov/16002912/).
- Diagnostic standard and guidelines for sarcoidosis—2006 [in Japanese]. *Jpn J Sarcoidosis and Granulomatous Disorders* 2007; 27: 89–102.
- Birnie DH, Sauer WH, Bogun F, et al. HRS expert consensus statement on the diagnosis and management of arrhythmias associated with cardiac sarcoidosis. *Heart Rhythm*. 2014; 11(7): 1305–1323, doi: [10.1016/j.hrthm.2014.03.043](https://doi.org/10.1016/j.hrthm.2014.03.043), indexed in Pubmed: [24819193](https://pubmed.ncbi.nlm.nih.gov/24819193/).
- Terasaki F, Azuma A, Anzai T, et al. JCS 2016 Guideline on Diagnosis and Treatment of Cardiac Sarcoidosis. *Circ J*. 2019; 83(11): 2329–2388, doi: [10.1253/circj.CJ-19-0508](https://doi.org/10.1253/circj.CJ-19-0508), indexed in Pubmed: [31597819](https://pubmed.ncbi.nlm.nih.gov/31597819/).
- Crouser ED, Maier LA, Wilson KC, et al. Diagnosis and detection of sarcoidosis. An official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2020; 201(8): e26–e51, doi: [10.1164/rccm.202002-0251ST](https://doi.org/10.1164/rccm.202002-0251ST), indexed in Pubmed: [32293205](https://pubmed.ncbi.nlm.nih.gov/32293205/).
- Kyperska K, Kuchynka P, Slovakova A, et al. Role of magnetic resonance in the detection of cardiac involvement in patients with newly diagnosed extracardiac sarcoidosis: Single center experience. *Kardiologia Polska*. 2022; 80(9): 897–901, doi: [10.33963/KP.a2022.0163](https://doi.org/10.33963/KP.a2022.0163), indexed in Pubmed: [35775447](https://pubmed.ncbi.nlm.nih.gov/35775447/).
- Nagai T, Kohsaka S, Okuda S, et al. Incidence and prognostic significance of myocardial late gadolinium enhancement in patients with sarcoidosis without cardiac manifestation. *Chest*. 2014; 146(4): 1064–1072, doi: [10.1378/chest.14-0139](https://doi.org/10.1378/chest.14-0139), indexed in Pubmed: [24853830](https://pubmed.ncbi.nlm.nih.gov/24853830/).
- Date T, Shinozaki T, Yamakawa M, et al. Elevated plasma brain natriuretic peptide level in cardiac sarcoidosis patients with preserved ejection fraction. *Cardiology*. 2007; 107(4): 277–280, doi: [10.1159/000095518](https://doi.org/10.1159/000095518), indexed in Pubmed: [16954683](https://pubmed.ncbi.nlm.nih.gov/16954683/).
- Barssoum K, Altibi AM, Rai D, et al. Speckle tracking echocardiography can predict subclinical myocardial involvement in patients with sarcoidosis: A meta-analysis. *Echocardiography*. 2020; 37(12): 2061–2070, doi: [10.1111/echo.14886](https://doi.org/10.1111/echo.14886), indexed in Pubmed: [33058271](https://pubmed.ncbi.nlm.nih.gov/33058271/).
- Cacoub P, Chapelon-Abrie C, Resche-Rigon M, et al. Cardiac sarcoidosis: A long term follow up study. *PLoS One*. 2020; 15(9): e0238391, doi: [10.1371/journal.pone.0238391](https://doi.org/10.1371/journal.pone.0238391), indexed in Pubmed: [32946452](https://pubmed.ncbi.nlm.nih.gov/32946452/).