

Open Research Online

The Open University's repository of research publications and other research outputs

Plio-Pleistocene productivity reconstructions in the Indian Monsoon region

Conference or Workshop Item

How to cite:

Gray, Emmeline; Anand, Pallavi; Badger, Marcus; Bolton, Clara; Leng, M. J. and Murayama, Masafumi (2022). Plio-Pleistocene productivity reconstructions in the Indian Monsoon region. In: GC10-Pliocene-44: The Warm Pliocene: Bridging the geological data and modelling communities, 23-26 Aug 2022, Leeds, UK.

For guidance on citations see [FAQs](#).

© Author(s) 2022.



<https://creativecommons.org/licenses/by/4.0/>

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.5194/egusphere-gc10-pliocene-44>, 2022

<https://meetingorganizer.copernicus.org/GC10-Pliocene/GC10-Pliocene-44.html>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

GC10-Pliocene-44

<https://doi.org/10.5194/egusphere-gc10-pliocene-44>

The warm Pliocene: Bridging the geological data and modelling communities

© Author(s) 2022. This work is distributed under

the Creative Commons Attribution 4.0 License.



Plio-Pleistocene productivity reconstructions in the Indian Monsoon region

Emmeline Gray¹, Pallavi Anand¹, Marcus Badger¹, Clara Bolton², Melanie Leng³, and Masafumi Murayama⁴

¹School of Environment, Earth and Ecosystem Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

²Aix Marseille University, CNRS, IRD, INRA, Coll France, CEREGE, Aix-en-Provence, France

³British Geological Survey, Keyworth, Nottingham, NG12 5GG, UK

⁴Kochi University, Kochi, Japan

The Indian Summer Monsoon (ISM) brings seasonal winds and rains to the Indian subcontinent. The winds cause surface ocean mixing in the southern Bay of Bengal (BoB), bringing nutrients to the surface that fuel ocean productivity and export of carbon to the sea floor. To improve the understanding of the role of monsoon winds in low-latitude surface ocean productivity and carbon export, the Plio-Pleistocene interval is ideal because boundary conditions were evolving. We aim to reconstruct variability in surface ocean and export productivity across the late-Pliocene and early-Pleistocene in response to ISM wind-driven mixing in the southern BoB.

Here we analyse sediments from IODP Site U1443 (Exp. 353) from ~1.9-2.8 million years ago. A new benthic oxygen isotope ($\delta^{18}\text{O}$) stratigraphy (at ~3-thousand-year resolution) and age model tied to the global benthic $\delta^{18}\text{O}$ stack are presented. We utilise these sediments to obtain bulk sediment X-ray fluorescence (XRF) core scanning elemental data and coccolithophore assemblages to infer changes in summer monsoon runoff and surface ocean productivity influenced by monsoon wind strength. We use % *Florisphaera profunda* (coccolithophore assemblage) along with bulk sediment XRF Br as productivity indicators and a “terrigenous” bulk sediment XRF elemental composition stack as a wind strength and runoff indicator. We observe increased productivity during glacials (MIS 96, 98, 100), coinciding with increased terrigenous input. This observation is coherent with previous low-latitude productivity records from the equatorial Pacific. However, in contrast to equatorial Pacific productivity records, influenced by obliquity, our southern BoB records show robust surface productivity (% *F. profunda*) and summer monsoon runoff (terrigenous stack) peaks in both the obliquity and precession bands. We will discuss linkages between monsoon wind and runoff across the Plio-Pleistocene in context with other monsoon records.