



UvA-DARE (Digital Academic Repository)

The need to calibrate standardized cannabis measurements across cultures

Kuhns, L.; Kroon, E.

DOI

[10.1111/add.15744](https://doi.org/10.1111/add.15744)

Publication date

2022

Document Version

Final published version

Published in

Addiction

License

CC BY-NC-ND

[Link to publication](#)

Citation for published version (APA):

Kuhns, L., & Kroon, E. (2022). The need to calibrate standardized cannabis measurements across cultures. *Addiction*, 117(6), 1518-1519. <https://doi.org/10.1111/add.15744>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

The need to calibrate standardized cannabis measurements across cultures

The iCann Toolkit is an important step towards standardized measurement of cannabis consumption in both clinical and research settings. However, there are still significant challenges caused by regional differences in cannabis potency and cultural differences in methods of administration that will need to be addressed.

Lorenzetti *et al.* [1] proposed the International Cannabis (iCann) Toolkit as a framework for international minimum standards for the measurement of cannabis use across multiple contexts based on agreement from international cannabis researchers. This is a crucial step forward to standardized measurements of cannabis consumption and has the potential to vastly improve comparability of evidence for the effects of cannabis consumption across contexts. As Lorenzetti *et al.* [1] point out, differences in legislation, cultural customs, cannabis products and patterns of consumption make the measurement of cannabis use particularly challenging. In a cannabis research setting, the proposed mid- and top layer measures are the most appropriate for quantifying cannabis use. However, challenges still lie ahead in how to integrate measurements across testing sites that often have distinct cannabis cultures.

The proposed top layer biological measures include quantifying cannabis consumption or abstinence through analysis of tetrahydrocannabinol (THC) and metabolites in urine and blood plasma. Although this will constitute a gold standard for cannabis use quantification, the related cost, practical and even legislative issues limit the feasibility of using these measures, highlighting the importance of harmonizing mid-layer assessments such as the Timeline Followback (TLFB). However, as Lorenzetti *et al.* [1] acknowledge, cross-cultural differences in potency, unit of measurement and method of administration result in extra challenges to consider when using the TLFB method.

The cannabinoid content of cannabis products is known to differ across regions [2] and can diverge substantially from the labelled amounts even in countries with legal dispensation [3]. Hence, patients and research participants may often not have an accurate perception of the potency of the cannabis products they typically use. This poses

significant challenges since the effects of cannabis consumption are known to be dependent on levels of THC (psychoactive), cannabidiol (CBD) (non-psychoactive) and their interaction (e.g. [4]). Although the TLFB is a relatively accurate and reliable self-report measure of cannabis exposure [5], it faces significant challenges in terms of detailed assessment of exposure to the main compounds in cannabis. This is particularly the case in cross-cultural cannabis research, in which regional differences in the typical THC:CBD ratio of cannabis products are common [2,6]. Moving towards a standard unit of THC can help in the effort to improve comparability of measurements and findings across studies from different regions. The proposed calculation of standard THC units in the mid-layer of the iCann Toolkit recommends the usage of visual aids to help participants estimate the type and quantity of product they use to calculate a proxy of potency. It is a priority to test whether these standard units will accurately reflect THC and metabolite exposure as measured via biological quantification, and furthermore, whether regional differences in the validity and reliability of standard THC units will emerge. Studies that calibrate top- and mid-level assessments of potency across regions should be of high priority since they can inform researchers with limited or no access to top-level measures.

Methods of administration differ substantially between individuals [7] as well as across countries and regions (e.g. [8]). These differences affect intoxication duration and sub-acute effects due to differences in bioavailability [9]. Thereby, it can differentially affect major research outcomes such as cognitive performance [10] and even affect cue-reactivity, a proposed mechanisms of heavy and dependent use [11]. As Lorenzetti *et al.* [1] point out a specific example of this is the combination of cannabis and tobacco. Extending the iCann toolkit, we strongly recommend the standard administration of the TLFB for tobacco to measure tobacco and cannabis co-use (e.g. blunts) as well as separate tobacco use (i.e. vaping, cigarettes).

The iCann Toolkit constitutes a valuable step forward. When possible, top-layer biological quantification should be used to provide insight into regional differences in cannabis content, as well as untangle the effects of potency on the effects of cannabis. Researchers that have the means to use biological quantification should collectively invest in validating and calibrating biological quantification relative to self-reports. Furthermore, cultural differences in methods of

Lauren Kuhns and Emese Kroon contributed equally to this work (shared first authorship).

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Addiction* published by John Wiley & Sons Ltd on behalf of Society for the Study of Addiction.

administration, including co-use of tobacco, should be systematically considered herein. Eventually, this will hopefully lead to high-quality accessible measures for the wider scientific community.

DECLARATION OF INTERESTS

None.

ACKNOWLEDGEMENTS


This article was supported by grant 1R01 DA042490-01A1 awarded to Janna Cousijn and Francesca Filbey from the National Institute on Drug Abuse/National Institutes of Health.

AUTHOR CONTRIBUTIONS

Lauren Kuhns: Conceptualization; writing - original draft; writing - review & editing. **Emese Kroon:** Conceptualization; writing - original draft; writing - review & editing.

KEYWORDS

Cannabis research, cultural differences, measurement, method of administration, potency, standardization

Lauren Kuhns 

Emese Kroon

*Developmental Psychology, University of Amsterdam, Amsterdam,
Noord-Holland, The Netherlands
Email: l.n.kuhns@uva.nl*

Lauren Kuhns and Emese Kroon contributed equally to this work (shared first authorship).

ORCID

Lauren Kuhns  <https://orcid.org/0000-0002-3156-8905>

REFERENCES

- Lorenzetti V, Hindocha F C, Petrilli K, Griffiths P, Brown J, Castillo-Carniglia Á, et al. The international cannabis toolkit (iCannToolkit): A multidisciplinary expert consensus on minimum standards for measuring cannabis use. *Addiction*. 2022;117:1510–1517. <https://doi.org/10.1111/add.15702>
- Chandra S, Radwan MM, Majumdar CG, Church JC, Freeman TP, ElSohly MA. New trends in cannabis potency in USA and Europe during the last decade (2008–2017). *Eur Arch Psychiatry Clin Neurosci* 2019 2691 [Internet]. 2019[cited 2021 Oct 28];269(1):5–15. Available from: <https://doi.org/10.1007/s00406-019-00983-5>
- Jikomes N, Zoorob M. The cannabinoid content of legal cannabis in Washington state varies systematically across testing facilities and popular consumer products. *Sci Rep*. 2018. [cited 2021 Oct 27];8(1): 1–15. Available from: <https://www.nature.com/articles/s41598-018-22755-2>
- Niesink RJM, van Laar MW. Does Cannabidiol protect against adverse psychological effects of THC? *Front Psych*. 2013;4:130.
- Robinson SM, Sobell LC, Sobell MB, Leo GI. Reliability of the timeline Followback for cocaine, cannabis, and cigarette use 2012
- United Nations Office on Drugs and Crime. Drug market trends: Cannabis Opioids [Internet]. *World Drug Report 2021* [cited 2021 Nov 2]. Available from: www.unodc.org/unodc/en/data-and-analysis/wdr2021.html
- Schauer GL, King BA, Bunnell RE, Promoff G, McAfee TA. Toking, vaping, and eating for health or fun: Marijuana use patterns in adults, U.S., 2014. *Am J Prev Med*. 2016;50(1):1–8.
- Hindocha C, Freeman TP, Ferris JA, Lynskey MT, Winstock AR. No smoke without tobacco: A global overview of cannabis and tobacco routes of administration and their association with intention to quit. *Front Psych*. 2016 [cited 2019 Apr 4];7:1–9. Available from: <https://doi.org/10.3389/fpsy.2016.00104/abstract>
- Sharma P, Murthy P, Bharath MMS. Chemistry, metabolism, and toxicology of cannabis: Clinical implications. *Iran J Psychiatry*. 2012 [cited 2021 Nov 1];7(4):149–56. Available from: [/pmc/articles/PMC3570572/](http://pmc/articles/PMC3570572/)
- Kroon E, Kuhns L, Cousijn J. The short-term and long-term effects of cannabis on cognition: Recent advances in the field. *Curr Opin Psychol*. 2021;38:49–55.
- Kuhns L, Kroon E, Filbey F, Cousijn J. Unraveling the role of cigarette use in neural cannabis cue reactivity in heavy cannabis users. *Addict Biol*. 2021 [cited 2021 Nov 2];26(3):e12941. Available from: <https://doi.org/10.1111/adb.12941>

How to cite this article: Kuhns L, Kroon E. The need to calibrate standardized cannabis measurements across cultures. *Addiction*. 2022;117:1518–1519. <https://doi.org/10.1111/add.15744>