

Roskilde **University**

Complete Genome Sequence of a Multidrug-Resistant Klebsiella pneumoniae Environmental Isolate from Zanzibar, Tanzania, Harboring Novel Insertion Elements and Two blaCTX-M-15 Genes

Lallement, Claire; Krogfelt, Karen A.; Skovgaard, Ole; Jelsbak, Lotte

Published in: Microbiology Resource Announcements

DOI:

10.1128/mra.00263-22

Publication date: 2022

Document Version Publisher's PDF, also known as Version of record

Citation for published version (APA):

Lallement, C., Krogfelt, K. A., Skovgaard, O., & Jelsbak, L. (2022). Complete Genome Sequence of a Multidrug-Resistant Klebsiella pneumoniae Environmental Isolate from Zanzibar, Tanzania, Harboring Novel Insertion Elements and Two *bla*CTX-M-15 Genes. *Microbiology Resource Announcements*, *11*(8 August), [e00263-22]. https://doi.org/10.1128/mra.00263-22

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- · Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
 You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 02. Dec. 2022







Complete Genome Sequence of a Multidrug-Resistant *Klebsiella* pneumoniae Environmental Isolate from Zanzibar, Tanzania, Harboring Novel Insertion Elements and Two *bla*_{CTX-M-15} Genes

Claire Lallement, a CKaren A. Krogfelt, a COle Skovgaard, a CLotte Jelsbaka

^aDepartment of Science and Environment, Roskilde University, Roskilde, Denmark

ABSTRACT Here, we report the annotated whole-genome sequence of *Klebsiella pneumoniae* strain KP_3b, isolated in Zanzibar, Tanzania, from plastic litter. The strain is extended-spectrum β -lactamase (ESBL) producing and multidrug resistant, encoding 17 resistance genes, most of which are located on a 230,544-bp plasmid. The isolate contains two copies of the $bla_{CTX-M-15}$ gene and novel insertion elements.

lebsiella pneumoniae strain KP_3b was isolated from plastic litter collected in Mtopepo (GPS coordinates, 6°08′46.0″S, 39°13′14.5″E), in Zanzibar, Tanzania, by cultivation on selective LB agar plates containing ciprofloxacin (1).

To investigate the context of the numerous drug resistance genes and insertion elements (IS) found in this strain (1), we report here the complete genomic sequence of Klebsiella pneumoniae strain KP_3b, assembled from newly obtained long reads combined with BGISEQ short reads obtained previously (1). Further details of the isolation site, cultivation, and DNA extraction can be found in reference 1. Long reads were obtained using an Oxford Nanopore MinION FLO-MIN106D R9.4.1 flow cell loaded with a library prepared using the SQK-LSK109 ligation kit, minimizing the shearing of DNA. Fast5 files were base called to fastg files using Guppy v5.0.7 with the dna_r9.4.1_450bps_sup model, yielding a total of 1,253,844,477 nucleotides (nt) in 128,252 reads (N_{50} , 14,277 nt). Reads longer than 20,000 nt were assembled into scaffolds with Flye v2.9 (2) or with iterations of the Minimap, Miniasm (3), and Racon (MMR) assembly (4). Both methods resulted in two contigs of 5,331 kbp and 230 kbp with a mean coverage of 64x. A blastn pairwise alignment of the assembled contigs indicated that they were congruent but deviated at 565 positions. The Flye assembly results indicated that these contigs are circular. The MMR assembly was polished with the BGISEQ short reads (BGI) (5,804,065 paired-end reads; length, 150 nt; coverage, 120×) using SPAdes v3.15, with the -isolate and -trusted-contigs parameters set (5). Finally, we resolved 13 indels, introduced by SPAdes and ranging from 76 bp to 211 bp, by alignment of the SPAdes polished sequence with the trusted MMR assembly. The contigs were manually rotated to encode DnaA (chromosome) and RepB (plasmid) as the first coding DNA sequences (CDS), respectively. The GC content is 57.4% for the chromosome and 51.6% for the plasmid. We annotated the sequences using PGAP v2021-07-01.build5508 (6) with default settings.

As reported previously (1), the isolate contains 16 resistance genes, including one extended-spectrum β -lactamase (ESBL) $bla_{\text{CTX-M-15}}$ gene. However, in the present assembly, PGAP annotation indicated that the isolate contains a $bla_{\text{CTX-M-15}}$ gene both on the chromosome and on the plasmid. Both the chromosomal and plasmid copies of $bla_{\text{CTX-M-15}}$ are associated with ISEcp1 (7). Point mutations in the efflux pump regulators acrR and ramR (8) and the porin genes ompK36 and ompK37 (9) were identified using ResFinder analyses (10). Thirty IS elements were annotated on the plasmid. Seven copies of IS26 (11) as well as two copies of IS5075 affiliated with antibiotic resistance genes were found on the plasmid. Three new IS candidates were identified (two plasmid borne, one chromosomal). These have been submitted

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2022 Lallement et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0

Address correspondence to Ole Skovgaard, olesk@ruc.dk, or Lotte Jelsbak, ljelsbak@ruc.dk.

The authors declare no conflict of interest.

Received 14 March 2022 Accepted 20 June 2022 Published 14 July 2022

TABLE 1 MIC data for the indicated antimicrobials

Antibiotic(s)	MIC $(\mu g/mL)^a$	Phenotype ^b
Amoxicillin + clavulanic acid	32	R
Cefepime	16	R
Ceftriaxone	>128	R
Imipenem	< 0.125	S
Ciprofloxacin	4	R
Cefuroxime	<0.25	1
Amikacin	4	S
Tigecycline	>16	R
Kanamycin	256	R
Streptomycin	>256	R
Gentamicin	8	R
Piperacillin/tazobactam (6 μ g/mL)	64	R

^a MICs were measured using broth microdilution according to the EUCAST guidelines.

to the ISfinder database (ISKpn91, ISKpn92, ISKpn93) (12). The isolate is multidrug resistant (Table 1).

Data availability. The complete genome sequence of *Klebsiella pneumoniae* strain KP_3b has been deposited at GenBank under accession numbers CP086724 (chromosome) and CP086725 (plasmid). The raw BGISEQ reads and the base-called Oxford Nanopore reads have been deposited in the NCBI SRA database under accession numbers SRX7403512 and SRX13090583, respectively, and BioProject accession number PRJNA596383.

REFERENCES

- Rasool FN, Saavedra MA, Pamba S, Perold V, Mmochi AJ, Maalim M, Simonsen L, Buur L, Pedersen RH, Syberg K, Jelsbak L. 2021. Isolation and characterization of human pathogenic multidrug resistant bacteria associated with plastic litter collected in Zanzibar. J Hazard Mater 405:124591. https://doi.org/10.1016/j .jhazmat.2020.124591.
- Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, errorprone reads using repeat graphs. Nat Biotechnol 37:540–546. https://doi .org/10.1038/s41587-019-0072-8.
- Li H. 2016. Minimap and miniasm: fast mapping and de novo assembly for noisy long sequences. Bioinformatics 32:2103–2110. https://doi.org/ 10.1093/bioinformatics/btw152.
- Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly from long uncorrected reads. Genome Res 27:737–746. https://doi.org/10.1101/gr.214270.116.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. https://doi.org/10.1089/cmb.2012.0021.
- Li W, O'Neill KR, Haft DH, DiCuccio M, Chetvernin V, Badretdin A, Coulouris G, Chitsaz F, Derbyshire MK, Durkin AS, Gonzales NR, Gwadz M, Lanczycki CJ, Song JS, Thanki N, Wang J, Yamashita RA, Yang M, Zheng C, Marchler-Bauer A, Thibaud-Nissen F. 2021. RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with protein family model curation. Nucleic Acids Res 49: D1020–D1028. https://doi.org/10.1093/nar/gkaa1105.

- 7. Poirel L, Decousser J-W, Nordmann P. 2003. Insertion sequence ISEcp1B is involved in expression and mobilization of a $blaCTX_M$ β -lactamase gene. Antimicrob Agents Chemother 47:2938–2945. https://doi.org/10.1128/AAC .47.9.2938-2945.2003.
- Lv F, Cai J, He Q, Wang W, Luo Y, Wang X, Mi N, Zhao Z, Li G, Luo W. 2021. Overexpression of efflux pumps mediate pan resistance of *Klebsiella pneumoniae* sequence type 11. Microb Drug Resist 27:1405–1411. https://doi.org/10.1089/mdr.2020.0395.
- Al Mana H, Sundararaju S, Tsui CKM, Perez-Lopez A, Yassine H, Al Thani A, Al-Ansari K, Eltai NO. 2021. Whole-genome sequencing for molecular characterization of carbapenem-resistant Enterobacteriaceae causing lower urinary tract infection among pediatric patients. Antibiotics (Basel) 10:972. https://doi.org/ 10.3390/antibiotics10080972.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother 67:2640–2644. https://doi.org/ 10.1093/jac/dks261.
- Hansen KH, Andreasen MR, Pedersen MS, Westh H, Jelsbak L, Schønning K. 2019. Resistance to piperacillin/tazobactam in *Escherichia coli* resulting from extensive IS26-associated gene amplification of *bla*_{TEM-1}. J Antimicrob Chemother 74:3179–3183. https://doi.org/10.1093/jac/dkz349.
- Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference centre for bacterial insertion sequences. Nucleic Acids Res 34:D32–D36. https://doi.org/10.1093/nar/gkj014.

^b Phenotypes (R, resistant; I, intermediate; S, sensitive) are based on clinical breakpoints according to the EUCAST guidelines.