

15^e Conférence Internationale Tournesol
15th International Sunflower Conference



12-15 juin 2000
12-15 June 2000

Toulouse - France
Toulouse - France

ACTES
PROCEEDINGS



Tome I

AVANT-PROPOS

Ces deux volumes des actes de la 15^{ème} Conférence Internationale Tournesol contiennent les textes des conférences données lors des séances plénières et ceux correspondants aux posters présentés lors des ateliers.

Le Comité Scientifique avait pensé qu'il serait utile de faire le point sur tous les aspects de la recherche sur le Tournesol. C'est pourquoi il a invité seize personnalités reconnues dans leurs domaines d'activité à présenter des conférences lors de quatre séances plénières consacrées à l'Economie, à la Technologie, à l'Agronomie et l'Environnement et à la Génétique, la Sélection et la Biotechnologie. Elles avaient pour tâche de permettre à tous les participants, spécialistes ou non des sujets traités, d'approfondir leurs connaissances dans tous les domaines de recherche touchant au tournesol.

Le Comité Scientifique remercie vivement les conférenciers pour leurs apports et les présidents des séances plénières qui ont accepté d'animer les débats avec la salle.

Par ailleurs, le Comité Scientifique avait aussi souhaité que les travaux de recherche consacrés au tournesol à travers le monde soient présentés lors d'ateliers posters et fassent l'objet de synthèses et de discussions. Les résumés des posters reçus par le Comité lui ont permis d'organiser 15 ateliers. Des Présidents ont été proposés pour présenter les synthèses et organiser les débats. Ils étaient assistés de correspondants français qui ont eu la lourde tâche de relire et éventuellement de corriger les textes publiés ici. Les Présidents ont reçus tous les textes de leurs ateliers avant la conférence.

Le Comité Scientifique tient à remercier les Présidents et les correspondants pour leur travail et leur engagement dans la réussite de la conférence et la qualité de la publication.

Le Comité Scientifique

Foreword.

These two volumes of the Proceedings of the 15th International Sunflower Conference contain the texts of the lectures given during the plenary sessions and those corresponding to the posters presented at the workshops. The scientific committee considered that it would be useful to have general updates on all aspects of research on sunflower and so invited sixteen acknowledged authorities in their domains of activity to give lectures during four plenary sessions devoted to Economics, to Technology, to Agronomy and Environment and to Genetics, Breeding and Biotechnology. Their task was to allow participants, specialists or not of the subjects dealt with, to widen their knowledge of all aspects of sunflower research.

The Scientific Committee warmly thanks the lecturers for their contribution and the plenary session chairmen who have accepted to lead the discussions with the audience.

The Committee also wanted to encourage the presentation of research programmes throughout the world during poster workshops, giving subjects for synthesis and debate. Poster summaries received by the Committee helped to decide the organisation of 15 workshops. Chairmen were proposed to present the syntheses and to organise the debates. They were assisted by french correspondents who had the hard task of reading and possibly correcting, the full texts published herein. The chairmen received all the texts for their workshops beforehand.

The scientific Committee wishes to thank both the chairmen and their correspondents for their work and their dedication to the success of the conference and the quality of this publication.

Séance plénière/Plenary session : A - Economie/Economics

- PLA- EVOLUTION COMPARÉE DE LA PRODUCTION ET DE LA COMPÉTITIVITÉ DU TOURNESOL DANS DIFFÉRENTES AIRES DE PRODUCTION PLA-1
André Pouzet, D. Delplancke
- PLA- LE TOURNESOL EN ARGENTINE : SITUATION ET DEFIS PLA-11
Rubén C. Devoto, Martine Guibert

Séance plénière/Plenary session : B - Technologie/Technology

- PLB- PHYSICAL REFINING OF SUNFLOWER OIL PLB-1
Katalin Kövari, Jean Denise, Asolt Kemény, Katalin Recseg
- PLB- QUELQUES EXEMPLES DE VALORISATION NON-ALIMENTAIRE DU TOURNESOL PLB-13
M.E. Borredon, A. Gaset
- PLB- INTÉRÊT NUTRITIONNEL DES TOURNESOLS PLB-15
B. Delpanque

Séance plénière/Plenary session : C - Agronomie et Environnement/Agronomy and Environment

- PLC- DISEASE CONTROL CONCERNS EVERY BODY: THE EXAMPLE OF SUNFLOWER DOWNY MILDEW PLC-1
Denis Tourvieille de Labrouhe
- PLC- PRINCIPLES AND APPLICATIONS OF SUNFLOWER CROP SIMULATION MODELS PLC-9
Francisco J. Villalobos
- PLC- ENVIRONMENTAL ISSUES OF SUNFLOWER PLC-18
Paolo Cecon, Philippe Delbaeke, Pedro González-Fernández
- PLC- SUNFLOWER ECOPHYSIOLOGY: SOME UNRESOLVED ISSUES PLC-28
Antonio J. Hall

Séance plénière/Plenary session : D - Génétique, Sélection et Biotechnologies/Genetics, Breeding and Biotechnology

- PLD- THE DEVELOPMENT OF TOOLS FOR MOLECULAR BREEDING AND GENOMICS RESEARCH IN CULTIVATED SUNFLOWER PLD-1
Steven J. Knapp, Mary B. Slabaugh, Shunxue Tang
- PLD- UNRAVELING THE MOLECULAR MECHANISMS OF PATHOGEN RESISTANCE IN SUNFLOWERS PLD-8
Said Mouzeyar
- PLD- PROBLEMS AND GOALS IN STUDYING OIL COMPOSITION VARIATION IN SUNFLOWER PLD-16
André Bervillé, Séverine Lacombe
- PLD- PHYLOGENY OF HELIANTHUS AND RELATED GENERA PLD-26
Edward E. Schilling

Atelier posters/Poster workshop : B - Nouveaux usages/New uses

- A-F33 - ANALYSIS OF DESATURASE TRANSCRIPT ACCUMULATION IN NORMAL AND IN HIGH OLEIC OIL SUNFLOWER DEVELOPMENT SEEDS
Lacombe Séverine, Berville André
- A-SPT - BIOCHEMICAL CONTROL OF HIGH PALMITIC ACID BIOSYNTHESIS
Rafael Garcés, Rosario Alvarez-Ortega, Sara Canhisan, Enrique Martínez-Force
- A-F31 - GENETIC AND MOLECULAR CHARACTERIZATION OF THE HIGH OLEIC CONTENT OF SUNFLOWER OIL IN PERENNETS
Lacombe Séverine, Guillot Hélène, Kaan François, Millet Christine, Berville André
- A-F29 - TRIACYLGLYCERIDE COMPOSITION IN F1 SEEDS USING FACTORIAL AND DIALEL CROSSSES BETWEEN SUNFLOWER LINES
Didier Vares, Jacky Cleomene, Séverine Lacombe, Yves Griveau, André Berville, François Kaan,
- A-19 - OIL COMPOSITION AND ACCUMULATION OF FATTY ACIDS IN NEW OLEIC SUNFLOWER (HELIANTHUS ANNUUS L.) HYBRIDS
Lagrèze Thierry, Lacombe Séverine, Surl Olivier, Kleiber Didier, Berville André, Dayde Jean
- A-SP3 - INHERITANCE OF OLEIC ACID CONTENT UNDER CONTROLLED ENVIRONMENT
Leonardo Velasco, Begoña Pérez-Vich, José M. Fernández-Martínez
- A-USA15 - COMPARISON OF OIL QUALITY CHARACTERISTICS OF AGENES FROM ORIGINAL AND REGENERATED POPULATIONS OF WILD SUNFLOWER SPECIES
Gerald J. Sailer, Mary E. Brothers
- A-YU12 - DEVELOPMENT OF INBRED LINES OF SUNFLOWER WITH VARIOUS OIL QUALITIES
Simsa Joci, Dragan Skoric, Nada Lecic, Igor Molnar
- A-B2 - PROGRESS IN THE DEVELOPMENT OF DNA-BASED MARKERS FOR HIGH STEARIC ACID CONTENT IN SUNFLOWER
Begoña Pérez-Vich, José María Fernández-Martínez, Juan Muñoz-Ruz, Steven J. Knapp, Simon T. Berry
- A-B3 - MOLECULAR MARKERS ASSOCIATED WITH SUNFLOWER OLEIC AND STEARIC ACID CONCENTRATIONS IN A HIGH STEARIC X HIGH OLEIC CROSS
Begoña Pérez-Vich, José María Fernández-Martínez, Steven J. Knapp, Simon T. Berry
- A-55 - EFFECT OF CULTURAL CONDITIONS ON YIELD, OIL CONTENT AND FATTY ACID COMPOSITION OF SUNFLOWER KERNEL
Piva G., Bouniols A., Mondés M
- A-F42 - EFFECT OF TEMPERATURE FROM FLOWERING TO MATURITY ON SEED COMPOSITION OF HIGH OLEIC SUNFLOWER INBREDS AND MID OLEIC HYBRIDS
Anne-Marie Thibot-Blondel, Bernard Bonnemoy, Robert Folcimgne, Michel Mortignac, Joëlle Messaoud, Jacqueline Philippou, Felicity Year
- A-F68 - EFFECTS OF TEMPERATURE VARIATIONS ON FATTY ACID COMPOSITION IN OLEIC SUNFLOWER OIL (HELIANTHUS ANNUUS L.) HYBRIDS
Lagrèze Thierry, Champolivier Luc, Lacombe Séverine, Kleiber Didier, Berville André, Dayde Jean
- A-117 - EFFECTS OF DIFFERENT WATER AVAILABILITY ON FATTY ACID COMPOSITION OF THE OIL IN STANDARD AND HIGH OLEIC SUNFLOWER HYBRIDS.
Marta Baldini, Romano Giovanardi, GianPaolo Vannozzi
- A-AR29 - STABILITY OF HIGH OLEIC SUNFLOWER HYBRIDS FOR GRAIN YIELD AND OIL AND OLEIC ACID CONTENTS IN THE SUNFLOWER SOWING REGION IN ARGENTINA A-85
Aguero M.E., Lúquez J., Pereyra Y.R., Aguirrezábal, L.A.N.
- A-AR41 - RENDIMIENTO Y ESTABILIDAD DE RENDIMIENTO EN HÍBRIDOS DE GIRASOL LINOLEICOS Y ALTO OLEICOS:
II. EFECTO DE FACTORES COMBINADOS O DEL AMBIENTE
Sergio Uhart, Mariano Frugone, Guillermo Pozzi, Ramón Correa, Claudio Simonella
- A-AR38 - EFFECT OF SUNFLOWER SEED DRYING ON SOME QUALITY PARAMETERS OF THE SEED AND OIL
Estela M. Santalla, Olinda E. Quiroga, María C. Gely
- A-MEX2 - CALIDAD FISIOLÓGICA Y SU RELACION CON LA COMPOSICIÓN QUÍMICA EN SEMILLAS DE GIRASOL (HELIANTHUS ANNUUS L.).
Velázquez Cagal Macario, Sevilla Panigua Evangelina, Hernández García, Leonardo y López Ramírez Javier
- A-103 - HETEROGENEITY AND NATURAL VARIABILITY OF PROTEINASE INHIBITORS IN SUNFLOWER (HELIANTHUS L.) AND OTHER COMPOSITAE
Alexander Konarev, Irina Anisimova, Vera Gavrilova, Tatyana Vachrusheva, Larisa Shashlova, Galina Korchnaya, Peter Shewry
- A-109 -

- B-AR52 - DEGOMADO DE ACEITES DE GIRASOL. INFLUENCIA DE LAS CONDICIONES DE PROCESO
L.G. Pan, A. Noli, A. Campaña, M. Barrera, M.C. Tomás, M.C. Añón
- B-F72 - IMPROVEMENT OF THE SUNFLOWER DEMAXING AT LABORATORY SCALE USING ULTRASOUNDS
Xavier Pagès, Carine Alfos
- B-7 -

BLF66 -	INTERET DES HUILES COMBINÉES À BASE DE TOURNESOL, TOURNESOL OLÉIQUE ET COLZA EN NUTRITION HUMAINE DÉFINITION DES LIMITES OPTIMALES DE L'APPORT EN ACIDE OLÉIQUE APRÈS STABILISATION DES APPORTS EN ACIDES LINOLÉIQUE ET ALPHA-LINOLÉIQUE . . .	B-14
	<i>Bernadette Delplanque, Brigitte Le Roy, Nicole Combe, François Mendy, Evelyne Fenart, Bernard Entressangles, Anissa Thaminy, Farisa Syeda, Stéphanie Tanguy</i>	
BLF77 -	SUPERCritical CARBON DIOXIDE EXTRACTION OF SUNFLOWER SEEDS AND RAPESEEDS OILS	B-20
	<i>Michelangelo Cammarata, Valerio Abbate, Tatiana Görner, Danielle Barth</i>	
BLF44 -	SUNFLOWERATES 1 : SYNTHESIS OF ALKYL OLEATES BY TRANSESTERIFICATION OF OLEIC SUNFLOWER OIL WITH C8-ALCOHOLS CATALYSED BY HYDRATED H ₃ PO ₄	B-26
	<i>Corrine Dufaure, Zéphirin Mouloungui</i>	
BLF45 -	TOURNESOLEATE 2 : PREPARATION DES MILIEUX REACTIONNELS DE FORMULATION DE GRAISSES LUBRIFIANTES A BASE D'HUILE DE TOURNESOL OLÉIQUE ADDITIONNE IN SITU D'OLEATE D'ALKYLE ET D'OLEIQUE METALLIQUE	B-31
	<i>Corrine Dufaure, Zéphirin Mouloungui, Luc Rigal</i>	
BLF46 -	TOURNESOLEATE 3 : STABILITÉ THERMIQUE, CHIMIQUE PROPRIETES TENSIO-ACTIVES ET ANTI-USURE DES OLEATES DE CARBONATE DE GLYCEROL	B-35
	<i>Pelet Sandrine, Zéphirin Mouloungui</i>	
BLF69 -	SYNTHESIS OF NEW DERIVATIVES FROM VEGETABLE SUNFLOWER OIL METHYL ESTERS VIA EPOXIDATION AND OXIRANE OPENING	B-41
	<i>Xavier Pagès, Carine Alfos</i>	
BLSP2 -	THERMOXIDATIVE STABILITY OF A NEW SUNFLOWER SEED OIL WITH HIGH LEVELS OF OLEIC AND PALMITIC ACIDS	B-48
	<i>Manuel Mancha, Rafael Garcés and Gloria Márquez-Ruiz</i>	
BLFB1 -	CREOL: A PILOT PLANT FOR OIL AND PROTEIN CROP PRODUCTS	B-54
	<i>Pierre Burghart, Jacques Evrard</i>	
BLF48 -	UTILISATION DE TOURTEAU DE TOURNESOL POUR LA FABRICATION DE POTS DE REPIQUAGE BIODEGRADABLES	B-58
	<i>A. Rouilly, F. Silvestre, L. Rigal, H. Caruel, E. Paux, J. Silvestre, P. Morard</i>	
BLF79 -	PREDICTION OF METABOLIZABLE ENERGY OF FULL-FAT SUNFLOWER SEEDS BY COCKERELS	B-64
	<i>Anne Bourdillon, Jacques Evrard, Marie-Pierre Le Guen</i>	
BLF80 -	EFFECTS OF CRUSHING CONDITIONS ON SUNMEAL QUALITY	B-69
	<i>Anne Bourdillon, Pierre Burghart, Jacques Evrard, Marie-Pierre Le Guen</i>	
BLF16 -	HYDROPHOBATION OF POLYSACCHARIDES WITH SUNFLOWER OIL AND ITS DERIVATIVES	B-73
	<i>S. Girardeau, C. Vaca-Garcia, J. Aburto-Anell, I. Alric, M.E. Borredon</i>	
BLF49 -	LES PROTÉINES DE TOURNESOL EN SOLUTION ACQUEUSE RHEOLOGIE ET PROPRIETES ADHESIVES	B-78
	<i>J. Leyris, F. Silvestre, L. Rigal</i>	
BLF50 -	PREPARATION OF FILMS FROM ISOLATE SUNFLOWER (HELIANTUS ANNUUS) PROTEINS	B-84
	<i>Fabiola Ayhllon-Meixueiro, François Silvestre</i>	
BLF51 -	OBTENTION D'AGROMATERIAUX BIODEGRADABLES A PARTIR DE TOURTEAU DE TOURNESOL PAR FORMAGE OU THERMOFORMAGE	B-90
	<i>J. Leyris, F. Silvestre, L. Rigal</i>	
BLF70 -	EVOLUTION DU DEGRÉ DE MÉTHYLATION DES PECTINES DE CAPITULE DE TOURNESOL EN FONCTION DE LA MATURITÉ	B-96
	<i>V. Marechal, L. Rigal, J. Raynal, G. Alibert</i>	
BLIT3 -	VARIABILITY FOR CHARACTERS RELATED IN INULIN ACCUMULATION IN TOPINAMBUR (HELIANTHUS TUBEROSUS L.)	B-102
	<i>Mario Baldini, Francesco Danuso, Maurizio Turi, GianPaolo Vannozzi</i>	
BLAR18 -	DRYING RATES ESTIMATION FOR DIFFERENT SUNFLOWER SEED VARIETIES GROWN IN ARGENTINA	B-108
	<i>M.C. Gely, E.M. Santalla</i>	

Ateliers posters/Poster workshops : C - Conduites des cultures/Crop management

CF6 -	TESTING CROP MANAGEMENT SYSTEMS FOR SUNFLOWER IN SOUTH-WEST FRANCE	C-1
	<i>Philippe Debaeke, Jean-Marie Nolot</i>	
CIT21 -	EVALUATION OF SUNFLOWER AGRO-ECONOMIC PERFORMANCES IN LOW-INPUT CROPPING SYSTEMS (EU REG. 2078/92) OF CENTRAL ITALY	C-7
	<i>Nicola Silvestri, Silvia Pampana, Enrico Bonari, Dario Conti</i>	
CYU2 -	EFFECT OF CULTURAL PRACTICES ON SUNFLOWER YIELDS IN A PERIOD OF SEVERAL YEARS IN ER YUGOSLAVIA	C-12

- C1720 - IMPORTANCE OF VARIETY CHOICE IN SUNFLOWER'S ANSWER TO DIFFERENT LEVELS OF CROPPING INTENSIFICATION
Silvia Pampana, Nicola Silvestri, Enrico Bonari, Giovan Maria Rodoni
- C173 - ASPHODEL : MODELE DE SIMULATION DES EPIDEMIES DE PHOMOPSIS DU TOURNE SOL (DIAPORTHE HELIANTHI)
Marc Delos, Jacques Moizard, Philippe Debœcke, André Pères
- C17 - RECHERCHE D'INDICATEURS DU COUVERT POUR LA PREVISION PRECOCE DU RISQUE PHOMOPSIS
Philippe Debœcke, André Estragnat, André Pères, Sophie Fiasentin
- CAR44 - EFECTO DE DIFERENTES SISTEMAS DE LARANZA SOBRE LA COMUNIDAD DE MALEZAS E INSECTOS DE SUELO, ASOCIADOS AL CULTIVO DE GIRASOL
Edgardo J. Zorza, Adhif López, Fernando Datta, Filomeno Sayago y Leonidas Chokky
- C49 - INTEGRATED SYSTEM OF PEST CONTROL AND BIOLOGICAL NUTRITION FOR SUNFLOWER CROP
Horia Iliescu, Ana Popescu, Alina Ionita, Simona Kupferberg, Ioana Prodan, Dumitru Manole
- C174 - EFFECT OF SOWING DENSITY ON THE DYNAMICS OF LEAF AREA DEVELOPMENT IN SUNFLOWER
Dusanic, N. and Milkic, V., Starevic, I., Crnobarac, J., Mihajlovic, D.
- C171 - TEMPORAL AND PLANT DENSITY EFFECTS OF INTERCROP COMPETITION ON SUNFLOWER
Sudhakar Babu, Shivashankar, K.
- C1725 - STUDIES ON AGRONOMIC MANIPULATIONS FOR IMPROVING THE SEED YIELD AND QUALITY OF KRS-1 SUNFLOWER HYBRID SEED PRODUCTION
K. Somasekhara, K.T. Krishne Gowda, V.R. Kalappa, Chikkadevaiah, K. Seenappa
- C173 - SUNFLOWER TOLERANCE TO PRE-EMERGENCE BROADLEAF WEED HERBICIDES
Alexandre M. Brighenti, Marcelo F. Oliveira, Dionisio L. P. Gazziero, José E. Pereira Elomar Voll
- C174 - CONTROL OF ANNUAL ASTERACEAE AND APIACEAE IN SUNFLOWER CROPS IN FRANCE: THE EXAMPLES OF AMBROSIA ARTEMISIIFOLIA AND AMMI MAJUS
Etienne Pilorge
- C177 - CONTROL OF *Cicum arvense* (Canada thistle) IN SUNFLOWER - CEREALS ROTATIONS
Etienne Pilorge, André Estragnat, Lionel Jouy, Jean-Luc Verdier
- C176 - EXAMPLE OF CROP MODEL APPLICATION TO EVALUATE IRRIGATION STRATEGIES IN SUNFLOWER
Michele Rinaldi
- C179 - SUNFLOWER IRRIGATION TREATMENTS IN INTERACTION WITH A PREVIOUS MULCH CROP IN SOUTHERN ITALY
Michele Rinaldi, Michele Maiorana
- C185 - PERSPECTIVES D'EVOLUTION DU TOURNE SOL DANS LES ASSOLEMENTS IRRIGUES DU SUD-OUEST DE LA FRANCE
Kevin Lemaire, Bernard Le Clech, Raymond Reau, Jean Kaimbavuli, Jean-Luc Lespina, Géraldine Paget
- C103 - IMPACT OF IRRIGATION ON SUNFLOWER PRODUCTIVITY
Dhillon A. Singh, Sidhu M. Singh
- C171 - UTILISATION PAR LE TOURNE SOL DE L'EAU STOCKEE AU SEMIS EN CLIMAT MEDITERRANEE SEMI-ARIDE
Abdelilah Aboudrare, Hassam Chekri, Ahmed Bouaziz, Philippe Debœcke
- C172 - VOIES D'AMELIORATION DE LA PRODUCTIVITE ET DE L'EFFICACITE D'UTILISATION DE L'EAU CHEZ LE TOURNE SOL
Sous climat méditerranéen semi-aride
Abdelilah Aboudrare, Ahmed Bouaziz, Philippe Debœcke
- C121 - RENDEMENT ET QUALITE DES GRAINES DU TOURNE SOL CULTIVE SOUS CONTRAINTE HYDRIQUE. RESULTATS DES ESSAIS EN COOPERATION AU MAROC ET DANS LE SUD-OUEST DE LA FRANCE
Mohammed El Asri, Abdelrahim Essahat, André Bouzols, Michel Mondès
- C127 - THE EFFECT OF CROP WATER STATUS ON THE YIELD, COMPOSITION AND PROCESSING QUALITY OF SUNFLOWER SEED
A.A. Nel, H.L. Loubser, P.S. Hammes
- C172 - EFFECT OF SUPPLEMENTARY IRRIGATION ON SEED YIELD AND OIL QUALITY OF SUNFLOWER (*Helianthus annuus* L.) GROWN IN A SUB-ARID ENVIRONMENT
Z. Flagella, T. Rotunno, R. Di Caterina, G. de Simone, L. Ciceriti, A. De Caro
- C182 - SUNFLOWER RESPONSE TO WATER STRESS AND BORON APPLICATION
César de Castro, Adonis Moreira, João Batista Rodrigues de Abreu
- C185 - INFLUENCE OF SOIL TILLAGE AND SOWING TECHNOLOGY ON SUNFLOWER PRODUCTION
Gh. Sin, Gh. Petcu, St. Ionita, M. Popa
- C172 - RESPONSE OF SUNFLOWER TO TILLAGE, STUBBLE AND WEED MANAGEMENT UNDER PADDY FALLOWS
B.C. Channakeshava, Chikkadevaiah, K. Somasekhara
- C156

C.160 - FERTILIZACION PROFUNDA DEL GIRASOL EN SIEMBRA DIRECTA EN LA REGION PAMPEANA ARGENTINA	C-159
<i>Carlos Ricardo Baumer, Nestor Gonzalez, Carlos Devito</i>	
C.165 - PRODUCCION DE GIRASOL BAJO DISTINTOS SISTEMAS DE LABRANZAS Y FERTILIZACION, EN UNA REGION SUB-HUMEDA	C-165
<i>Elena M. Fernandez, Leonidas Cholaky, Edgardo J. Zorza, Gabriel P. Espósito, Jorge J. Gesumaria, Walter E. Asnal,</i>	
C.169 - DIAGNOSTICO DE FERTILIZACION NITROGENADA PARA EL CULTIVO DE GIRASOL BAJO DIFERENTES SISTEMAS DE LABRANZA EN EL SUD ESTE DE LA PROVINCIA DE BUENOS AIRES (ARGENTINA)	C-171
<i>Jorge González Montaner, Marcelo Di Napoli</i>	
C.170 - INTEGRATED PHOSPHORUS MANAGEMENT IN SOYBEAN - SUNFLOWER CROPPING SYSTEM	C-177
<i>D.M. Hegde</i>	
C.180 - RESPONSE OF SUNFLOWER TO FOLIAR SPRAYING WITH EPSOM SALT (MgSO4.7H2O) SOLUTION	C-182
<i>Miroslav Krizmanic, Zorica Jurkovic, Anto Mijic, Eberhard Uebel</i>	

Ateliers posters/Poster workshops : D - Agro-physiologie/Crop physiology

D.1A10 - RESPUESTAS A LA TEMPERATURA DEL PESO FINAL, DURACION Y TASA DE LLENADO DEL EMBRION EN GIRASOL	D-1
<i>Claudio Chimenti, Antonio J. Hall, María S. López</i>	
D.1A14 - EFFECT OF INTERCEPTED SOLAR RADIATION DURING THE FRUIT FILLING PERIOD ON SUNFLOWER OIL YIELD	D-7
<i>Guillermo A. A. Dosio, Victor R. Pereyra, Fernando H. Andrade, Luis A. N. Aguirrezabal</i>	
D.1A27 - ANATOMÍA Y DESARROLLO DEL PERICARPIO DE DISTINTOS GENOTIPOS DE GIRASOL (<i>Helianthus annuus</i> L.)	D-13
<i>Lilja Ivone Lindström, Cecilia Noemi Pellegrini, Luis Francisco Hernández</i>	
D.1A31 - RELATIONSHIPS BETWEEN SEED OIL PERCENTAGE AND KERNEL OIL PERCENTAGE & KERNEL PERCENTAGE (w/w) AS AFFECTED BY PLANT DENSITY & SOLAR RADIATION INTERCEPTED DURING GRAIN FILLING IN SUNFLOWER	D-19
<i>Nolasco S.M., Dosio G.A.A., Riccobene I.C., Santalla E.M., Aguirrezabal L.A.N.</i>	
D.1A11 - RESPUESTAS DEL NUMERO DE GRANOS A LA TEMPERATURA DURANTE LA DIFERENCIACION FLORAL	D-25
<i>Claudio Chimenti, Antonio J. Hall, María S. López</i>	
D.1A34 - PATRON DE VANE0 EN CAPITULOS DE GIRASOL (<i>Helianthus annuus</i> L.) CRECIDOS EN CONDICIONES NATURALES	D-31
<i>P.J. Paoloni, L.F. Hernández, C.N. Pellegrini</i>	
D.1A8 - REDUCTION IN THE NUMBER OF FILLED SEED IN SUNFLOWER (<i>Helianthus annuus</i> L.) BY LIGHT STRESS	D-35
<i>Julio E. Cantagallo, Antonio J. Hall</i>	
D.1F17 - SPATIAL AND TEMPORAL PATTERNS OF CELL DIVISION AND TISSUE EXPANSION AS AFFECTED BY TEMPERATURE AND WATER DEFICIT IN SUNFLOWER LEAVES	D-41
<i>Christine Granier, François Tardieu</i>	
D.1F28 - STABILITY OF SUNFLOWER LEAF DEVELOPMENT IN FRANCE AND ARGENTINA	D-47
<i>Guillermo Dosio, Thierry Lefeuvre, Benoît Suard, Natalia Izquierdo, Luis Aguirrezabal, Olivier Turc</i>	
D.1J18 - INFLUENCE DE LA TEMPÉRATURE ET DE LA PHOTOPÉRIODE SUR LA PHÉNOLOGIE ET LES CARACTÉRISTIQUES DU RENDEMENT DU TOURNESOL	D- 52
<i>Rosalena Tuttobene, Umberto Anastasi, Orazio Sortino, Michelangelo Cammarata</i>	
D.1J4 - RESPONSE OF SUNFLOWER TO PROGRESSIVE WATER STRESS	D-58
<i>Stefano Bona, Giuliano Mosca, Antonio Cantale, Teofilo Vamerli</i>	
D.1S5 - EFICIENCIA EN EL USO DEL AGUA EN EL GIRASOL DE SIEMBRA INVERNAL Y PRIMAVERAL DETERMINADA MEDIANTE LISIMETROS DE DRENAJE	D-64
<i>E. Sobrino, M.D. Curt</i>	
D.1A37 - LEAF PHOTOSYNTHETIC CAPACITY, CHLOROPHYLL CONTENT, AND SPECIFIC LEAF NITROGEN RELATIONSHIPS IN SUNFLOWER CROPS PRIOR TO ANTHESIS	D-70
<i>M. Cecilia Rousseaux, A. J. Hall, R. A. Sánchez, E. I. Ploschuk</i>	
D.1J11 - THE SUNFLOWER (<i>HELIANTHUS ANNUUS</i> L.) MUTANT XAN1 IS IMPAIRED IN PHOTOSYNTHETIC ACTIVITY	D-76
<i>Marco Fambrini, Lucia Guidi, Antonella Castagna, Paolo Vernieri, Francesco Ferraro, Mauro Durante, Claudio Pugliesi</i>	
D.1A2 - LEAF TEMPERATURE OF SUNFLOWER AS AFFECTED BY NITROGEN AND WATER STATUS	D-82
<i>Menon Parameswaran</i>	
D.1A4 - MODELLING AND MEASURING TRANSPIRATION EFFICIENCY IN COMMERCIAL AND EXOTIC SUNFLOWER	D-87
<i>Scott C. Chapman, Christopher J. Lambrides, Bodapati P. Naidu, Ray Shorter</i>	
D.1E9 - NUMERICAL SUNFLOWER PLANTS RESPONDING TO LIGHT AND TEMPERATURE	

D-106	PHYSIOLOGICAL BASES OF GENOTYPE BY ENVIRONMENT INTERACTIONS FOR SOWING DATE IN SUNFLOWER <i>Abelardo J. de la Vega, Antonio J. Hall</i>
D-112	ENVIRONMENTAL ATTRIBUTES UNDERLYING ENVIRONMENTAL MAIN-EFFECTS AND GENOTYPE BY ENVIRONMENT INTERACTIONS IN SUNFLOWER <i>Abelardo J. de la Vega, Scott C. Chapman</i>
D-117	CORRELATION BETWEEN PHYSIOLOGICAL INDEX, SUNFLOWER PLANT HEIGHT AND DRY MATTER IN DIFFERENT PHENOLOGICAL STAGES <i>Nuria F. Calvet, Maria Regina G. Ungaro</i>
D-123	PHYSIOLOGICAL CHARACTERISTICS OF THE SUNFLOWER CROP (<i>HELIANTHUS ANNUUS</i> L.) AS AFFECTED BY PLANTING GEOMETRY <i>Wulf Diepenbrock, Mu-Hua Long, Boy Fell</i>
D-129	RENDIMIENTO Y ESTABILIDAD DE RENDIMIENTO EN HÍBRIDOS DE GIRASOL LINOLEICOS Y ALTO OLEICOS: I. EFECTO DE LA VARIACION EN RADIACION SOLAR <i>Sergio Uhart, Mariano Frugone, Guillermo Pozzi, Ramón Correa, Claudio Simonella</i>
D-136	SUNFLOWER CULTIVAR RESPONSES TO NON-UNIFORMITY OF THE STAND: EFFECTS ON YIELD PER PLANT AND AT THE CROP LEVEL <i>N. Trapani, M. Lopez Pereira, A.J. Hall</i>
D-140	EFFECT OF THE LOSS OF PLANTS OR HEADS AT DIFFERENT PHENOLOGICAL STAGES ON SUNFLOWER YIELD <i>Pereyra V.R., Aguirrezabal L.A.N., Pedraza M.V.</i>
D-144	RELATIONSHIP BETWEEN HYBRID PRODUCTIVITY AND MAIN PHYSIOLOGICAL INDEX UNDER SUMMER CROPPING SUNFLOWER <i>Cui Langji, Xin Huaqun, Liang Guozhan, Wang dexing</i>
D-150	STUDIES ON MANIPULATION OF SOURCE-SINK RELATIONSHIP FOR IMPROVING SEED YIELD AND QUALITY OF MALE PARENTAL LINE (RHA 6D-1) OF KWSH-1 SUNFLOWER HYBRID <i>K. Somasekhara, K.T. Krishne Gowda, V.P. Kalappa, P. Balakrishna, D. Nuthan, S.D. Nehru, and K. Seenappa</i>
D-156	AGRONOMIC RESPONSE OF SUNFLOWER TO WATERLOGGING SUWATCHAI CHUENCHOM AND PAISAN LAOSUAN <i>Suwatjai Chuenchom, Paisan Laosuan</i>

TOME II	Ateliers posters/Poster workshops : E - Productivité : composants, aptitude à la combinaison, interaction génotype x environnement <i>Productivity: components, combining ability, genotype x environment interaction</i>
TOME II	Atelier posters/Poster workshop : F - Qualité des semences/Seed Quality
TOME II	Atelier posters/Poster workshop : G - Traitement des semences/Seed treatment
TOME II	Atelier posters/Poster workshop : H - Entomologie/Entomology
TOME II	Atelier posters/Poster workshop : I - Epidémiologie, description des maladies/Epidemiology, description of diseases
TOME II	Atelier posters/Poster workshop : J - Résistance verticale/Vertical resistance
TOME II	Atelier posters/Poster workshop : K - Résistance horizontale/Horizontal resistance
TOME II	Atelier posters/Poster workshop : L - Biologie cellulaire, Biotechnologie/Cellular biology, Biotechnology
TOME II	Atelier posters/Poster workshop : M - Cartographie, marqueurs/Mapping, markers
TOME II	Atelier posters/Poster workshop : N - Perspectives du développement du tournesol dans différents pays <i>Future development of the sunflower around the world</i>
TOME II	Atelier posters/Poster workshop : O - Ressources génétiques/Genetic resources

DEVELOPMENT OF INBRED LINES OF SUNFLOWER WITH VARIOUS OIL QUALITIES

Siniša Jocić, *Institute of Field and Vegetable Crops*
Maksima Gorkog 30, 21000 Novi Sad, Yugoslavia
Tel/fax: 00 381 21 413 833, E-mail: sinjocic@ifvcns.ns.ac.yu

Dragan Škoric, *Institute of Field and Vegetable Crops*
Maksima Gorkog 30, 21000 Novi Sad, Yugoslavia
Tel/fax: 00 381 21 413 833, E-mail: skoricd@ifvcns.ns.ac.yu

Nada Lecic, *Institute of Field and Vegetable Crops*
21000 Novi Sad, M. Gorkog 30, Yugoslavia
Tel/fax: 00 381 21 413 833

Igor Molnar, *Institute of Field and Vegetable Crops*
Maksima Gorkog 30, 21000 Novi Sad, Yugoslavia
Tel/fax: 00 381 21 413 833, E-mail: igor-fam@Eunet.yu

SUMMARY

The cultivated sunflower (*Helianthus annuus*) is one of most important oil crops in the world. Although sunflower is primarily grown for extraction of oil there is a limited production of non-oilseed types used for confections or as a bird feed.

The objective of this research was the development of B- and R-lines with stable and high oleic acid content and modified tocopherol composition, with high GCA and SCA values for the two most important agronomic characters (seed yield and oil yield) and high tolerance to Phomopsis.

B-lines [Lg-21 (tph1, Ol), Lg-24 (tph1, tph2) Lg-25 (tph2, Ol)] and R-lines (near-isogenic lines for three genes, i.e., Tph1, Tph2 and Ol-VK 66) were used as donors of tph1 and tph2 recessive alleles and Ol gene, in a backcross program with lines for high GCA and SCA values and high tolerance to Phomopsis: B-lines (Ha-74, Ha-98 CMS-III-8) and R-lines (RHA-583, RHA-576, RHA-SEL, RHA-SNRF). Tocopherol composition was determined by thin-layer chromatography followed by the Emmerie-Engel reaction and densitometer quantification. The screening for oleic acid content was performed by gas liquid chromatography. The program of incorporation of tph1 and tph2 alleles was based on the backcross method for the recessive traits of sunflower seeds, which includes the selfing of backcrossed plants and simultaneous individual crossing with the recurrent line.

As a result of this investigation we made a set of new inbred lines with high or normal content of oleic acid and modified tocopherol composition which may be used for developing materials with certain levels of oleic and linoleic acids and a desirable content of alpha, beta, gamma and delta-tocopherol. As a final result, a set of specialty oils might be obtained for human consumption and industrial purposes.

KEY WORDS: sunflower, tocopherol composition, oleic acid, oil quality

INTRODUCTION

The cultivated sunflower (*Helianthus annuus*) is counted among most important oil crops in the world. Sunflower oil is in a group of plant oils of highest quality. Composition of higher fatty acids and contents of tocopherol, sterol, carotenoids and other compounds determine oil quality. Sunflower is considered a promising species for genetic modification of oil quality (Scharp, 1986).

Fatty acids composition and mutual ratios are basic parameters determining oil quality. Standard sunflower oil contains linoleic acid, oleic acid, palmitic acid, stearic acid, and small amounts of other higher fatty acids. Linoleic and oleic acid take 90% of the total content, 55-65% the former and 20-30% the latter (Škoric *et al.*, 1994). The first source of high oleic acid was developed by Soldatov (1976) who treated seeds of the variety VNIMK 8931 with a 0.5% solution of dimethylsulphate (DMS). In the M₃ generation, Soldatov selected individual plants that contained over 50% of oleic acid which, after further selection, gave rise to the variety Pervenetz containing 80-90% oleic acid in oil. Pervenetz had been extensively used for the development of high oleic inbred lines and hybrids although genetic studies have not provided a definite answer on the mode of inheritance and the number of genes that control this characteristic. Fick (1984) reported that the high oleic acid content is controlled by a single gene for partial dominance (Ol), Urie (1985) and Miler *et al.* (1987) reported one dominant (Ol) and one recessive modifier gene (ml), while Fernandez-Martinez *et al.* (1989) reported three complementary genes (Ol₁, Ol₂, Ol₃).

Not only the content of higher fatty acids but also the composition of tocopherol may be modified by genetic manipulations. The standard sunflower oil contains about 95% of alpha-tocopherol, 3% of beta-tocopherol, 2% of gamma-tocopherol and delta-tocopherol in traces. By selfing the variety VNIIMK 8931, Demurin (1988) obtained an inbred line that contained 50% of alpha-tocopherol and 50% of beta-tocopherol. The high content of beta-tocopherol was controlled by a single recessive gene, tph₁. Further searching through the VIR gene bank, the same author came across the population No. 44 containing the recessive gene tph₂ which controls high content of gamma-tocopherol (5% of alpha-tocopherol and 95% of gamma-tocopherol). By combining these two recessive genes, Demurin (1993) obtained an inbred line with a high content of delta-tocopherol (8% of alpha-tocopherol, 84% of gamma-tocopherol and 8% of delta-tocopherol).

The objective of this study was to develop B- and Rf-inbred lines with a high content of oleic acid and modified tocopherol composition, good GCA and SCA for seed and oil yields and high tolerance to Phomopsis. These lines may be subsequently used for the development of hybrids with modified oil content.

MATERIAL AND METHOD

In this study, the donor lines for oil quality were inbred lines developed at VNIIMK, Krasnodar, Russia:

1. B-lines: Lg-21 (Ol, tph₁), Lg-25 (Ol, tph₂) and Lg-24 (tph₁, tph₂)
2. Rf-lines: VK-66-1 (tph₁, tph₂), VK-66-2 (Ol, tph₁) and VK-66-3 (Ol, tph₂)

The donors for good GCA and SCA and high tolerance to Phomopsis had been developed at the Institute of Field and Vegetable Crops, Novi Sad, Yugoslavia:

1. B-lines: Ha-74, Ha-981 and CMS-3-8
2. Rf-lines: RHA-583, RHA-576, RHA-SEL and RHA-SNRF-b

In the first year of the experiment (1993), each B-line tolerant to Phomopsis was crossed to all B-lines - donors for quality, and each Rf-line tolerant to Phomopsis to all Rf-lines - donors for quality. During the autumn/winter of 1993/94, backcross with the lines tolerant to

Phomopsis (F_1BC_{P1}) and selfing of F_1 hybrids to produce the F_2 generation were completed in a greenhouse. The female plants were emasculated by hand, in early morning, before anthers open up. In 1994, 1995, 1996 and 1997, the selfing program was continued for development of new inbred lines. The conversion of selected B-lines into the cms-sterile form started in the summer of 1995. Three generations were produced each year, first in the period January-April in the greenhouse, second in the period May-September in the field and third in the period September-January in the greenhouse.

After harvest of each generation, individual plants were analyzed for the composition of tocopherols, oleic acid content and oil content in seed. Field generations were also assessed for resistance to Phomopsis. The composition of tocopherols was determined by the half-seed method (Demurin, 1994) and thin-layer chromatography (TLC), the composition of higher fatty acids by gas chromatography (GC), the oil content in seed by the method of nuclear magnetic resonance (NMR). Phomopsis tolerance of the newly developed material was rated on the scale 0-5 (0-resistant, 5-susceptible).

RESULTS AND DISCUSSION

The inbred lines from VNIIMK, Krasnodar (Russia) used in this study as donors for oil quality, i.e., donors of genes Ol , tph_1 and tph_2 , have poor agronomic characteristics, in the first place high susceptibility to Phomopsis. This is why they had to be crossed to inbred lines from the Institute of Field and Vegetable Crops in Novi Sad which have good agronomic characteristics and high tolerance to Phomopsis. A part of the obtained F_1 generations was selfed for the production of the F_2 generation, the other part was backcrossed to the donors for Phomopsis resistance. In this way we made an initial population for the development of inbred lines with modified oil quality and high tolerance to Phomopsis.

After each generation of selfing, the materials were analyzed for the content of oleic acid in oil in order to identify the material with completely stable and high oleic acid content, i.e., the material heterozygous for this characteristic. Also, each generation selfed in the field was selected for Phomopsis resistance. Analyses of the composition of tocopherols and oil content in seed were done too. Inbred lines with increased content of oleic acid and high tolerance to Phomopsis were then selected on the basis of the obtained results.

Among the high oleic B-lines (Table 1), particularly promising are the inbreds UK-18 and UK-26, with high oil content in seed (> 50%), high tolerance to Phomopsis and the ratio alpha vs. beta tocopherol 50:50. These inbreds also have a stable and high content of oleic acid (> 80%). They have been converted to the sterile form, which is the basic prerequisite for the development of high oleic sunflower hybrids. Among the high oleic Rf-lines, there is a group of restorers distinguished for the oil content in seed over 50 %, high tolerance to Phomopsis and the ratio alpha vs. beta tocopherol 50:50 (Table 2). Some of these Rf-lines have a high and stable content of oleic acid, as indicated by low values of standard deviation. There is also a large set of high oleic restorers possessing medium tolerance to Phomopsis and various levels of oil content in seed (Table 3). In some of them, the oil content exceeds 55%, and in the restorer Rus-rf-01-154 it reaches 63.9 %. Regarding tocopherol, the set may be subdivided into a subset with dominant Tph_1 gene (100% alpha-tocopherol) and a subset with recessive tph_1 gene (50% alpha- tocopherol and 50% beta-tocopherol).

The newly developed B- and Rf-lines may be used for the development of high oleic sunflower hybrids possessing a high oil content in seed and different compositions of tocopherols. Taking in consideration the high tolerance to Phomopsis in some of these lines, it seems reasonable to expect of them to render resistant hybrids. Combining ability tests that started this year produced preliminary results which indicate that we are close to developing high oleic sunflower hybrids which match the conventional hybrids regarding the important

agronomic traits. According to *Friedt et al.* (1994), agronomic inferiority has been the main obstacle to a large-scale commercial production of high oleic sunflower hybrids.

The results of *Demurin et al.* (1994) and *Škorića et al.* (1996) indicate that the stability of high oleic sunflower oil is threefold that of the conventional oil. Combining high oleic content with increased contents of beta-, gamma- and delta-tocopherol may further increase oil stability. *Demurin* (1996) states that the presence of gamma-tocopherol in oil and a high proportion of oleic acid increase the stability of sunflower oil 16.4 times over that in the conventional sunflower oil.

CONCLUSION

Following conclusion were drawn on the basis of the obtained results.

- We developed a large set of high oleic B-lines that are tolerant to Phomopsis and that possess various compositions of tocopherols.
- Also, we developed high oleic and high oil Rf-lines that are highly tolerant to Phomopsis and that possess various compositions of tocopherols.
- The newly developed B- and Rf-lines will be used for the development of new sunflower hybrids with modified oil quality.

LITERATURE

- Demurin Ya. N.* 1988 Genetički analiz sastava tocopherolov v semenah podsolnečnika. Avtoreferat disertacije za KBN: 1-12.
- Demurin Ya. N.* 1993 Genetic variability of tocopherol composition in sunflower seeds. *Helia* 16: 59-62.
- Demurin Ya. N., Škoric D., Popov P., Efimenko S. and Bochkovoy A.* 1994.: Tocopherol genetics in sunflower breeding for oil quality. Proc. of EUCARPIA- Symposium on Breeding of Oil and Protein Crops, Albena, Bulgaria, 193-197.
- Demurin Ya. N i Lecic Nada* 1994.: Određivanje sastava tocopherol metodom "pola zrna" Zbornik savetovanjao unapre cenju uljarstva Jugoslavije, 24-27 maj, Bar, 255-258.
- Demurin Ya. N., Škoric D. and Karlovic D.* 1996.: Genetic variability of tocopherol composition in sunflower seeds as a basis of breeding for improved oil quality. *Plant Breeding* 115, 33-36.
- Fernandez-Martinez J., Jimenez A., Dominguez J., Garcia J.M., Garces R. and Mancha M.* 1989.: Genetic analysis of the high oleic acid content in cultivated sunflower (*Helianthus annuus* L). *Euphytica*: 41, 39-51.
- Fick G.N.* 1984.: Inheritance of high oleic acid in the seed oil of sunflower. P. 9 Proc.Sunflower Research Workshop. Bismarck, ND. National Sunflower Association
- Friedt W., Ganssmann M. and Korell M.,* 1994.: Improvement of sunflower oil quality. Proc of EUCARPIA – Symposium on Breeding of Oil and Protein Crops, Albena, Bulgaria. 1-29.
- Miler F.J., Zimmerman C.D. and Vick A.B.* 1987.: Genetic control of high oleic acid content i sunflower oil. *Crop Science*, Vol. 27, Septembar-October, 923-926.
- Scharp W.R.,* 1986: Opportunities for biotechnology in the development of new edible vegetable oil products. *YAOCS*, 63, 5: 594-600
- Soldatov K.J.,* 1976.: Chemical mutagenesis for sunflower breeding. Proc. of the 7th Int. Sunflower Conference, Krasnodar, Russia: 352-357.
- Škoric D., Demurin Ya. N., Jovic S., Lecic Nada i Verešbaranji I.* 1994.: Stvaranje hibrida suncokreta sa različitim kvalitetom ulja. Zbornik savetovanjao unapredenju uljarstva Jugoslavije, 24-27 maj, Bar, 8-15.
- Škoric D., Demurin Ya. N. and Jovic S.* 1996.: Development of hybrids with various oil quality. Proc. of 14th Intern. Sunflower Conference, Beijing/Shenyang, China, 54-59.
- Škoric D., Jovic S. and Lečić Nada,* 1998.: Sunflower breeding for different oil quality. Proc. of the 15th EUCARPIA Congress, Viterbo, Italy, 339-346.
- Urie L. A.,* 1985.: Inheritance of high oleic acid in sunflower. *Crop Science*, Vol. 25, November-December, 986-989.

Table 1. B-lines with increased oleic acid content and various compositions of tocopherols

No.	Line	Fatty acids composition (%)		Tocopherols (%)		Oil content (%)	Phomopsis resistance (scale 0-5)
		oleic	linoleic	alpha	beta		
1.	UK-4	88	3	50	50	43.86	4
2.	UK-8	85	5	100	/	46.90	3
3.	UK-9	86	3	50	50	48.36	4
4.	UK-18	86	4	40	60	52.72	2
5.	UK-26	88	3	50	50	53.05	2
6.	UK-29	86	3	50	50	41.64	2
7.	UK-56	88	2	50	50	42.80	2
8.	UK-57	89	2	50	50	47.96	2
9.	UK-58	86	4	50	50	43.92	2
10.	UK-87	88	3	100	/	47.10	2
11.	UK-88	86	4	50	50	/	2
12.	UK-89	87	3	50	50	/	2
13.	UK-90	88	3	50	50	40.90	2
14.	UK-91	88	2	50	50	40.17	2
15.	UK-96	86	4	100	/	49.10	2

Table 2. Selected high oleic Rf-lines having high tolerance to Phomopsis and oil content in seed over 50%

No.	Line	Oleic acid content (%)			
		Average	Max	Min	Std.Dev.
1.	Rf-kv-1	85.4	87	80	3.05
2.	Rf-kv-2	88.8	89	88	0.45
2.	Rf-kv-3	88.6	91	88	1.34
4.	Rf-kv-4	88.0	90	86	1.41
5.	Rf-kv-5	85.6	88	78	4.28
6.	Rf-kv-6	87.6	89	85	1.67
7.	Rf-kv-7	89.2	91	88	1.30
8.	Rf-kv-8	88.2	89	87	0.84
9.	Rf-kv-9	81.2	90	49	18.01
10.	Rf-kv-10	83.0	88	70	7.55
11.	Rf-kv-11	87.8	89	87	0.84
12.	Rf-kv-12	87.4	89	86	1.34
13.	Rf-kv-13	71.8	89	41	19.59
14.	Rf-kv-14	82.4	87	71	6.54
15.	Rf-kv-15	85.2	89	73	6.87
16.	Rf-kv-16	88.2	89	87	0.84

Table 3. High oleic Rf-lines with various compositions of tocopherols

No.	Line	Fatty acids composition (%)		Tocopherols (%)		Oil content (%)	Phomopsis resistance (scale 0-5)
		oleic	linoleic	alpha	beta		
1.	Rus-rf-tph-ol-8	86	4	50	50	51.21	4
2.	Rus-rf-tph-ol-10	86	3	90	10	40.50	2
3.	Rus-rf-tph-ol-14	90	3	50	50	54.37	4
4.	Rus-rf-tph-ol-15	82	10	50	50	54.05	4
5.	Rus-rf-tph-ol-16	87	3	50	50	54.22	4
6.	Rus-rf-tph-ol-23	91	2	100	/	53.87	4
7.	Rus-rf-tph-ol-25	89	2	50	50	55.38	3
8.	Rus-rf-tph-ol-26	90	2	50	50	56.68	3
9.	Rus-rf-tph-ol-35	87	3	50	50	50.05	3
10.	Rus-rf-tph-ol-37	84	6	50	50	52.08	3
11.	Rus-rf-tph-ol-39	89	2	50	50	42.32	4
12.	Rus-rf-tph-ol-43	88	3	50	50	/	5
13.	Rus-rf-tph-ol-51	87	5	50	50	49.59	4
14.	Rus-rf-tph-ol-69	85	7	50	50	39.54	4
15.	Rus-rf-tph-ol-81	83	5	50	50	55.92	4
16.	Rus-rf-tph-ol-83	86	6	50	50	49.93	4
17.	Rus-rf-tph-ol-87	81	9	50	50	51.70	4
18.	Rus-rf-tph-ol-93	84	4	50	50	51.40	4
19.	Rus-rf-tph-ol-96	87	4	50	50	33.73	4
20.	Rus-rf-ol-27	85	4	100		47.68	4
21.	Rus-rf-ol-38	89	4	100		54.70	4
22.	Rus-rf-ol-39	87	5	100		44.87	4
23.	Rus-rf-ol-54	90	2	100		51.71	3
24.	Rus-rf-ol-67	89	2	100		49.41	4
25.	Rus-rf-ol-68	87	4	100		47.88	4
26.	Rus-rf-ol-70	85	5	100		43.71	4
27.	Rus-rf-ol-77	91	1	100		58.39	4
28.	Rus-rf-ol-78	88	2	100		56.00	4
29.	Rus-rf-ol-80	87	4	100		45.98	4
30.	Rus-rf-ol-91	88	3	100		55.26	3
31.	Rus-rf-ol-94	86	3	100		43.55	4
32.	Rus-rf-ol-134	85	6	100		43.46	4
33.	Rus-rf-ol-140	84	4	100		46.43	4
34.	Rus-rf-ol-142	88	2	100		59.13	3
35.	Rus-rf-ol-154	84	4	100		63.90	3
36.	Rus-rf-ol-206	88	3	100		42.02	4
37.	Rus-rf-ol-207	90	2	100		53.46	4
38.	Rus-rf-ol-209	90	1	100		52.83	4
39.	Rus-rf-ol-222	90	2	100		52.57	4
40.	Rus-rf-ol-242	91	3	100		52.57	4