

Isolation of Phytochemical Constituents from *Hunteria umbellata* K. SchumIftikhar Ali<sup>1\*</sup>, Abiodun Falodun<sup>2</sup>, Baraa Siyo<sup>3</sup>, Bankeu K. J. Jules<sup>4</sup>, Hidayat Hussain<sup>5</sup>, Peter Langer<sup>6,7</sup><sup>1</sup>Department of Chemistry, Karakoram International University, 15100-Gilgit, Pakistan.<sup>2</sup>Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.<sup>3</sup>Department of Chemistry, Tishreen University, Latakia, Syria.<sup>4</sup>Department of Chemistry, Faculty of Science, University of Bamenda, P.O. Box 39 Bambili-Cameroon, Cameroon.<sup>5</sup>Leibniz Institute of Plant Biochemistry, Department of Bioorganic Chemistry, Weinberg 3, D-06120 Halle (Salle), Germany.<sup>6</sup>Institut für Chemie, Universität Rostock, Albert-Einstein-Str. 3a, 18059 Rostock, Germany.<sup>7</sup>Leibniz-Institut für Katalyse an der Universität Rostock e.V., Albert Einstein Str. 29a, 18059 Rostock, Germany.

## ARTICLE INFO

## ABSTRACT

## Article history:

Received 17 January 2019

Revised 10 March 2019

Accepted 19 March 2019

Published online 04 April 2019

**Copyright:** © 2019 Ali *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Hunteria umbellata* K. Schum has been reported for the treatment of diabetes in Nigeria. In the present study, the ethanolic extract of dried leaves of *Hunteria umbellata* K. Schum was investigated for the chemical principles. The isolated pure compounds were characterized by NMR, IR and Mass spectral studies. Ursolic acid (1), oleanolic acid (2) and squalene (3) were the main constituents isolated from the extract.

**Keywords:** *Hunteria umbellata*, Ursolic acid, Oleanolic acid, Squalene

## Introduction

Herbal medicine has been considered the best source of naturally produced compounds that account for drug discovery that is an expensive, long, and complex process.<sup>1</sup> The polyherbal formulations have been reported for the treatment of type 2 diabetes, a chronic metabolic disorder resulting from either insulin insufficiency or insulin dysfunction.<sup>2, 3</sup> As recently reported, approximately one-third of all stroke patients have diabetes<sup>4</sup>, and diabetes mellitus is one of the risk factors associated with increased breast cancer mortality.<sup>6, 7</sup> The genus *Hunteria* belongs to the family Apocynaceae. *Hunteria umbellata* K. Schum (Syn: *Carpodinus umbellatus* K. Schum.) has been reported to be used in the treatment of diabetes and obesity in Nigeria.<sup>8</sup> *Hunteria umbellata* K. Schum (*H. umbellata*) has been reported for its aphrodisiac effect,<sup>9</sup> and as phosphodiesterase type 5 (PDE-5) and arginase inhibitor,<sup>10</sup> antioxidant, hypolipidemic,<sup>8</sup> analgesic,<sup>11</sup> antiinflammatory<sup>12</sup> etc. *H. umbellata* has also been reported to decrease hepatic glucose-6-phosphatase activity.<sup>13</sup> Phytochemically the plant *H. umbellata* has been reported to contain saponins, steroids, and alkaloids.<sup>14</sup>

The main aim of this research was to investigate the ethanolic extract of *H. umbellata* for various chemical constituents. As a result of our investigation, ursolic acid (1), oleanolic acid (2) and squalene (3) were isolated and identified. However, compound (2) and (3) are being reported for the first time from *H. umbellata*.

\*Corresponding author. E mail: [iftikhar.ali@kiu.edu.pk](mailto:iftikhar.ali@kiu.edu.pk)  
Tel: +92-3339665739

**Citation:** Ali I, Falodun A, Siyo B, Jules BKJ, Hussain H, Langer P.

Isolation of Phytochemical Constituents from *Hunteria umbellata* K. Schum. Trop J Nat Prod Res. 2019; 3(3):95-98. [doi.org/10.26538/tjnpr/v3i3.6](https://doi.org/10.26538/tjnpr/v3i3.6)

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

## Materials and Methods

## Instrumentation and chromatography

The IR spectra were recorded on JASCO FT/IR-5300 spectrophotometer. <sup>1</sup>H (300 MHz), <sup>13</sup>C (300 MHz; 62 MHz), and 2D NMR spectra were recorded using JEOL JNM-A500 and/or Varian INOVA-500 spectrometers. MS spectra were recorded using JEOL JMS-700 instrument. Silica-gel 60 (Merck) column chromatography was employed for isolation and purification, and pre-coated silica gel plates (Merck) were used for TLC study. The spots were visualized by spraying with ceric sulphate followed by heating.

## Plant material

Fresh leaves of *H. umbellata* were collected from the Ugbowo Campus of the University of Benin, Nigeria in January 2011. The plant material was authenticated at the Forestry Research Institute of Nigeria (FRIN), Ibadan and an herbarium specimen with the voucher number FHI-107678 was deposited.

## Extraction and fractionation

The fresh leaves of *H. umbellata* were collected, dried in shade and powdered (500 g), macerated with 2L EtOH at room temperature for 32 hours. The crude extract was obtained by evaporating the filtrate solvent under reduced pressure at 40°C with the help of rotary evaporator. The combined ethanolic extract (82 g) was further fractionated with n-heptane, chloroform, and ethyl acetate. The n-heptane soluble fraction was employed to medium pressure liquid chromatography (MPLC) to get 100% n-heptane and 100% CHCl<sub>3</sub> sub fractions.

## Phytochemical isolation

The 100% n-heptane sub-fraction was subjected to column chromatography over silica gel eluting with n-heptane (100%), EtOAc:n-heptane (10% to 90%), EtOAc (100%) in increasing order of polarity. The fraction obtained from (100%) EtOAc elution gave ursolic acid (1, 110 mg). The 100% CHCl<sub>3</sub> sub fraction was subjected

to CC over silica gel eluting with n-heptane (100%), EtOAc:n-heptane (10% to 90%), EtOAc (100%), MeOH:EtOAc (1% to 10%), MeOH (100%) in increasing order of polarity. The fraction obtained from (60%) EtOAc elution gave oleanolic acid (**2**, 32 mg) and in further elution the fractions obtained from (70% EtOAc:n-heptane to 5% MeOH:EtOAc) were combined and this yielded squalene (**3**, 7.3 mg). The structures of the isolated compounds as shown in figure 1, were determined using a combination of spectral techniques i.e. IR, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, and MS, the compound structures were further verified by direct comparison with the literature.

## Results and Discussion

### Ursolic acid (1)

Pure white solid; 110 mg; IR (ATR-Messung)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3400, 2924, 2869, 1687, 1455, 1386, 1028, 996, 661. <sup>1</sup>H-NMR (DMSO, 300 MHz)  $\delta$ : 5.12 (brs, 1H, H-12), 4.26-4.28 (m, 1H, H-3), 3.49-3.37 (m, 1H), 1.09-1.55 (m, 7H), 1.44-1.29 (m, 11H), 0.64-1.03 (m, 27H). <sup>13</sup>C-NMR (DMSO-d<sub>6</sub>, 300 MHz): See Table 1. HR-ESI-MS (negative ion mode) m/z: 455.3527. HR-EI-MS m/z: 456 (1), 262 (5), 248 (100), 203 (60), 133 (49), 69 (19), 55 (17), 44 (29), 39 (4). Compound (**1**) was identified as ursolic acid.

### Oleanolic acid (2)

Colorless solid; 32 mg; IR (ATR-Messung)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3400, 2923, 2855, 1683, 1651, 1455, 1028, 996, 661. <sup>1</sup>H-NMR (CDCl<sub>3</sub>/CD<sub>3</sub>OH, 300 MHz)  $\delta$ : 5.01 (brs, 1H, H-12), 4.00-3.98 (m, 1H, H-3), 3.49-3.37 (m, 1H, H-18), 2.05-1.68 (m, 7H), 1.44-1.29 (m, 11H), 1.11-0.55 (m, 27H). <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 300 MHz): See Table 1. HR-ESI-MS (-ve ion mode) m/z: 455.3530.

### Squalene (3)

Oily substance; 7.3 mg; IR (ATR-Messung,  $\text{cm}^{-1}$ ): 2962, 2919, 2852, 1722, 1666, 1444, 1376, 1106, 834; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): 1.52 (18H, s, 6CH<sub>3</sub>, H-24, H-25, H-26, H-27, H-28, H-29), 1.60 (6H, s, 2CH<sub>3</sub>, H-1, H-30), 1.90-2.02 (20H, m, 10CH<sub>2</sub>, H-4, H-5, H-8, H-9, H-12, H-13, H-16, H-17, H-20, H-21), 4.99-5.07 (6H, m, 6CH, H-3, H-7, H-11, H-15, H-19, H-22). <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 300 MHz): See Table 2; GC MS m/z (%): 410 (0.97), 341 (1.83), 149 (7.01), 81 (51.10), 69 (100), 41 (29.29); ESI MS (+ve ion mode): 410, 377, 338, 325, 301, 279 (100), 125.

Compound **1** was obtained as pure white solid. Based on HR-ESI-MS (negative ion mode) m/z: 455.3527 and <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectral data, the molecular formula of **1** was deduced to be C<sub>30</sub>H<sub>48</sub>O<sub>3</sub>. The <sup>1</sup>H-NMR spectrum showed the presence of one broad singlet at  $\delta$  5.12. The <sup>13</sup>C-NMR spectrum exhibited 30 well resolved signals. The carbonyl carbon for the acidic group was observed at  $\delta$  179.1 (C-28), and the two olefinic carbon signals were observed at  $\delta$  125.5 (CH, C-12), and 139.0 (C, C-13). The oxygenated methine carbon signal was observed at  $\delta$  77.7 (CH, C-3). Similarly seven methyl, nine methylene, seven methine, and seven quaternary carbons were observed (Table 1). These data showed that compound **1** is a triterpene derivative. By comparison with the NMR spectral data in the literature,<sup>15, 16</sup> the structure of the compound **1** was characterized and identified as ursolic acid.


Compound **2** was obtained as colourless solid. Based on HR-ESI-MS (negative ion mode) m/z: 455.3530 and <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectral data, the molecular formula of **2** was deduced to be C<sub>30</sub>H<sub>48</sub>O<sub>3</sub>. The <sup>1</sup>H-NMR spectrum indicated the presence of one broad singlet at  $\delta$  5.01 (1H, H-12). The <sup>13</sup>C-NMR spectrum exhibited 30 carbon signals. The carbonyl carbon for the acidic group was observed at  $\delta$  180.2 (C-28), and the two olefinic carbon signals were observed at  $\delta$  125.1 (CH, C-12), and 137.8 (C, C-13). The oxygenated methine carbon signal was observed at  $\delta$  78.3 (CH, C-3). Similarly seven methyl, ten methylene, five methine, and eight quaternary carbons were observed (Table 1). The spectral data showed that compound **2** is a triterpene derivative. By comparison with the previous NMR spectral data in the literature,<sup>15, 16</sup> the structure of the compound **2** was characterized and identified as oleanolic acid.

Compound **3** was obtained as oily substance. Based on GC-MS m/z: 410 and <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectral data, the molecular formula of **3** was deduced to be C<sub>30</sub>H<sub>50</sub>. The <sup>1</sup>H-NMR spectrum indicated the presence of one singlet at  $\delta$  1.52 for six methyl protons (18H, H-24, H-25, H-26, H-27, H-28, H-29). Another singlet, for two methyl protons, was observed at  $\delta$  1.60 (6H, H-1, H-30). Similarly two multiplet signals were observed at  $\delta$  1.90-2.02, and 4.99-5.07 for ten methylene protons (20H; H-4, H-5, H-8, H-9, H-12, H-13, H-16, H-17, H-20, H-21) and six methine protons (6H; H-3, H-7, H-11, H-15, H-19, H-22), respectively. The <sup>13</sup>C-NMR spectrum exhibited 15 carbon signals. A signal at  $\delta$  15.9 was assigned to two methyl carbons (C-27, C-28), and another signal at  $\delta$  16.0 was assigned to two other methyl carbons (C-26, C-29). Two methyl carbons (C-25, C-30) were observed at  $\delta$  17.6, and methyl carbons (C-1, C-24) were observed at  $\delta$  25.6. The methylene carbons were observed at  $\delta$  26.6 (C-8, C-17), 26.7 (C-4, C-21), 28.2 (C-12, C-13), 39.7 (C-9, C-16), and 39.7 (C-5, C-20). While the methine carbons were resonating at  $\delta$  124.2 (C-7, C-18), 124.2 (C-11, C-14), and 124.3 (C-3, C-22). The quaternary carbons were observed at  $\delta$  131.2 (C-2, C-23), 134.8 (C-6, C-19), and 135.0 (C-10, C-15). By comparison with the previous NMR spectral data in literature,<sup>17</sup> the structure of the compound **3** was characterized and identified as squalene.

*H. umbellata* has been investigated for preliminary analysis of saponins and its glycosides, alkaloids, steroid, tannins, oils, etc.<sup>14</sup> Other available reports<sup>18-20</sup> have focused on the phytochemical analysis of *H. umbellata* for the isolation of alkaloids. The triterpenoid saponin derivatives i.e. ursolic acid (**1**) and oleanolic acid (**2**) have already exhibited antimicrobial properties and its mechanism is established.<sup>21, 22</sup> Oleanolic acid has also exhibited anti-tubercular activity.<sup>23</sup> The derivatives of oleanolic acid have been reported for the treatment of pancreatic cancer,<sup>24</sup> and oleanolic acid saponins have shown significant antimicrobial properties,<sup>25</sup> and the combination of oleanolic acid with hypocrellin A has been reported for the treatment of hepatocellular carcinoma diseases,<sup>26</sup> and oleanolic acid derivative SZC014 has exhibited anticancer activity.<sup>27</sup> Oleanolic acid has shown antidiabetic,<sup>28, 29</sup> and hepatoprotective activities,<sup>30,31</sup> which may establish the use of *H. umbellata* as antidiabetic by the local inhabitants in Nigeria.

## Conclusion

The present study reveals that the plant extract contains certain bioactive compounds. The triterpenoids including ursolic acid and oleanolic acid isolated from this plant have previously been shown to have hepatoprotective and antiinflammatory potentials.



**Figure 1:** Molecular structures of chemical constituents isolated from *H. umbellata*.

**Table 1:** <sup>13</sup>C-NMR chemical shifts (δ, ppm) of Compounds (1) in DMSO-d<sub>6</sub>, 300 MHz and (2) in CDCl<sub>3</sub>, 300 MHz.

C No	Compound (1)		Literature		Compound (2)		Literature	
	δ C		15	16	δ C		15	16
1	39.1	CH <sub>2</sub>	39.2	40.3	38.3	CH <sub>2</sub>	39.0	38.7
2	27.9	CH <sub>2</sub>	28.2	29.4	27.6	CH <sub>2</sub>	28.1	27.8
3	77.7	CH	78.2	79.4	78.3	CH	78.2	77.8
4	39.3	C	39.6	40.6	39.0	C	39.4	39.1
5	55.7	CH	55.9	57.1	54.9	CH	55.9	55.5
6	18.9	CH <sub>2</sub>	18.8	19.9	17.9	CH <sub>2</sub>	18.8	18.7
7	33.6	CH <sub>2</sub>	33.7	34.8	32.6	CH <sub>2</sub>	33.4	33.0
8	40.0	C	40.1	41.2	38.2	C	39.8	39.5
9	47.7	CH	48.1	49.3	52.4	CH	48.2	49.4
10	37.2	C	37.5	38.5	36.5	C	37.4	37.1
11	23.7	CH <sub>2</sub>	23.7	24.9	23.8	CH <sub>2</sub>	23.8	23.6
12	125.5	CH	125.7	126.9	125.1	CH	122.6	122.3
13	139.0	C	139.3	140.5	137.8	C	144.8	144.6
14	40.4	C	42.6	43.7	41.6	C	42.2	41.9
15	28.4	CH <sub>2</sub>	28.8	29.9	29.2	CH <sub>2</sub>	28.4	28.1
16	24.7	CH <sub>2</sub>	25.0	26.2	22.8	CH <sub>2</sub>	23.8	23.6
17	42.5	C	48.1	49.3	47.3	C	46.7	47.9
18	53.2	CH	53.6	54.8	38.7	CH	42.1	41.8
19	39.4	CH	39.5	40.7	36.4	CH <sub>2</sub>	46.6	46.2
20	39.2	CH	39.4	40.6	29.2	C	31.0	30.7
21	31.1	CH <sub>2</sub>	31.1	32.3	30.2	CH <sub>2</sub>	34.3	34.0
22	37.4	CH <sub>2</sub>	37.4	38.7	26.3	CH <sub>2</sub>	33.2	32.9
23	29.1	CH <sub>3</sub>	28.8	30.1	27.5	CH <sub>3</sub>	28.8	28.5
24	16.9	CH <sub>3</sub>	16.5	17.5	16.4	CH <sub>3</sub>	16.5	16.6
25	16.1	CH <sub>3</sub>	15.7	16.4	14.9	CH <sub>3</sub>	15.6	15.4
26	17.8	CH <sub>3</sub>	17.5	18.5	16.5	CH <sub>3</sub>	17.5	17.7
27	24.1	CH <sub>3</sub>	24.0	25.2	20.5	CH <sub>3</sub>	26.2	25.9
28	179.1	C	179.7	181.6	180.2	C	180.0	180.0
29	17.9	CH <sub>3</sub>	17.5	18.6	15.1	CH <sub>3</sub>	33.4	33.0
30	21.9	CH <sub>3</sub>	21.4	22.3	23.0	CH <sub>3</sub>	23.8	23.6

**Table 2:** <sup>13</sup>C-NMR chemical shifts (δ, ppm) of Compound (3) in CDCl<sub>3</sub>, 300 MHz.

C No.	Compound (3) δ C		Literature <sup>17</sup>
1, 24	25.67	CH <sub>3</sub>	25.67
2, 23	131.22	C	131.22
3, 22	124.39	CH	124.40
4, 21	26.76	CH <sub>2</sub>	26.77
5, 20	39.74	CH <sub>2</sub>	39.75
6, 19	134.87	C	134.88
7, 18	124.26	CH	124.20
8, 17	26.65	CH <sub>2</sub>	26.66
9, 16	39.72	CH <sub>2</sub>	39.72
10, 15	135.08	C	135.08
11, 14	124.29	CH	124.30
12, 13	28.26	CH <sub>2</sub>	28.27
25, 30	17.66	CH <sub>3</sub>	17.65
26, 29	16.02	CH <sub>3</sub>	16.02
27, 28	15.98	CH <sub>3</sub>	15.98

**Conflict of interest**

The authors declare no conflict of interest.

**Authors' Declaration**

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

**Acknowledgments**

The authors thank Prof. Dr. Peter Langer for financial support. We thank the plant taxonomist Prof. MacDonald Idu Department of Plant Biology and Biotechnology, University of Benin for the identification of the plant sample.

## References

- Zhang A, Sun H, Wang X. Mass spectrometry-driven drug discovery for development of herbal medicine. *Mass Spectrom Rev.* 2018; 37(3):307-320.
- Telapolu S, Kalachavedu M, Punnoose AM, Bilikere D. MD-1, a poly herbal formulation indicated in diabetes mellitus ameliorates glucose uptake and inhibits adipogenesis - an in vitro study. *BMC Compl Altern Med.* 2018; 18(1):113.
- Khalili N, Fereydoonzadeh R, Mohtashami R, Mehrzadi S, Heydari M, Huseini HF. Silymarin, Olibanum, and Nettle, A Mixed Herbal Formulation in the Treatment of Type II Diabetes: A Randomized, Double-Blind, Placebo-Controlled, Clinical Trial. *J Evid Based Compl Altern Med.* 2017; 22(4):603-608.
- Lau LH, Lew J, Borschmann K, Thijs V, Ekinci EI. The prevalence of Diabetes and its effects on Stroke Outcomes; a meta-analysis and literature review. *J Diabet Investig.* 2018. [Epub ahead of print]
- Mehmood K and Junaid N. Prevalence of undiagnosed type 2 diabetes mellitus in Pakistan: Results of screen-diabetes disease registry. *J Pak Med Assoc.* 2018; 68(8):1171-1178.
- Chang YL, Sheu WH, Lin SY, Liou WS. Good glycaemic control is associated with a better prognosis in breast cancer patients with type 2 diabetes mellitus. *Clin Exp Med.* 2018; 18(3):383-390.
- Maskarinec G, Fontaine A, Torfadottir JE, Lipscombe LL, Lega IC, Figueroa J, Wild S. The Relation of Type 2 Diabetes and Breast Cancer Incidence in Asian, Hispanic and African American Populations-A Review. *Can J Diabet.* 2018; 42(1):100-105.
- Ajiboye TO, Hussaini AA, Nafiu BY, Ibitoye OB. Aqueous seed extract of *Hunteria umbellata* (K. Schum.) Hallier f. (Apocynaceae) palliates hyperglycemia, insulin resistance, dyslipidemia, inflammation and oxidative stress in high-fructose diet-induced metabolic syndrome in rats. *J Ethnopharmacol.* 2017; 198:184-193.
- Oboh G, Adebayo AA, Ademosun AO, Abegunde OA. Aphrodisiac effect of *Hunteria umbellata* seed extract: Modulation of nitric oxide level and arginase activity in vivo. *Pathophysiol.* 2019; 26(1):39-47.
- Oboh G, Adebayo AA, Ademosun AO, Boligon AA. In vitro inhibition of phosphodiesterase-5 and arginase activities from rat penile tissue by two Nigerian herbs (*Hunteria umbellata* and *Anogeissus leiocarpus*). *J Basic Clin Physiol Pharmacol.* 2017; 28(4):393-401.
- Adeyemi OO, Adeneye AA, Alabi TE. Analgesic activity of the aqueous seed extract of *Hunteria umbellata* (K. Schum.) Hallier f. in rodents. *Indian J. Exp. Biol.*, 2011; 49(9):698-703.
- Igbe I, Ching FP, Eromon A. Anti-inflammatory activity of aqueous fruit pulp extract of *Hunteria umbellata* K. Schum in acute and chronic inflammation. *Acta Pol Pharm.* 2010; 67(1):81-5.
- Adeneye AA, Adenekan SO, Adeyemi OO, Agbaje EO. Glucose utilization and anti-oxidative mechanisms of the aqueous *Hunteria umbellata* seed extract in alloxan-induced diabetic rats. *Nig J Physiol Sci.* 2014; 29(1):71-9.
- Falodun A, Nworgu ZA, Ikponmwonsa MO. Phytochemical components of *Hunteria umbellata* (K. Schum) and its effect on isolated non-pregnant rat uterus in oestrus. *Pak J Pharm Sci.* 2006; 19(3):256-8.
- Seebacher W, Simic N, Weis R, Saf R, Kunert O. Spectral Assignments and Reference Data: Complete assignments of <sup>1</sup>H and <sup>13</sup>C-NMR resonances of oleanolic acid, 18a-oleanolic acid, ursolic acid and their 11-oxo derivatives. *Magn Reson Chem.* 2003; 41:636-638.
- Woo KW, Han JY, Choi SU, Kim KH, Lee KR. Triterpenes from *Perilla frutescens* var. *acuta* and Their Cytotoxic Activity. *Nat Prod Sci.* 2014; 20(2):71-75.
- Alam K, Harun-or-Rashid, Rahman AA, Baki A, Sadik MG. Chemical constituents of *Hemigraphis hirta* T. anders (Acanthaceae). *Pak J Biol Sci.* 2002; 5(11):1264-1266.
- Morita Y, Hesse M, Schmid H. [Concerning the alkaloid eripine from *Hunteria umbellata*]. *Helv Chim Acta.* 1968; 51(6):1438-443.
- Bevan CW, Patel MB, Rees AH. The seed alkaloids of *Hunteria umbellata*. *Tetrahedron* 1967; 23(9):3809-21.
- Bevan CW, Patel MB, Rees AH. The seed alkaloids of *Hunteria umbellata*: the x-ray crystal structure of corymine hydrobromide monohydrate. *Chem Ind.* 1965; 14:603-604.
- Park SN, Lim YK, Choi MH, Cho E, Bang IS, Kim JM, Ahn SJ, Kook JK. Antimicrobial Mechanism of Oleanolic and Ursolic Acids on *Streptococcus mutans* UA159. *Curr Microbiol.* 2018; 75(1):11-19.
- Zhou T, Li Z, Kang OH, Mun SH, Seo YS, Kong R, Shin DW, Liu XQ, Kwon DY. Antimicrobial activity and synergism of ursolic acid 3-O-alpha-L-arabinopyranoside with oxacillin against methicillin-resistant *Staphylococcus aureus*. *Int J Mol Med.* 2017; 40(4):1285-1293.
- Ebeh Messanga R, Dominique Serge NB, Abouem AZA, Norbert MNI, Esther Del Florence MN, Patrick Herve BD, Maximilienne Ascension N, Alex De Theodore A, Dieudonne Emmanuel P, Christian GB, Koert U. Rauvolfianine, a new antimycobacterial glyceroglycolipid and other constituents from *Rauvolfia caffra*. *Sond (Apocynaceae).* *Nat Prod Res.* 2018; 32(16):1971-1976.
- Song YL, Gao L, Tang ZY, Li HL, Sun B, Chu P, Qaed E, Ma XD, Peng JY, Wang SS, Hu M, Tang ZY. Anticancer effect of SZC015 on pancreatic cancer via mitochondria-dependent apoptosis and the constitutive suppression of activated nuclear factor B and STAT3 in vitro and in vivo. *Journal of Cellular Physiology*, 2019; 234(1):777-788.
- Tagousop CN, Tamokou JD, Kengne IC, Ngnokam D, Voutquenne-Nazabadioko L. Antimicrobial activities of saponins from *Melanthera elliptica* and their synergistic effects with antibiotics against pathogenic phenotypes. *Chem Cent J.* 2018; 12(97):1-9.
- Wang L, Wang J, Cao YY, Li WJ, Wang Y, Xu JL, Xu GL. Molecular evidence for better efficacy of hypocrellin A and oleanolic acid combination in suppression of HCC growth. *Eur J Pharmacol.* 2019; 842:281-290.
- Chu P, Li H, Luo R, Ahsan A, Qaed E, Shopit A, Ma X, Lin Y, Peng J, Zhang J, Wang S, Tang Z. Oleanolic acid derivative SZC014 inhibit cell proliferation and induce apoptosis of human breast cancer cells in a ROS-dependent way. *Neoplasma*, 2017; 64(5):681-692.
- Wang X, Li YL, Wu H, Liu JZ, Hu JX, Liao N, Peng J, Cao PP, Liang X, Hai CX. Antidiabetic effect of oleanolic acid: a promising use of a traditional pharmacological agent. *Phytother Res.* 2011; 25(7):1031-1040.
- Castellano JM, Guinda A, Delgado T, Rada M, Cayuela JA. Biochemical basis of the antidiabetic activity of oleanolic acid and related pentacyclic triterpenes. *Diabet.* 2013; 62(6):1791-1799.
- Gutierrez-Rebolledo GA, Siordia-Reyes AG, Meckes-Fischer M, Jimenez-Arellanes A. Hepatoprotective properties of oleanolic and ursolic acids in antitubercular drug-induced liver damage. *Asian Pac J Trop Med.* 2016; 9(7):644-51.
- Wang W, Chen K, Xia Y, Mo W, Wang F, Dai W, Niu P. The Hepatoprotection by Oleanolic Acid Preconditioning: Focusing on PPARalpha Activation. *PPAR Res.* 2018; 2018:3180396.