

Comparative assessment of the pattern of anti-hypertensive drugs prescribed in medicine and cardiology outpatient department

Bhargav Gaikwad, Sagar Bhagat*, Ketaki Patil

Department of Pharmacology,
Grant Medical College, Mumbai,
Maharashtra, India

Received: 27 June 2016

Accepted: 05 August 2016

***Correspondence to:**

Dr. Sagar Bhagat,

Email: sagarbhagat04@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Hypertension or high blood pressure is one of the strongest modifiable risk factors for cardiovascular and renal diseases and is a condition that afflicts almost 1 billion people worldwide. The objective of the present study was to compare the prescribing pattern of antihypertensive drugs in patient attending medicine and cardiology OPD.

Methods: It was an observational, cross sectional study conducted in the department of pharmacology in collaboration with the medicine and the cardiology department.

Results: The present study which was conducted for a period of 6 months in the medicine and cardiology OPD showed unequal distribution of the patients in the medicine and cardiology OPD, with 223 patients attended medicine OPD while 434 patients attended cardiology OPD. Amongst the patients attending medicine OPD, majority of the patients (61.88%) were stage I hypertensive (>140/90 mmHg) according to JNC VII classification whereas majority (69.38%) of those attending cardiology OPD were stage II hypertensive (>160/100). Calcium channel blockers (CCB) were the most commonly prescribed drug in medicine OPD followed by angiotensin converting enzyme inhibitors (ACE I) whereas beta blockers (BB) were most commonly prescribed in cardiology OPD followed by ACE I.

Conclusions: The study concludes that there is a wide variation in the prescription of the anti-hypertensive drugs among medicine and cardiology OPD. CCBs were most commonly used in medicine OPD whereas beta blockers were preferred in cardiology OPD. Monotherapy was preferred in medicine OPD, whereas combination therapy was preferred by cardiologist. Double and triple drug combination therapies were used more in comparison to monotherapy.

Keywords: OPD, Hypertension, High Blood pressure, Medicine, Cardiology, Prescription

INTRODUCTION

Hypertension or high blood pressure is one of the strongest modifiable risk factors for cardiovascular and renal diseases and is a condition that afflicts almost 1 billion people worldwide.¹ It is usually asymptomatic until the damaging effects of hypertension such as stroke, myocardial infarction, renal dysfunction, visual problems etc. are observed.¹ It is labeled as 'silent killer' because in considerable proportion of cases the disease tends to be asymptomatic for prolonged time.² It is also considered as an 'Iceberg' disease because unknown morbidity far exceeds the known morbidity.³ It ranks fourth in the world by its prevalence.⁴

Hypertension is an important public health challenge in both economically developing and developed countries. WHO health statistics 2012 revealed that, the prevalence of hypertension in India was 23.1% in men and 22.6% in women in equal or more than 25 years age.⁵ According to a recent review on "Global Burden of hypertension", the estimated prevalence of hypertension (in aged 20 years and older) in India is expected to increase by approximately 60% by 2025.⁶

It is well documented that, in addition to lifestyle modifications, convenient antihypertensive drug therapy substantially reduces the risk of hypertension-related morbidity and mortality.⁶ The treatment of hypertension

is constantly evolving in response to new evidence published every now and then and a plethora of new drugs are being added at a rapid pace.

Various international committees have published guidelines on the treatment of hypertension. The JNC 7 (Joint National Committee on prevention, detection, evaluation and treatment of high blood pressure) recommends the use thiazides type diuretics as the first choice when used alone or in combination with drugs from other classes of anti-hypertensive in uncomplicated essential hypertension.⁷ For >20/10 mmHg above goal BP, combination of two agents is recommended with one of them is usually being thiazides diuretic.⁷ But in recent JNC 8 guidelines it do not consider diuretics as the first choice, rather, considers first-line and later-line treatments to be limited to 4 classes of medications: thiazides-type diuretics, calcium channel blockers (CCBs), angiotensin converting enzyme inhibitors (ACEI) and angiotensin receptor blockers (ARBs) followed by second- and third-line alternatives included higher doses or combinations of ACE inhibitors, ARBs, thiazides-type diuretics and CCBs.⁸ The National Institute for Health and Clinical Excellence (NICE) guideline recommend angiotensin-converting enzyme (ACE) inhibitor or angiotensin-II receptor blocker (ARB) as first choice antihypertensive drug under 55 years, whereas calcium channel blockers (CCBs) are preferred first choice over 55 years.⁹ A combination therapy has recently been recommended as first-line intervention, particularly in patients with severe hypertension,^{7,8} but despite the undisputed importance to treat hypertension aggressively, control rates are still unsatisfactory.

At present, hypertensive patients are treated in medicine as well as the cardiology department. There is no clear guideline which says about the treatment of hypertensive patient should be done in a particular department. The pattern of prescription may vary between the two departments which may affect the compliance of patients and rationality of prescription. Therefore, the present study was planned with the objective to compare the prescribing pattern of antihypertensive drugs in patient attending medicine and cardiology OPD and to analyse the factors affecting the difference in prescribing pattern of antihypertensive in the above mentioned OPD, if any.

METHODS

The present study was an observational, cross sectional study conducted in the department of pharmacology in collaboration with the medicine and the cardiology department of Grant Govt. medical college and Sir JJ group of hospitals for a period of 6 months from January 2014 to June 2014. The study commenced after approval of Institutional Ethics Committee (IEC) and the concerned department. Confidentiality with respect to identity of participating patients and the data obtained was maintained at all levels. The personal right to with-

draw from the study at any moment was ensured. Patients of either sex or age diagnosed with hypertension by the consulting physicians of our tertiary care hospital after recording blood pressure for frequent time and at frequent intervals, taking at least one anti-hypertensive drug and willing to participate by giving written consent were included in the study while pregnant females, mentally compromised, unconscious patients and patients unable to respond to verbal questions were excluded from the study.

Data was collected in a pre-designed proforma which included patient's demographic details and details of the drugs prescribed. Data was compiled in Microsoft office excel 2010 version and a descriptive statistical analysis was carried out. Observations were presented as simple percentages of different variables.

RESULTS

The present study which was conducted for a period of 6 months in the medicine and cardiology OPD showed unequal distribution of the patients in the medicine and cardiology OPD, with 223 patients attended medicine OPD while 434 patients attended cardiology OPD. Demographic characteristic of the patients showed females were more compared to males in medicine OPD whereas male were more in the cardiology OPD. Age wise distribution showed majority of the patients belong to 41-50 years in medicine OPD whereas 51-60 year old patients were in majority in cardiology OPD as shown in Figure 1.

Amongst the patients attending medicine OPD, majority of the patients (61.88%) were stage 1 hypertensive (>140/90 mmHg) according to JNC 7 classification whereas majority (69.38%) of those attending cardiology OPD were stage II hypertensive (>160/100).⁷ Diabetes mellitus was the most common co-morbid condition associated with hypertension in both the OPDs followed by asthma in medicine OPD as shown in Figure 2 and myocardial infarction in cardiac OPD as in Figure 3.

Analysis of the anti-hypertensive drugs prescribed showed, calcium channel blockers (CCB) were the most commonly prescribed drug in medicine OPD followed by angiotensin converting enzyme inhibitors (ACEI) whereas beta blockers (BB) were most commonly prescribed in cardiology OPD followed by ACEI as given in Figure 4.

Anti-hypertensive drugs in the form of fixed drug combinations (FDC) were prescribed more in cardiology OPD (18.93%) in comparison to medicine OPD (7.72%). Single drug prescription was the most preferred in medicine OPD followed by combination of two drugs whereas combination of two drugs was preferred in cardiology OPD followed by combination of three drugs as in Figure 5.




Figure 1: Age and sex distribution of study population.




Figure 2: Co-morbid condition associated with hypertension in medicine OPD.




Figure 3: Co-morbid condition associated with hypertension in cardiology OPD.




Figure 4: Percentage wise prescription of drugs in medicine and cardiology OPD.




Figure 5: Anti-hypertensive drugs per prescription.

Amongst the single drug prescription CCBs were most commonly prescribed in medicine OPD followed by BB whereas BBs were the most commonly prescribed in cardiology OPD followed by CCBs as presented in Table 1. Combination of CCB and BB were more preferred in

medicine OPD followed by combination of CCBs and diuretics (D) whereas combination of ACEIs and BB were preferred in cardiology OPD followed by combination of CCBs and Angiotensin receptor blockers (ARBs) as given in Table 2.

Table 1: Single drug per prescription in medicine and cardiology OPD.

Medicine OPD			Cardiology OPD		
Drugs	No of drugs	Total (%)	Drugs	No of drugs	Total
CCBs			CCBs		
Amlodipine	52	54 (55.10)	Amlodipine	18	19(28.78)
Diltiazem	2		Diltiazem	1	
BB			BB		
Metoprolol	4	19 (19.38)	Metoprolol	19	28 (42.42)
Atenolol	15		Atenolol	9	
α + β Blocker-			α + β Blocker-		
Carvedilol	2	2(2.04)	Carvedilol	1	1 (1.51)
ACEIs			ACEIs		
Enalapril	4	15(15.30)	Enalapril	3	6(9.09)
Ramipril	9		Ramipril	1	
Captopril	2		Captopril	2	
ARBs			ARBs		
Telmisartan	3	6(6.12)	Telmisartan	2	9(13.63)
Losartan	3		Losartan	7	
Diuretics			Diuretics		
Frusemide	2	2(2.04)	Frusemide	2	2(3.03)
α-Blocker-			α-Blocker-		
Prazosin	0	0(0)	Prazosin	1	1(1.51)
TOTAL	98	98(100)	TOTAL	66	66 (100)

When three anti-hypertensive drugs were prescribed, combination of CCB, ARB and D were most common in medicine OPD whereas combination of D, ACEIs and BBs were preferred in cardiology OPD as shown in Table 3. Combination of 2 BB, CCB and D was preferred in medicine OPD when 4 drugs were needed for prescription, whereas combination of CCBs, BB, ARBs

and D was preferred in cardiology OPD as given in Table 4. Combination of 5 drugs was never prescribed in medicine OPD whereas it was prescribed to 2 patients in cardiology OPD, ACEIs, ACEIs, D, CCBs and BB. FDCs of ARBs and D was most commonly prescribed in medicine OPD followed by CCB and BB whereas FDC of ARB and CCB was most commonly prescribed in

cardiology OPD followed by that of BB and ACEIs.

Majority of the drugs were prescribed by their brand names in both the OPDs.

Table 2: Two drug per prescription in medicine and cardiology OPD.

Medicine OPD		Cardiology OPD	
Drugs	Total (%)	Drugs	Total (%)
CCBs ,BB	34 (40.96)	CCBs ,BB	18(9.09)
CCBs ,ACEIs	9(10.84)	CCBs , ACEIs	3(1.51)
CCBs, ARBs	8(0.96)	CCBs, ARBs	24(12.12)
CCBs, D	13(15.66)	CCBs, D	1(0.50)
BB, D	2(2.40)	BB, D	3(1.51)
ACEIs, D	5(6.02)	ACEIs, D	4(2.02)
ACEIs, BB	7(8.43)	ACEIs, BB	105(53.03)
ARBs, D	4(4.81)	ARBs, D	16(8.08)
CCBs, α - Blocker	1(1.20)	ACEIs, α + β Blocker	1(0.50)
		ARBs, BB	21(10.60)
TOTAL	83(100)	D + D	2(1.01)
		TOTAL	198 (100)

Table 3: Three drugs per prescription in medicine and cardiology OPD.

Medicine OPD		Cardiology OPD	
Antihypertensive Drugs	Total (%)	Antihypertensive Drugs	Total (%)
CCBs , ARBs+ D	11(34.37)	CCBs , ARBs+ D	4(3.03)
CCBs , ARBs, BB	8(25)	CCBs + BB, ARBs	27(20.45)
CCBs + ARBs, D	3(9.37)	CCBs + ARBs+ D	2(1.51)
α + β Blocker, D, ACEIs	3(9.37)	α + β Blocker, D, ACEIs	14(10.60)
CCBs, D, α -Blocker	2(6.25)	CCBs, D, α -Blocker	2(1.51)
ARBs + D, BB	2(6.25)	ARBs, D, BB	11 (8.33)
BB, ACEIs, ARBs	1(3.12)	BB, ACEIs, ARBs	6(4.54)
α + β Blocker, D, ARBs	1(3.12)	α + β Blocker, D, ARBs	2(1.51)
α + β Blocker, ARBs + D	1(3.12)	α + β Blocker, ARBs + D	1(0.75)
α + β Blocker, D, K ⁺ channel openers	0	α + β Blocker, D, K ⁺ channel openers	3(2.27)
BB, D, K ⁺ channel openers	0	BB, D, K ⁺ channel openers	2(1.51)
ARBs, D+ D	0	ARBs, D+ D	1(0.75)
CCBs, D + D	0	CCBs, D + D	1(0.75)
α + β Blocker, D, D	0	α + β Blocker, D, D	4(3.03)
CCBs , ACEIs, BB	0	CCBs , ACEIs, BB	17(12.87)
CCBs , ACEIs, D	0	CCBs , ACEIs, D	4(3.03)
CCBs + BB, ACEIs	0	CCBs + BB, ACEIs	5(3.78)
D , ACEIs, BB	0	D , ACEIs, BB	62(46.96)
BB + ACEIs, K ⁺ channel openers	0	BB + ACEIs, K ⁺ channel openers	3(2.27)
BB, BB , ACEIs	0	BB, BB , ACEIs	1(0.75))
CCBs, ACEIs, ACEIs	0	CCBs, ACEIs, ACEIs	1(0.75)
TOTAL	32	TOTAL	129

DISCUSSION

Hypertension is one of the leading causes of mortality and morbidity around the world. The prevalence of this disorder is expected to increase to 1.15 billion by 2025 in developing countries alone including India. Irrespective

of the fact that many medicines are present at present to prevent the mortality and morbidity associated with hypertension the incidence is still on rise. Hypertensive patients can be treated in medicine and cardiology OPD. So, the prescription pattern may vary between the departments.

Table 4: Four drugs per prescription in medicine and cardiology OPD.

Medicine OPD		Cardiology OPD	
Antihypertensive Drugs	Total (%)	Antihypertensive Drugs	Total (%)
CCBs, α + β Blocker, D, α -Blocker	3 (30)	CCBs+ARBs, D, α Blocker	5(12.19)
CCBs, BB, ARBs + D	2(20)	CCBs, BB, ARBs, D	14(34.14)
CCBs, BB, D, BB	4(40)	CCBs + BB, ARBs + BB	2(4.87)
CCBs, ARBs, D + D	1(10)	CCBs, BB, D, D	6(14.63)
		BB, D, ARBs+D	3(7.31)
		CCBs+ARBs, BB+ACEIs	1(2.43)
		CCBs+ARBs, BB, K ⁺ channel openers	2(4.87)
		CCBs+ARBs, D, Central Sympatholytic	1(2.43)
		CCBs+ARBs, Diuretics, CCBs	2(4.87)
		α + β Blocker, D, ACEIs, K ⁺ channel openers	1(2.43)
		ACEIs, BB, ARBs+D	2(4.87)
TOTAL	10	TOTAL	39

So the present study was planned for 6 month to compare the prescription pattern of the two departments. Our study showed unequal distribution of the patients among the two departments with majority of the patients 434 (66.05%) opting for cardiology department and the remaining 223 (33.94%) for medicine OPD. Overall result showed males 333 (50.68%) and females 324 (49.31%) were in almost equal proportion, but, females were in majority in the medicine OPD while males were in majority in cardiology OPD. The results of our study were in accordance with the study conducted by Jhaj and Goel where male and females were in almost equal proportion but majority of the studies conducted in India and abroad, Hong Kong, Germany.¹⁰⁻¹⁴ On the contrary males were in majority in a study conducted by Malhotra et al in North India.¹⁵ Age wise distribution of the patients showed patients belongs to 51-60 years of age presented more with complaints of hypertension, but, majority of the patients belongs to 41-50 years in the medicine OPD and to 51-60 years in cardiology OPD. The above result did prove that as the age increases the risk of hypertension also increase especially after 40 years. Results of our study were comparable to studies done in past where the maximum prevalence of hypertension was in the age group of 51-60years.^{16,17} But a study conducted by Khurshid et al, reported the prevalence to be more among 31-40 years.¹¹ The above results prove that apart from advancing age, a multi factorial inheritance of the disorder.

Diabetes mellitus was the most common co-morbid condition associated with hypertension in both medicine and cardiology OPD. Similar pattern of co-morbidity was seen in the study conducted by Kaur et al and Beg et al.^{18,19} The other co-morbidity present in our study was angina, asthma, arthritis, stroke, myocardial infarction. The presence of co-morbidity decides mainly the pattern of drugs prescribes to patients, as, some drugs

are contraindicated in particular condition and some drugs are favored in other. The average number of antihypertensive drug prescribed in medicine OPD was 1.79 (\pm 0.85) with maximum single drug prescribed to 43.94% patients and the average antihypertensive drug prescribed in cardiology OPD was 2.33 (\pm 0.85) with maximum 2 drugs per prescription prescribed to 45.62% patients. The above results can be compared to study conducted by Khurshid et al and Malhotra et al where the average anti-hypertensive drugs were 1.8 and 1.9 respectively.^{11,15}

A combination of two or more drugs was prescribed to 56.05% patients in medicine OPD whereas it was prescribed to 84.79% patients in cardiology OPD. The above results can be compared to several other studies which demonstrated that combination therapy was needed in atleast 70% of patients to attain optimal blood pressure control.^{20,21} But, many other studies have demonstrated the preference of single drug therapy by the physician.^{12,22,23} Overall assessment of the anti-hypertensive drugs prescribed showed, CCBs (40.26%) were the most common drug prescribed in the medicine OPD followed by BBs (20.52%), whereas BBs (30.28%) were the most preferred drug in cardiology OPD followed by ACEIs (21.13%).

The pattern of anti-hypertensive drugs prescribed differs from study to study, may be because of the different demographic characteristic and the associated co morbid condition. ARBs and ACEIs were the most commonly prescribed drugs in the study conducted by Beg et al, Elliot et al.^{19,25} A common observation in all the above mentioned studies, including ours, was the under-utilization of diuretics, inspite of the JNC 7 report recommendation that in the absence of any specific indications, a diuretic should be selected as the initial therapy for hypertension.⁷

Amongst the single drug prescription, Amlodipine, a calcium channel blocker, 52 (53.06%) was the most commonly prescribed single drug in medicine OPD followed by atenolol, a beta blocker, 15 (15.30%). whereas, metoprolol, a beta blocker, 19 (28.78%), was preferred as a single drug in cardiology OPD followed by amlodipine, 18 (27.27%). This result can be compared to study done in India where CCBs was the most commonly used drug as single drug therapy followed by ACEIs, ARBs, diuretics and β blockers.²⁶ Other Indian studies showed ARBs to be the first choice monotherapy.^{26,27} Nigerian study showed diuretics were prescribed more commonly.²⁸

In the combination drug therapy, dual drug therapy was most preferred. Combination of CCBs and BBs were most commonly prescribed in medicine OPD, whereas combination of ACEI and BBs was preferred in cardiology OPD. Similar to our study, CCBs and BBs was most common combination used in a study conducted in other part of India, whereas, combination of ACEI with BBs was most commonly prescribed in a study conducted by Solanki et al, ARB with diuretic was the most frequently prescribed in a study conducted by Pai et al, Sivakumar et al and Pavani et al.^{11,12,19,29,30,26,27} Etuk (28) reported combination of ACEI and diuretic to be the most common two drug combination used.²⁸ CCB with diuretics 24 (30%) were most commonly prescribed in a study conducted by Konwar et al.²⁵

When combination of three drugs were analysed, combination of CCB, ARB and diuretics was most commonly prescribed in medicine OPD whereas that of ACEI, BB and diuretics were preferred in cardiology OPD. Various other study have reported the use of various other combination agent in the treatment of hypertension such as, combination of ARB, BB and diuretics were preferred in a study conducted by Pavani.²⁷ ARB, CCB and diuretics was most commonly prescribed in a study conducted by Sivakumar et al.²⁶ α -Methyl dopa, ACEI and diuretics was preferred in Nigerian study.²⁸ ACEI, BB and diuretics were preferred in study conducted by Konwar et al.²⁵ ACEI, CCB and diuretics was most commonly prescribed in study by Solanki et al Whereas, ACEI, BB and CCB was most commonly prescribed in study conducted by Khurshid et al.^{29,11} Thus, all the four major drugs are use in combination in various studies and it was also seen that diuretic was involved in almost all combination. Combination of CCBs, α - β Blocker, D and α -Blocker was preferred as four drug combination therapy in medicine OPD, whereas combination of CCBs, BB, ARBs and D was preferred in cardiology OPD.

Fixed drug combination (FDC) of ARB and diuretic was preferred in medicine OPD while that of ARB and CCB was preferred in cardiology OPD followed by that of BB and ACEI. FDCs were prescribed more by cardiologist in comparison to the physicians. Majority of the drugs were prescribed according to their brand name in both medicine

and cardiology OPD. Thus, there is a wide variation in the prescription of anti-hypertensive drugs among various physicians and it was also seen that variation does exist between the medicine and cardiology OPD. This difference might be due to physician's choice with relation to the characteristics of patients, their concurrent illness, as well as the availability of medicines in the hospital. There are very limited data available where, comparison of prescription of a particular disorder in two department of a tertiary care hospital is available. Such studies will definitely help policy makers in generating policies for the disorder.

CONCLUSION

The study concludes that there is a wide variation in the prescription of the anti-hypertensive drugs among medicine and cardiology OPD. CCBs were most commonly used in medicine OPD whereas beta blockers were preferred in cardiology OPD. Monotherapy was preferred in medicine OPD, whereas combination therapy was preferred by cardiologist. Double and triple drug combination therapies were used more in comparison to monotherapy.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. Klabunde RE. Text book of Cardiovascular Physiology Concepts. 2nd edition. Lippincott Williams & Wilkins; 2011.
2. Kulkarni AT. 'Hypertension – A Silent Killer'. *Indian Medical Gazette.*1998;32(3):73-7.
3. Park K. Text book of Preventive and Social Medicine. 16th edition. M. S. Banarsidas Bhanot Publishers; 2000: 277-80.
4. The World Health Report 1998 – WHO, Geneva, 1998
5. WHO health statistics 2012, (cited 2013 Dec 28) available from http://www.who.int/gho/publications/world_health_statistics/2012/en/. Accessed on 01 August 2016.
6. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *The Lancet.* 2005(365):217-23.
7. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL et al. The seventh report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. The JNC 7 report. *JAMA.* 2003;289: 2560-72.
8. JAMES PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. Evidence-Based Guideline for the Management of High Blood Pressure in Adults: Report From the Panel Members

- Appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014;311(5):507-20.
9. National Institute for Health and Care Excellence 2011, Hypertension: clinical management of primary hypertension in adults. CG127. London: National Institute for Health and Care Excellence.
 10. Jhaj R, Goel NK, Gautam CS, Hota D, Sangeeta B, Sood A, et al. Prescribing patterns and cost of antihypertensive drugs in an internal medicine clinic. *Indian Heart J*. 2001;53:323-7.
 11. Khurshid F, Aqil M, Alam MS, Kapur P, Pilla KK: Antihypertensive Medication Prescribing Patterns in a University Teaching Hospital In South Delhi. *Int J Pharm Sci Res*. 2012;3(7):2057-63.
 12. Tiwari H, Kumar A, Kulkarni SK. Prescription monitoring of anti-hypertensive drug utilisation at the Panjab University Health Centre in India. *Singapore Med J*. 2004;45:117-20.
 13. Lee PK, Li RK, Chan JC, Chang S, Lee SC, Tomlinson B, et al. A prescription survey in a hospital hypertension outpatient clinic. *Br J Clin Pharmacol*. 1997;44:577-82.
 14. Pittrow D, Kirch W, Bramlage P, Lehnert H, Hofler M, Unger T, et al. Patterns of antihypertensive drug utilization in primary care. *Eur J Clin Pharmacol*. 2004;60:135-42.
 15. Malhotra S, Karan RS, Pandhi P, Jain S. Pattern of use and pharmaco-economic impact of antihypertensive drugs in a north Indian referral hospital. *Eur J Clin Pharmacol*. 2001;57:535-40.
 16. Shobha JC, Kumar TR, Raju BS, Kamath S, Rao M, Harwal, et al. Evaluation of efficacy and safety of Losartan potassium in the treatment of mild to moderate hypertension as compared to Enalapril maleate. *J Assoc Physicians Ind*. 2000;48:497-500.
 17. Fogari R, Zoppi A. Comparative effects of Lisinopril and Losartan on insulin sensitivity in the treatment of non diabetic hypertensive patients. *Br J Clin Pharmacol*. 1998;46:467-71.
 18. Sharminder Kaur et al. Prescribing pattern of antihypertensive drugs in a tertiary care hospital of northern india. *IJPBS*. 2013;3(2):304-9.
 19. Beg MA, Dutta S, Varma A, Kant R, Bawa S, Anjoom M, et al. Study on drug prescribing pattern in hypertensive patients in a tertiary care teaching hospital at Dehradun, Uttarakhand. *Int J Med Sci Public Health*. 2014;3:922-6.
 20. Chalmers J. The ABCD of anti-hypertensive therapy. *J Hypertens*. 2002;20:615-6.
 21. Yusuff KB, Balogun O. Physicians' prescribing of anti-hypertensive combinations in a tertiary care setting in southwestern Nigeria. *J Pharm Pharm Sci*. 2005;8:235-42.
 22. Mori H, Ukai H, Yamamoto H, Saitou S, Hirao K, Yamauchi M, et al. Current status of antihypertensive prescription and associated blood pressure control in Japan. *Hypertens Res*. 2006;29:143-51.
 23. Arief Md, Harika B, Satyanarayana B, Pasha S, Poloju D, Pokkula S, et al. Evaluation Of Prescribing Pattern Of Antihypertensive Drugs In A Tertiary Care Hospital. *Acta Chim. Pharm. Indica*. 2013;3(2):172-81.
 24. Elliott WJ, Meyer PM. Incident diabetes in clinical trials of antihypertensive drugs: a network meta-analysis. *Lancet*. 2007;369:201-7.
 25. Konwar M, Paul PK, Das S. Prescribing Pattern Of Antihypertensive Drugs In Essential Hypertension In Medicine Out Patients Department In A Tertiary Care Hospital Asian J Pharm Clin Res. 2014;7(2):142-4.
 26. Sivakumar A, Venkateswaramurthy N, Sambathkumar R. Study of drug prescription pattern of anti-hypertensives in a tertiary care hospital. *Der Pharmacia Lettre*. 2014;6(4):86-8.
 27. Pavani V, Cidda M, Krishna TR, Parmar MY, Nalini M. Study Of Prescribing Patterns Of Antihypertensive Drugs. *IJPBS*. 2012;2(2):317-27.
 28. Etuk E, Isezuo Sa, Chika A, Akuche J, Ali M. Prescription Pattern Of Anti-Hypertensive Drugs In A Tertiary Health Institution In Nigeria *Annals Of African Medicine*. 2008;7(3):128-32.
 29. Solanki KC, Mistry RA, Singh AP, Jadav SP, Patel NM, Trivedi HR. Drug utilization study of anti-hypertensive drugs and their adverse effects in patients of a tertiary care hospital. *J Clin Exp Res*. 2013;1:58-67.
 30. Pai PG, Shenoy J, Sanji N. Prescribing patterns of antihypertensive drugs in a South Indian tertiary care hospital. *Drug Invention Today*. 2011;3:38-40.

Cite this article as: Gaikwad B, Bhagat S, Patil K. Comparative assessment of the pattern of anti-hypertensive drugs prescribed in medicine and cardiology outpatient department. *Int J Basic Clin Pharmacol* 2016;5:1958-65.