

Research Article

Anatomical variations and clinical relevance of dorsal sacral foramina in North Indians

Jugesh Khanna*, Renu Chauhan

Department of Anatomy, UCMS and GTB Hospital, Dilshad Garden, Delhi-110095, India

Received: 28 June 2013

Accepted: 4 July 2013

*Correspondence:

Dr. Jugesh Khanna,

E-mail: jugeshkhanna@rediffmail.com

© 2013 Khanna J et al. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Lateral side of each intermediate crest has four dorsal sacral foramina for the passage of posterior division of the sacral nerves. Not only they serve as an important landmark in placement of spinal instrumentation but are also of great importance in transsacral block of sacral nerves for producing analgesia and anesthesia in pelvic surgeries. Therefore a study on the variation in number of dorsal sacral foramina was conducted in 60 adult dry North Indian sacra. Three dorsal sacral foramina were seen on the lateral side of each intermediate crest in four sacra (6.6%) and five in 6 sacra (10%) amongst all the sacra examined. Either a single pair, double pair or totally absent foramina were not observed in any of the sacra seen. This study revealed that there is variation in the number of dorsal sacral foramina. Anaesthetists should be aware of this disparity to be able to perform trans-sacral nerve block successfully. Radiologists must also know about this variation to interpret the radiographs of lumbosacral region. Neurologist and orthopedist should also be cognizant of this fact to diagnose patients presenting with bizarre clinical findings.

Keywords: Dorsal sacral foramina, Intermediate crest, Spinal instrumentation, Sacrum

INTRODUCTION

Sacrum is the most variable portion of spine as a number of anatomical variations occur in this region quite frequently.¹ Lateral to the intermediate crest on the dorsal aspect of sacrum are four dorsal sacral foramina (Figure 1), which transmit the posterior division of sacral nerves.² Occasionally, the first sacral vertebra is not fused to other four segments of sacrum (lumbarization of first sacral vertebra) resulting in three sacral foramina (Figure 2, 3) or the fifth lumbar vertebra fuses with the first sacral vertebra (sacralization of fifth lumbar vertebra) leading to formation of five pairs of dorsal sacral foramina (Figure 4). As they serve as an important landmark in spinal instrumentation and can also be used for blocking sacral nerves trans-sacrally in order to produce analgesia and anesthesia in the surgeries involving pelvic region therefore, it becomes essential to know the frequency of variation in the number of dorsal sacral foramina. Deficient literature is available on dorsal sacral foramina

in North Indians, hence this study was conducted. Embryological origin and clinical significance of its variable number have been discussed in the paper.

METHODS

Eighty adult dry human sacra were obtained from the Anatomy Department of U.C.M.S and G.T.B. Hospital, Dilshad Garden, Delhi. The sacra were grossly examined. Twenty sacra were damaged therefore were excluded from the study. Ultimately, sixty sacra were seen. The number of dorsal sacral foramina in each of the remaining sacrum was counted. The result obtained was analyzed and discussed.

RESULTS

Sixty adult dry human sacra were examined. Three pairs of dorsal sacral foramina were present lateral to intermediate crest in 4 sacra (6.6%) (Figure 3) and five

pairs of dorsal sacral foramina were seen in 6 sacra (10%) (Figure 4). Rest of the sacra showed the normal four pair of foramina. No sacrum showed either a single or double pair or a total absence of dorsal sacral foramina.



Figure 1: Showing dorsal aspect of adult dry sacrum. Normally, four pairs of dorsal sacral foramina are seen lateral to intermediate crest.



Figure 2: Showing ventral view of the adult dry sacrum. Three pairs of ventral sacral foramina are seen.



Figure 3: Showing dorsal aspect of the adult dry sacrum having three pairs of dorsal sacral foramina lateral to intermediate crest of sacrum.



Figure 4- Dorsal aspect of the adult dry sacrum showing five pairs of dorsal sacral foramina lateral to intermediate crest of sacrum.

DISCUSSION

Normally, sacrum is formed by fusion of five sacral vertebrae which results in formation of four pairs of sacral foramina (Figure 1).³ The upper four sacral nerve roots exit the sacrum through paired dorsal and ventral sacral foramina. Fifth sacral and coccygeal roots and the filum terminale exit the sacrum caudally through the sacral hiatus.⁴ A couple of abnormalities of the sacrum influence the free functioning of the spine.

Lumbarization of the first sacral vertebra is said to occur when the uppermost segment of the sacrum is not fused to other four segments of sacrum. Rather the first sacral vertebra is mobile and participates along with the neighbouring lumbar vertebrae in spinal activities.⁵ The resultant sacrum will have four vertebrae instead of five as in the normal sacrum and will have three sacral foramina along the pelvic and dorsal aspect of sacrum. Although few authors have mentioned presence of three dorsal sacral foramina in their study^{6,7} but no literature was available on the variation in number of the dorsal sacral foramina in North Indian population. In the present study, three dorsal sacral foramina were observed in four out of a total number of sixty sacra examined (Figure 3). Radiographic studies conducted by Paul (1938) on one thousand patients who complained of pain in the lower part of the back and legs suggested that approximately 30 per cent showed an anomaly of the lumbo-sacral spine.⁸ It is possible that the patients who have three dorsal sacral foramina could present with pain in lower part of the back and legs as suggested by Paul. If such an anomaly exists then there is all possibility that the fourth sacral nerve roots also passes through the sacral hiatus along with the fifth sacral, coccygeal roots and the filum terminale exiting through it. In that case, any sacrococcygeal tumour may compress upon fourth sacral roots also, leading to neurological involvement of the bladder and rectum. Physicians, Surgeons and Uro-surgeons must be aware of the presence of such variation while dealing with unusual clinical findings. However,

Pamela suggested that the patients are frequently told that a spinal anomaly observed in their radiograph is the cause of their back problems. But the anomalies in the lumbar and sacral spine are simply variants of the normal bony architecture and are typically of no consequence. It would be very rare for an abnormality such as a sixth lumbar vertebra or extra sacral segment to cause back problems.⁹



Figure 5: Ventral aspect of the adult dry sacrum showing sacralization of fifth lumbar vertebra on the left side resulting in five pairs of ventral sacral foramina on that side only. Incomplete process of sacralization of fifth lumbar vertebra seen on the right side.

Five pairs of dorsal sacral foramina were also seen in six sacra in the present study. When fifth lumbar vertebra fuses with the first sacral vertebra, it is known as sacralization of lumbar vertebra and when first coccygeal vertebra fuses with the apex of sacrum, it is known as sacralization of coccygeal vertebra. The processes described above leads to formation of an additional fifth pair of sacral foramina.¹⁰ Either unilateral (Figure 5) or bilateral (Figure 6) sacralization of fifth lumbar vertebra was seen in the specimen studied. Unilateral sacralization was observed in one specimen only which showed five dorsal sacral foramina on the left side (Figure 7). Rajani reported a case with five pairs of dorsal sacral foramina while taking osteology demonstration of MBBS students of Medical University, Lucknow. The fifth pair developed due to sacralization of coccyx. This pair of foramina gives passage to fifth pair of sacral and coccygeal nerves. This variant is of importance to surgeons and obstetricians dealing with these nerves. Rajani (2011) is of the opinion that due to fusion of sacrum and coccyx as was seen in her study, antero-posterior diameter of pelvic outlet does not increase during second stage of labour which may lead to prolonged labour and perineal tears.¹¹ In the present study, all the sacra seen with five pairs of sacral foramina were male specimen (Figure 4, 5, 6). No sacrum belonging to a female possessed five pairs of dorsal

sacral foramina. The variation seen in this study is in contrast to that reported by Rajani (2011). No sacralization of the coccygeal vertebra was observed in any of the sacra seen in the present study as was seen by Platzer (2008) and Rajani (2011) in their study.



Figure 6: Ventral aspect of the adult dry sacrum showing bilateral complete sacralization of fifth lumbar vertebra resulting in five pairs of ventral sacral foramina.



Figure 7: Dorsal aspect of the adult dry sacrum showing 5 complete dorsal sacral foramina on the left side only.

Simultaneous fusion of fifth lumbar vertebra with the cranial end of the first sacral vertebra and fusion of first coccygeal vertebra with the caudal end of fifth sacral vertebra may lead to formation of six pairs of sacral foramina as suggested by Bergman, Afifi and Miyauchi.¹² In the present study no sacrum was seen to possess 6 pairs of foramina.

Sadler is of the opinion that HOX gene is responsible for patterning of the shapes of vertebra.¹³ Sacralization of the

fifth lumbar vertebra or lumbarization of the first sacral vertebra as seen in the present study could have been due to mutation of HOX gene as was suggested by Sadler. Other less common etiologies may be arthritic changes, traumatic injury and intentional spinal fusion surgery.

The observations made by renowned anatomists and surgeons several years ago, still holds true. But one needs to learn a lot to keep pace with recent advances in surgical methods and instrumentation in the lumbosacral region. Obstetricians, radiologists, anaesthetists, neurologists and orthopedic surgeons must know about the existence of this variation to be able to correctly investigate, diagnose and treat the patients presenting with unusual signs and symptoms.

Funding: None

Competing interests: None declared

Ethical approval: Not required

REFERENCES

1. Esses SE, Botsford DJ. Surgical Anatomy and Operative Approaches to the Sacrum. In: Frymoyer JW, Ducker TB, Hadler NM, et al eds. The Adult Spine: Principles and Practice. 2nd ed. Philadelphia, Lippincott-Raven; 1997: Vol 2, 2329-2341.
2. Joseph SC, John KS. Anatomy of the Sacrum. Neurosurg Focus 2003;15(2).
3. Standring S. Anatomical Basis of Clinical Practice. In: Gray's Anatomy ed. 39th ed. London, Churchill Livingstone; 2005: 749–750.
4. Spinal Anatomy. Available at <http://www.scientificspine.com> Accessed 24 June 2013.
5. Available at <http://www.simplebackpain.com/lumbarisation-and-sacralisation.html#ixzz2Wk4IE8Lr>. Accessed 24 June 2013.
6. Panjakash S, Londhe S, Kori R. Lumbarization of first sacral vertebra: A case Report. International Journal of Basic and Applied Medical Sciences 2012; 2 (2): 154-157.
7. Dharati M Kubavat, Shaileshkumar K Nagar, Chintan Lakhani, Srushti S Ruparelia, Shilpa Patel, Padma Varlekar. A Study of Sacrum with Three Pairs of Sacral Foramina in Western India. International journal of medical science and public health 2012; 1(2): 127-131.
8. Paul CW. Anomalies of the Lumbo-Sacral Spine. Radiology 1938; 30: 361-369.
9. Spinal Abnormalities Rarely Cause Back Problems. Available at <http://www.spine-health.com> Accessed 23 June 2013.
10. Platzer W. Locomotor System. In: Color Atlas of Human Anatomy. 6th ed. Vol 1. Stuttgart, Thieme; 2008:11.
11. Rajani S. Sacrum with five pairs of sacral foramina. International Journal of Anatomical Variations 2011; 4: 139-140.
12. Bergman RA, Afifi AK, Miyauchi R. Illustrated Encyclopedia of Human Anatomic Variation. <http://www.anatomyatlases.org/AnatomicVariants/SkeletalSystem/Text/SacrumCoccyx.html>. Accessed 11 June 2013.
13. Sadler TW. In: Langman's Medical Embryology. 8th ed. Chapel Hill, North Carolina, Lippincott Williams and Wilkins; 2000:183.

DOI: 10.5455/2320-6012.ijrms20130820
Cite this article as: Khanna J, Chauhan R.
Anatomical variations and clinical relevance of dorsal sacral foramina in North Indians. Int J Res Med Sci 2013;1:263-6.