



This is a repository copy of *On Siegel eigenvarieties at Saito–Kurokawa points*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/193293/>

Version: Published Version

Article:

Berger, T. orcid.org/0000-0002-5207-6221 and Bertina, A. (2022) On Siegel eigenvarieties at Saito–Kurokawa points. *Annales de l'Institut Fourier*, 72 (3). pp. 901-961. ISSN 0373-0956

<https://doi.org/10.5802/aif.3482>

Reuse


This article is distributed under the terms of the Creative Commons Attribution-NoDerivs (CC BY-ND) licence. This licence allows for redistribution, commercial and non-commercial, as long as it is passed along unchanged and in whole, with credit to the original authors. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



ANNALES DE L'INSTITUT FOURIER

Tobias BERGER & Adel BETINA

On Siegel eigenvarieties at Saito–Kurokawa points

Tome 72, n° 3 (2022), p. 901-961.

<https://doi.org/10.5802/aif.3482>

Article mis à disposition par ses auteurs selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

e-ISSN : 1777-5310

ON SIEGEL EIGENVARIETIES AT SAITO–KUROKAWA POINTS

by Tobias BERGER & Adel BETINA (*)

ABSTRACT. — We study the geometry of the p -adic Siegel eigenvariety \mathcal{E} of paramodular tame level at certain Saito–Kurokawa points having a critical slope. For $k \geq 2$ let f be a cuspidal new eigenform of $S_{2k-2}(\Gamma_0(N))$ ordinary at a prime $p \nmid N$ with sign $\epsilon_f = -1$ and write α for the p -adic unit root of the Hecke polynomial of f at p . Let π_α be the semi-ordinary p -stabilization of the Saito–Kurokawa lift of the cusp form f to $\mathrm{GSp}(4)$ of weight (k, k) and paramodular tame level. Under the assumption that the dimension of the Selmer group $H_{f, \mathrm{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ attached to f is at most one and some mild assumptions on the automorphic representation attached to f , we show that \mathcal{E} is smooth at the point corresponding to π_α , and that the irreducible component of \mathcal{E} specializing to π_α is not globally endoscopic. Finally we give an application to the Bloch–Kato conjecture, by proving under some mild assumptions that the smoothness failure of \mathcal{E}_Δ at π_α yields that $\dim H_{f, \mathrm{unr}}^1(\mathbb{Q}, \rho_f(k-1)) \geq 2$.

RÉSUMÉ. — On étudie la géométrie de la variété de Hecke-Siegel \mathcal{E}_Δ de niveau paramodulaire Δ associée à un entier N en certains points de Saito-Kurokawa dont la pente est critique. Pour $k \geq 2$, soit f une forme cuspidale de $S_{2k-2}(\Gamma_0(N))$, ordinaire en un nombre premier $p \nmid N$, de signe $\epsilon_f = -1$, et notons α l'unité p -adique racine du polynôme de Hecke de f en p . Soit π_α la p -stabilisation semi-ordinaire du relèvement de Saito–Kurokawa de f à GSp_4 de poids (k, k) et de niveau modéré Δ . Sous l'hypothèse que la dimension du groupe de Selmer $H_{f, \mathrm{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ associé à f est au plus égale à 1 et certaines hypothèses supplémentaires sur la représentation p -adique ρ_f associée à f , on démontre que l'espace analytique rigide \mathcal{E}_Δ est lisse au point correspondant à π_α . Cela signifie qu'il existe une unique composante irréductible de \mathcal{E}_Δ passant par π_α et on démontre que cette composante n'est pas globalement endoscopique. Pour finir, on donne une application à la conjecture de Bloch-Kato, en prouvant, sous certaines hypothèses raisonnables, qu'une singularité de \mathcal{E}_Δ en π_α entraîne que $\dim H_{f, \mathrm{unr}}^1(\mathbb{Q}, \rho_f(k-1)) \geq 2$.

Keywords: Eigenvarieties for GSp_4 , Saito–Kurokawa lift, Selmer groups and pseudo-deformation theory.

2020 *Mathematics Subject Classification:* 11F33, 11F46, 11F80, 14G22.

(*) Both authors acknowledge support from the EPSRC Grant EP/R006563/1.

1. Introduction

Let p be a prime number. Eigenvarieties are p -adic rigid analytic spaces interpolating the Hecke eigenvalues of automorphic representations of a particular reductive group G of finite slope eigenvalues for Hecke operators at p , fixed tame level away from p and varying weights. Following the seminal works of Hida [21] and Coleman–Mazur [15] their geometry has been studied by many people, e.g. Bellaïche–Chenevier [6], Majumdar [28], Bellaïche–Dimitrov [8], Betina–Dimitrov–Pozzi and Betina–Dimitrov [11, 12] for $G = \mathrm{GL}_2(\mathbb{Q})$, and by Bellaïche–Chenevier [3, 7] for unitary groups.

Andreatta, Iovita and Pilloni constructed in [1] an eigenvariety parametrizing locally analytic overconvergent cuspidal Siegel eigenforms of genus two, principal level N and finite slope, and they proved that the Siegel eigenvariety of tame level 1 is étale over the weight space at certain classical non-critical points of regular cohomological weights with Iwahoric level at p . The proof uses the classicality criteria for overconvergent Siegel cusp forms of Hida [22, Proposition 3.6], Tilouine and Urban [45, Theorem 3.2], Pilloni [36, Theorem 2] and the multiplicity one theorem of Arthur’s classification for GSp_4 [2].

We investigate in this work the geometry of the Siegel eigenvariety \mathcal{E}_Δ of paramodular level N at the points corresponding to Saito–Kurokawa lifts of ordinary cusp forms for $\mathrm{GL}_2(\mathbb{Q})$ (which have a critical slope), including the case of the non-cohomological weight $(2, 2)$.

In order to state our results, we recall some facts and fix some notations: Let $N \geq 1$ be an integer prime to p . For a prime ℓ the paramodular subgroup of $\mathrm{GSp}_4(\mathbb{Q}_\ell)$ is defined as $\Delta_\ell = \gamma \mathrm{M}_4(\mathbb{Z}_\ell) \gamma^{-1} \cap \mathrm{GSp}_4(\mathbb{Q}_\ell)$ for $\gamma = \mathrm{diag}[1, 1, \ell, 1]$. We write $\Delta := \prod_{\ell|N} \Delta_\ell \cap \mathrm{GSp}_4(\mathbb{Q})$ for the paramodular congruence subgroup of level N . If $N = 1$ we put $\Delta = \mathrm{GSp}_4(\mathbb{Z})$.

Let $f \in \mathrm{S}_{2k-2}(\Gamma_0(N), K_f)$ be a weight $2k - 2$ cuspidal N -new eigenform for $\mathrm{GL}_2(\mathbb{Q})$ with coefficients in a number field K_f (and trivial central character). Assume that f has an ordinary p -stabilization and denote it by f_α , where $U_p(f_\alpha) = \alpha \cdot f_\alpha$.

The L-function $L(f, s)$ attached to f satisfies the following functional equation:

$$L(f, s) = \epsilon_f L(f, 2k - 2 - s).$$

We have that $\epsilon_f = (-1)^{\mathrm{ord}_s = k-1} L(f, s)$. Assume until the end of this paper that $\epsilon_f = -1^{(1)}$, which means that there exists a lift $\mathrm{SK}(f)$ to a holomorphic weight (k, k) cuspform of level Δ called the Saito–Kurokawa lift of f .

(1) When $N = 1$, one has $\epsilon_f = (-1)^{k-1}$.

It satisfies

$$L^N(\text{SK}(f), \text{spin}, s) = \zeta^N(s - k + 1)\zeta^N(s - k + 2)L^N(s, f).$$

When $N = 1$ this lift was constructed by Maass, Andrianov and Zagier; Gritsenko generalized it to any level N . A representation theoretic approach building on results of Piatetski–Shapiro and Waldspurger is discussed in [38] (see also [40, Theorem 5.3]).

In order to p -adically deform $\text{SK}(f)$, one must first choose a semi-ordinary⁽²⁾ p -stabilization of $\text{SK}(f)$, that is an eigenform of tame level the paramodular group Δ and sharing the same eigenvalues as $\text{SK}(f)$ away from p and of finite slope. Denote by π_α the p -stabilization of $\text{SK}(f)$ such that $U_0(\pi_\alpha) = \alpha \cdot \pi_\alpha$, and $U_1(\pi_\alpha) = p \cdot \alpha \cdot \pi_\alpha$ where U_0, U_1 are the Hecke operators attached to $\text{diag}[1, 1, p, p]$ (U_0 is often denoted by U_p), $\text{diag}[1, p, p^2, p]$, and U_1 has been renormalized to have a good p -adic interpolation (see for example [41, Theorem 2.4.14]).

Let \mathcal{E}_Δ be Siegel eigenvariety of tame paramodular level Δ (see appendix Section B.4). It is reduced and equidimensional of dimension 2, and endowed with a morphism

$$\kappa : \mathcal{E}_\Delta \rightarrow \mathcal{W}$$

called the weight map (which is locally finite and torsion-free), where the weight space \mathcal{W} is the rigid analytic space over \mathbb{Q}_p such that $\mathcal{W}(\mathbb{C}_p) = \text{Hom}_{\text{cont}}((\mathbb{Z}_p^\times)^2, \mathbb{C}_p^\times)$.

The cuspidal eigenform π_α defines a point of \mathcal{E}_Δ which we denote again (by a slight abuse of notation) by π_α . Write L for the residue field of the point $\pi_\alpha \in \mathcal{E}_\Delta$, a finite extension of \mathbb{Q}_p . Note that the slopes of U_0 and U_1 are locally constant on \mathcal{E}_Δ , and equal to 0 for U_0 and 1 for U_1 locally at π_α . This means that the cuspform π_α has a critical slope since it does not satisfy the small slope condition of [1, Theorem 7.3.1].

One can show that there exists a pseudo-character $\text{Ps} = \text{Ps}_{\mathcal{E}_\Delta} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta)$ of dimension 4 such that the specialization $\text{Ps}(y)$ of Ps at a classical point $y \in \mathcal{E}_\Delta(\overline{\mathbb{Q}}_p)$ is the trace of the semi-simple p -adic Galois representation $\rho_y : G_{\mathbb{Q}} \rightarrow \text{GL}_4(\overline{\mathbb{Q}}_p)$ of dimension 4 attached to a cuspidal Siegel eigenform g_y corresponding to y (i.e. $L(g_y, \text{spin}, s) = L(\rho_y, s)$). For $y = \pi_\alpha$ we have

$$\text{Ps}(\pi_\alpha) = \epsilon_p^{1-k} + \epsilon_p^{2-k} + \text{Tr } \rho_f,$$

⁽²⁾Semi-ordinary means that the eigenvalue for the Hecke operator U_0 is a p -adic unit. Following Tilouine–Urban this is also called Siegel ordinary.

where $\rho_f : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(L)$ is the p -adic Galois representation attached to f (i.e. $L(f, s) = L(\rho_f, s)$) and ϵ_p is the p -adic cyclotomic character (L contains the Hecke field K_f of f).

Let \mathcal{T} be the local ring of \mathcal{E}_{Δ} at π_{α} for the rigid topology, \mathfrak{m} the maximal ideal of \mathcal{T} and Λ the local ring of \mathcal{W} for the rigid topology at the weight $\kappa(\pi_{\alpha})$ of π_{α} (they are both Henselian rings). Note that \mathcal{T} is an equidimensional ring of dimension 2.

DEFINITION 1.1. — *We say that an irreducible affinoid $\mathcal{Z} \subset \mathcal{E}_{\Delta}$ of dimension 2 is stable if and only if the reducibility locus of the pseudo-character $\mathrm{Ps}_{\mathcal{Z}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{Z})$ given by the composition of $\mathrm{Ps}_{\mathcal{E}_{\Delta}}$ with the natural morphism $\mathcal{O}(\mathcal{E}_{\Delta}) \rightarrow \mathcal{O}(\mathcal{Z})$ is strictly contained in \mathcal{Z} (i.e. of dimension less or equal to 1).*

Let $\pi_f = \bigotimes_{\ell} \pi_{f,\ell}$ be the automorphic representation attached to f . We list below the main assumptions that we need in this work:

(BK) The Bloch–Kato Selmer group

$$\begin{aligned} & H_{f,\mathrm{unr}}^1(\mathbb{Q}, \mathrm{ad}^0 \rho_f) \\ &= \ker \left(H^1(\mathbb{Q}, \mathrm{ad}^0 \rho_f) \rightarrow H^1(\mathbb{Q}_p, \mathrm{ad}^0 \rho_f \otimes B_{\mathrm{crys}}) \bigoplus_{\ell \neq p} H^1(I_{\ell}, \mathrm{ad}^0 \rho_f) \right) \end{aligned}$$

is trivial (B_{crys} is Fontaine’s crystalline period ring).

(Reg) If $k = 2$ assume $\alpha \neq 1$ (α is the p -adic unit root of the Hecke polynomial at p of f).

(St) For any prime $\ell \mid N$, $\rho_f|_{I_{\ell}}$ is special and the sign $\epsilon_{\ell}(f)$ in the functional equation of f at ℓ is always equal to 1 and the local sign $\epsilon_{\infty}(f)$ at the archimedean place ∞ is equal to -1 (which means that k is even). In that case, for any prime $\ell \mid N$, $\pi_{f,\ell} \simeq \mathrm{St} \otimes \xi$, where ξ is the unramified character with $\xi(\ell) = -1$.

We highlight that the sign $\epsilon_f = \prod_{\ell \mid N \cup \{\infty\}} \epsilon_{\ell}(f)$ and $\epsilon_{\infty}(f) = (-1)^{k-1}$. Moreover, if f satisfies **(St)**, then the integer N is necessarily squarefree.

Andreatta–Iovita–Pilloni pose the following question in [1, Section 8]:

OPEN PROBLEM. — *Let $x(g)$ be a classical point of the Siegel eigenvariety \mathcal{E}_N of tame level $\Gamma(N)$, the principal congruence subgroup of level N . Is the map $\kappa : \mathcal{E}_N \rightarrow \mathcal{W}$ unramified at $x(g)$?*

Let \mathfrak{m}_{Λ} be the maximal ideal of Λ , the completed local ring of \mathcal{W} at $\kappa(\pi_{\alpha})$, $\mathcal{T}' = \mathcal{T}/\mathfrak{m}_{\Lambda}\mathcal{T}$ be the local ring of the fiber $\kappa^{-1}(\kappa(\pi_{\alpha})) \subset \mathcal{E}_{\Delta}$ at π_{α} (since κ is locally finite, \mathcal{T}' is an Artinian algebra), and let $\mathfrak{t}_{\pi_{\alpha}}$ (resp. $\mathfrak{t}_{\pi_{\alpha}}^0$)

be the Zariski tangent space of \mathcal{T} (resp. \mathcal{T}' , i.e the relative tangent space of $\kappa^\# : \Lambda \rightarrow \mathcal{T}$).

Let $\omega_p : G_{\mathbb{Q}} \rightarrow \mathbb{Z}_p^\times$ be the Teichmüller character and $L_p(f_\alpha, \omega_p^{-1}, \cdot) \in \Lambda := \overline{\mathbb{Z}}_p[[T]]$ be the Manin–Vishik p -adic L -function attached to $f_\alpha \otimes \omega_p^{-1}$ (see e.g. [23, 16.2]), and let

$$\begin{aligned} & H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) \\ &= \ker \left(H^1(\mathbb{Q}, \rho_f(k-1)) \rightarrow H^1(\mathbb{Q}_p, \rho_f(k-1) \otimes B_{\text{crys}}) \bigoplus_{\ell \nmid p} H^1(I_\ell, \rho_f(k-1)) \right) \end{aligned}$$

be the Bloch–Kato Selmer group attached to f .

Recall that \mathcal{E}_Δ is equidimensional of dimension 2. Our main result is the following theorem describing the local geometry at π_α of the rigid analytic space \mathcal{E}_Δ :

THEOREM 1.2 (see Sections 2.5 and 7.2). — *Put*

$$s = \dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)).$$

- (1) *Assume that $k \geq 2$ and **(Reg)**. Then all the irreducible affinoids of \mathcal{E}_Δ of dimension 2 specializing to π_α are stable.*
- (2) *Assume that $k \geq 2$, **(Reg)**, **(BK)** and **(St)**, and assume also when $k = 2$ that $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$, then*

$$2 \leq \dim \mathfrak{t}_{\pi_\alpha} \leq 1 + s^2 \quad \text{and} \quad \dim \mathfrak{t}_{\pi_\alpha}^0 \leq s^2.$$

Moreover, if $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$, then \mathcal{E}_Δ is smooth at π_α , and the reducibility locus of the pseudo-character $\text{Ps}_{\mathcal{T}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta) \rightarrow \mathcal{T}$ is the closed irreducible smooth subscheme of $\text{Spec } \mathcal{T}$ of dimension 1 associated to the Saito–Kurokawa lift of the ordinary Hida family \mathcal{F} passing through f_α , and it is even a principal Weil divisor of $\text{Spec } \mathcal{T}$.

A key step in the proof is the determination of the schematic reducibility locus of the pseudo-character $\text{Ps}_{\mathcal{T}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta) \rightarrow \mathcal{T}$ carried by \mathcal{E}_Δ at π_α , and our approach uses pseudo-representations of p -adic families of cuspidal Siegel eigenforms and p -adic Hodge theory. We provide a more detailed sketch of the proof in Section 1.1.

A direct consequence of (2) of the above theorem is that under these assumptions there exists a unique irreducible component of \mathcal{E}_Δ specializing to π_α when the Selmer group $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ is 1-dimensional.

The smoothness of the eigencurve at critical points is a crucial ingredient for the construction of a family of p -adic L -functions on an open neighborhood of these points, see e.g. [5]. Our result on the smoothness of

\mathcal{E}_Δ opens up the possibility of constructing a family of p -adic L-functions in a neighbourhood of π_α , a challenging question in Iwasawa theory.

Remark 1.3. — We require assumption **(BK)** for applying an $R = T$ theorem in characteristic zero (see Proposition 7.6). Assumption **(BK)** is conjectured to always hold by the Bloch–Kato conjectures as $L(\text{ad}^0(\rho_f), 1) \neq 0$. It has been proven under various assumptions, in particular, by Weston [49, Theorem 5.5] if f is not CM, and special or supercuspidal at all $\ell \mid N$, and by Kisin [25] under mild assumptions on the residual representation $\bar{\rho}_f$, e.g. if $\bar{\rho}_f|_{G_L}$ is absolutely irreducible for any quadratic extension L/\mathbb{Q} with $L \subset \mathbb{Q}(\zeta_{p^3})$ or if $\bar{\rho}_f^{\text{ss}}$ over an algebraic closure is given by the sum of two characters that are distinct on $G_{\mathbb{Q}(\zeta_{p^\infty})}$.

Using results about Λ -adic Selmer groups we exhibit many examples where the Selmer group $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ is 1-dimensional (see Appendix C). We also have an example of an elliptic curve satisfying all the assumptions of (2) of the above theorem (see Section C).

COROLLARY 1.4. — *Assume that $k \geq 2$, **(Reg)**, **(BK)** and **(St)**, and assume also when $k = 2$ that $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$. If the rigid analytic space \mathcal{E}_Δ is singular at π_α then $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) \geq 2$.*

Hence we have a geometric criterion to detect if $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) \geq 2$. Thus, the question of finding a lower bound of the dimension of the Selmer group $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ can be reduced to certain computations of spaces of semi-ordinary p -adic modular cuspforms for GSp_4 .

It turns out that the geometry of \mathcal{E}_N at π_α depends on the tame level. When we change the tame level to the principal Siegel congruence subgroup $\Gamma(N)$ it is in general non-smooth. In particular, the answer to the question in [1] is negative if N is not prime.

COROLLARY 1.5 (see Corollary 8.5). — *Assume that $\ell_1, \ell_2 \mid N$, where $\{\ell_i\}_{\{1,2\}}$ are prime numbers and assume that f is Steinberg at both these primes. Then the eigenvariety \mathcal{E}_N of level $\Gamma(N)$ is singular at π_α and has at least two irreducible endoscopic components specializing to π_α .*

1.1. Sketch of the proof of Theorem 1.2

Using [41, Theorem 3.2.9] we show that any irreducible affinoid $\mathcal{Z} \subset \mathcal{E}_\Delta$ of dimension 2 specializing to π_α such that its pseudo-character $\text{Ps}_{\mathcal{Z}}$ is reducible contains necessarily a classical point corresponding to an endoscopic cuspidal eigenform of trivial central character, Iwahoric level and old

at p . In fact, it follows from Arthur’s classification and results of Roberts and Schmidt that no such Yoshida lift exists for level Δ . This establishes that all irreducible components of \mathcal{E}_Δ containing π_α are stable.

Hence by localizing the pseudo-character $\text{Ps}_{\mathcal{E}_\Delta} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta)$ of dimension 4 at the local Henselian ring \mathcal{T} , we get a pseudo-character $\text{Ps}_{\mathcal{T}} : G_{\mathbb{Q}} \rightarrow \mathcal{T}$ deforming $\text{Ps}(\pi_\alpha)$ which is generically irreducible on each irreducible component containing π_α . Following the results of [7], we obtain a GMA matrix $S = \mathcal{T}[G_{\mathbb{Q}}]/\ker(\text{Ps}_{\mathcal{T}})$ with orthogonal idempotents lifting the natural idempotents of the semi-simple representation $\varrho = \epsilon_p^{2-k} \oplus \rho_f \oplus \epsilon_p^{1-k}$.

The total reducibility ideal \mathcal{I}^{tot} of $\text{Ps}_{\mathcal{T}}$ is defined to be the smallest ideal I of \mathcal{T} such that

$$\text{Ps}_{\mathcal{T}} \bmod I = T_1 + T_2 + T_3$$

for pseudocharacters T_i with $T_i \bmod \mathfrak{m} = \text{Tr}(\rho_i)$ for $\rho_1 = \epsilon_p^{2-k}$, $\rho_2 = \rho_f$, $\rho_3 = \epsilon_p^{1-k}$. By results of [7] it is controlled by the entries of the GMA S (see Proposition 3.4). These in turn give rise to S -extensions of ρ_i by ρ_j for $i \neq j$. We prove in Theorem 6.8 when $s := \dim H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$ that \mathcal{I}^{tot} is principal (or more generally we bound the number of its generators by s^2) by proving that these extension satisfy the required local properties to lie in the corresponding Selmer groups $H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-2)) = \{0\}$ (a deep result of Kato [23]), $H_{f,\text{unr}}^1(\mathbb{Q}, \epsilon_p) \stackrel{\text{Kummer}}{\simeq} \mathbb{Z}^\times \otimes L = \{0\}$ and $H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$.

This local analysis forms the technical heart of the paper. At p we use that any representation ρ_z attached to a classical point z of a sufficiently small open admissible affinoid $\mathcal{Z} \subset \mathcal{E}_\Delta$ containing π_α is semi-ordinary (i.e. $\dim \rho_z^{I_p} \geq 1$). Using this we prove in Sections 3 and 5 that any S -extension W (resp. W') occurring in the cohomology group $H^1(\mathbb{Q}, \rho_f(k-1))$ (resp. $H^1(\mathbb{Q}, \rho_f(k-2))$) is in fact ordinary at p , in the sense that $W^{I_p} \neq 0$, $(W')^{I_p} \neq 0$ and Frob_p acts on them by α . Therefore, W (resp. W' when $k \geq 3$) is ordinary in the sense of Fontaine–Perrin-Riou (so de Rham, see Theorem 2.2), and hence crystalline since $H_g^1(\mathbb{Q}_p, \rho_f(k-i)) = H_f^1(\mathbb{Q}_p, \rho_f(k-i))$ for $i \in \{1, 2\}$.

To prove the crystallinity of the S -extensions in $\text{Ext}_{G_{\mathbb{Q}}}^1(\epsilon_p^{1-k}, \epsilon_p^{2-k})$ we apply in Section 4 the results of [7, Section 4] on the analytic continuation of crystalline periods for the smallest Hodge–Tate weight in families of p -adic $G_{\mathbb{Q}_p}$ -representations occurring in a torsion free coherent module of generic 3. To this end we establish in Section B that classical points which are old at p are very Zariski dense in \mathcal{E}_Δ . To be able to study the period we are interested in we need to consider the quotient by the line fixed by inertia due to semi-ordinarity. At a classical point $z \in \mathcal{Z}$ of cohomological

weight (l_1, l_2) the smallest Hodge–Tate weight of the 3-dimensional $G_{\mathbb{Q}_p}$ -representation $\rho_z/\rho_z^{I_p}$ is $l_2 - 2$ and $\dim \mathcal{D}_{\text{crys}}(\rho_z/\rho_z^{I_p})_{U_1/U_0(z)} p^{l_2-2} = 1$.

This allows us to prove that the S -extensions occurring in $\text{Ext}_{G_{\mathbb{Q}}}^1(\epsilon_p^{1-k}, \epsilon_p^{2-k})$ have a crystalline period equal to

$$\lim_{z_n \in \mathcal{E}_{\Delta}, z_n \rightarrow \pi_{\alpha}} U_1/U_0(z) p^{l_2(z)-2} = U_1/U_0(\pi_{\alpha}) p^{k-2} = p^{k-1}.$$

This means that for any S -extension $V \in \text{Ext}_{L[G_{\mathbb{Q}}^N p]}^1(\epsilon_p^{1-k}, \epsilon_p^{2-k})$, we have $\mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(V) \neq 0$ so that $\dim \mathcal{D}_{\text{crys}}(V) = 2$, i.e. that V is crystalline at p .

For $\ell \mid N$, [41, Lemma 4.1.3] shows that the S -extensions occurring in the cohomology group $H^1(\mathbb{Q}, \rho_f(k-1))$ (resp. $H^1(\mathbb{Q}, \rho_f(k-2))$) are unramified outside p . For proving that the S -extensions occurring in $H^1(\mathbb{Q}, \epsilon_p)$ are unramified at $\ell \mid N$ (under the hypothesis **(St)**), we use the semi-continuity of the rank of the monodromy operator attached to the Weil–Deligne representation at ℓ of p -adic families and that the rank is generically one for families of paramodular tame level.

Having bounded the number of generators of \mathcal{I}^{tot} by s^2 we determine in Section 7 the local ring $A := \mathcal{T}/\mathcal{I}^{\text{tot}}$ by proving under the hypothesis **(BK)** that the completion \widehat{A} of A with respect to its maximal ideal is isomorphic to the universal ring representing the p -ordinary minimally ramified deformations of ρ_f , which is a discrete valuation ring. Since \mathcal{T} is equidimensional of dimension 2, $\mathcal{T}/\mathcal{I}^{\text{tot}} = A$ is regular of dimension one (implied by \widehat{A} being regular) and \mathcal{I}^{tot} is principal when $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$ (or more generally generated by at most s^2 elements), it follows that the generator of \mathcal{I}^{tot} is a regular local parameter of \mathcal{T} when $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$ (or more generally, we obtain the desired bound of the Zariski tangent space of \mathcal{T}).

This means that the tangent space of \mathcal{T} is of dimension 2 when

$$\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$$

and \mathcal{T} is regular of dimension 2. Thus the rigid analytic space \mathcal{E}_{Δ} is smooth at π_{α} , and as a consequence, \mathcal{E}_{Δ} has a unique irreducible component (of dimension 2) specializing to π_{α} .

However, for the case when $k = 2$ (i.e Theorem 1.2(2)), we need to prove in addition that the S -extensions occurring in $H^1(\mathbb{Q}, \rho_f)$ are crystalline at p . This seems difficult to establish (see Remark 5.2). But we know that these extensions are ordinary in the sense that they have an unramified line on which Frob_p acts by α , and so they belong to a Greenberg’s type Selmer group $\text{Sel}_{\mathbb{Q}, f, \alpha}$ attached to $\rho_f^{\vee}(-1)$ (see Section 5.1). Moreover, we know from the Iwasawa main conjecture for GL_2 that the Pontryagin dual of the Λ -adic

Greenberg’s Selmer group of f_α is a torsion Λ -module, and its characteristic ideal contains the p -adic L function $L_p(f_\alpha, \omega_p^{-1}, \cdot)$ (see [42, Theorem 3.25]). Hence, the condition that $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$ is sufficient for the vanishing of $\text{Sel}_{\mathbb{Q}, f_\alpha}$.

By a general principle of Hida the adjoint L -function of modular forms controls congruences with other modular forms. This demonstrates that our assumptions are natural for understanding the geometry at π_α since $H_{f, \text{unr}}^1(\mathbb{Q}, \text{ad}^0(\varrho))$ contains the Selmer groups

$$H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k - 2)), H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k - 1)), \text{ and } H_{f, \text{unr}}^1(\mathbb{Q}, \text{ad}^0(\rho_f)),$$

where $\varrho = \epsilon_p^{1-k} \oplus \epsilon_p^{2-k} \oplus \rho_f$ is the semi-simple p -adic representation attached to π_α . As we show the first two groups control the generators of \mathcal{I}^{tot} (which correspond to congruences with stable components) and $H_f^1(\mathbb{Q}, \text{ad}^0(\rho_f))$ controls $\mathcal{T}/\mathcal{I}^{\text{tot}}$ (corresponding to the number of ordinary families of p -adic representations specializing to ρ_{f_α}).

1.2. Relationship to other results in the literature

Bellaïche–Chenevier studied in [7] the geometry of some eigenvarieties X attached to unitary Shimura varieties at points with reducible Galois representation and gave applications to the Bloch–Kato conjecture. They focus on points $z \in X$ with Galois representation given by $\mathbb{1} \oplus \epsilon_p \oplus \rho_z$, where ρ_z is an irreducible n -dimensional representation anti-ordinary at p . They proved that at $z \in X$, the local Galois deformation at p is irreducible on every Artinian thickening of z (the reducibility locus at z of the pseudo-character carried by X is the maximal ideal of $\mathcal{O}_{X, z}$). It should be pointed out that our setting is quite different since the reducibility locus at π_α of the pseudo-character $\text{Ps}_{\mathcal{E}_\Delta}$ is given by a principal Weil divisor of the 2-dimensional affine scheme $\text{Spec } \mathcal{T}$ and corresponds on the modular side to the Saito–Kurokawa lift of the Hida family passing through f_α . A further difference between these settings lies in the position of the Hodge–Tate weights and their distribution between the different pieces of the reducible Galois representations $\mathbb{1} \oplus \epsilon_p \oplus \rho_z$ and $\rho_{\pi_\alpha} := \epsilon_p^{1-k} \oplus \epsilon_p^{2-k} \oplus \rho_f$. More precisely, while the smallest Hodge–Tate of ρ_{π_α} is zero and occurs in the 2-dimensional representation ρ_f , the smallest Hodge–Tate weight of $\mathbb{1} \oplus \epsilon_p \oplus \rho_z$ is -1 and occurs in the one dimensional sub-representation ϵ_p , and ρ_z has no Hodge–Tate weights equal to $\{0, -1\}$, and this difference makes the proof of the crystallinity of the $S := \mathcal{T}[G_{\mathbb{Q}}]/\ker(\text{Ps}_{\mathcal{T}})$ -extensions occurring in $\text{Ext}_{G_{\mathbb{Q}}}^1(\epsilon^{1-k}, \epsilon^{2-k})$ (in our setting) more subtle than [7, Proposition 8.2.14]

(see Section 4). In addition, we investigate also in this paper the geometry of \mathcal{E}_Δ at Saito–Kurokawa points π_α of non-cohomological weights (i.e when $k = 2$) and in that case ρ_{π_α} has only two Hodge–Tate weights $\{0, 1\}$ (with multiplicity two).

Skinner–Urban constructed in [41, Theorem 2.4.10] a semi-ordinary eigenvariety as an admissible open of \mathcal{E}_N . Using a deep automorphic argument they established the existence of a stable semi-ordinary p -adic cuspidal component \mathcal{Y} of dimension 2 specializing to π_α (see [41, Theorem 4.2.7]), with fewer assumptions on the level and the local representation ρ_f at $\ell \mid Np$ than us (they assumed only that f is ordinary at p). They then applied the lattice construction of [46] (generalizing Ribet’s Lemma to higher dimensions) to obtain non-trivial extensions in $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$.

In [9] short crystalline, minimal, essentially self-dual deformations of non-semisimple mod p Galois representations $\bar{\rho}$ with $\bar{\rho}^{\text{ss}} = \bar{\rho}_{\pi_\alpha} = \bar{\epsilon}_p^{2-k} \oplus \bar{\rho}_f \oplus \bar{\epsilon}_p^{1-k}$ are studied. In this analysis the principality of the total reducibility ideal of the universal pseudodeformation of $\text{Tr}(\bar{\rho})$ to \mathcal{O}_L -algebras also played a crucial role.

Hernandez constructed in [20] a three dimensional p -adic eigenvariety for the group $\text{U}(2, 1)(E)$, where E is a quadratic imaginary field in which p is inert (the Picard modular surface has an empty ordinary locus in that case), and gave an application by reproving particular cases of the Bloch–Kato conjecture for Galois characters of E .

Acknowledgement

We would first like to thank the referee for their careful reading, helpful comments and suggestions that helped to improve the exposition and weaken the hypotheses in our result. We would like also to thank Riccardo Brasca, Valentin Hernandez, Kris Klosin, Vincent Pilloni, Jacques Tilouine, Chris Skinner and Eric Urban for helpful communications related to the topics of this article.

Notation and some remarks

- (1) Throughout this article, we fix a prime number p , as well as an algebraic closure $\bar{\mathbb{Q}}$ of \mathbb{Q} and $\bar{\mathbb{Q}}_p$ of \mathbb{Q}_p and an embedding $\iota_p : \bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_p$. Observe that ι_p yields a canonical embedding $G_{\mathbb{Q}_p} \hookrightarrow G_{\bar{\mathbb{Q}}}$ through which we will see $G_{\mathbb{Q}_p}$ as a decomposition group of $G_{\bar{\mathbb{Q}}}$ at p . We denote by I_p the inertia subgroup of $G_{\mathbb{Q}_p}$.

- (2) We fix an embedding $\overline{\mathbb{Q}} \subset \mathbb{C}$ and a complex conjugation $c \in G_{\mathbb{Q}}$.
- (3) Let $\mathbb{Q}_p(1)$ denote the $G_{\mathbb{Q}}$ representation of dimension 1 on which $G_{\mathbb{Q}}$ acts by the p -adic cyclotomic character $\epsilon_p : G_{\mathbb{Q}} \rightarrow \mathbb{Z}_p^{\times} \hookrightarrow \mathbb{Q}_p^{\times}$.
- (4) The Hodge–Tate–Sen weight of $\mathbb{Q}_p(1)$ is -1 and its Sen polynomial is $X + 1$ (we are following the geometric convention).
- (5) Let B_{crys} denote the crystalline period ring endowed with the semi-linear Frobenius Φ and the natural $G_{\mathbb{Q}_p}$ -action.
- (6) Let $t \in B_{\text{crys}}$ be the element on which $G_{\mathbb{Q}_p}$ -acts by ϵ_p and $\Phi(t) = p.t$. Note that t generates the maximal ideal of the integral de Rham periods ring B_{dR}^+ ; i.e $B_{\text{dR}}^+/t.B_{\text{dR}}^+ \simeq \mathbb{C}_p$ as $G_{\mathbb{Q}_p}$ -modules.
- (7) Let $B_{\text{crys}}^+ \subset B_{\text{crys}}$ denote the ring of period defined in [18, Exposé II, Section 2.3].
- (8) Let V be a $G_{\mathbb{Q}_p}$ -representation of finite dimension over a p -adic field L . Let $\mathcal{D}_{\text{crys}}(V)$ denote the L -vector space $(B_{\text{crys}} \otimes_{\mathbb{Q}_p} V)^{G_{\mathbb{Q}_p}}$ of dimension at most $\dim_L V$. And we denote again by Φ for the L -linear action given by $\Phi \otimes \text{Id}_V$ on $\mathcal{D}_{\text{crys}}(V)$. Denote also by $\mathcal{D}_{\text{crys}}^+(V)$ for $(B_{\text{crys}}^+ \otimes_{\mathbb{Q}_p} V)^{G_{\mathbb{Q}_p}}$.
- (9) Let $z \in \mathcal{E}_N$ be a classical point such that the Galois representation V attached to z is crystalline. Then the (Φ, Γ) -module attached to V is trianguline in the sense of Colmez. However, the triangulation can be given by non-étale (Φ, Γ) -submodules, and hence $V|_{G_{\mathbb{Q}_p}}$ is not necessarily ordinary at p .
- (10) Remark that $\mathcal{D}_{\text{crys}}^+(\epsilon_p) = 0$, $\mathcal{D}_{\text{crys}}(\epsilon_p) = \mathbb{Q}_p.t^{-1}$ (t^{-1} is not in B_{crys}^+), and $\mathcal{D}_{\text{crys}}^+(\epsilon_p^{-1}) = \mathbb{Q}_p.t$.
- (11) Let $\mathbb{1}$ be the trivial representation of dimension 1.
- (12) We shall always write Frob_{ℓ} for the geometric Frobenius at the prime ℓ .
- (13) Let $\alpha \in \mathbb{Q}$, we shall denote \mathcal{E}_N^{α} for the admissible open locus of \mathcal{E}_N defined by $|U_0 U_1|_p = \alpha$.
- (14) We write $G_{\mathbb{Q}}^{Np}$ for the Galois group of the maximal extension of \mathbb{Q} unramified outside of Np and ∞ . For any $G_{\mathbb{Q}}$ -geometric representation V we define the Bloch–Kato Selmer groups

$$H_{f, \text{unr}}^1(\mathbb{Q}, V) = \ker \left(H^1(\mathbb{Q}, V) \rightarrow H^1(\mathbb{Q}_p, V \otimes B_{\text{crys}}) \bigoplus_{\ell \nmid p} H^1(I_{\ell}, V) \right)$$

and

$$H_f^1(\mathbb{Q}, V) = \ker \left(H^1(\mathbb{Q}, V) \rightarrow H^1(\mathbb{Q}_p, V \otimes B_{\text{crys}}) \right).$$

- (15) Let A be a ring and M be a finite length A -module. We shall always denote by $l(M)$ the length of M as A -module.

2. Some properties of automorphic p -adic representations

In this section we recall some facts about the Galois representations associated to classical and Siegel modular forms. The end of this section is devoted to the proof of the first part of Theorem 1.2.

2.1. Ordinary Galois representations

We recall in the following definition the notion of ordinary p -adic representations.

DEFINITION 2.1. — *Let V be finite-dimensional L -vector space endowed with a continuous $G_{\mathbb{Q}_p}$ -action. We say that V is ordinary, if it has a decreasing filtration by $G_{\mathbb{Q}_p}$ -subrepresentations $\dots F^{i+1} \subset F^i \dots$ satisfying $\bigcap_i F^i = \{0\}$ and $\bigcup_i F^i = V$, and such that all graded factors have of the following form*

$$F^i/F^{i+1} = V_i(i), \text{ with } V_i \text{ unramified at } p, \text{ and } V_i(i) \text{ is the twist by } \epsilon_p^i.$$

Nekovar proved the following useful theorem.

THEOREM 2.2 ([33, Theorem 1.30]). — *Any ordinary p -adic representation is semistable.*

2.2. Elliptic modular forms

Recall that $(p, N) = 1$, $f \in S_{2k-2}(\Gamma_0(N), L)$ is an elliptic cuspidal eigenform with coefficients in the p -adic field L , and let $\rho_f : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(L)$ be the p -adic Galois representation attached to f in the sense that $L(\rho_f, s) = L(f, s)$. We note that $\rho_f^\vee \simeq \rho_f(2k-3)$ by the duality of 2-dimensional representations. It is known that ρ_f is de Rham and that its Hodge–Tate–Sen weights are $(2k-3, 0)$. Moreover, ρ_f is crystalline at p since $p \nmid N$.

Since f_α is ordinary at p , $(\rho_f)|_{G_{\mathbb{Q}_p}} \sim \begin{pmatrix} \psi & * \\ 0 & \psi^{-1} \epsilon_p^{3-2k} \end{pmatrix}$, where $\psi : G_{\mathbb{Q}_p} \rightarrow L^\times$ is the unramified character such that $\psi(\mathrm{Frob}_p) = \alpha$ (where $U_p(f_\alpha) = \alpha f_\alpha$) and $\det \rho_f = \epsilon_p^{3-2k}$. Note that the characteristic polynomial of the L -linear Frobenius Φ acting on $\mathcal{D}_{\mathrm{crys}}(\rho_f)$ is equal to the p -th Hecke polynomial of f .

PROPOSITION 2.3. — *Let $\ell \mid N$ be a prime number.*

(1) *Assume that $\pi_{f,\ell}$ is special at ℓ , then*

$$\begin{aligned} \dim \text{Ext}_{G_{\mathbb{Q}_\ell}}^1(\rho_f, \epsilon_p^{2-k}) &= \dim \text{Ext}_{G_{\mathbb{Q}_\ell}}^1(\epsilon_p^{1-k}, \rho_f) \\ &= \dim H^1(\mathbb{Q}_\ell, \rho_f(k-1)) = 0. \end{aligned}$$

(2) *Assume that $\pi_{f,\ell} \simeq \text{St} \otimes \xi$ (i.e. $a_\ell(f) = -\ell^{k-2}$), then*

$$\begin{aligned} \dim \text{Ext}_{G_{\mathbb{Q}_\ell}}^1(\rho_f, \epsilon_p^{1-k}) &= \dim \text{Ext}_{G_{\mathbb{Q}_\ell}}^1(\epsilon_p^{2-k}, \rho_f) \\ &= \dim H^1(\mathbb{Q}_\ell, \rho_f(k-2)) = 0. \end{aligned}$$

Remark 2.4. — When $k = 2$, the assumption that $a_\ell = -1$ when $\ell \mid N$ is a prime holds if and only if the abelian variety A_f attached to the weight 2 cuspidal eigenform f has non-split multiplicative reduction at ℓ .

Proof. — We know, in fact, that $(\rho_f)|_{G_{\mathbb{Q}_\ell}} = \begin{pmatrix} \psi_\ell^{-1} & * \\ 0 & \psi_\ell^{-1} \epsilon_p^{-1} \end{pmatrix}$ with infinite image of inertia, where ψ_ℓ is an unramified character such that $\psi_\ell(\text{Frob}_\ell) = a_\ell(f)$. Note that by [30, Theorem 4.6.17 (2)] $a_\ell^2(f) = \ell^{2k-4}$. Our assumption on a_ℓ implies that $H^0(\mathbb{Q}_\ell, \rho_f(k-1)) = H^0(\mathbb{Q}_\ell, \rho_f(k-2)) = 0$.

By applying the Euler characteristic formula and Tate duality, we obtain:

$$\begin{aligned} \dim H^1(\mathbb{Q}_\ell, \rho_f(k-1)) \\ = \dim H^0(\mathbb{Q}_\ell, \rho_f(k-1)) + \dim H^0(\mathbb{Q}_\ell, (\rho_f(k-1))^\vee(1)). \end{aligned}$$

Since $\rho_f^\vee = \rho_f(2k-3)$ (the duality for 2-dimensional representations), the above equality yields that

$$(2.1) \quad \dim H^1(\mathbb{Q}_\ell, \rho_f(k-1)) = 0.$$

The other cases are proved similarly. □

2.3. Siegel modular forms

We define the abstract Hecke algebra \mathcal{H}_N as the \mathbb{Z} -algebra generated by the Hecke operators $T_{\ell,1}, T_{\ell,2}, S_\ell$ for $\ell \nmid Np$ and the Hecke operators U_0, U_1 at p , where $T_{\ell,1}$ (resp. $T_{\ell,2}, S_\ell$) is the Hecke operator attached to $\text{diag}[1, 1, \ell, \ell]$ (resp. $\text{diag}[1, \ell, \ell^2, \ell], \text{diag}[\ell, \ell, \ell, \ell]$); and U_0, U_1 are the Hecke operators attached respectively to $\text{diag}[1, 1, p, p], \text{diag}[1, p, p^2, p]$ (U_1 is renormalized as in [41, Theorem 2.4.14]).

We recall the p -adic properties of Galois representation arising from Siegel modular eigenforms. The following theorem has been proved by Laumon and Weissauer (see [26, 48]).

THEOREM 2.5. — *Let π be a Siegel modular eigenform of central character ω_π of level $\Gamma(N)$ and of cohomological weight $k = (l_1, l_2)$ with corresponding Hecke character $\lambda_\pi : \mathcal{H}_N \rightarrow L_\pi$, where $L_\pi \subset \overline{\mathbb{Q}}_p$ is a p -adic field generated over \mathbb{Q}_p by the systems of Hecke eigenvalues of π . Then there exist a finite extension L'_π of L_π and a p -adic representation $\rho_\pi : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_4(L'_\pi)$ unramified outside Np and such that for all $\ell \nmid Np$,*

$$\det(X \cdot \mathrm{Id} - \rho_\pi(\mathrm{Frob}_\ell)) = P_{\pi, \ell}(X),$$

where $P_{\pi, \ell}(X)$ is the Hecke–Andrianov polynomial at ℓ attached to π . Moreover, we have the symplectic relation :

$$(2.2) \quad \rho_\pi \simeq \rho_\pi \otimes \chi_\pi^{-1},$$

and $\det \rho_\pi = \chi_\pi^2$. We also have the following relation between the similitude character χ_π and the central character:

$$\omega_\pi \epsilon_p^{3-l_1-l_2} = \chi_\pi.$$

We have the following properties at p of ρ_π following from the works of Chai–Faltings, Laumon, Taylor, Urban and Weissauer (see [17, 26, 44, 47, 48]).

THEOREM 2.6. — *Under the notations of the above theorem we have :*

- (1) *The Galois representation ρ_π is de Rham and its Hodge–Tate weights are*

$$\{0, l_2 - 2, l_1 - 1, l_1 + l_2 - 3\}.$$

- (2) *If π is old at p and semi-ordinary (i.e U_0 acts on π by a p -adic unit), then the p -adic representation ρ_π is crystalline at p , and the characteristic polynomial of Φ acting on $\mathcal{D}_{\mathrm{crys}}(\rho_\pi)$ is the Hecke polynomial at p . The eigenvalues of the L'_π -linear Frobenius Φ acting on $\mathcal{D}_{\mathrm{crys}}(\rho_\pi)$ are*

$$\{\lambda_\pi(U_0), \lambda_\pi(U_1 \cdot U_0^{-1})p^{l_2-2}, \lambda_\pi(U_0 \cdot U_1^{-1})^{-1}p^{l_1-1}, \lambda_\pi(U_0)^{-1}p^{l_1+l_2-3}\}.$$

- (3) *Assume that π is semi-ordinary at p and of finite slope for $\mathbb{U} = U_0U_1$, then*

$$(2.3) \quad (\rho_\pi)|_{G_{\mathbb{Q}_p}} \sim \begin{pmatrix} \phi_\pi & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & \phi_\pi^{-1} \epsilon_p^{-l_1-l_2+3} \end{pmatrix},$$

where $\phi_\pi : G_{\mathbb{Q}_p} \rightarrow L_\pi^\times$ is the unramified character having $\lambda_\pi(U_0)$ as value at Frob_p .

COROLLARY 2.7. — Assume that π is old at p , semi-ordinary, non-endoscopic and cohomological. Let $\rho_\pi^{I_p}$ be the $G_{\mathbb{Q}_p}$ -stable line of $(\rho_\pi)_{|G_{\mathbb{Q}_p}}$ on which $G_{\mathbb{Q}_p}$ acts by the unramified character ϕ_π , then the 2-dimensional $G_{\mathbb{Q}_p}$ -stable subspace W_π of the quotient of $(\rho_\pi)_{|G_{\mathbb{Q}_p}}$ by $\rho_\pi^{I_p}$ (see (2.3)) is crystalline with Hodge–Tate weight $(l_1 - 1, l_2 - 2)$. Moreover, the eigenvalues of the L'_π -linear Frobenius Φ acting on $\mathcal{D}_{\text{crys}}(W_\pi)$ are $\lambda_\pi(U_1 U_0^{-1})p^{l_2-2}$ and $\lambda_\pi(U_0 U_1^{-1})p^{l_1-1}$.

Remark 2.8. — Note that the p -adic Galois representation attached to a cuspidal Siegel eigenform is not necessarily irreducible. Schmidt makes the consequences of Arthur’s classification for GSp_4 explicit in [39]. All cuspidal automorphic representations are either of type (G), (Y), (B), (Q), or (P). The latter three are CAP representations, with type (P) for the Siegel parabolic including the Saito–Kurokawa lifts. Type (Y) representations are endoscopic representations (“of Yoshida type”). Type (G) representations are “stable” in the sense that their transfer to GL_4 stays cuspidal, and therefore their Galois representations are expected to be irreducible.

Recall that we write $\Delta := \prod_{\ell|N} \Delta_\ell \cap \text{GSp}_4(\mathbb{Q})$ for the paramodular congruence subgroup of level N . If $N = 1$ we put $\Delta = \text{GSp}_4(\mathbb{Z})$.

PROPOSITION 2.9. — Let π be a holomorphic Siegel modular eigenform of cohomological weight (l_1, l_2) , trivial central character ω_π and of paramodular level Δ for $N \geq 1$. Then π is either of type (G) or a Saito–Kurokawa lift (in particular it is not endoscopic).

Proof. — This is proven in [39, Lemma 2.5] and [40, Proposition 5.2]. \square

2.4. Properties at $\ell \neq p$ of a p -adic representation arising from a Siegel cusp form

We have the following result on the local properties of ρ_π at the primes $\ell \mid N$ (compare [41, Conjecture 3.1.7]) proved by [32, Theorem 3.5] (local-global compatibility up to Frobenius semi-simplification for type (G) representations) and [43, Corollary 1] (monodromy rank 1). Mok [32] used Arthur’s classification for GSp_4 , whose proof was completed by Gee–Taibi in [19]⁽³⁾.

⁽³⁾Note the comments at [19, p. 472] about this result still being conditional on unpublished work of Arthur and cases of the twisted weighted fundamental lemma.

PROPOSITION 2.10. — *Under the notations of Theorem 2.5, and assuming that π is non-CAP and non-endoscopic and $\pi^{\Delta_\ell} \neq 0$ for $\ell \mid N$, the rank of the monodromy operator of the Weil–Deligne representation attached to the Galois representation $(\rho_\pi)|_{G_{\mathbb{Q}_\ell}}$ is at most one.*

2.5. Non-existence of endoscopic components of \mathcal{E}_Δ specializing to π_α

Recall that f is an elliptic eigenform of $S_{2k-2}(\Gamma_0(N), L)$, $\text{SK}(f)$ is the Saito–Kurokawa lift of f to a weight (k, k) cuspform of paramodular level Δ , and π_α is a semi-ordinary p -stabilization of $\text{SK}(f)$ (the eigenvalue of π_α for U_0 is a p -adic unit). We recall also that π_α defines a point of the eigenvariety \mathcal{E}_Δ which we denote again by π_α . The following theorem is a consequence of [41, 3.2.9] and Proposition 2.9.

THEOREM 2.11. — *Assume (Reg) and $k \geq 2$, then any irreducible affinoid \mathcal{Z} of \mathcal{E}_Δ of dimension two containing π_α is stable.*

Proof. — Assume that there exists such an irreducible affinoid \mathcal{Z} that is not stable, i.e. that the pseudo-character $\text{Ps}_{\mathcal{Z}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{Z})$ is reducible. Since \mathcal{Z} specializes to π_α and ρ_f is absolutely irreducible, a subconstituent of the pseudo-character $\text{Ps}_{\mathcal{Z}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{Z}) \rightarrow \mathcal{O}_{\mathcal{Z}, \pi_\alpha}$ is a pseudo-character of dimension 2 whose reducibility locus is of dimension at most one. (One can rule out the existence of a 3-dimensional irreducible constituent by specializing at sufficiently regular classical weights and applying the argument from the proof of Case A (iii) in [41, Theorem 3.2.1].) Hence one can find a sufficiently small affinoid neighborhood $\mathcal{X} = \text{Spm } R$ of π_α with an odd representation $\varrho_1 : G_{\mathbb{Q}} \rightarrow \text{GL}_2(R)$ specializing to the 2-dimensional odd representation ρ_f and such that any classical specialization of ϱ_1 is irreducible, and a 2-dimensional pseudo-character $\text{Ps}_2 : G_{\mathbb{Q}} \rightarrow R$ specializing to $\epsilon^{1-k} \oplus \epsilon^{2-k}$, and such that $\text{Tr } \varrho_1 + \text{Ps}_2 = \text{Ps}_{\mathcal{X}}$ (Ps_2 is the trace of a 2-dimensional $G_{\mathbb{Q}}$ -representation ϱ_2 valued in $\text{Frac } R$). Moreover, the p -regularity assumption on π_α (when $k = 2$) and [41, Proposition 3.3.6] yield (after shrinking again \mathcal{X} to a smaller affinoid which we denote again by \mathcal{X}) that ϱ_1 is ordinary at p (in the sense that $\varrho_1^{I_p}$ is a direct summand in ϱ_1 of rank 1). Hence, Theorem [41, 3.2.9] implies that any specialization of \mathcal{X} at a classical point $z \in \mathcal{X}$ of a cohomological weight is CAP or endoscopic. Note that π_α is of algebraic weight, has trivial central character, and is old at p . This implies that the old (at p) classical points of trivial central character and non-parallel very regular weights of \mathcal{X} are Zariski

dense (see Corollary B.5). By [46, Proposition 3.3] the specialization of \mathcal{X} at these points cannot be a CAP form and hence necessarily endoscopic by Theorem [41, 3.2.9]. Since such forms do not exist by Proposition 2.9 we get the desired contradiction. \square

3. The GMA S and ordinarity of S -extensions occurring in $H^1(\mathbb{Q}, \rho_f(k-1))$

Recall that Theorem 2.11 implies that all irreducible components of \mathcal{E}_Δ passing through π_α are stable, and that \mathcal{T} , the local ring of \mathcal{E}_Δ at π_α , is reduced and equidimensional of dimension 2 since \mathcal{E}_Δ is reduced and equidimensional of dimension 2. Let \mathfrak{m} be the maximal ideal of \mathcal{T} and L be the residue field of \mathcal{T} .

Let A be a reduced Noetherian ring. Recall that the total fraction ring of A is the fraction ring $Q(A) := \mathcal{S}^{-1}A$ where $\mathcal{S} \subset A$ is the multiplicative subset of non-zero-divisors of A . We check at once that the natural map $A \rightarrow \mathcal{S}^{-1}A$ is injective and flat, and that the non-zero-divisors of A are invertible in $\mathcal{S}^{-1}A$. Moreover, since A is Noetherian the zero divisors of A are the elements of the union of the (finitely many) minimal prime ideal of A , so $\mathcal{S}^{-1}A = \prod_{\mathcal{P}_i} A_{\mathcal{P}_i}$, where \mathcal{P}_i runs over the minimal prime ideals of A . Moreover, each $A_{\mathcal{P}_i}$ is a field, since it is reduced, local and of Krull dimension equal to zero. Let $K = \prod K_i$ be the total field of fractions of the reduced equidimensional ring \mathcal{T} , where K_i is the localisation of \mathcal{T} at a minimal prime ideal.

DEFINITION 3.1 (Definition/Proposition). — *The pseudo-character⁽⁴⁾*

$$\text{Ps}_{\mathcal{T}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta) \rightarrow \mathcal{T}$$

is residually multiplicity free and the corresponding Cayley–Hamilton faithful algebra

$$S := \mathcal{T}[G_{\mathbb{Q}}] / \ker \text{Ps}_{\mathcal{T}}$$

can by [7, Theorem 1.4.4(i)] be equipped with the structure of a GMA (in the sense of [7, Definition 1.3.1]). It is of finite type and torsion-free as \mathcal{T} -module. Since \mathcal{T} is reduced we further have an associated Galois representation $\rho_K : G_{\mathbb{Q}} \rightarrow \text{GL}_4(K)$ by [7, Theorem 1.4.4(ii)]. Note that $\rho_K : G_{\mathbb{Q}} \rightarrow \text{GL}_4(K)$ is absolutely irreducible, since all the minimal prime ideals of \mathcal{T} correspond to stable irreducible components of \mathcal{E}_Δ passing

⁽⁴⁾The pseudo-character $\text{Ps}_{\mathcal{T}}$ is obtained by composing $\text{Ps}_{\mathcal{E}_\Delta}$ with the localization map $\mathcal{O}(\mathcal{E}_\Delta) \rightarrow \mathcal{T}$.

through π_α (so each Galois representation $\rho_{K_i} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_4(K_i)$ is irreducible).

Assume until the end of this paper that $\alpha \neq 1$ when $k = 2$ (which we will refer to as “ p -adic regularity”). Recall that $\varrho = \begin{pmatrix} \epsilon_p^{2-k} & 0 & 0 \\ 0 & \rho_f & 0 \\ 0 & 0 & \epsilon_p^{1-k} \end{pmatrix}$ is the

Galois representation attached to π_α in a basis such that $\varrho(\tau) \sim \begin{pmatrix} \mu & 0 & 0 & 0 \\ 0 & \alpha & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & \gamma \end{pmatrix}$, where the eigenvalues of $\tau \in G_{\mathbb{Q}_p}$ are all distinct (since $\alpha \neq 1$ when $k = 2$)

(necessarily in this basis $\varrho(G_{\mathbb{Q}_p}) \sim \begin{pmatrix} \epsilon_p^{2-k} & 0 & 0 & 0 \\ 0 & \psi & * & 0 \\ 0 & 0 & \psi^{-1} \epsilon_p^{3-2k} & 0 \\ 0 & 0 & 0 & \epsilon_p^{1-k} \end{pmatrix}$).

Remark 3.2. — Note that the character ϕ_{π_α} of Theorem 2.6 (3) equals ψ since $U_p(f_\alpha) = \alpha \cdot f_\alpha$ and $U_0(\pi_\alpha) = \alpha \cdot \pi_\alpha$.

Since all reducible components of ϱ have multiplicity one, [7, Theorem 1.4.4] implies that there exist orthogonal idempotents $\tilde{e}_1, \tilde{e}_2, \tilde{e}_3$ of $S = \mathrm{Im}(\mathcal{T}[G_{\mathbb{Q}}] \rightarrow M_4(K))$ lifting the idempotents e_1, e_2, e_3 of ϱ , and corresponding respectively to $\epsilon_p^{2-k}, \rho_f, \epsilon_p^{1-k}$. Moreover, we can see S as

$$S = \begin{pmatrix} \mathcal{T} & M_{1,2}(\mathcal{T}_{1,2}) & \mathcal{T}_{1,3} \\ M_{2,1}(\mathcal{T}_{2,1}) & M_2(\mathcal{T}) & M_{2,1}(\mathcal{T}_{2,3}) \\ \mathcal{T}_{3,1} & M_{1,2}(\mathcal{T}_{3,2}) & \mathcal{T} \end{pmatrix},$$

where $\mathcal{T}_{i,j}$ are fractional ideals of K ($\mathcal{T}_{i,j}$ are finite type \mathcal{T} -modules).

Put $\rho_1 = \epsilon_p^{2-k}, \rho_2 = \rho_f$ and $\rho_3 = \epsilon_p^{1-k}$. We recall Bellaïche and Chenevier’s definition of reducibility ideals:

DEFINITION 3.3 ([7] Definition 1.5.2, Proposition 1.5.1). — *Let $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_s)$ be a partition of the set $\mathcal{I} = \{1, 2, 3\}$. The ideal of reducibility $I^{\mathcal{P}}$ (associated to $\mathrm{Ps}_{\mathcal{T}}$ and the partition \mathcal{P}) is the smallest ideal I of \mathcal{T} with the property that there exist pseudocharacters $T_1, \dots, T_s : \mathcal{T}/I[G_{\mathbb{Q}}] \rightarrow \mathcal{T}/I$ such that*

- (1) $\mathrm{Ps}_{\mathcal{T}} \otimes \mathcal{T}/I = \sum_{l=1}^s T_l$,
- (2) for each $l \in \{1, \dots, s\}$, $T_l \otimes L = \sum_{i \in \mathcal{P}_l} \mathrm{trace} \rho_i$.

PROPOSITION 3.4 ([7, Proposition 1.5.1], [9, Corollary 6.5]). — *One has*

$$I^{\mathcal{P}} = \sum_{\substack{(i,j) \\ i, j \text{ not in the same } \mathcal{P}_l}} \mathcal{T}_{i,j} \mathcal{T}_{j,i}.$$

For $\mathcal{P} = \{\{1\}, \{2\}, \{3\}\}$ we write

$$\mathcal{I}^{\mathrm{tot}} := I^{\mathcal{P}} = \mathcal{T}_{3,1} \mathcal{T}_{1,3} + \mathcal{T}_{2,3} \mathcal{T}_{3,2} + \mathcal{T}_{1,2} \mathcal{T}_{2,1}.$$

Let $\mathcal{T}'_{i,j} = \mathcal{T}_{i,k}\mathcal{T}_{k,j}$ for i, j, k distinct. Since the maximal ideal \mathfrak{m} of \mathcal{T} contains the total reducibility ideal \mathcal{I}^{tot} [7, Theorem 1.5.5] implies that for $i \neq j \in \{1, 2, 3\}$ there exists an injective homomorphism of L -modules

$$(3.1) \quad \text{Hom}(\mathcal{T}_{i,j}/\mathcal{T}'_{i,j}, L) \hookrightarrow H^1(\mathbb{Q}_{Np}, \rho_i^\vee \otimes \rho_j \otimes L).$$

THEOREM 3.5. — Assume that $\alpha \neq 1$ when $k = 2$. For $(i, j) = (1, 2)$ the injective homomorphism of L -modules of (3.1) gives rise to

$$(3.2) \quad \text{Hom}(\mathcal{T}_{1,2}/\mathcal{T}'_{1,2}, L) \hookrightarrow H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)).$$

Proof. — The proof of [7, Theorem 1.5.5] tells us that the homomorphism (3.1) is given by

$$(3.3) \quad \begin{aligned} \text{Hom}(\mathcal{T}_{1,2}/\mathcal{T}'_{1,2}, L) &\hookrightarrow H^1(\mathbb{Q}, \rho_f(k-1)) \\ h &\mapsto (g \rightarrow h(\bar{b}_1(g), \bar{b}_2(g))\rho_f^{-1}(g)), \end{aligned}$$

where $(\bar{b}_1(g), \bar{b}_2(g))$ is the class of $t_{1,2}(g) = (b_1(g), b_2(g)) \in M_{1,2}(\mathcal{T}_{1,2})$ in $M_{1,2}(\mathcal{T}_{1,2}/\mathcal{T}'_{1,2})$. The classical points old at p and of regular weights form a very Zariski dense set Σ in every irreducible component of \mathcal{E}_Δ specializing to π_α (see Lemma B.3 and [41, Proposition 3.3.6]). By Theorem 2.6, the set of Hodge–Tate–Sen weights of the semi-simple representation ρ_y attached to any point $y \in \mathcal{E}_\Delta$ corresponding to a classical cuspidal Siegel eigenform old at p of weight (l_1, l_2) is $\{0, l_2 - 2, l_1 - 1, l_1 + l_2 - 3\}$ and ρ_y is crystalline at p .

On the other hand, for any $y \in \Sigma$, let us denote by $\rho_y : G_\mathbb{Q} \rightarrow \text{GL}_4(L_y)$ the semi-simple p -adic Galois representation attached to the Siegel eigenform corresponding to y (i.e. $\text{Tr } \rho_y$ is the specialization of the universal pseudo-character $\text{Ps}_{\mathcal{E}_\Delta} : G_\mathbb{Q} \rightarrow \mathcal{O}(\mathcal{E}_\Delta)$ at y). Theorem 2.6 implies that $\dim \mathcal{D}_{\text{crys}}^+(\rho_y)^{\Phi=U_0(y)} = 1$, and then $(\text{Ps}_{\mathcal{E}_\Delta}, \Sigma, U_0, \{\kappa_i\})$ is a weakly refined family (in the sense of [7, Definition 4.2.7]) since $U_0 \in \mathcal{O}(\mathcal{E}_\Delta)^\times$. Note also that condition $(*)$ of [7, Definition 4.2.7] is satisfied since we have a torsion free morphism $\kappa : \mathcal{E}_\Delta \rightarrow \mathcal{W}$; condition (v) of [7, Definition 4.2.7] is satisfied by Lemma A.6, Lemma B.3 and Corollary B.5 (so the classical points of \mathcal{E}_Δ which are old at p accumulate to π_α).

Moreover, $\dim \mathcal{D}_{\text{crys}}^+(\varrho)^{\Phi=\alpha=U_0(\pi_\alpha)} = 1$ by regularity assumption on ϱ at p . Hence, [7, Theorem 4.3.6] implies that any $G_\mathbb{Q}$ -representation V corresponding to a cohomology class in the image of the morphism (3.3) satisfies

$$\dim \mathcal{D}_{\text{crys}}^+(V)^{\Phi=\alpha} = 1.$$

We use this to first prove that V is crystalline at p . One can see V as the following $G_\mathbb{Q}^{Np}$ -extension:

$$0 \rightarrow \epsilon_p^{2-k} \rightarrow V \rightarrow \rho_f \rightarrow 0.$$

Let $\tilde{\rho} = \begin{pmatrix} \epsilon_p^{2-k} & * \\ 0 & \rho_f \end{pmatrix}$ be the realization of V by a matrix. The restriction to $G_{\mathbb{Q}_p}$ of $\tilde{\rho}$ has the form $\begin{pmatrix} \epsilon_p^{2-k} & b & c \\ 0 & \psi & * \\ 0 & 0 & \psi^{-1}\epsilon_p^{3-2k} \end{pmatrix}$. Hence, we have an extension of $G_{\mathbb{Q}_p}$ -modules

$$0 \rightarrow \begin{pmatrix} \epsilon_p^{2-k} & b \\ 0 & \psi \end{pmatrix} \rightarrow \tilde{\rho}|_{G_{\mathbb{Q}_p}} \rightarrow \psi^{-1}\epsilon_p^{3-2k} \rightarrow 0.$$

Let $V^0 \subset V$ be the L -vector space of dimension 2 on which $G_{\mathbb{Q}_p}$ acts by $\begin{pmatrix} \epsilon_p^{2-k} & b \\ 0 & \psi \end{pmatrix}$. By applying the left exact functor $\mathcal{D}_{\text{crys}}^+(\cdot)^{\Phi=\alpha}$ to the above exact sequence, we obtain

$$\mathcal{D}_{\text{crys}}^+(V^0)^{\Phi=\alpha} \simeq \mathcal{D}_{\text{crys}}^+(\tilde{\rho}|_{G_{\mathbb{Q}_p}})^{\Phi=\alpha}.$$

Since $\dim \mathcal{D}_{\text{crys}}(V)^{\Phi=\alpha} = 1$, we get $\dim \mathcal{D}_{\text{crys}}^+(V^0)^{\Phi=\alpha} = 1$. Hence, $V_0 = \begin{pmatrix} \epsilon_p^{2-k} & b \\ 0 & \psi \end{pmatrix}$ is crystalline at p which implies that the cohomology class of b in $\text{Ext}_{G_{\mathbb{Q}_p}}^1(\psi, \epsilon_p^{2-k})$ is trivial (i.e. $\tilde{\rho}|_{G_{\mathbb{Q}_p}} \simeq \begin{pmatrix} \epsilon_p^{2-k} & 0 & c \\ 0 & \psi & * \\ 0 & 0 & \psi^{-1}\epsilon_p^{3-2k} \end{pmatrix}$). Thus, $\tilde{\rho}$ is ordinary in the sense of Definition 2.1 and then semi-stable (hence de Rham) at p by Theorem 2.2. Therefore the extension V gives a cohomology class in

$$\mathrm{H}_g^1(G_{\mathbb{Q}}^{Np}, \rho_f(k-1)) = \ker\left(\mathrm{H}^1(\mathbb{Q}, \rho_f(k-1)) \rightarrow \mathrm{H}^1(\mathbb{Q}_p, \rho_f(k-1) \otimes B_{\text{dR}})\right).$$

Since $\mathrm{H}_g^1(G_{\mathbb{Q}}^{Np}, \rho_f(k-1)) \simeq \mathrm{H}_f^1(G_{\mathbb{Q}}^{Np}, \rho_f(k-1))$ (see e.g. [41, Lemme 4.1.3]) we deduce that V is crystalline at p .

Finally, the restriction of the map

$$\mathrm{H}^1(\mathbb{Q}, \rho_f(k-1)) \rightarrow \mathrm{H}^1(I_\ell, \rho_f(k-1)),$$

when $\ell \mid N$ is trivial by Proposition [41, Lemme 4.1.3]. □

3.1. Symplectic relation and the anti-involution τ on S

Recall that

$$(3.4) \quad \text{Ps}_{\mathcal{E}_\Delta} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta)$$

are pseudo-characters of dimension 4 and since the classical points of \mathcal{E}_Δ are Zariski dense, the relation (2.2) implies that the pseudo-character Ps_τ is invariant under the anti-involution

$$\tau : \mathcal{T}[G_{\mathbb{Q}}] \rightarrow \mathcal{T}[G_{\mathbb{Q}}] \text{ sending } g \rightarrow \epsilon_p^{-\kappa_1 - \kappa_2 + 3} \cdot g^{-1},$$

where $\epsilon_p^{-\kappa_1 - \kappa_2 + 3}$ is the character $G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{U})^\times$ interpolating the similitude character of the $G_{\mathbb{Q}}$ -semi-simple representations whose trace correspond to the classical specializations of the pseudo-character $\text{Ps}_{\mathcal{U}} : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_{\Delta}) \rightarrow \mathcal{O}(\mathcal{U})$ for a small enough connected affinoid neighborhood \mathcal{U} of π_{α} .

Hence τ yields an anti-automorphism on S given by $\rho_K \circ \tau$ and it follows from [7, Lemma 1.8.3] that we can choose our idempotent $\tilde{e}_1, \tilde{e}_2, \tilde{e}_3$ of S lifting the idempotents e_1, e_2, e_3 attached respectively to $\epsilon_p^{2-k}, \rho_f, \epsilon_p^{1-k}$, and such that $\tilde{e}_{\tau(1)} = \tilde{e}_3$ (τ preserves the idempotent corresponding to ρ_f , and switches the idempotents corresponding to $\epsilon_p^{1-k}, \epsilon_p^{2-k}$).

By (3.1) there exists an injection

$$(3.5) \quad \text{Hom}(\mathcal{T}_{2,3}/\mathcal{T}'_{2,3}, L) \hookrightarrow \text{Ext}_{G_{\mathbb{Q}}^{Np}}^1(\epsilon_p^{1-k}, \rho_f) \simeq H^1(\mathbb{Q}, \rho_f(k-1)).$$

Proposition [7, Proposition 1.8.6] yields immediately the following corollary.

COROLLARY 3.6. — *Assume that $\alpha \neq 1$ when $k = 2$. Then the image of (3.5) lands in $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ and has dimension equal to the dimension of the image of the morphism (3.2).*

4. Crystallinity of the S -extensions occurring in $H^1(\mathbb{Q}, \epsilon_p)$

In this section we show using the analytic continuation of the crystalline periods in a family of p -adic $G_{\mathbb{Q}_p}$ -representations of generic rank 3 interpolating $\{\rho_z/\rho_z^I, z \in \mathcal{E}_{\Delta}^{|U_{0,p}|=1}\}$ the crystallinity of the S -extensions occurring in $H^1(\mathbb{Q}, \epsilon_p)$. Assume in this section **(Reg)**, **(St)** and that $k \geq 2$.

By (3.1) we have a natural injection

$$(4.1) \quad \begin{aligned} \text{Hom}(\mathcal{T}_{1,3}/\mathcal{T}'_{1,3}, L) &\hookrightarrow \text{Ext}_{G_{\mathbb{Q}}^{Np}}^1(\epsilon_p^{1-k}, \epsilon_p^{2-k}) \simeq H^1(G_{\mathbb{Q}}^{Np}, \epsilon_p) \\ h &\mapsto \left(g \rightarrow \frac{h(\tilde{t}_{1,3}(g))}{\epsilon_p^{1-k}(g)} \right), \end{aligned}$$

where $\tilde{t}_{1,3}(g)$ is the class of $t_{1,3}(g) \in \mathcal{T}_{1,3}$ in $\mathcal{T}_{1,3}/\mathcal{T}'_{1,3}$.

Now we have to determine the exact image of the injective morphism (4.1). In [7, Section 1.5.4], Bellaïche–Chenevier introduce a left ideal $M_3 = S.E_3$ of $S \subset M_4(K)$ which is the third column of the GMA matrix S and hence it is a projective left S -module (see [7, 1.3.3] for the definition of E_3), and they proved in [7, Theorem 1.5.6] and [7, Lemma 4.3.9] the following results:

(i) There exists an exact sequence of S -left modules

$$(4.2) \quad 0 \rightarrow E \rightarrow M_3/\mathfrak{m}M_3 \rightarrow \epsilon_p^{1-k} \rightarrow 0$$

- (ii) Any simple S -subquotient of E occurs in the set $\{\rho_f, \epsilon_p^{2-k}\}$ (in particular it is not isomorphic to ϵ_p^{1-k}).
- (iii) The image of the morphism (4.1) consists of extensions occurring as quotient of the $S/\mathfrak{m}S$ -module $M_3/\mathfrak{m}M_3 \oplus \epsilon_p^{2-k}$ by an S -submodule \mathcal{Q} such that the S -simple subquotient of \mathcal{Q} occurs in $\{\rho_f, \epsilon_p^{2-k}\}$ (in particular it is not isomorphic to ϵ_p^{1-k}).

We will need the following additional property:

LEMMA 4.1. — *Let S_p be the subring generated by the image of $G_{\mathbb{Q}_p}$ in S . Then the S_p -simple subquotients of \mathcal{Q} occur in $\{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{3-2k}\}$.*

Proof. — Let Ps_p be the restriction of $\text{Ps}_{\mathcal{T}}$ to S_p . By [7, Lemma 1.2.7] we have $S/\text{rad}(S) \cong \overline{S}/\ker \overline{\text{Ps}}$ and $S_p/\text{rad}(S_p) \cong \overline{S}_p/\ker \overline{\text{Ps}}_p$, hence $\text{rad}(S) \cap S_p \subset \text{rad}(S_p)$, and we obtain a morphism $S_p/\text{rad}(S) \cap S_p \twoheadrightarrow S_p/\text{rad}(S_p) = \overline{S}_p/\ker \overline{\text{Ps}}_p \cong \prod_{i=1}^4 \text{End}_L(\rho_i)$, where $\rho_i \in \{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{2k-3}, \epsilon_p^{1-k}\}$. In particular, one can see that all $\{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{2k-3}, \epsilon_p^{1-k}\}$ are simple S_p -modules. Now, we claim that any simple S_p -representation occurs in

$$\{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{2k-3}, \epsilon_p^{1-k}\},$$

and it follows immediately from the injection $S_p/\text{rad}(S) \cap S_p \hookrightarrow S/\text{rad}(S) \simeq \overline{S}/\ker \overline{\text{Ps}} \cong \prod_{i=1}^3 \text{End}_L(\rho_i)$ whose image is $\prod_{i=1}^3 \text{End}_L((\rho_i)|_{G_{\mathbb{Q}_p}})$ (so $S_p/\text{rad}(S_p)$ is a semi-simple quotient of $\prod_{i=1}^3 \text{End}_L((\rho_i)|_{G_{\mathbb{Q}_p}})$).

The rest of the lemma follows from the fact (see (iii) above) that the S -module \mathcal{Q} has a Jordan–Holder sequence, all subquotients of which are isomorphic to either ρ_f or ϵ_p^{2-k} , and it has a refinement as S_p -module for which the S_p -simple subquotients occur in $\{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{2k-3}\}$. \square

We recall that a torsion-free A -module is a module over a ring A such that 0 is the only element annihilated by a regular element (i.e non-zero-divisor of A) of the ring. A coherent sheaf \mathcal{F} over a rigid analytic space X is a sheaf of $\mathcal{O}_X^{\text{rig}}$ -modules such that there exists an admissible covering of X by affinoid subdomains $\{U_i = \text{Spm } R_i\}$ of X for which the restriction $\mathcal{F}|_{U_i}$ is associated to \widetilde{M}_i and M_i is a finite type R_i -module.

The sheaf \mathcal{F} is said to be torsion-free if all those modules M_i are torsion-free over their respective rings. Alternatively, \mathcal{F} is torsion-free if and only if it has no local torsion sections.

LEMMA 4.2.

- (1) *One has $M_3 \subset K^4$ and $M_3.K = K^4$. Moreover, M_3 is a \mathcal{T} -torsion-free lattice of the representation $\rho_K \rightarrow \text{GL}_4(K)$.*

(2) *The natural morphism $M_3 \rightarrow M_3 \otimes_{\mathcal{T}} K$ is injective and the natural morphism*

$$M_3 \otimes_{\mathcal{T}} K \rightarrow M_3.K$$

is an isomorphism.

Proof.

(1). — Note that the finite type \mathcal{T} -module M_3 corresponds to the third column of the GMA matrix $S \subset M_4(K)$, hence $M_3 \subset K^4$. Since $\rho_K : G_{\mathbb{Q}} \rightarrow S^{\times} \subset \mathrm{GL}_4(K)$ is irreducible, then $M_3.K$ is necessarily of rank 4 over K .

(2). — Recall that $M_3 \otimes_{\mathcal{T}} K = M_3 \otimes_{\mathcal{T}} \mathcal{S}^{-1}\mathcal{T}$, where \mathcal{S} is the set of non-zero divisors of \mathcal{T} . Hence, $M_3 \otimes_{\mathcal{T}} K = \mathcal{S}^{-1}M_3$ and the injection follows from the fact that M_3 is torsion-free. It yields also that the natural surjection $M_3 \otimes_{\mathcal{T}} K \twoheadrightarrow M_3.K = K^4$ is in fact an isomorphism. \square

THEOREM 4.3. — *Assume **(Reg)** and **(St)**. Then the image of the injective morphism of L -modules*

$$\mathrm{Hom}(\mathcal{T}_{1,3}/\mathcal{T}'_{1,3}, L) \hookrightarrow \mathrm{Ext}_{G_{\mathbb{Q}}^{N_p}}^1(\epsilon_p^{1-k}, \epsilon_p^{2-k}) \simeq H^1(G_{\mathbb{Q}}^{N_p}, \epsilon_p)$$

lands in $H_{f,\mathrm{unr}}^1(\mathbb{Q}, \epsilon_p) = \{0\}$.

Proof. — To simplify notation, let M denote the finite type S -module M_3 . We recall that M is a torsion-free finite type \mathcal{T} -module, because S is of finite type over \mathcal{T} and $M \subset S \subset M_4(K)$. According to [7, Lemma 4.3.7], there exists an open affinoid neighborhood $\mathcal{U} = \mathrm{Spm} A$ of π_{α} inside \mathcal{E}_{Δ} such that we can extend M to an analytic torsion-free coherent sheaf $\widetilde{\mathcal{M}}$ over \mathcal{U} (\mathcal{M} is the A -module associated to $\widetilde{\mathcal{M}}$) and such that:

- $Q(A) \otimes \mathcal{M} = Q(A)^4$ (i.e the generic rank of \mathcal{M} is 4 ⁽⁵⁾)
- $\mathcal{M} \otimes_A \mathcal{T} = M$ (i.e the stalk of $\widetilde{\mathcal{M}}$ at π_{α} is M).
- The A -module \mathcal{M} carries a continuous action of $G_{\mathbb{Q}}$ compatible with the action of $G_{\mathbb{Q}}$ on its localization M at π_{α} , and the generic representation $G_{\mathbb{Q}} \rightarrow \mathrm{GL}_4(Q(A))$ is semi-simple and its trace is just the trace given by $\mathrm{Ps}_A : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_{\Delta}) \rightarrow A$.

On the other hand, by semi-ordinarity at p , the action of I_p on $Q(A)^4$ stabilizes a line $(Q(A)^4)^{I_p}$ on which Frob_p acts by U_0 . Let $\widetilde{\mathcal{L}}$ be the subsheaf of \mathcal{M} given by $(Q(A)^4)^{I_p} \cap \mathcal{M}$ (i.e the sections of $\widetilde{\mathcal{L}}$ are the sections of $\widetilde{\mathcal{M}}$ on which I_p acts trivially and Frob_p acts by U_0). Moreover, $\widetilde{\mathcal{L}}$ is the coherent sheaf associated to the A -submodule \mathcal{L} of \mathcal{M} given by the elements which

⁽⁵⁾ We have to choose $\mathrm{Spm} A$ small enough in the aim that it is connected and it contains no more irreducible components than $\mathrm{Spec} \mathcal{T}$, to have a natural inclusion $Q(A) \subset K$.

are invariant under the actions of the inertia I_p and on which Frob_p acts by U_0 .

Let $\widetilde{\mathcal{M}}'_+$ be the quotient presheaf $\widetilde{\mathcal{M}}/\widetilde{\mathcal{L}}$ and $\widetilde{\mathcal{M}}'$ be the sheaf associated to the presheaf $\widetilde{\mathcal{M}}'_+$, and it is $\widetilde{\mathcal{M}}/\widetilde{\mathcal{L}}$ since \mathcal{U} is an affinoid, and is endowed naturally with an action of $G_{\mathbb{Q}_p}$.

Let $M' := \mathcal{M}' \otimes_A \mathcal{T}$. Since $M_K := M \otimes_{\mathcal{T}} K = M \cdot K = K^4$, it is obvious that M' is a \mathcal{T} -submodule of $K^4/(K^4)^{I_p}$, where $(K^4)^{I_p}$ means the I_p -invariant subspace on which Frob_p acts by U_0 . Hence, M' is a finite type torsion-free \mathcal{T} -module of generic rank 3 over K , and the regularity assumption when $k = 2$ yields that the $G_{\mathbb{Q}_p}$ -semi-simplification $M' \otimes_{\mathcal{T}} L$ does not contain ψ .

Similarly, since $Q(A) \subset K$ and $(K^4)^{I_p} \cap \mathcal{M} = \mathcal{M}^{I_p}$, we obtain that \mathcal{M}' injects into $K^4/(K^4)^{I_p}$ and \mathcal{M}' is torsion-free over A and with generic rank equal to 3. Moreover, the regularity assumption yields that the $G_{\mathbb{Q}_p}$ -semi-simplification of its specialization at $\pi_\alpha = x$ does not contain the character $\psi|_{G_{\mathbb{Q}_p}}$. In fact, Corollary 2.7 implies the characteristic polynomial of the Frobenius Φ acting on the crystalline module of almost of the classical specializations y of \mathcal{M}' has no root equal to $U_0(y)$.

Let $Z = V(\mathcal{I}) \subset \mathcal{U}$ be the Zariski closed set defined by the ideal \mathcal{I} generated by the 4-th Fitting ideal Fitt_4 of the A -module \mathcal{M} and by the 3-rd Fitting ideal Fitt_3 of the A -module \mathcal{M}' , then any point y lies in $V(\text{Fitt}_4)$ (resp. $V(\text{Fitt}_3)$) if and only if $\dim_{k(y)}(\mathcal{M}(y)) \geq 5$ (resp. $\dim_{k(y)}(\mathcal{M}'(y)) \geq 4$), where $\mathcal{M}(y)$ (resp. $\mathcal{M}'(y)$) is the fiber of \mathcal{M} (resp. \mathcal{M}') at y and $k(y)$ is the residue field at y .

Thus $\mathcal{U} - V(\mathcal{I})$ is the biggest admissible open subset of \mathcal{U} on which \mathcal{M} (resp. \mathcal{M}') can be locally generated (on stalks) by 4 elements (resp. 3 elements). Moreover, since the coherent \mathcal{M} (resp. \mathcal{M}') is generically of rank 4 (resp. 3) and torsion-free then one can deduce that the coherent sheaf \mathcal{M} (resp. \mathcal{M}') is locally free of rank 4 (resp. 3) on the admissible open $\mathcal{U} - Z = \mathcal{U}'$ (\mathcal{U}' does not necessarily contain π_α). Thus, the direct summand \mathcal{M}' of \mathcal{M} is also locally free of rank 3 on \mathcal{U}' . Hence one can deduce that the Hodge–Tate weights of the specialization of \mathcal{M}' at classical points of \mathcal{U}' of weight $l_1 > l_2 + 1$ and having crystalline representation (they form a very Zariski dense set) are $l_2 - 2, l_1 - 1, l_1 + l_2 - 3$; and then $l_2 - 2$ is the smallest Hodge–Tate weight (see Corollary 2.7).

In addition, if $\mathcal{M}'(y)$ (resp. $\mathcal{M}(y)$) denotes the specialization of \mathcal{M}' (resp. \mathcal{M}) at a very classical point $y \in \mathcal{U}'$. We can enlarging Z if it is necessary to have that for any $y \in \mathcal{U}'$, $\mathcal{M}(y)^{\text{ss}} = \mathcal{M}(y)$. Now, if $y \in \mathcal{U}'$ is a classical point of weight (l_1, l_2) and ρ_y is a crystalline representation at p , then the

eigenvalues of the Frobenius Φ acting on $\mathcal{D}_{\text{crys}}(\mathcal{M}'(y))$ are $\lambda_y(U_1U_0^{-1})p^{l_2-2}$, $\lambda_y(U_0U_1^{-1})p^{l_1-1}$ and $\lambda_y(U_0^{-1})p^{l_2+l_1-3}$, where $\lambda_y : \text{Spm } L_y \rightarrow \mathcal{E}_\Delta$ is the morphism corresponding to y . When $y = x$, we have $\lambda_x(U_1U_0^{-1}) = p$ ($\underline{k} = (k, k)$) is the weight of π_α).

The exact sequence (4.2) (i.e. ϵ_p^{1-k} occurs with multiplicity one in $\mathcal{M}'/\mathfrak{m}\mathcal{M}'$), the regularity assumption (i.e. $\alpha \neq 1$) of ϱ at p when $k = 2$, and the fact that $\mathcal{M}' \otimes_A \mathcal{T} = M'$ (since $\mathcal{M}^{I_p} \otimes_A \mathcal{T} = M^{I_p}$), yield that

$$(4.3) \quad \dim \mathcal{D}_{\text{crys}}^+(\mathcal{M}'(\pi_\alpha)^{\text{ss}})^{\Phi=p^{k-1}} = \dim \mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(\epsilon_p^{1-k}) = 1.$$

Hence, one has (after a twist by ϵ^{k-2})

$$(4.4) \quad \dim \mathcal{D}_{\text{crys}}^+(\mathcal{M}'(\pi_\alpha)^{\text{ss}}(k-2))^{\Phi=p} = 1.$$

Since the set Σ of classical points of \mathcal{E}_Δ of cohomological weights and old at p (i.e having a crystalline representation) of \mathcal{E}_Δ are very Zariski dense (see Corollary B.5), it follows from Lemma A.7 that $\Sigma \cap \mathcal{U} - (\Sigma \cap Z)$ is Zariski dense in \mathcal{U} , and hence we obtain a refined family

$$(G_{\mathbb{Q}_p} \rightarrow \text{Aut}_{\mathcal{U}}(\mathcal{M}'), \Sigma \cap \mathcal{U} - (\Sigma \cap Z), \{\kappa_i\}, U_0/U_1 \in \mathcal{O}(\mathcal{E}_\Delta)^\times)$$

of generic rank equal to 3 over K . Note also that condition $(*)$ of [7, Definition 4.2.7] is satisfied since we have a torsion free morphism $\kappa : \mathcal{E}_\Delta \rightarrow \mathcal{W}$; the condition (v) of [7, Definition 4.2.7] is satisfied by Lemma A.6, Lemma B.3 and Corollary B.5 (so $\Sigma \cap \mathcal{U} - (\Sigma \cap Z)$ accumulate to π_α).

Since $\mathcal{M}' \otimes_A \mathcal{T} = M'$, it follows from [7, Theorem 3.4.1] that

$$\dim \mathcal{D}_{\text{crys}}^+(M'/\mathfrak{m}M'(k-2))^{\Phi=p} = 1.$$

Then

$$\dim \mathcal{D}_{\text{crys}}^+(M'/\mathfrak{m}M')^{\Phi=p^{k-1}} = 1.$$

Finally, by [7, Theorem 1.5.6] any $S/\mathfrak{m}S$ -extension V of ϵ_p^{1-k} by ϵ_p^{2-k} (i.e occurring in the image of the morphism (4.1)) is a quotient of $M/\mathfrak{m}M \oplus \epsilon_p^{2-k}$ by an S -submodule \mathcal{Q} (see **(iii?)** of (4.5)).

However, by the regularity assumption at p the non-trivial unramified character $\psi|_{G_{\mathbb{Q}_p}}$ does not occur in $V|_{G_{\mathbb{Q}_p}} \in \text{Ext}_{G_{\mathbb{Q}_p}}^1(\epsilon_p^{1-k}, \epsilon_p^{2-k})$, which implies that $V|_{G_{\mathbb{Q}_p}}$ is a quotient of $M'/\mathfrak{m}M' \oplus \epsilon_p^{2-k}$. Thus we obtain a surjection of $G_{\mathbb{Q}_p}$ -modules

$$(4.5) \quad M'/\mathfrak{m}M' \oplus \epsilon_p^{2-k} \xrightarrow{\pi'} V|_{G_{\mathbb{Q}_p}},$$

with kernel isomorphic to a quotient of the $G_{\mathbb{Q}_p}$ -module \mathcal{Q} .

Since the semi simplification of $M_3/\mathfrak{m}M_3$ is isomorphic to the representation

$$\rho_f^{n_1} \oplus (\epsilon_p^{2-k})^{n_2} \oplus \epsilon_p^{1-k}$$

by (4.2), the regularity assumption at p on ϱ when $k = 2$ (i.e. $\alpha \neq 1$), and the fact that the S_p -simple subquotients of \mathcal{Q} do not equal ϵ_p^{1-k} by Lemma 4.1 (they occur in $\{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{3-2k}\}$), one has

$$\mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(\ker(\pi')) = 0.$$

Thus the surjective morphism (4.5) yields the following injection

$$\mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(M'/\mathfrak{m}M') \hookrightarrow \mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(V),$$

and implies that $\mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(V) \neq 0$ (since $\dim \mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(M'/\mathfrak{m}M') = 1$).

On the other hand, by applying the left exact functor $\mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(\cdot)$ to the exact sequence

$$0 \rightarrow \epsilon_p^{2-k} \rightarrow V \rightarrow \epsilon_p^{1-k} \rightarrow 0,$$

and using the fact that $\mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(\epsilon_p^{2-k}) = 0$ and $\dim \mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(\epsilon_p^{1-k}) = 1$, we obtain that $\dim \mathcal{D}_{\text{crys}}^{\Phi=p^{k-1}}(V) = 1$ (since it is non-zero by the above discussion). Hence the characteristic polynomial of Φ has two roots $\{p^{k-2}, p^{k-1}\}$ yielding that $\dim \mathcal{D}_{\text{crys}}(V) = 2$ and that V is crystalline, so $V \otimes \epsilon_p^{k-2}$ is also crystalline at p .

It remains to proof that the image of the map

$$\text{Hom}(\mathcal{T}_{1,3}/\mathcal{T}'_{1,3}, L) \hookrightarrow H_f^1(G_{\mathbb{Q}}^{Np}, \epsilon_p)$$

consists of extensions which are unramified outside p . Let ℓ denote a prime number dividing N (so prime to p), note that any $G_{\mathbb{Q}_\ell}$ -extension of ϵ_p^{-1} by $\mathbb{1}$ is trivial or its restriction to the inertia has the following form

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

and hence the monodromy operator of the Weil–Deligne representation attached to its 2-dimensional $G_{\mathbb{Q}_\ell}$ -representation is of rank 1 (i.e a Steinberg type).

We know that the rank over L of the monodromy operator attached to the Weil–Deligne representation corresponding to $(\rho_f)_{|G_{\mathbb{Q}_\ell}}$ is one (since we assumed that ρ_f is a twisted Steinberg at every prime $\ell \mid N$).

Recall that the $G_{\mathbb{Q}}$ -coherent sheaf \mathcal{M} is locally free of rank 4 on the admissible open $\mathcal{U} - Z = \mathcal{U}'$ and it admits a Weil–Deligne representation $(r_{\mathcal{U}}, N_{\mathcal{U}})$ by [7, Lemma 7.8.14] at ℓ (for which $N_{\mathcal{U}} \in \text{End}_A(\mathcal{M})$). Since the rank of the monodromy operator of the Weil–Deligne representation attached to the specializations of $(r_{\mathcal{U}}, N_{\mathcal{U}})$ at classical points of non-endoscopic, non-CAP points \mathcal{U}' is at most 1 by Theorem 2.10, [7, Proposition 7.8.19(ii)] implies that the generic rank over K of the monodromy operator of the Weil–Deligne representation attached $(r_{\mathcal{U}}, N_{\mathcal{U}})$ is

also 1 (since it is non-trivial at π_α). Therefore, the generic rank of the monodromy $N_K = N_{\mathcal{U}} \otimes K$ operator of the Weil–Deligne representation attached to $(\rho_K)|_{G_{\mathbb{Q}_\ell}}$ is one.

Let S_ℓ be the image of $\mathcal{T}[G_{\mathbb{Q}_\ell}]$ inside S . Thanks to Proposition 2.3, one can apply [7, Lemma 8.2.11] ⁽⁶⁾ to $\mathcal{P} = \{\epsilon_p^{1-k}, \epsilon_p^{2-k}\}$, and we obtain that there exists idempotents $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3)$ of S lifting the idempotents attached respectively to $\epsilon_p^{2-k}, \epsilon_p^{1-k}, \rho_f$ and such that $\tilde{e} = \tilde{e}_1 + \tilde{e}_2$ is in the center of S_ℓ (see [7, Lemma 8.2.12]), and hence S_ℓ is block diagonal of type $(2, 2)$ in S . Thus,

$$S_\ell/\mathfrak{m}S_\ell = \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & \rho_f \end{pmatrix}, \quad \text{and} \quad S_\ell = \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & M_{2,2}(\mathcal{T}) \end{pmatrix}.$$

By [7, Lemma 7.8.14] one can see N_K as element of S_ℓ . By (iii) it is enough to prove that $\tilde{e}N_K \in \tilde{e}S_\ell$ is trivial for showing that the image of $\text{Hom}(\mathcal{T}_{1,3}/\mathcal{T}'_{1,3}, L) \hookrightarrow H^1(G_{\mathbb{Q}}^{Np}, \epsilon_p)$ gives rise to classes unramified at ℓ .

As an element of S_ℓ we know that $N_K = \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & * \end{pmatrix}$ and it is of rank 1 as discussed before. By [7, Proposition 7.8.8] applied to $(1 - \tilde{e})S_\ell(1 - \tilde{e})$ we further know that the rank of $(1 - \tilde{e})N_K$ is one, using that ρ_f is a twisted Steinberg at ℓ (the rank of the monodromy operator of $WD_\ell(\rho_f)$ is one) and the surjection $(1 - \tilde{e}).S/\mathfrak{m}S.(1 - \tilde{e}) \twoheadrightarrow \rho_f$. Hence, $\tilde{e}_i N_K = 0$ for $i \in \{1, 2\}$, which yields that $\tilde{e}N_K = 0$. □

The proof of Theorem 4.3 yields the following corollary.

COROLLARY 4.4. — *There exists a representation $\rho_{\mathcal{M}'} : G_{\mathbb{Q}_p} \rightarrow \text{Aut}_{\mathcal{U}}(\mathcal{M}')$, where \mathcal{M}' is a torsion-free coherent sheaf on an admissible open affinoid $\mathcal{U} = \text{Spm } A \subset \mathcal{E}_\Delta$ containing π_α and $\rho_{\mathcal{M}'}$ is of generic rank 3 over the total ring of fractions K of \mathcal{U} such that:*

- (1) *There exists a very Zariski dense set $\Sigma' \subset \mathcal{U}$ such that the specialization of the representation $\rho_{\mathcal{M}'}$ at any point z of Σ' gives rise to a crystalline $G_{\mathbb{Q}_p}$ -representation ρ'_z of dimension 3, with Hodge–Tate–Sen weights given by $(\kappa_2 - 2, \kappa_1 - 1, \kappa_1 + \kappa_2 - 3)$.*
- (2) *The smallest Hodge–Tate weight of ρ'_z is $\kappa_2(z) - 2$ and $U_1/U_0 \in \mathcal{O}(\mathcal{E}_\Delta)^\times$ interpolates the crystalline period of the smallest Hodge–Tate weight. In other words, one has*

$$\dim \mathcal{D}_{\text{crys}}(\rho'_z)^{\Phi=U_1/U_0(z)p^{\kappa_2(z)-2}} = 1.$$

⁽⁶⁾The assumption that $\pi_{f,\ell} \simeq \text{St} \otimes \xi$ is crucial to prove the vanishing of $H^1(G_{\mathbb{Q}_\ell}, \rho_f(k-2))$.

(3) Let $M' := \mathcal{M}' \otimes_A \mathcal{T}$, then for any cofinite ideal \mathcal{J} of \mathcal{T} one has that

$$l(\mathcal{D}_{\text{crys}}^+(M'/\mathcal{J}M' \otimes (\epsilon_p^{\kappa_2-2}))^{\Phi=U_1/U_0}) = l(\mathcal{T}/\mathcal{J}).$$

(4) The Sen operator of $\mathcal{D}_{\text{sen}}(M'/\mathcal{J}M')$ is annihilated by the polynomial

$$(T - (\kappa_2 - 2))(T - (\kappa_1 - 1))(T - (\kappa_1 + \kappa_2 - 3)).$$

Proof. — (1) and (2) follows directly from the proof of Theorem 4.3 and [7, Theorem 1.5.6]. Thus, it remains to show (3) and (4), which follows immediately from similar arguments to those already used to prove of Theorem 4.3, [7, Theorem 3.4.1] and [7, Lemma 4.3.3(i)]. \square

5. Crystallinity of the S -extensions occurring in

$$H^1(\mathbb{Q}, \rho_f(k - 2))$$

By (3.1) we have a natural injection

$$(5.1) \quad \text{Hom}(\mathcal{T}_{3,2}/\mathcal{T}'_{3,2}, L) \hookrightarrow \text{Ext}_{G_{\mathbb{Q}}^{Np}}^1(\rho_f, \epsilon_p^{1-k}) \simeq H^1(G_{\mathbb{Q}}^{Np}, \rho_f(k - 2)).$$

Now we have to determine the exact image of the injective morphism (5.1). As in Section 4 we apply the results of [7, Theorem 1.5.6] and [7, Lemma 4.3.9] for the left ideal $M_2 = S.E_2$ of S given by the second column of the GMA matrix S :

(i) There exists an exact sequence of S -left modules

$$(5.2) \quad 0 \rightarrow E' \rightarrow M_2/\mathfrak{m}M_2 \rightarrow \rho_f \rightarrow 0$$

(ii) Any simple S -subquotients of E' is not isomorphic to ρ_f and they occur in the set $\{\epsilon_p^{1-k}, \epsilon_p^{2-k}\}$.

(iii) The image of the morphism (5.1) consists of extensions occurring as quotient of the $S/\mathfrak{m}S$ -module $M_2/\mathfrak{m}M_2 \oplus \epsilon_p^{1-k}$ by an S -submodule \mathcal{Q}' whose S -simple subquotients occur in the set $\{\epsilon_p^{1-k}, \epsilon_p^{2-k}\}$.

Since ρ_K is absolutely irreducible and M_2 is a finite type torsion free \mathcal{T} -module we again have $M_2.K = K^4$.

THEOREM 5.1. — Assume **(Reg)**. Let $\mathcal{T}'_{3,2}$ be the \mathcal{T} -module $\mathcal{T}_{3,1}\mathcal{T}_{1,2} \subset \mathcal{T}_{3,2}$, then:

(1) There exists an injective homomorphism of L -modules

$$(5.3) \quad \text{Hom}(\mathcal{T}_{3,2}/\mathcal{T}'_{3,2}, L) \hookrightarrow \ker \left(H^1(\mathbb{Q}, \rho_f(k - 2)) \rightarrow H^1(\mathbb{Q}_p, \rho_f/\rho_f^{I_p}(k - 2)) \bigoplus_{\ell \neq p} H^1(I_\ell, \rho_f(k - 2)) \right).$$

(2) Assume that $k \geq 3$, then

$$(5.4) \quad \text{Hom}(\mathcal{T}_{3,2}/\mathcal{T}'_{3,2}, L) \hookrightarrow H^1_{f, \text{unr}}(\mathbb{Q}, \rho_f(k-2)).$$

Proof.

(1). — By (5.2) we have a surjective morphism of S -modules $\pi : M_2/\mathfrak{m}M_2 \twoheadrightarrow \rho_f$ whose kernel does not contain ρ_f and whose semi-simplification contains only $G_{\mathbb{Q}}$ -representations lying in the set $\{\epsilon_p^{1-k}, \epsilon_p^{2-k}\}$. Moreover, our assumptions yield that the irreducible constituents of the semi-simplification of $\varrho|_{G_{\mathbb{Q}_p}}$ are without multiplicity, hence $M_2^{I_p} := \{x \in M_2, \forall g \in I_p, g.x = x \text{ and } \text{Frob}_p.x = U_0.x\}$ is not contained in $\mathfrak{m}M_2$. Let $V \in \text{Ext}^1_{G_{\mathbb{Q}}}(\rho_f, \epsilon_p^{1-k}) = H^1(G_{\mathbb{Q}}^{N_p}, \rho_f(k-2))$ be in the image of (5.4). By (iii) above we have an exact sequence of left S -modules

$$0 \rightarrow \mathcal{Q}' \rightarrow M_2/\mathfrak{m}M_2 \oplus \epsilon_p^{1-k} \rightarrow V \rightarrow 0.$$

Similar to Lemma 4.1 we can show that \mathcal{Q}' has no $L[G_{\mathbb{Q}_p}]$ -simple subquotients equal to ψ or $\psi^{-1}\epsilon_p^{2k-3}$. This shows that the image of $M_2^{I_p}$ in V is non-zero. It follows that

$$V^{I_p} \neq 0.$$

Moreover, since Frob_p acts on $M_2^{I_p}$ by U_0 , the action of Frob_p on V^{I_p} is given by ψ . If the realization of V is given by $\tilde{\rho} = \begin{pmatrix} \epsilon_p^{1-k} & * \\ 0 & \rho_f \end{pmatrix}$ then the restriction of $\tilde{\rho}$ to $G_{\mathbb{Q}_p}$ is given by $\begin{pmatrix} \epsilon_p^{1-k} & 0 & c \\ 0 & \psi & * \\ 0 & 0 & \psi^{-1}\epsilon_p^{3-2k} \end{pmatrix}$ since $V^{I_p} \neq 0$. Finally, it remains to show that the extensions V are unramified at every prime $\ell \mid N$, and this fact follows immediately from Proposition [41, Lemme 4.1.3].

(2). — The fact that $k \geq 3$ implies that $3 - 2k < 1 - k$ and hence $\tilde{\rho}$ is ordinary in the sense of Definition 2.1 and then semi-stable (hence de Rham) at p by Theorem 2.2. Therefore the extension V gives a cohomology class in $H^1_g(G_{\mathbb{Q}}^{N_p}, \rho_f(k-2))$ which is isomorphic to $H^1_f(G_{\mathbb{Q}}^{N_p}, \rho_f(k-2))$ by [41, Lemme 4.1.3]. \square

Remark 5.2. — For $k = 2$ ordinarity/crystallinity of the extension would require us to prove additionally that $\tilde{\rho}/\tilde{\rho}^{I_p} \cong \begin{pmatrix} \epsilon^{-1} & \\ 0 & \psi^{-1}\epsilon^{-1} \end{pmatrix}$ is a trivial extension. This would follow, e.g. if one could prove that the generator of $H^1(G_{\mathbb{Q}}^{N_p}, \rho_f)$ (which is conjectured to be 1-dimensional by Jannsen) has no line fixed by inertia at p . See Section 5.1 below for an alternative approach in this case.

Similarly to Corollary 3.6, [7, Proposition 1.8.6] yields immediately the following corollary.

COROLLARY 5.3. — Assume **(Reg)**. Then the image of the natural injective morphism of L -modules

$$\text{Hom}(\mathcal{T}_{2,1}/\mathcal{T}_{2,3}\mathcal{T}_{3,1}, L) \hookrightarrow H^1(\mathbb{Q}, \rho_f(k - 2))$$

is isomorphic to the image of (5.1) (which is described in Theorem 5.1).

5.1. On the vanishing of the Greenberg’s Selmer group attached to f_α

Assume in this subsection that $k = 2$ and let

$$\text{Sel}_{\mathbb{Q}, f_\alpha} = \ker \left(H^1(\mathbb{Q}_{Np}, \rho_f) \rightarrow H^1(\mathbb{Q}_p, \rho_f/\rho_f^{I_p}) \bigoplus_{\ell \nmid p} H^1(I_\ell, \rho_f) \right)$$

be the Greenberg-type Selmer group we used in Theorem 5.1(1) attached to the ordinary elliptic cuspform f_α . In the literature, Greenberg’s Selmer group is often defined using the representation $\rho_f^\vee(-1)$ (arithmetic Frobenius convention). The p -adic representation ρ_f^\vee corresponds to the Tate module $T_p(A_f)$ of the abelian variety A_f , and ρ_f is the Galois representation obtained from the p -adic étale cohomology of A_f . We remark also that for $k = 2$ the condition at p is weaker than the “usual” condition for the ordinary representation ρ_f (which would require the class to be split at p). Our condition of having an I_p -fixed quotient for the extension $\left(\begin{smallmatrix} \rho_f & * \\ 0 & 1 \end{smallmatrix}\right)$ (or dually an I_p -fixed line for $\left(\begin{smallmatrix} \epsilon_p^{-1} & * \\ 0 & \rho_f \end{smallmatrix}\right)$) is the one that would normally be required for $\rho_f(1) \cong \rho_f^\vee$.

Note that ρ_f is not critical in the sense of Deligne. We use Iwasawa theory for the cyclotomic \mathbb{Z}_p -extension to bound $\text{Sel}_{\mathbb{Q}, f_\alpha}$: It follows from Kato [23] that the Pontryagin dual of the Selmer group $\text{Sel}_{\mathbb{Q}_\infty, f_\alpha}$ is a torsion Λ -module with characteristic ideal $g(T) \in \Lambda$. Furthermore, according to the Iwasawa main conjecture (Kato’s bound, see e.g. [42, Theorem 3.25]), $g(T) \mid L_p(f, \omega^{-1}, \cdot)$. Hence $\dim \text{Sel}_{\mathbb{Q}, f_\alpha} = 0$ when $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$ (see [9, Proposition 2.10] and [9, Theorem 2.11] for more details). Moreover, it follows from the control theorem for the Λ -adic Greenberg’s Selmer group $\text{Sel}_{\mathbb{Q}_\infty, f_\alpha}$ (see also [35]) that $g(T = p) \neq 0$ is a necessary condition for the vanishing of $\text{Sel}_{\mathbb{Q}, f_\alpha}$.

6. Schematic reducibility locus of the pseudo-character $\text{Ps}_{\mathcal{T}}$ on $\text{Spec } \mathcal{T}$ and applications to the Bloch–Kato conjecture

Recall that we view S as the generalized matrix attached to the pseudo-character

$$\text{Ps}_{\mathcal{T}} : G_{\mathbb{Q}} \rightarrow \mathcal{T}$$

with respect to a set of idempotents compatible with the anti-involution τ and have

$$S = \begin{pmatrix} \mathcal{T} & M_{1,2}(\mathcal{T}_{1,2}) & \mathcal{T}_{1,3} \\ M_{2,1}(\mathcal{T}_{2,1}) & M_2(\mathcal{T}) & M_{2,1}(\mathcal{T}_{2,3}) \\ \mathcal{T}_{3,1} & M_{1,2}(\mathcal{T}_{3,2}) & \mathcal{T} \end{pmatrix},$$

where $\mathcal{T}_{i,j}$ are fractional ideals of K that satisfy $\mathcal{T}_{i,j}\mathcal{T}_{j,k} \subset \mathcal{T}_{i,k}$ and $\mathcal{T}_{i,j}\mathcal{T}_{j,i} \subset \mathfrak{m}$.

In this section we will compute the total reducibility ideal $\mathcal{I}^{\text{tot}} \subset \mathcal{T}$ (see Definition 3.3). By Proposition 3.4 it is given by

$$(6.1) \quad \mathcal{I}^{\text{tot}} = \mathcal{T}_{3,1}\mathcal{T}_{1,3} + \mathcal{T}_{2,3}\mathcal{T}_{3,2} + \mathcal{T}_{1,2}\mathcal{T}_{2,1}.$$

The following lemma follows directly from the anti-involution $\tau : S \rightarrow S$ and the fact that $\text{Ps}_{\mathcal{T}}$ is invariant under the action of τ .

LEMMA 6.1. — *One always has:*

$$\mathcal{T}_{2,3}\mathcal{T}_{3,2} = \mathcal{T}_{1,2}\mathcal{T}_{2,1}.$$

Proof. — This is proved exactly as in Lemma [7, 8.2.16] using the anti-involution τ . □

Hence, the above lemma implies that

$$(6.2) \quad \mathcal{I}^{\text{tot}} = \mathcal{T}_{3,1}\mathcal{T}_{1,3} + \mathcal{T}_{1,2}\mathcal{T}_{2,1}.$$

LEMMA 6.2. — *Assume (Reg) and (St). Then one has*

$$\mathcal{I}^{\text{tot}} = \mathcal{T}_{1,2}\mathcal{T}_{2,1} = \mathcal{T}_{2,3}\mathcal{T}_{3,2}.$$

Proof. — We first show that $\mathcal{T}_{1,3} = \mathcal{T}'_{1,3} = \mathcal{T}_{1,2}\mathcal{T}_{2,3}$. By Theorem 4.3 we have an injective map

$$\text{Hom}(\mathcal{T}_{1,3}/\mathcal{T}'_{1,3}, L) \hookrightarrow H^1_{f,\text{unr}}(\mathbb{Q}, \epsilon_p).$$

Note that the Kummer map provides an isomorphism

$$H^1_{f,\text{unr}}(\mathbb{Q}, \epsilon_p) \simeq \mathbb{Z}^{\times} \otimes L.$$

Hence $H_{f, \text{unr}}^1(\mathbb{Q}, \epsilon_p)$ is trivial, and then $\mathcal{T}_{1,3}/\mathcal{T}'_{1,3} = 0$ by Nakayama's lemma ($\mathcal{T}_{1,3}$ is of finite type over \mathcal{T} since S is). Thus, we have

$$(6.3) \quad \mathcal{T}_{1,3} = \mathcal{T}_{1,2}\mathcal{T}_{2,3}.$$

It is easy to see that

$$(6.4) \quad \begin{aligned} \mathcal{I}^{\text{tot}} &= \mathcal{T}_{3,1}\mathcal{T}_{1,3} + \mathcal{T}_{1,2}\mathcal{T}_{2,1} \\ &= \mathcal{T}_{1,2}\mathcal{T}_{2,3}\mathcal{T}_{3,1} + \mathcal{T}_{1,2}\mathcal{T}_{2,1} \\ &= \mathcal{T}_{1,2}\mathcal{T}_{2,1}, \end{aligned}$$

since $\mathcal{T}_{2,3}\mathcal{T}_{3,1} \subset \mathcal{T}_{2,1}$ □

COROLLARY 6.3. — *One has*

$$\mathcal{T}'_{1,2} = \mathcal{I}^{\text{tot}}.\mathcal{T}_{1,2}.$$

Proof. — Since $\mathcal{T}_{1,3} = \mathcal{T}_{1,2}\mathcal{T}_{2,3}$ by relation (6.3) we get $\mathcal{T}'_{1,2} = \mathcal{T}_{1,3}\mathcal{T}_{3,2} = \mathcal{T}_{1,2}\mathcal{T}_{2,3}\mathcal{T}_{3,2}$. On the other hand, we have by Lemma 6.1 that $\mathcal{T}_{2,3}\mathcal{T}_{3,2} = \mathcal{T}_{1,2}\mathcal{T}_{2,1}$, and we have also by Lemma 6.2 $\mathcal{I}^{\text{tot}} = \mathcal{T}_{1,2}\mathcal{T}_{2,1} = \mathcal{T}_{2,3}\mathcal{T}_{3,2}$. Thus $\mathcal{T}'_{1,2} = \mathcal{T}_{1,3}\mathcal{T}_{3,2} = \mathcal{T}_{1,2}\mathcal{T}_{2,3}\mathcal{T}_{3,2} = \mathcal{I}^{\text{tot}}\mathcal{T}_{1,2}$. □

6.1. Application to Bloch–Kato conjecture

Since we have assumed that the sign ϵ_f of $L(f, s)$ is -1 , the functional equation

$$L(f, s) = -L(f, 2k - 2 - s)$$

yields that $L(f, s)$ vanishes at the central value $k - 1$. The Selmer group $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k - 1))$ classifies the extensions with everywhere good reduction and one can think of the Bloch–Kato conjecture as a generalization of the Birch and Swinnerton–Dyer conjecture for the motive M_f corresponding to f of weight $2k - 2 \geq 2$. One has the following application related to the Bloch–Kato conjecture:

COROLLARY 6.4. — *Assume that $k \geq 2$, **(Reg)** and **(St)**. Then there exists an injection*

$$(6.5) \quad \text{Hom}(\mathcal{T}_{1,2}/\mathfrak{m}.\mathcal{T}_{1,2}, L) \hookrightarrow H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k - 1)),$$

and $\dim \mathcal{T}_{1,2}/\mathfrak{m}.\mathcal{T}_{1,2} \geq 1$.

Proof. — The following injection follow from Theorem 3.5 and Corollary 6.3:

$$(6.6) \quad \text{Hom}(\mathcal{T}_{1,2}/\mathfrak{m}.\mathcal{T}_{1,2}, L) \simeq \text{Hom}(\mathcal{T}_{1,2}/\mathcal{I}^{\text{tot}}.\mathcal{T}_{1,2}, L) \hookrightarrow H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k - 1))$$

Moreover, $\mathcal{T}_{1,2}/\mathfrak{m}\mathcal{T}_{1,2} \neq \{0\}$ since $\rho_K : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_4(K)$ is absolutely irreducible (so $\mathcal{I}^{\mathrm{tot}} = \mathcal{T}_{1,2}\mathcal{T}_{2,1} \neq (0)$). □

PROPOSITION 6.5.

$$\dim H^1(G_{\mathbb{Q}}^{Np}, \epsilon_p^{-1}) = 1.$$

Proof. — This follows from [28, Proposition 2.2]. □

PROPOSITION 6.6. — Assume that $k \geq 3$ and **(Reg)**. The image of the natural injection

$$(6.7) \quad \mathrm{Hom}(\mathcal{T}_{1,2}/\mathcal{T}'_{1,2}, L) \hookrightarrow H^1_{f, \mathrm{unr}}(\mathbb{Q}, \rho_f(k-1)),$$

is non-trivial (i.e. $\mathcal{T}_{1,2}/\mathcal{T}'_{1,2} \neq \{0\}$).

Proof. — Since $\dim H^1_{f, \mathrm{unr}}(\mathbb{Q}, \rho_f(k-2)) = 0$ (by [23]), Theorem 5.1 and Corollary 5.3 yield that $\mathrm{Hom}(\mathcal{T}_{3,2}/\mathcal{T}'_{3,2}, L) = \{0\}$, and hence Nakayama’s lemma implies that $\mathcal{T}_{3,2} = \mathcal{T}'_{3,2} = \mathcal{T}_{3,1}\mathcal{T}_{1,2}$. We will proceed by contradiction. Assume that $\mathcal{T}_{1,2}/\mathcal{T}'_{1,2} = \{0\}$ and so $\mathcal{T}_{1,2} = \mathcal{T}'_{1,2} = \mathcal{T}_{1,3}\mathcal{T}_{3,2}$. Thus, $\mathcal{T}_{1,2} = \mathcal{T}_{1,3}\mathcal{T}_{3,1}\mathcal{T}_{1,2}$. Moreover, $\mathcal{T}_{1,3}\mathcal{T}_{3,1} \subset \mathfrak{m} \subset \mathcal{T}$ by [7, Theorem 1.4.4] and hence $\mathcal{T}_{1,2} = \mathfrak{m}\mathcal{T}_{1,2}$. Therefore, Nakayama’s lemma yields that $\mathcal{T}_{1,2} = 0$. The involution introduced in Section 3.1 implies by [7, Lemma 1.8.5] that $\mathcal{T}_{1,2} = \mathcal{T}_{2,3}$ and $\mathcal{T}_{2,1} = \mathcal{T}_{3,2}$, and so also $\mathcal{T}_{2,3} = 0$. Furthermore, Theorem 5.1 yields that $\mathcal{T}_{2,1} = \mathcal{T}'_{2,1} = \mathcal{T}_{2,3}\mathcal{T}_{3,1}$. Finally, this yields that $\mathcal{T}_{3,2} = \mathcal{T}_{2,1} = \mathcal{T}_{2,3}\mathcal{T}_{3,1} = 0$ and also $\mathcal{T}_{1,2} = \mathcal{T}_{2,3} = 0$, and it follows that the generic representation $\rho_K : G_{\mathbb{Q}} \rightarrow S^{\times} \subset \mathrm{GL}_4(K)$ attached to $K = Q(\mathcal{T})$ is reducible, which contradicts Theorem 2.11. □

Assume now that $\bar{\rho}_f$ is absolutely irreducible. Let \mathbb{I} be the finite flat integral extension of the Iwasawa algebra $\mathbb{Z}_p[[T]]$ generated by the coefficients of a Hida family \mathcal{F} specializing to f_{α} (\mathcal{F} is unique up to Galois conjugacy) and let $\rho_{\mathcal{F}} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\mathbb{I})$ be the p -adic Galois representation attached to \mathcal{F} . Let $\chi_{\mathrm{univ}} : G_{\mathbb{Q}} \rightarrow \mathbb{Z}_p[[T]]^{\times}$ be the universal character given by the composition of the p -adic cyclotomic character $\epsilon_p : G_{\mathbb{Q}} \rightarrow 1 + p^{\nu}\mathbb{Z}_p$ with the tautological character $1 + p^{\nu}\mathbb{Z}_p \rightarrow \mathbb{Z}_p[[1 + p^{\nu}\mathbb{Z}_p]]^{\times} \simeq \mathbb{Z}_p[[T]]^{\times}$, where $\nu = 1$ if $p \geq 3$ and $\nu = 2$ if $p = 2$. It follows from the work of Nekovar [34, Proposition 4.2.3] that the \mathbb{I} -adic Selmer group $H^1_{f, \mathrm{unr}}(\mathbb{Q}, \rho_{\mathcal{F}} \otimes \chi_{\mathrm{univ}}^{-1/2})$ is of finite type over the Iwasawa algebra $\mathbb{Z}_p[[T]]$, and so over \mathbb{I} since \mathbb{I} is finite flat over $\mathbb{Z}_p[[T]]$.

COROLLARY 6.7. — Assume that $\bar{\rho}_f$ is absolutely irreducible, then the generic rank of the \mathbb{I} -adic Selmer group $H^1_{f, \mathrm{unr}}(\mathbb{Q}, \rho_{\mathcal{F}} \otimes \chi_{\mathrm{univ}}^{-1/2})$ is at least one (i.e. $H^1_{f, \mathrm{unr}}(\mathbb{Q}, \rho_{\mathcal{F}} \otimes \chi_{\mathrm{univ}}^{-1/2})$ has a non-torsion class over \mathbb{I}).

Proof. — It follows from Corollary 6.6 that the \mathbb{I} -adic Selmer group $H_{f,\text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}} \otimes \chi_{\text{univ}}^{-1/2})$ specializes at infinitely many classical points of $\text{Spm} \mathbb{I}[1/p]$ to a non-trivial Selmer group. Hence, the generic rank of $H_{f,\text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}} \otimes \chi_{\text{univ}}^{-1/2})$ over \mathbb{I} is non-zero. \square

6.2. Bounding the number of generators of \mathcal{I}^{tot}

THEOREM 6.8. — Assume **(Reg)**, **(St)** and $\dim H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$.

- (1) There exists idempotents $\{e'_1, e'_2, e'_3\}$ of S lifting the idempotents of ϱ attached to $\{\epsilon_p^{2-k}, \rho_f, \epsilon_p^{1-k}\}$ such that S has the following form

$$S = \begin{pmatrix} \mathcal{T} & M_{1,2}(\mathcal{T}) & \mathcal{T} \\ M_{2,1}(\mathcal{I}^{\text{tot}}) & M_2(\mathcal{T}) & M_{2,1}(\mathcal{T}) \\ \mathcal{T}_{3,1} & M_{1,2}(\mathcal{I}^{\text{tot}}) & \mathcal{T} \end{pmatrix},$$

where $\mathcal{T}_{3,1} = \mathcal{J} \subset \mathcal{I}^{\text{tot}}$ is an ideal.

- (2) Assume $k \geq 3$. Then $\mathcal{I}^{\text{tot}} = \mathcal{J} = \mathcal{T}_{3,1}$ and $\mathcal{I}^{\text{tot}} = \mathcal{T}.g + (\mathcal{I}^{\text{tot}})^2$ for an element g in \mathcal{I}^{tot} , which yields that the reducibility ideal \mathcal{I}^{tot} is principal and generated by g .
- (3) Assume that $k = 2$ and $\dim \text{Sel}_{\mathbb{Q}, f_\alpha} = 0$, then $\mathcal{I}^{\text{tot}} = \mathcal{J} = \mathcal{T}_{3,1}$ and $\mathcal{I}^{\text{tot}} = \mathcal{T}.g + (\mathcal{I}^{\text{tot}})^2$ for an element g in \mathcal{I}^{tot} , and the reducibility ideal \mathcal{I}^{tot} is principal and generated by g .

Remark 6.9. — Using results about Λ -adic Selmer groups we exhibit many examples where the Selmer group $H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ is 1-dimensional (see Appendix C).

Proof.

- (1). — By Theorem 3.5 and Corollary 6.3, we have the following:

$$(6.8) \quad \text{Hom}(\mathcal{T}_{1,2}/\mathcal{I}^{\text{tot}}.\mathcal{T}_{1,2}, L) \hookrightarrow H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1)).$$

Moreover, since $\mathcal{I}^{\text{tot}} \subset \mathfrak{m}$, we have an injection

$$\text{Hom}(\mathcal{T}_{1,2}/\mathfrak{m}.\mathcal{T}_{1,2}, L) \hookrightarrow \text{Hom}(\mathcal{T}_{1,2}/\mathcal{I}^{\text{tot}}.\mathcal{T}_{1,2}, L).$$

By the assumption on the dimension of $H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1))$ we get

$$\dim \text{Hom}(\mathcal{T}_{1,2}/\mathfrak{m}.\mathcal{T}_{1,2}, L) \leq 1.$$

On the other hand, the fact that $\rho_K : G_{\mathbb{Q}} \rightarrow \text{GL}_4(K)$ is irreducible implies that $\mathcal{I}^{\text{tot}} = \mathcal{T}_{1,2}\mathcal{T}_{2,1} \neq 0$ and hence

$$\dim \text{Hom}(\mathcal{T}_{1,2}/\mathfrak{m}.\mathcal{T}_{1,2}, L) = 1.$$

Thus, Nakayama’s lemma implies that the \mathcal{T} -modules $\mathcal{T}_{1,2}$ is a monogenic \mathcal{T} -module.

Since $\mathcal{T}_{1,2}$ is a fractional ideal of K and each component ρ_{K_i} of ρ_K is absolutely irreducible, the annihilator of the generator of $\mathcal{T}_{1,2}$ over \mathcal{T} is trivial. Hence, $\mathcal{T}_{1,2}$ is a free rank one \mathcal{T} -module. Moreover, the symmetry under the anti-involution implies that $\mathcal{T}_{1,2} \simeq \mathcal{T}_{2,3}$ and hence $\mathcal{T}_{2,3}$ is also a free \mathcal{T} -module of rank one.

Let $\alpha \in K$ (resp. $\beta \in K$) be a generator of $\mathcal{T}_{1,2}$ (resp. of $\mathcal{T}_{2,3}$) as \mathcal{T} -module. A direct computation shows that one can choose $e'_1 = \alpha.e_1, e'_2 = e_2, e'_3 = \beta^{-1}.e_3$ as a suitable basis of idempotents.

Moreover, we recall that we have an injection by Theorem 4.3

$$\text{Hom}(\mathcal{T}_{1,3}/\mathcal{T}_{1,2}\mathcal{T}_{2,3}, L) = \text{Hom}(\mathcal{T}_{1,3}/\mathcal{T}, L) \hookrightarrow H^1_{f,\text{unr}}(\mathbb{Q}, \epsilon_p) = \{0\}.$$

Hence, Nakayama’s lemma implies that $\mathcal{T}_{1,3} = \mathcal{T}$. Now, we can conclude from the fact that $\mathcal{T}_{1,2}\mathcal{T}_{2,1} = \mathcal{T}_{2,3}\mathcal{T}_{3,2} = \mathcal{I}^{\text{tot}}$ that $\mathcal{T}_{2,1} = \mathcal{T}_{3,2} = \mathcal{I}^{\text{tot}}$.

(2). — By (3.1) applied with $(i, j) = (3, 2)$ and $(3, 1)$, respectively, applying Theorem 5.1 and Corollary 5.3 for $(i, j) = (3, 2)$ and using that $\mathcal{T}_{3,2} = \mathcal{I}^{\text{tot}}, \mathcal{T}'_{3,2} = \mathcal{T}_{3,1}\mathcal{T}_{1,2} = \mathcal{J}, \mathcal{T}_{3,1} = \mathcal{J}$, and $\mathcal{T}'_{3,1} = \mathcal{T}_{3,2}\mathcal{T}_{2,1} = (\mathcal{I}^{\text{tot}})^2$ we get injective morphisms

$$(6.9) \quad \begin{aligned} \text{Hom}(\mathcal{I}^{\text{tot}}/\mathcal{J}, L) &\hookrightarrow H^1_{f,\text{unr}}(\mathbb{Q}, \rho_f(k-2)) \\ \text{Hom}(\mathcal{J}/(\mathcal{I}^{\text{tot}})^2, L) &\hookrightarrow H^1(G_{\mathbb{Q}}^{Np}, \epsilon_p^{-1}). \end{aligned}$$

One has $\dim H^1_{f,\text{unr}}(\mathbb{Q}, \rho_f(k-2)) = 0$ (by a deep result of Kato [23]), hence Nakayama’s lemma applied to $\mathcal{I}^{\text{tot}}/\mathcal{J}$ yields that $\mathcal{I}^{\text{tot}} = \mathcal{J}$. Moreover, the ideal \mathcal{I}^{tot} is non-zero since ρ_K is irreducible. Thus, the fact that $\dim H^1(G_{\mathbb{Q}}^{Np}, \epsilon_p^{-1}) = 1$ (by Proposition 6.5) yields that $\mathcal{I}^{\text{tot}} = \mathcal{T}.g + (\mathcal{I}^{\text{tot}})^2$ and g is a generator of the ideal \mathcal{I}^{tot} .

(3). — The assertion follows from similar arguments to those already used to prove (1), (2) and the fact that $\text{Hom}(\mathcal{I}^{\text{tot}}/\mathcal{J}, L) \hookrightarrow \text{Sel}_{\mathbb{Q}, f_\alpha}$ by Theorem 5.1 and Corollary 5.3. □

One has the following general bound of the number of generators of \mathcal{I}^{tot} :

COROLLARY 6.10. — *Let $s := \dim H^1_{f,\text{unr}}(\mathbb{Q}, \rho_f(k-1))$. Assume **(Reg)**, **(St)**, and that $\dim \text{Sel}_{\mathbb{Q}, f_\alpha} = 0$ if $k = 2$. Then \mathcal{I}^{tot} is generated by at most s^2 elements.*

Proof. — It follows from (3.5) and Corollary 6.3 that $\mathcal{T}_{1,2}$ (resp. $\mathcal{T}_{2,3}$) is generated by at most s elements. Moreover, it follows from Theorem 5.1 and Corollary 5.3 that $\mathcal{T}_{2,1} = \mathcal{T}_{2,3}\mathcal{T}_{3,1}$ and $\mathcal{T}_{3,1} = \mathcal{T}_{3,2}\mathcal{T}_{2,1} + g.\mathcal{T}$. Thus, $\mathcal{T}_{2,1} = (\mathcal{T}_{3,2}\mathcal{T}_{2,1} + g.\mathcal{T})\mathcal{T}_{2,3} = \mathcal{I}^{\text{tot}}\mathcal{T}_{2,1} + g.\mathcal{T}_{2,3}$. Hence, $\mathcal{T}_{2,1}$ is generated also

by at most s elements and then $\mathcal{I}^{\text{tot}} = \mathcal{T}_{1,2}\mathcal{T}_{2,1}$ is generated by at most s^2 elements. □

7. Smoothness of \mathcal{E}_Δ at π_α

The goal of this section is to prove under the assumptions **(Reg)**, **(St)** and **(BK)** that $A := \mathcal{T}/\mathcal{I}^{\text{tot}}$ is a regular ring of dimension one (so it is a discrete valuation ring) and deduce that \mathcal{T} is a regular ring of dimension two when $\dim H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$.

7.1. Ordinary deformations of ρ_f

Recall that $\rho_f : G_{\mathbb{Q}}^{Np} \rightarrow \text{GL}_2(L)$ is the irreducible odd p -adic representation attached to f and f_α is the p -ordinary p -stabilisation of $f \in S_{2k-2}(\Gamma_0(N), L)$. We assume until the end of this subsection that **(St)** holds for f and consider the following deformation problem attached to ρ_f : for B any local L -Artinian algebra with maximal ideal \mathfrak{m}_B and residue field $B/\mathfrak{m}_B = L$, we define $\mathcal{D}(B)$ as the set of strict equivalence classes of representations $\rho_B : G_{\mathbb{Q}}^{Np} \rightarrow \text{GL}_2(B)$ lifting ρ_f (that is $\rho_B \bmod \mathfrak{m}_B \simeq \rho_f$) and which are ordinary at p in the sense that:

$$(7.1) \quad \rho_B|_{G_{\mathbb{Q}_p}} \simeq \begin{pmatrix} \psi_{1,B} & * \\ 0 & \psi_{2,B} \end{pmatrix},$$

where $\psi_{1,B} : G_{\mathbb{Q}_p} \rightarrow B^\times$ is an unramified character, and such that they are minimally ramified at every $\ell \mid N$ (i.e $\rho_B^{I_\ell}$ is free of rank one over B for any $\ell \mid N$). Let \mathcal{D}' be the subfunctor of \mathcal{D} of deformation with constant determinant (so equal to $\det \rho_f = \epsilon_p^{3-2k}$).

Since ρ_f is absolutely irreducible, we know from [29] that \mathcal{D} is prorepresentable by a complete local noetherian ring \mathcal{R}^{ord} together with its universal ordinary deformation $\rho^{\text{ord}} : G_{\mathbb{Q}} \rightarrow \text{GL}_2(\mathcal{R}^{\text{ord}})$. We know also from [29] that the tangent space $\mathcal{D}(L[\epsilon])$ (resp. $\mathcal{D}'(L[\epsilon])$) is isomorphic to a subspace $H_{\text{ord}}^1(\mathbb{Q}, \text{ad } \rho_f)$ (resp. $H_{\text{ord}}^1(\mathbb{Q}, \text{ad}^0 \rho_f)$) of $H^1(\mathbb{Q}, \text{ad } \rho_f)$ (resp. $H^1(\mathbb{Q}, \text{ad}^0 \rho_f)$).

The determinant of ρ^{ord} is a deformation of $\det \rho_f$, and endows \mathcal{R}^{ord} with a structure of A_1 -algebra, where A_1 is the completed local ring of $\mathbb{Z}_p[[\mathbb{Z}_p^\times]] = \bigoplus_{0 \leq l \leq p-1} \mathbb{Z}_p[[X]]^{(7)}$ at the height one prime ideal corresponding to $\det \rho_f$ ($A_1 = \mathbb{Q}_p[[X]]$).

(7) One has $\text{Spec } \mathbb{Z}_p[[\mathbb{Z}_p^\times]](\mathbb{C}_p) = \text{Hom}_{\text{cont}}(\mathbb{Z}_p^\times, \mathbb{C}_p^\times)$.

PROPOSITION 7.1. — Assume **(St)**. Then the L -vector space $H_{\text{ord}}^1(\mathbb{Q}, \text{ad}^0 \rho_f)$ is contained in $H_{f, \text{unr}}^1(\mathbb{Q}, \text{ad}^0 \rho_f)$.

Proof. — Let $\rho_\epsilon = (1 + \epsilon\rho_1)\rho_f$ be a representative of a strict equivalence class in $\mathcal{D}'(L[\epsilon])$, so ρ_1 is a 1-cocycle with respect to the adjoint action $\text{ad}^0 \rho_f$, and its cohomology class lies in $H_{\text{ord}}^1(\mathbb{Q}, \text{ad}^0 \rho_f)$. We will show first that $[\rho_1] \in H_g^1(\mathbb{Q}, \text{ad}^0 \rho_f) := \ker(H^1(\mathbb{Q}, \text{ad}^0 \rho_f) \rightarrow H^1(\mathbb{Q}_p, \text{ad}^0 \rho_f \otimes B_{\text{dR}}))$, where B_{dR} is Fontaine’s de Rham period ring. So it is enough so show that ρ_ϵ is ordinary in the sense of Definition 2.1 (so semi-stable by Theorem 2.2) as a 4-dimensional p -adic representation over L (since the L -representation ρ_ϵ corresponds to an element of $\text{Ext}_{G_{\mathbb{Q}}}^1(L, \text{ad}^0 \rho_f) = H^1(\mathbb{Q}, \text{ad} \rho_f^0)$). Since ρ_ϵ is ordinary at p , one can find a basis (e_1, e_2) of $L[\epsilon]^2$ such that the line $L[\epsilon] \cdot e_1$ is $G_{\mathbb{Q}_p}$ -stable and invariant under the action of the inertia I_p . Let $v_1 := e_1, v_2 = \epsilon \cdot e_1, v_3 := e_2, v_4 = \epsilon \cdot e_2$ be a basis of $L[\epsilon]^2$ as L -vector space.

By ordinarity at p one has

$$(\rho_\epsilon)|_{G_{\mathbb{Q}_p}} = ((1 + \epsilon\rho_1)\rho_f)|_{G_{\mathbb{Q}_p}} = \begin{pmatrix} \psi' & * \\ 0 & \psi'' \end{pmatrix},$$

where ψ' is unramified and $\psi''|_{I_p} = \epsilon_p^{3-2k}$ (since $\det \rho_\epsilon = \epsilon_p^{3-2k}$). Thus, the realization of ρ_ϵ by matrices in the basis (v_1, v_2, v_3, v_4) of $L[\epsilon]^2$ has the following form on I_p :

$$(\rho_\epsilon)|_{I_p} = \begin{pmatrix} 1 & 0 & * & * \\ 0 & 1 & * & * \\ 0 & 0 & \epsilon_p^{3-2k} & 0 \\ 0 & 0 & 0 & \epsilon_p^{3-2k} \end{pmatrix}.$$

Hence ρ_ϵ is ordinary and it follows from Theorem 2.2 that ρ_ϵ is semi-stable, and then $[\rho_1] \in H_g^1(\mathbb{Q}, \text{ad}^0 \rho_f)$. It remains to show that $\rho_1 \in H_{f, \text{unr}}^1(\mathbb{Q}, \text{ad}^0 \rho_f)$. Since ρ_ϵ is minimally ramified at any $\ell \mid N$ (i.e. $(\rho_\epsilon)|_{I_\ell} = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$), the restriction of the cohomology class of ρ_1 to I_ℓ is trivial (because $\text{Hom}(I_\ell, L)$ is 1-dimensional). To be able to conclude, we need to show that $H_g^1(\mathbb{Q}_p, \text{ad}^0 \rho_f) = H_f^1(\mathbb{Q}_p, \text{ad}^0 \rho_f)$. This follows immediately from the fact that

$$\dim H_g^1(\mathbb{Q}_p, \text{ad}^0 \rho_f) = \dim H_f^1(\mathbb{Q}_p, \text{ad}^0 \rho_f) + \dim \mathcal{D}_{\text{crys}}((\text{ad}^0 \rho_f)|_{G_{\mathbb{Q}_p}}(1))^{\phi=1}$$

and

$$\dim \mathcal{D}_{\text{crys}}((\text{ad}^0 \rho_f)|_{G_{\mathbb{Q}_p}}(1))^{\phi=1} = 0. \quad \square$$

7.2. Regularity of $\mathcal{T}/\mathcal{I}^{\text{tot}}$

Recall that $(\kappa_1, \kappa_2) \subset (\mathcal{O}(\mathcal{W}))^2$ are the universal weights interpolating k_1, k_2 (they are the derivative at 1 of $\epsilon_p^{\kappa_1}, \epsilon_p^{\kappa_2}$). Hence one can see κ_i as

global section in $\mathcal{O}(\mathcal{E}_\Delta)$ via the weight map $\kappa : \mathcal{E}_\Delta \rightarrow \mathcal{W}$. Recall also that $\epsilon_p^{-\kappa_1}$ and $\epsilon_p^{-\kappa_2}$ specialize at $\underline{k} = (k_1, k_2)$ to the characters $\epsilon_p^{-k_1}, \epsilon_p^{-k_2}$, respectively.

Let A be the local quotient ring $\mathcal{T}/\mathcal{I}^{\text{tot}}$ of dimension ≤ 2 . Note that A is Henselian, since \mathcal{T} is Henselian (the local ring of a rigid analytic space for the rigid topology is always Henselian).

Let $\text{Ps}_A : G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{E}_\Delta) \rightarrow A$ be the natural pseudo-character of dimension 4. Moreover, $\text{Ps}_A = \Psi_1 + \Psi_2 + \text{Tr}_A$ such that $\text{Tr}_A : G_{\mathbb{Q}} \rightarrow A$ is a pseudo-character lifting the pseudo-character $\text{Tr}(\rho_f)$ and $\{\Psi_i\}_{i=1,2} : G_{\mathbb{Q}} \rightarrow A^\times$ are characters lifting respectively ϵ_p^{2-k} and ϵ_p^{1-k} . Moreover, since ρ_f is absolutely irreducible, $\text{Tr}_A : G_{\mathbb{Q}} \rightarrow A$ is the trace of a deformation $\rho_A : G_{\mathbb{Q}} \rightarrow \text{GL}_2(A)$ of ρ_f . The deformation $\det \rho_A$ of $\det \rho_f$ yields a natural local morphism of $\overline{\mathbb{Q}}_p$ -algebras $\Lambda_1 \rightarrow A$ (see [8, Section 6]).

THEOREM 7.2. — Assume **(Reg)**. For any cofinite ideal $\mathcal{J} \subset \mathcal{T}$ containing \mathcal{I}^{tot} the \mathcal{T}/\mathcal{J} -module $\mathcal{D}_{\text{crys}}^+(M'/\mathcal{J}M' \otimes (\epsilon_p^{\kappa_2-2}))^{\Phi=U_1/U_0}$ is free of rank one and $\kappa_1 - \kappa_2 \in \mathcal{I}^{\text{tot}}$. Moreover, one has that $\Psi_2 \equiv \epsilon_p^{1-\kappa_2} \pmod{\mathcal{I}^{\text{tot}}}$.

Proof. — Recall that in the proof of Theorem 4.3 and Corollary 4.4, we have constructed a family of p -adic representations $\rho_{\mathcal{M}'} : G_{\mathbb{Q}_p} \rightarrow \text{GL}_{\mathcal{U}}(\mathcal{M}')$ over an affinoid $\mathcal{U} := \text{Spm } B \subset \mathcal{E}_\Delta$ containing π_α , and such that $\mathcal{M}' = \mathcal{M}/\mathcal{M}^{I_p}$ is a torsion-free quotient of \mathcal{M} of generic rank 3 (the generic rank of \mathcal{M} over \mathcal{U} is 4). By [7, Theorem 1.5.6] we have surjections

$$M = \mathcal{M} \otimes_B \mathcal{T} \twoheadrightarrow M/\mathcal{J}M \twoheadrightarrow \Psi_2 \pmod{\mathcal{J}},$$

such that any semi-simple S -subquotient of the S -module $\ker(M/\mathcal{J}M \rightarrow \Psi_2 \pmod{\mathcal{J}})$ occurs in $\{\rho_f, \epsilon_p^{2-k}\}$ (any S -simple module is necessarily an $S/\mathfrak{m}S$ -module).

On the other hand, since $\mathcal{M}/\mathcal{M}^{I_p} = \mathcal{M}'$, the surjection $M/\mathcal{J}M \rightarrow \Psi_2 \pmod{\mathcal{J}}$ must factor through

$$(7.2) \quad M'/\mathcal{J}M' \twoheadrightarrow \Psi_2 \pmod{\mathcal{J}}$$

for $M' = \mathcal{M}' \otimes_B \mathcal{T}$.

We recall from Corollary 4.4 that

$$l(\mathcal{D}_{\text{crys}}^+(M'/\mathcal{J}M' \otimes (\epsilon_p^{\kappa_2-2}))^{\Phi=U_1/U_0}) = l(\mathcal{T}/\mathcal{J}).$$

On the other hand, it follows from the fact that the semi-simple subquotients of

$$\ker(M'/\mathcal{J}M' \rightarrow \Psi_2 \pmod{\mathcal{J}})$$

occur in $\{\epsilon_p^{2-k}, \psi, \psi^{-1}\epsilon_p^{3-2k}\}$ that

$$\mathcal{D}_{\text{crys}}(\ker(M'/\mathcal{J}M' \rightarrow \Psi_2 \pmod{\mathcal{J}}) \otimes \epsilon_p^{\kappa_2-2})^{\Phi=U_1/U_0} = \{0\}.$$

Therefore, $l(\mathcal{D}_{\text{crys}}^+(\Psi_2 \otimes (\epsilon_p^{\kappa_2-2}) \bmod \mathcal{J})^{\Phi=U_1/U_0}) = l(\mathcal{T}/\mathcal{J})$. Thus, [7, Lemma 3.3.9] yields that

$$(7.3) \quad \mathcal{D}_{\text{crys}}^+(\Psi_2 \otimes (\epsilon_p^{\kappa_2-2}) \bmod \mathcal{J})^{\Phi=U_1/U_0} \text{ is a free rank one } \mathcal{T}/\mathcal{J}\text{-module,}$$

and then

$$\mathcal{D}_{\text{crys}}^+(M'/\mathcal{J}M' \otimes (\epsilon_p^{\kappa_2-2}))^{\Phi=U_1/U_0} \text{ is a free rank one } \mathcal{T}/\mathcal{J}\text{-module.}$$

At the same time [7, Proposition 2.5.4] (i.e the “constant weight lemma”) yields that $\Psi_2 \otimes \epsilon_p^{\kappa_2-2}$ has a constant weight given by 1 (i.e by the weight of $\epsilon_p^{-1} \equiv \Psi_2 \otimes \epsilon_p^{\kappa_2-2} \bmod \mathfrak{m}$) and that the Sen operator acts on Ψ_2 by multiplication by 1. It follows from Corollary 4.4 that 1 is a root of

$$(T)(T - (\kappa_1 - \kappa_2 + 1))(T - (\kappa_1 - 1)).$$

Thus, $(\kappa_2 - \kappa_1)(2 - \kappa_1) \equiv 0 \bmod \mathcal{J}$. Assume now that $k \geq 3$, then $(2 - \kappa_1)$ is invertible in (the local ring) \mathcal{T}/\mathcal{J} . Therefore, $\kappa_2 - \kappa_1 \in \mathcal{J}$. Assume now that $k = 2$, we can consider by Theorem 2.6(3) the sub-representation $\rho_{\mathcal{M}''} : G_{\mathbb{Q}_p} \rightarrow \text{Aut}_{\mathcal{U}}(\mathcal{M}'')$ generically of dimension 2 of $\rho_{\mathcal{M}'}$ (\mathcal{M}'' is a torsion-free $\mathcal{O}_{\mathcal{U}}$ -module) and it specialises at every point z of the Zariski dense set $\Sigma' \subset \mathcal{U}$ to a crystalline $G_{\mathbb{Q}_p}$ -representation ρ''_z of dimension 2 such that:

- (1) The Hodge–Tate weights of ρ''_z are $\{\kappa_1(z) - 1, \kappa_2(z) - 2\}$ with $\kappa_2(z) - 2 < \kappa_1(z) - 1$.
- (2) $\dim \mathcal{D}_{\text{crys}}(\rho''_z)^{\Phi=U_1/U_0(z)p^{p^{\kappa_2(z)}-2}} = 1$.

Let $M'' := \mathcal{M}'' \otimes_B \mathcal{T} = \mathcal{M}''_{\pi_\alpha}$ be the stalk of \mathcal{M}'' at π_α . Similar arguments to those already used to prove [7, Lemma 4.3.3(i)] yield that the Sen operator of $\mathcal{D}_{\text{sen}}(M''/\mathcal{J}M'')$ is annihilated by the polynomial $(T - (\kappa_2 - 2))(T - (\kappa_1 - 1))$. The assumption **(Reg)** and the fact that $G_{\mathbb{Q}_p}$ acts by $U_0 \epsilon_p^{-\kappa_1 - \kappa_2 + 3}$ on M'/M'' (U_0 means the unramified character with value U_0 at Frob_p) yield that the composition $M''/\mathcal{J}M'' \rightarrow M'/\mathcal{J}M' \rightarrow \Psi_2 \bmod \mathcal{J}$ is surjective. Thus, $T(T - (\kappa_1 - \kappa_2 + 1))$ annihilates the Sen operator of $\Psi_2 \otimes \epsilon_p^{\kappa_2-2}$, and since 1 is a root of that polynomial, we deduce immediately that $\kappa_2 - \kappa_1 \in \mathcal{J}$.

Meanwhile, the fact that $\mathcal{D}_{\text{crys}}^+(\Psi_2 \otimes (\epsilon_p^{\kappa_2-2}) \bmod \mathcal{J})^{\Phi=U_1/U_0}$ is a free rank one \mathcal{T}/\mathcal{J} -module yields that the character $\Psi_2 \otimes \epsilon_p^{\kappa_2-2} \bmod \mathcal{T}/\mathcal{J}$ is a crystalline $L[G_{\mathbb{Q}_p}]$ -representation with Hodge–Tate weight 1. Thus, $\Psi_2 \otimes (\epsilon_p^{\kappa_2-2}) \otimes \epsilon_p \bmod \mathcal{J}$ is of Hodge–Tate weight 0 and crystalline, therefore unramified. Thus, by class field theory we deduce that $\Psi_2 \otimes (\epsilon_p^{\kappa_2-2}) \otimes \epsilon_p \bmod \mathcal{J}$ is the trivial character (since \mathbb{Q} has a unique \mathbb{Z}_p -extension). Therefore, $\Psi_2 \bmod \mathcal{J} = \epsilon_p^{1-\kappa_2} \bmod \mathcal{J}$.

Finally, we conclude that $\kappa_2 - \kappa_1 \in \mathcal{I}^{\text{tot}}$ and that $\Psi_2 = \epsilon_p^{1-\kappa_2} \bmod \mathcal{I}^{\text{tot}}$ since the ideal \mathcal{I}^{tot} is the intersection of all cofinite length containing it (by Krull’s theorem). \square

COROLLARY 7.3. — Assume **(Reg)**. Then the local ring A is topologically generated by the image of $\text{Tr}(\rho_A)$ over Λ_1 .

Proof. — Let A' be the subring of A topologically generated by the image of the trace $\text{Tr}(\rho_A)$ over Λ_1 . Since $\kappa_1 - \kappa_2 \in \mathcal{I}^{\text{tot}}$ by Theorem 7.2, the polarisation of $\text{Ps}_{\mathcal{T}}$ described in Section 3.1 and the fact that the subconstituents of $\text{Ps}_{\mathcal{T}} \bmod \mathcal{I}^{\text{tot}}$ are uniquely determined (see [7, Proposition 1.5.1]) yield that $\text{Tr} \rho_A = \text{Tr}(\rho_A^{\vee} \otimes \epsilon_p^{3-2\kappa_1})$, $\Psi_1 = \Psi_2^{-1} \otimes \epsilon_p^{3-2\kappa_1} = \epsilon_p^{2-\kappa_1}$ (the last equality follows from Theorem 7.2). Thus, $\det \rho_A$ is given by the character $\epsilon_p^{3-2\kappa_1}$. As the determinant can be expressed in terms of the trace the image of $\epsilon_p^{2\kappa_1}$ lies in A' , and so by Hensel’s lemma the image of the character $\epsilon_p^{\kappa_1} = \epsilon_p^{\kappa_2} \bmod \mathcal{I}^{\text{tot}}$ (and therefore also of Ψ_1 and Ψ_2) lies in A' . Finally, as Ps_A is surjective onto A by construction of \mathcal{E}_{Δ} this establishes the proposition. \square

PROPOSITION 7.4. — Assume **(Reg)** and **(St)**. Then the representation ρ_A is p -ordinary and minimal.

Proof. — According to [7, Theorem 1.5.6] and [7, Lemma 4.3.9], there exists a \mathcal{T} -module $M \subset K^4$ of generic rank 4 endowed with a $G_{\mathbb{Q}}$ -continuous action which is generically given by the semi-simple representation

$$\rho_K : G_{\mathbb{Q}} \rightarrow S^{\times} \subset \text{GL}_4(K),$$

and equipped with a surjection $\pi : M/\mathcal{I}^{\text{tot}}M \twoheadrightarrow \rho_A$ such that the S -simple subquotients of its kernel are either ϵ_p^{1-k} or ϵ_p^{2-k} .

Since \mathcal{T} is reduced and ρ_K is semi-ordinary ($\rho_K^{I_p}$ is of dimension one and Frob_p acts on it by U_0) and $\alpha \neq 1$ when $k = 2$, we again have (as in Section 5) that M^{I_p} is not contained in $\mathfrak{m}M$. Since the S -simple subquotients of $\ker \pi$ do not contain ρ_f and contain only the representations in the set $\{\epsilon_p^{1-k}, \epsilon_p^{2-k}\}$, the regularity assumption further implies that the image of M^{I_p} under the surjection $\pi' : M/\mathfrak{m}M \twoheadrightarrow \rho_f$ is non-zero and hence the image of M^{I_p} under the surjection $\pi : M/\mathcal{I}^{\text{tot}}M \twoheadrightarrow \rho_A$ is non-zero and it is not contained in $\mathfrak{m}A^2$.

Thus, we have an exact sequence of $A[G_{\mathbb{Q}_p}]$ -modules:

$$(7.4) \quad 0 \rightarrow \rho_A^{I_p} \rightarrow \rho_A \rightarrow \rho_A/\rho_A^{I_p} \rightarrow 0.$$

Since $\rho_A/\rho_A^{I_p} \otimes_A L$ is of rank one Nakayama’s lemma implies that $\rho_A/\rho_A^{I_p}$ and $\rho_A^{I_p}$ are monogenic A -modules and generated respectively by y_1, y_2 .

Therefore y_1, y_2 generate A^2 and they must even form a basis of A^2 . Hence the exact sequence (7.4) splits as A -modules and yields that ρ_A is p -ordinary.

We shall now prove that ρ_A is minimally ramified at every $\ell \mid N$. Let ℓ be a prime number dividing N . From the proof of Theorem 4.3 we know that there exist idempotents $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3)$ of S lifting the idempotents attached respectively to $\epsilon_p^{2-k}, \epsilon_p^{1-k}, \rho_f$ such that $\tilde{e} = \tilde{e}_1 + \tilde{e}_2$ is in the center of $S_\ell = \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & M_{2,2}(\mathcal{T}) \end{pmatrix}$, the image of $\mathcal{T}[G_{\mathbb{Q}_\ell}]$ inside S . We also recall that N_K , the monodromy operator corresponding to the Weil–Deligne representation attached to $G_{\mathbb{Q}_\ell} \rightarrow S_\ell^\times$, can be viewed as an element of S_ℓ , has rank 1 by Proposition 2.10 and satisfies $\tilde{e}_3 N_K \tilde{e}_3 \neq 0$. For $N := \tilde{e}_3 N_K \tilde{e}_3 \in M_2(\mathcal{T})$ we know that N is non-trivial modulo $\mathfrak{m}_\mathcal{T}$ (since the rank of the monodromy operator of $WD_\ell(\rho_f)$ is one) and so the morphism $\rho_A|_{G_{\mathbb{Q}_\ell}} : G_{\mathbb{Q}_\ell} \rightarrow \mathrm{GL}_2(\mathcal{T}) \rightarrow \mathrm{GL}_2(A)$ is also minimally ramified. \square

Remark 7.5. — The assumption **(Reg)** is crucial to ensure the existence of a line in M fixed by inertia on which Frob_p acts by U_0 (i.e. $M^{I_p} \not\subset \mathfrak{m}M$). We have many examples for $\mathrm{GL}_2(\mathbb{Q})$ [11, Theorem C] where the 2-dimensional p -adic Galois representation $\rho_{\mathfrak{h}}$ attached to the local ring \mathfrak{h} of the eigencurve at p -irregular weight one forms is not ordinary, but the representation becomes ordinary when we extend the scalar to the field of fractions of \mathfrak{h} (i.e. $\rho_{\mathfrak{h}}^{I_p} \neq 0$ but $\rho_{\mathfrak{h}}^{I_p} \subset \mathfrak{m}_{\mathfrak{h}}\rho_{\mathfrak{h}}^{I_p}$).

PROPOSITION 7.6. — *Let \widehat{A} denote the completion of A with respect to its maximal ideal and assume **(St)**, **(Reg)** and **(BK)**. Then the natural morphism $\mathcal{R}^{\mathrm{ord}} \rightarrow \widehat{A}$ associated to $\rho_A \otimes_A \widehat{A}$ is an isomorphism of discrete valuation rings. Moreover, $\mathcal{R}^{\mathrm{ord}}$ is étale over A_1 .*

Proof. — According to Proposition 7.3, \widehat{A} is topologically generated by $\mathrm{Tr} \rho_A(G_{\mathbb{Q}})$. It follows that $\mathcal{R}^{\mathrm{ord}}$ surjects on \widehat{A} . We claim that the Krull dimension of \widehat{A} is at least one, and it is a consequence of the fact that A surjects onto the local ring $\mathcal{O}_{SK(\mathcal{F}),\alpha}$ at π_α of the 1-dimensional irreducible analytic subspace corresponding to the Saito–Kurokawa family $SK(\mathcal{F})$ specializing to π_α , where \mathcal{F} is the Hida family specialising to f_α (see [41, Proposition 4.2.5]).

On the other hand, it follows from Proposition 7.1 that the relative tangent space of $\Lambda \rightarrow \mathcal{R}^{\mathrm{ord}}$ (i.e. $\mathcal{D}'_{\rho_f}(L[\epsilon])$) is contained in $H_{f,\mathrm{unr}}^1(\mathbb{Q}, \mathrm{ad}^0 \rho_f)$, which is trivial under the assumption **(BK)**. Thus, the tangent space of $\mathcal{R}^{\mathrm{ord}}$ is at most 1-dimensional and $\mathcal{R}^{\mathrm{ord}}$ is unramified over A_1 . However, the existence of the surjection $\mathcal{R}^{\mathrm{ord}} \twoheadrightarrow \widehat{A}$ yields that the tangent space of $\mathcal{R}^{\mathrm{ord}}$ is necessarily 1-dimensional and that $\mathcal{R}^{\mathrm{ord}} \twoheadrightarrow \widehat{A}$ is an isomorphism of

1-dimensional regular rings. Since $\det \rho_A = \epsilon_p^{3-2\kappa_1}$, A_1 injects in $\mathcal{R}^{\text{ord}} \simeq A$ and \mathcal{R}^{ord} is necessarily flat over A_1 . Thus, \mathcal{R}^{ord} is étale over A_1 because it is unramified and flat. \square

THEOREM 7.7. — Assume **(BK)**, **(Reg)**, **(St)**, $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1))=1$ and when $k = 2$ also assume that $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$. Then the local ring \mathcal{T} is regular of dimension 2, i.e. \mathcal{E}_Δ is smooth at π_α . Moreover, the reducibility ideal of the pseudo-character $\text{Ps}_\mathcal{T}$ corresponds to the principal Weil divisor (closed subset of dimension one) of $\text{Spec } \mathcal{T}$ corresponding to the Saito–Kurokawa family $SK(\mathcal{F})$ specializing to π_α , where \mathcal{F} is the Hida family passing through f_α .

Proof. — We have to show that the tangent space of \mathcal{T} is of dimension 2. Since the Krull dimension is always less or equal to the dimension of the tangent space, we have to show that the maximal ideal \mathfrak{m} of \mathcal{T} has at most two generators. Note that $\mathcal{I}^{\text{tot}} = (g)$ (see Theorem 6.8 and Section 5.1) and $A = \mathcal{T}/(g)$ is regular of dimension 1. Hence \mathfrak{m} has at most two generators. Thus \mathcal{T} is regular. The rest of the assertion follows from the fact that $\mathcal{I}^{\text{tot}} = (g)$ and $\mathcal{O}_{SK(\mathcal{F}), x} = A = \mathcal{T}/(g)$ (since A is a discrete valuation ring). \square

One has the following general bound of the Zariski tangent space of $\pi_\alpha \in \mathcal{E}_\Delta$.

COROLLARY 7.8. — Assume **(BK)**, **(Reg)**, **(St)** and when $k = 2$ also assume that $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$. Then we have

$$2 \leq \dim \mathfrak{t}_{\pi_\alpha} \leq 1 + (\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)))^2$$

and

$$\dim \mathfrak{t}_{\pi_\alpha}^0 \leq (\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(k-1)))^2.$$

Proof. — The assertion follows immediately from Corollary 6.10 (i.e. \mathcal{I}^{tot} is generated by at most s^2 elements) and from Theorem 7.7 (i.e. $A = \mathcal{T}/\mathcal{I}^{\text{tot}}$ is étale over $A_1 \simeq \Lambda/(\kappa_1 - \kappa_2)$). \square

8. Smoothness failure of \mathcal{E}_N at π_α when N is square free and not prime

We prove in this subsection that our main results fail when we change the tame level to $\Gamma(N)$. In this subsection we can remove the assumption on the global root number ϵ_f being -1 as there exists a Saito–Kurokawa lift of level $\Gamma(N)$ for either sign (see [38]).

Coleman, Gouvea and Jochnowitz proved in [14] that the p -adic modular form for $\mathrm{GL}_2(\mathbb{Q})$ $G_2(q) = \frac{\zeta(-1)}{2} + \sum_{n=1}^\infty \sigma(n)q^n$, where $\sigma(n) = \sum_{d|n} d$ is not overconvergent, however the p -ordinary p -stabilization $E_2^{\mathrm{ord}_p}(q) = G_2(q) - p \cdot G_2(q^p)$ of $G_2(q)$ is classical, hence the critical p -stabilization $E_2^{\mathrm{crit}_p} = G_2(q) - G_2(q^p)$ of $G_2(q)$ is not overconvergent. On the other hand, any ordinary ℓ -stabilization $E_2^{\mathrm{crit}_p, \mathrm{ord}_\ell}$ of $E_2^{\mathrm{crit}_p}$ is an overconvergent modular form of weight two and level $\Gamma_0(\ell p)$. Note that $a_{\ell'}(E_2^{\mathrm{crit}_p, \mathrm{ord}_\ell}) = 1 + \ell'$ where $\ell' \nmid \ell, p$, and $a_\ell(E_2^{\mathrm{crit}_p, \mathrm{ord}_\ell}) = 1$, $a_p(E_2^{\mathrm{crit}_p, \mathrm{ord}_\ell}) = p$.

$E_2^{\mathrm{crit}_p, \mathrm{ord}_\ell}$ is a cuspidal overconvergent form of tame level $\Gamma_0(\ell)$ since each constant term of its q -expansion is trivial at each cusp of the multiplicative ordinary locus of the rigid curve attached to the semi-stable modular curve $X_1(\Gamma_1(4\ell) \cap \Gamma_0(p))/\mathbb{Z}_p$ (these cusps are in the $\Gamma_0(p)$ -orbit of the standard cusp ∞ , see [12, Sections 3.1 and 3.2]).

Let \mathcal{C}_N be the reduced eigencurve of tame level N constructed using the Hecke operators T_ℓ for $\ell \nmid Np$ and U_p (we omit the Hecke operators U_ℓ for $\ell \mid N$). Recall that there exists a flat and locally finite morphism $w : \mathcal{C}_N \rightarrow \mathcal{V}$, called the weight map, where \mathcal{V} is the weight space ($\mathcal{V}(\mathbb{C}_p) = \mathrm{Hom}_{\mathrm{cont}}(\mathbb{Z}_p^\times, \mathbb{C}_p^\times)$).

PROPOSITION 8.1. — *Let \mathcal{Y} be an irreducible component of the p -adic Eigencurve \mathcal{C}_N of tame level N specializing to a point y corresponding to the system of Hecke eigenvalues of $E_2^{\mathrm{crit}_p}$. Denote by $\rho_{\mathcal{U}} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(K_{\mathcal{U}})$ the Galois representation attached to \mathcal{U} , where $K_{\mathcal{U}}$ is the field of fractions of some connected affinoid subdomain \mathcal{U} of \mathcal{Y} containing y , then $\rho_{\mathcal{U}}$ is Steinberg at least one prime $\ell \mid N$ (hence $N \neq 1$).*

Proof. — Let $A := \mathcal{O}_{\mathcal{Y}, y}$ be the local ring of $\mathcal{Y} \subset \mathcal{C}_N$ at point y . One has a universal pseudo-character carried by \mathcal{C}_N

$$(8.1) \quad G_{\mathbb{Q}} \rightarrow \mathcal{O}(\mathcal{C}_N)$$

sending Frob_q to the Hecke operator T_q , where $q \nmid Np$ is a prime number. The localization of the pseudo-character (8.1) at A gives rise to a pseudo-character

$$\mathrm{Ps}_A : G_{\mathbb{Q}} \rightarrow A$$

of dimension 2 specializing to $\epsilon_p^{-1} \oplus \mathbb{1}$ modulo the maximal ideal of A . Moreover, Ps_A is the trace of a 2-dimensional irreducible Galois representation $\rho_A : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(Q(A))$ (since \mathcal{Y} corresponds to a cuspidal Coleman family). Hence, we obtain from ρ_A a non-trivial cohomology class c_y in $H^1(\mathbb{Q}, \epsilon_p)$ (see [7, Section 1.5]). The cohomology class c_y corresponds to an extension $V = \mathbb{Q}_p^2$ of ϵ_p^{-1} by $\mathbb{1}$ unramified outside Np . It is known that for

any classical point y' in \mathcal{C}_N , the semi-simple p -adic Galois representation $\rho_{y'} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}(V_{y'})$ of dimension 2 attached to the modular form corresponding to y' (i.e. $\mathrm{Tr} \rho_{y'}$ is the specialization of (8.1) at y') has a crystalline periods equal to $U_p(y')$ (see [24]) and it corresponds to its smaller Hodge–Tate weight which is zero (i.e. $\mathcal{D}_{\mathrm{crys}}(V_{y'})^{\Phi=U_p(y')} \neq 0$), hence by using the analytic continuation of the crystalline periods U_p on the Eigencurve \mathcal{C}_N (see [7, Theorem 4.3.6]), one has $\mathcal{D}_{\mathrm{crys}}(V)^{\Phi=U_p(y)} = \mathcal{D}_{\mathrm{crys}}(V)^{\Phi=p} \neq 0$ (note that $U_p(y) = U_p(E_2^{\mathrm{crit}_p}) = p$). Thus, c_y is crystalline extension of ϵ_p^{-1} by $\mathbb{1}$, and it belongs to

$$H_f^1(G_{\mathbb{Q}}^{Np}, \epsilon_p) = \ker(H^1(G_{\mathbb{Q}}^{Np}, \epsilon_p) \rightarrow H^1(\mathbb{Q}_p, \epsilon_p \otimes B_{\mathrm{crys}})).$$

Let us proceed now by contradiction. Assume that $\rho_{\mathcal{U}}$ is not Steinberg at any $\ell \mid N$ (i.e the rank of the monodromy operator of the Weil–Deligne representation attached to $\rho_{\mathcal{U}}$ by [7, Lemma 7.8.14] at any ℓ is zero), hence $\rho_{\mathcal{U}}$ is principal series or supercuspidal, which implies that for any $\ell \mid N$, the image of the inertia group I_{ℓ} by $\rho_{\mathcal{U}}$ is finite (we also have a natural inclusion $K_{\mathcal{U}} \subset Q(\mathcal{O}_{y,y})$), and then semi-simple and reducible. Moreover, $\epsilon_p^{-1} \oplus \mathbb{1}$ is trivial on I_{ℓ} when $\ell \nmid p$, hence $\rho_{\mathcal{U}}$ is unramified outside p .

Thus, the extension c_y is not Steinberg at any $\ell \mid N$ (hence unramified outside p) and it belongs necessarily to $H_{f,\mathrm{unr}}^1(\mathbb{Q}, \epsilon_p)$ which is trivial. Finally, the cohomology class c_y is trivial, contradicting the fact that ρ_y is absolutely irreducible. \square

Remark 8.2.

- (1) The Atkin–Lehner eigenvalue of the classical specializations of \mathcal{Y}_{ℓ} at ℓ is constant and equal to -1 .
- (2) Let $\ell \mid N$ be a prime for which f is special, then the Hida family \mathcal{F} specializing to f_{α} is special at ℓ and the Atkin–Lehner eigenvalue of the classical specializations of \mathcal{F} at ℓ is constant.
- (3) According to [28], the weight map $w : \mathcal{C}_{\ell} \rightarrow \mathcal{V}$ is étale at $E_2^{\mathrm{crit}_p, \mathrm{ord}_{\ell}}$, and since w is locally finite, one can shrink any affinoid neighborhood of $E_2^{\mathrm{crit}_p, \mathrm{ord}_{\ell}}$ to ensure that it will be étale over \mathcal{V} (see Proposition A.5).

We can use this to construct endoscopic components containing π_{α} . We first note the following result about classical Yoshida lifts.

PROPOSITION 8.3 ([37, Proposition 3.1]). — *Let $f_1 \in S_{k_1}(N_1)$, $f_2 \in S_{k_2}(N_2)$ be newforms of squarefree level with even integers $k_1 \geq k_2 \geq 2$ and $M := \mathrm{gcd}(N_1, N_2) > 1$. Assume that the Atkin–Lehner eigenvalues of f_1 and f_2 for $\ell \mid M$ coincide. Put $N = \mathrm{lcm}(N_1, N_2)$. Then there exists a*

non-zero holomorphic Yoshida lift of level $\Gamma(N)$ and weight $((k_1 + k_2)/2, (k_1 - k_2 + 4)/2)$ with corresponding Galois representation $\rho_{f_1} \oplus \rho_{f_2}(\frac{k_1 - k_2}{2})$. For $p \nmid N$ there exists a p -stabilisation of this lift (of Iwahori level at p) with U_0 -eigenvalue α_1 and U_1 -eigenvalue⁽⁸⁾ $\alpha_1\alpha_2p^{\frac{k_1 - k_2 - 2}{2}}$, where α_i are roots of the Hecke polynomial of f_i at p for $i = 1, 2$.

Proof. — For the existence of the lift of level $\Gamma(N)$ see [37, Proposition 3.1]. For the p -stabilisation of the principal unramified series see [31, Section 7.1.1], but we use the normalization of [41, Section 2.4.16]. \square

THEOREM 8.4. — *Let $\ell \mid N$ be a prime number for which f is Steinberg, \mathcal{U}^1 be an affinoid subdomain of the p -adic eigencurve $w_2 : \mathcal{C}_\ell \rightarrow \mathcal{V}$ of tame level ℓ containing $E_2^{\text{crit}_p, \text{ord}_\ell}$, corresponding to a Coleman family $G = \sum_{n=1}^\infty a(n, G)q^n$, and such that it is étale over the weight space \mathcal{V} . Let \mathcal{U}^0 be an affinoid subdomain of the ordinary locus $\mathcal{C}_N^{\text{ord}}$ of the p -adic eigencurve \mathcal{C}_N of tame level N containing f_α and corresponding to the Hida family $\mathcal{F} = \sum_{n=1}^\infty a(n, \mathcal{F})q^n$, and such that it is étale⁽⁹⁾ over the weight space \mathcal{V} .*

There exists a Zariski closed immersion $\lambda_{Y_0} : \mathcal{U}^0 \times_{\mathbb{Q}_p} \mathcal{U}^1 \hookrightarrow \mathcal{E}_N$ with image denoted by $Y_0(\mathcal{F}, G)$ and such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{U}^0 \times_{\mathbb{Q}_p} \mathcal{U}^1 & \xrightarrow{\lambda_{Y_0}} & \mathcal{E}_N \\ \downarrow w_1 \times w_2 & & \downarrow \kappa \\ \mathcal{V} \times \mathcal{V} & \xrightarrow{\lambda_\kappa} & \mathcal{W} \end{array}$$

where $\lambda_\kappa(2k_1, 2k_2) = (k_1 + k_2, k_1 - k_2 + 2)$ and the morphism λ_{Y_0} corresponds to the morphism

$$\lambda_{Y_0}^* : \mathcal{O}(\mathcal{E}_N) \rightarrow \mathcal{O}(\mathcal{U}^0) \widehat{\otimes}_{\mathbb{Q}_p} \mathcal{O}(\mathcal{U}^1)$$

defined by

$$\begin{aligned} \lambda_{Y_0}^*(P_\ell(X)) &= (X^2 - a(\ell, \mathcal{F})X + \ell^{-3}\kappa_1\kappa_2(\ell)) \\ &\quad \times (X^2 - \kappa_2(\ell)\ell^{-2}a(\ell, G)X + \ell^{-3}\kappa_2(\ell)\cdot\kappa_1(\ell)), \end{aligned}$$

for any $\ell \nmid Np$, where $P_\ell(X) \in \mathcal{O}(\mathcal{E}_N)[X]$ is the Hecke–Andrianov polynomial at $\ell \nmid Np$, and $\lambda_{Y_0}^*(U_0) = a(p, \mathcal{F})$, and $\lambda_{Y_0}^*(U_1) = a(p, \mathcal{F}) \times a(p, G)$.

⁽⁸⁾For the different normalisation (B.2) of the U_1 operator on the eigenvariety this corresponds to the constant eigenvalue $\alpha_1\alpha_2$.

⁽⁹⁾According to Hida’s control theorem, the weight map $w_1 : \mathcal{C}_N \rightarrow \mathcal{V}$ is étale at f_α , and since w is locally finite, one can shrink any affinoid neighborhood of f_α to ensure that it will be étale over \mathcal{V} .

Proof. — One can choose the affinoids $\mathcal{U}^0 \subset \mathcal{C}_N$ and $\mathcal{U}^1 \subset \mathcal{C}_\ell$ étale over the weight space and small enough such that there exist $\epsilon, v \in \mathbb{R}$ and the Banach sheaf ω_ϵ^κ on $\overline{X}(v) \times W$ of locally analytic v -overconvergent p -adic families (see Section B.3), where $W = \text{Spm } R$ is an affinoid of the weight space \mathcal{W} given by $w_1(\mathcal{U}^0) \times_{\mathbb{Q}_p} w_2(\mathcal{U}^1)$. Let $\mathcal{T}_{W,1}$ be the affinoid \mathbb{Q}_p -algebra generated over R by the image of the abstract Hecke algebra \mathcal{H}_N in the space of endomorphisms of the sections of $\varinjlim_{v \rightarrow 0} H^0(\overline{X}(v) \times W, \omega_\epsilon^\kappa(-D))$ with slope ≤ 1 . By construction of \mathcal{E}_N (see Section B.3), $\mathcal{E}_{N,W}^1 = \text{Spm } \mathcal{T}_{W,1}$ is an affinoid subdomain of \mathcal{E}_N . Let $\theta : \mathcal{H}_N \rightarrow \mathcal{T}_{W,1}$ be the natural surjection and J be the kernel of θ generated by g_1, \dots, g_n .

On the other hand, let λ be the morphism

$$\lambda : \mathcal{H}_N \rightarrow \mathcal{O}(\mathcal{U}^0) \widehat{\otimes}_{\mathbb{Q}_p} \mathcal{O}(\mathcal{U}^1)$$

defined by

$$\begin{aligned} \lambda_{Y_o}^*(P_\ell(X)) &= (X^2 - a(\ell, \mathcal{F})X + \ell^{-3}\kappa_1\kappa_2(\ell)) \\ &\quad \times (X^2 - \kappa_2(\ell)\ell^{-2}a(\ell, G)X + \ell^{-3}\kappa_2(\ell) \cdot \kappa_1(\ell)), \end{aligned}$$

for $\ell \nmid Np$, where $P_\ell(X) \in \mathcal{H}_N[X]$ is the Hecke–Andrianov polynomial at $\ell \nmid Np$, and $\lambda_{Y_o}^*(U_0) = a(p, \mathcal{F})$, and $\lambda_{Y_o}^*(U^1) = a(p, \mathcal{F}) \times a(p, G)$.

It is enough to prove that for every $1 \leq i \leq n$, $\lambda(g_i) = 0$. Note that the classical points old at p of $\mathcal{U}^0, \mathcal{U}^1$ form a dense set Σ of $\mathcal{U}^0 \times_{\mathbb{Q}_p} \mathcal{U}^1$. It follows from Proposition 8.3 that the points Σ lift to a set $\widetilde{\Sigma}^{(10)}$ of points of $\mathcal{E}_{N,W}^1$. Hence, for any $1 \leq i \leq n$, the specialization of $\lambda(g_i)$ at the points of the dense subset Σ of $\mathcal{U}^0 \times_{\mathbb{Q}_p} \mathcal{U}^1$ is trivial, yielding that

$$(8.2) \quad \lambda(g_i) = 0 \text{ for any } 1 \leq i \leq n.$$

Hence, we obtain a surjective homomorphism

$$\mathcal{O}(\mathcal{E}_{N,W}^1) \twoheadrightarrow \mathcal{O}(\mathcal{U}^0) \widehat{\otimes}_{\mathbb{Q}_p} \mathcal{O}(\mathcal{U}^1),$$

yielding a morphism $\mathcal{U}^0 \times_{\mathbb{Q}_p} \mathcal{U}^1 \rightarrow \mathcal{E}_{N,W}^1$, and its image is an irreducible component of $\mathcal{E}_{N,W}^1$. □

COROLLARY 8.5. — *Assume $N > 1$ is squarefree and not prime. Assume that f is Steinberg for at least two primes $\ell_i \mid N, i = 1, 2$. Then the Siegel eigenvarieties \mathcal{E}_N of tame level N is singular at π_α and there exists at least two p -adic families specializing to π_α .*

⁽¹⁰⁾ Any point of Σ corresponds to a 2-tuple of old forms (f_1, g_1) at p . Hence, f_1 (resp. g_1) is the p -ordinary (resp. p -critical) p -stabilization of a classical form of level $\Gamma_0(N)$ (resp. $\Gamma_0(\ell)$) denoted by f_1^{old} (resp. g_1^{old}). So we can consider the Yoshida lift of (f_1^{old}, g_1^{old}) and take its semi-ordinary p -stabilization which gives a point of $\widetilde{\Sigma} \subset \mathcal{E}_N^1$.

Proof. — If f is Steinberg at ℓ_1 and ℓ_2 , then by the previous theorem we get two irreducible components of \mathcal{E}_N (they are endoscopic) specializing to π_α by taking \mathcal{U}^1 arising from \mathcal{Y}_{ℓ_i} . \square

A direct consequence of the above corollary is that $\kappa : \mathcal{E}_N \rightarrow \mathcal{W}$ is ramified at π_α . Let $S_k^\dagger(N)^{|\mathbb{U}|_p=1}[\pi_\alpha]$ be the generalized eigenspace attached to π_α inside the L -vector space of locally analytic overconvergent Siegel cusp forms $S_k^\dagger(N)^{|\mathbb{U}|_p=1}$ of tame level $\Gamma(N)$ and slope 1 for \mathbb{U} .

COROLLARY 8.6. — *One has $\dim_L S_k^\dagger(N)^{|\mathbb{U}|_p=1}[\pi_\alpha] \geq 2$.*

Proof. — Since \mathcal{W} is smooth at $\kappa(\pi_\alpha)$ and \mathcal{E}_N is singular at π_α , the local ring $\mathcal{T}_0 = \mathcal{O}_{\mathcal{E}_N, \pi_\alpha} / \mathfrak{m}_{\mathcal{O}_{\mathcal{W}, \kappa(\pi_\alpha)}} \mathcal{O}_{\mathcal{E}_N, \pi_\alpha}$ of the fiber of $\kappa^{-1}(\kappa(\pi_\alpha))$ at π_α is Artinian with a non-trivial tangent space (since κ is necessarily ramified at π_α in this case). On the other hand, it follows from the construction of eigenvarieties that the local ring \mathcal{T}_0 at π_α of the fiber $\kappa^{-1}(\kappa(\pi_\alpha))$ acts faithfully on $S_k^\dagger(N)^{|\mathbb{U}|_p=1}[\pi_\alpha]$. Hence, $\dim S_k^\dagger(N)^{|\mathbb{U}|_p=1}[\pi_\alpha] \geq 2$ (since $\dim_L \mathcal{T}_0 \geq 2$). \square

Appendix A. Some basic facts about rigid analytic geometry

We shall recall in this section the notions of “very Zariski dense” subset of a rigid analytic space and discuss accumulation points of a Zariski dense set and irreducible components of rigid analytic spaces. Moreover, we will recall some basic properties of finite and torsion-free morphisms of affinoid spaces.

The following proposition is an analogue to [10, Proposition 2.1.6] for \mathbb{Q}_p -rigid analytic spaces.

PROPOSITION A.1. — *Let $g : X \rightarrow Y$ be a finite morphism between two \mathbb{Q}_p -affinoid spaces, $y \in f(X) \subset Y$, and $g^{-1}(y) = \{x_1, x_2, \dots, x_n\}$, then there exists a small affinoid neighborhood \mathcal{U}_{i_0} of y in Y such that $g^{-1}(\mathcal{U}_{i_0}) = \bigcup_{1 \leq k \leq n} V_k^{i_0}$, and $V_k^{i_0} \cap V_j^{i_0} = \emptyset$, when $k \neq j$. Moreover, for any $1 \leq k \leq n$, the domains $\{V_k^i, i \in I \text{ and } i \leq i_0\}$ form a basis of neighborhood of x_k when \mathcal{U}_i varies in a family $\{\mathcal{U}_i, i \in I \text{ and } i \leq i_0\}$ of basis of affinoids containing y .*

Proof. — Let B (resp. A) be the affinoid \mathbb{Q}_p -algebra corresponding to X (resp. Y), and $\varphi : A \rightarrow B$ be the finite morphism corresponding to g . Let B_y be the finite $\mathcal{O}_{Y,y}$ -algebra $B \otimes_A \mathcal{O}_{Y,y}$; thanks to [10, Theorem 2.1.5] the

local ring $\mathcal{O}_{Y,y}$ is Henselian, hence

$$B_y = {}^{(11)} \prod_{x_i \in g^{-1}(y)} \mathcal{O}_{X,x_i}.$$

On the other hand, one has

$$B_y = B \otimes_A \mathcal{O}_{Y,x} = B \otimes_A \varinjlim_{\mathcal{U}_i} \mathcal{O}_Y(\mathcal{U}_i) = \varinjlim_{\mathcal{U}_i} B \otimes_A \mathcal{O}_Y(\mathcal{U}_i),$$

where $\{\mathcal{U}_i, i \in I\}$ runs over the affinoid neighborhood of y .

Hence, we have

$$\begin{aligned} \varinjlim_{\mathcal{U}_i, i \in I} B \otimes_A \mathcal{O}_Y(\mathcal{U}_i) &= {}^{(12)} \varinjlim_{\mathcal{U}_i, i \in I} B \widehat{\otimes}_A \mathcal{O}_Y(\mathcal{U}_i) \\ &= \varinjlim_{\mathcal{U}_i, i \in I} \mathcal{O}_X(g^{-1}(\mathcal{U}_i)) = \prod_{x_i \in g^{-1}(y)} \mathcal{O}_{X,x_i}. \end{aligned}$$

Thus, each local component \mathcal{O}_{X,x_j} of $\prod_{x_i \in g^{-1}(y)} \mathcal{O}_{X,x_i}$ corresponds to an idempotent e_j of B_y . So there exist an $i_0 \in I$ and orthogonal idempotents $\{\tilde{e}_j, 1 \leq j \leq n\}$ of $\mathcal{O}_X(g^{-1}(\mathcal{U}_{i_0}))$ whose image in B_y is $\{e_j, 1 \leq j \leq n\}$ and corresponding respectively to $\{x_1, \dots, x_n\}$. Thus, $\mathcal{O}_X(g^{-1}(\mathcal{U}_{i_0})) = \prod_{\tilde{e}_j, 1 \leq j \leq n} \tilde{e}_j \cdot \mathcal{O}_X(g^{-1}(\mathcal{U}_{i_0}))$, and hence each affinoid subdomain $\text{Spm } \tilde{e}_k \cdot \mathcal{O}_X(g^{-1}(\mathcal{U}_{i_0}))$ of X corresponds to a connected component $V_k^{i_0}$ of $g^{-1}(\mathcal{U}_{i_0})$ containing x_k . Hence, $g^{-1}(\mathcal{U}_{i_0}) = \bigcup_{1 \leq k \leq n} V_k^{i_0}$, and $V_l^{i_0} \cap V_k^{i_0} = \emptyset$, when $l \neq k$.

Finally, the rest of the assertion follows from the fact that

$$\varinjlim_{i \leq i_0} \mathcal{O}_X(V_k^i) = \mathcal{O}_{X,x_k},$$

and the inductive limit is taken on the connected component V_k^i of $g^{-1}(\mathcal{U}_i)$ containing x_k , when \mathcal{U}_i varies over the affinoid neighborhoods of x_k inside \mathcal{U}_{i_0} . □

We recall that F is an irreducible component of a \mathbb{Q}_p -separated reduced rigid analytic space X , if F is the image of a connected component of the normalization X^{nor} of X via the normalization morphism $X^{\text{nor}} \rightarrow X$ (see [16]). Moreover, when X is a reduced affinoid $\text{Spm } A$, then the irreducible components of X correspond to $\text{Spm } A/\mathcal{P}$, where \mathcal{P} is a minimal prime ideal of A .

We recall also that a subset Z of a reduced \mathbb{Q}_p -rigid analytic space X is said to be Zariski-dense if the only analytic subset of X containing Z is X itself.

(11) Since B_y is finite over the Henselian ring $\mathcal{O}_{Y,y}$, it is necessarily a product of local Henselian rings.

Remark A.2. — The set $S = \{(1/p^n, 1/p^m), \text{ where } n \in \mathbb{Z}, m \in \mathbb{N}\}$ of the rigid affine plane $\mathbb{A}_2^{\text{rig}}$ of dimension 2 is Zariski dense but for any open affinoid subdomain $\mathcal{U} \subset \mathbb{A}_2^{\text{rig}}$, the set $\mathcal{U} \cap S$ is not Zariski dense in \mathcal{U} (it follows from the maximum modulus principle).

This example motivates the notion of a very Zariski dense set of a rigid analytic space (see also [4, Definition II.5.1]):

DEFINITION A.3.

- (1) Let X be a \mathbb{Q}_p -separated reduced rigid analytic space over \mathbb{Q}_p , and $\Sigma \subset X$ be a Zariski dense subset. We say that Σ is very Zariski-dense in X if for every $z \in \Sigma$ there is a basis of open affinoid neighborhoods \mathcal{U} of z in X such that $\Sigma \cap \mathcal{U}$ is Zariski-dense in \mathcal{U} .
- (2) We say that a subset Z of a \mathbb{Q}_p -separated rigid analytic space Y accumulates at $y \in Y$ if there is a basis of affinoid neighborhoods $U \subset Y$ of y such that $U \cap Z$ is Zariski-dense in U .

Remark A.4. — Let X be a separated \mathbb{Q}_p -rigid space, $\{F_i\}$ be the irreducible components of X and \mathcal{U} be an admissible open of X . Then it follows from [16, Corollary 2.2.9] that each irreducible component of \mathcal{U} is contained in a unique F_i and for any i , $\mathcal{U} \cap F_i$ is empty or the union of irreducible components of \mathcal{U} .

PROPOSITION A.5.

- (1) Let $g : X \rightarrow Y$ be a finite flat morphism between two \mathbb{Q}_p -affinoid spaces such that X is equidimensional and Y is irreducible. Assume that g is étale at a Zariski dense set Σ of points of X , then after shrinking X to a smaller admissible open X' of X , the restriction $g : X' \rightarrow g(X')$ is étale and $g(X')$ is an admissible open of Y .
- (2) Let $g : X \rightarrow Y$ be a finite morphism between rigid analytic spaces, then for any irreducible component F of X , $g(X)$ is a closed irreducible component of Y .

Proof.

(1). — It is known that g is étale outside of the support of the relative differential sheaf $\Omega_{X/Y}$. Moreover, since g is étale at a Zariski dense set of points of X , the support Z of $\Omega_{X/Y}$ is a Zariski closed set of X of dimension $< \dim X$ (since Σ is Zariski dense in all irreducible components of X by [16, Proposition 2.2.8]). Hence, $g_{|X-Z} : X - Z \rightarrow Y$ is étale, and the image of

the Zariski open ⁽¹³⁾ $X - Z$ under g is a Zariski open of Y (a flat morphism is Zariski open).

(2). — The assertion follows from the fact that a finite morphism is Zariski closed and [16, Proposition 2.2.3]. \square

The following proposition was proved by Chenevier in [13] using base change arguments. We give in the following a more direct proof:

PROPOSITION A.6. — *Let $g : X \rightarrow Y$ be a finite torsion-free morphism between two reduced \mathbb{Q}_p -affinoid spaces and such that Y is irreducible. Then :*

- (1) X is equidimensional of dimension equal to $\dim Y$ and the image of each irreducible component of X under g is Y .
- (2) Let Σ be a Zariski dense set of Y , then $g^{-1}(\Sigma)$ is Zariski dense in X .

Proof.

(1). — Let B (resp. A) be the affinoid algebra corresponding to X (resp. Y) and $g^* : A \rightarrow B$ be the finite torsion-free morphism corresponding to g . Since Y is irreducible and reduced, A is a domain. Let \mathcal{P} be a minimal prime ideal of B corresponding to an irreducible component F of X , it follows from the fact that B is a torsion-free finite A -algebra that the morphism $A \rightarrow B/\mathcal{P}$ is injective (since the zero divisors of a reduced Noetherian ring are the union of its minimal prime ideals). Moreover, the image of the natural composition $F \rightarrow X \rightarrow Y$ is dense, because $A \rightarrow B/\mathcal{P}$ is injective (so the image of the morphism $\text{Spec } B/\mathcal{P} \rightarrow \text{Spec } A$ is Zariski dense) and $\text{Spec } A$ and $\text{Spec } B$ are Jacobson schemes (so $\text{Spm } A$ is Zariski dense in $\text{Spec } A$, and the same for B).

However, g is also finite, and then Zariski closed. Hence, the irreducible component F of X surjects onto Y , and since the morphism $A \rightarrow B/\mathcal{P}$ is injective and finite, then $\dim F = \dim Y$ (it follows from the Going-up theorem), and hence X is equidimensional of dimension equal to $\dim Y$.

(2). — A subset $\Sigma' \subset X$ is a Zariski dense set of a reduced affinoid X if and only if for any irreducible component F of X (see [16, Proposition 2.2.8]), $\Sigma' \cap F$ is a Zariski dense set of F . Thus, it is enough to prove the assertion when X is reduced and irreducible. Assume that X is irreducible and that Σ is a Zariski dense set of Y . Let $\Sigma' \subset X$ denote the subset

⁽¹³⁾ Note that a Zariski open U of rigid analytic space X is not necessarily an affinoid subdomain of X . Take $X = \text{Spm } \mathbb{Q}_p \langle T \rangle$ and $U = D(T)$ the locus where T is invertible; it is clear that U doesn't satisfy the maximal modulus principle for the function $1/T$, and hence U is not an affinoid. However, any Zariski open is an admissible open for the rigid topology.

$g^{-1}(\Sigma)$ of X . Since g is finite and torsion-free, then g is closed for the Zariski topology and surjective, and then the Zariski closure of Σ' is necessarily an analytic subspace $Z \subset X$ of dimension equal to $\dim Y = \dim X$, because $g(Z)$ is a Zariski closed set of Y containing Σ (so $g(Z)$ contains Y the closure of Σ). Hence, Z is finite and surjects on Y and it follows that $Z = X$, since they have the same dimension and X is irreducible. \square

LEMMA A.7. — *Let $\mathcal{U} = \text{Spm } A$ be an equidimensional affinoid of dimension 2, F be a Zariski closed subset of \mathcal{U} of dimension ≤ 1 , \mathcal{U}' be the admissible open given by $\mathcal{U} - F$. Let Σ be Zariski dense set of \mathcal{U} , then $\Sigma' = \Sigma \cap \mathcal{U}'$ is Zariski dense in \mathcal{U} and in \mathcal{U}' .*

Proof. — Note that $\Sigma = \Sigma' \cup (\Sigma \cap F)$. Hence, the Zariski closure $\bar{\Sigma}$ of Σ is equal to the union of the Zariski closure $\bar{\Sigma}'$ of Σ' with the closure $\bar{\Sigma} \cap \bar{F}$ of $\Sigma \cap F$. On the other hand, $\bar{\Sigma} = \mathcal{U}$ and it is equidimensional of dimension 2, and $\bar{\Sigma} \cap \bar{F} \subset F$ is of dimension at most one. Hence, $\bar{\Sigma}' = \mathcal{U}$, yielding that Σ' is dense in \mathcal{U} and so in \mathcal{U}' . \square

Appendix B. On the very Zariski density of classical points in the Eigenvariety \mathcal{E}_Δ

The goal of this section is to recall quickly the construction of the Siegel eigenvarieties and to prove that classical points which are old at p and of cohomological weights are very Zariski dense in them.

B.1. The Weight space \mathcal{W}

Recall that the connected components of \mathcal{W} are naturally indexed by $\mathcal{W}^{a,b}$, where $(a, b) \in (\mathbb{Z}/(p - 1)\mathbb{Z})^2$. The classical weights $(k_1, k_2) \in \mathcal{W}^{a,b}$ are congruent to $(a, b) \pmod{p - 1}$, in other words, the discrete part of the restriction of any character of $\mathcal{W}^{a,b}(\mathbb{C}_p)$ to $\mathbb{Z}/p\mathbb{Z}^\times$ is (ω_p^a, ω_p^b) , where ω_p is the Teichmüller character. In addition, the formal scheme $\text{Spf } \mathbb{Z}_p[[T_1, T_1]]$ is a Raynaud’s formal model of any connected component⁽¹⁴⁾ $\mathcal{W}^{a,b}$ of the weight space \mathcal{W} .

Remark B.1. — The category of admissible \mathbb{Z}_p -formal schemes (which includes locally topologically of finite type formal \mathbb{Z}_p -schemes) is localised

⁽¹⁴⁾Note that $\text{Spm } \mathbb{Z}_p[[T_1, T_1]][1/p] = \mathcal{W}^{a,b}$, and $\text{Spm } \mathbb{Z}_p[[T_1, T_1]][1/p]$ is the open disk of dimension 2 and radius 1.

with respect to blowups in the special fibre and the Raynaud generic fibre functor defines an equivalence of categories between the localised category of admissible formal \mathbb{Z}_p -schemes and the category of rigid analytic spaces over \mathbb{Q}_p .

Now, let $\underline{k} = (k_1, k_2) \in \mathbb{Z}^2$, any morphism $\underline{k} : (\mathbb{Z}_p^\times)^2 \rightarrow \mathbb{Q}_p^\times$ sending $(z_1, z_2) \rightarrow z_1^{k_1} \cdot z_2^{k_2}$ give a point of $\mathcal{W}(\mathbb{Q}_p)$ and which denote again by \underline{k} , and we call it “an algebraic weight”. More generally, any character of $\mathcal{W}(\mathbb{C}_p)$ which is a product of a character $\underline{k} \in \mathbb{Z}^2 \subset \mathcal{W}(\mathbb{Q}_p)$ with a finite character $\chi : (\mathbb{Z}_p^\times)^2 \rightarrow \overline{\mathbb{Q}}_p^\times$ is called ”an arithmetic character” and denoted by (\underline{k}, χ) .

LEMMA B.2. — *The classical weights \mathbb{Z}^2 of $\mathcal{W}(\mathbb{Q}_p)$ are very Zariski dense in the weight space \mathcal{W} .*

Proof. — It follows from the Weierstrass preparation theorem that the set \mathbb{Z}^2 of integral weights is Zariski-dense in \mathcal{W} . Moreover, the p -adic topology on the union of open discs \mathcal{W} induces by restriction the topology on \mathbb{Z}^2 for which we have a natural basis of neighborhood of $\underline{k} \in \mathbb{Z}^2$ given by the congruence classes modulo $p^n(p - 1)$ for all n . Hence \mathbb{Z}^2 is very Zariski dense. □

B.2. Geometric Siegel cuspforms

Let G denote the algebraic group GSp_4 and $\Gamma(N)$ be the open compact subgroup of $G(\widehat{\mathbb{Z}})$ of level N given by $\{\gamma \in G(\widehat{\mathbb{Z}}) \mid \gamma = \mathbb{1}_4 \pmod N\}$.

Assume now that $N \geq 5$, and let $X/\mathbb{Z}_p^{(15)}$ be the Siegel scheme of level $\Gamma(N) \cap I_1$, where I_r is the standard Iwahoric at p of G given by $\{\gamma \in \mathrm{GSp}_4(\mathbb{Z}_p) \mid \gamma \pmod{p^r} \in B(\mathbb{Z}/p^r\mathbb{Z})\}$ and B is the Borel of GSp_4 . There exists a universal abelian scheme A/X with identity section e and we let $\omega := e^*(\Omega_{A/X})$ be the conormal sheaf. Note that ω is a locally free sheaf of rank 2 over X . Let \overline{X} denote a toroidal compactification of X (it is not unique and depends on a combinatorial choice, see [17]), \overline{A} be the semi-abelian scheme extending A to \overline{X} and $D = \overline{X}/X$ be the normal crossing divisor at infinity. The sheaf ω extends to a locally free sheaf of rank 2 over \overline{X} , which we again denote by ω .

The classical cuspidal Siegel forms of level $\Gamma(N) \cap I_1$ and weight $k = (k_1, k_2)$ and coefficients in a p -adic field L (we have $k_1 \geq k_2$) are the elements of $H^0(\overline{X}_L, \omega_L^k(-D))^{(16)}$, where ω^k is the locally free sheaf

⁽¹⁵⁾The generic fiber X/\mathbb{Q}_p is smooth, and the special fiber X/\mathbb{F}_p is singular, and it even has vertical components.

⁽¹⁶⁾It follows from Koecher principle that $H^0(\overline{X}_L, \omega_L^k)$ does not depend on the choice of the toroidal compactification \overline{X} of X .

$\text{Sym}^{k_1-k_2} \omega \otimes \det \omega^{k_2}$, and ω_L^k is the base change of ω^k to \overline{X}_L . Let $\overline{X}^{\text{rig}}/\mathbb{Q}_p$ be the rigid analytic space given by taking the *generic fiber* of the formal scheme given by the completion of \overline{X} along its special fiber, and writing again ω for the analytification of ω , and let $\overline{X}^{\text{ord}}$ be the multiplicative ordinary locus of $\overline{X}^{\text{rig}}$ (it is not an affinoid), then the p -adic (resp. v -overconvergent) cuspidal Siegel modular forms of tame level $\Gamma(N)$ weight $k = (k_1, k_2)$ and coefficients in L are $H^0(\overline{X}_L^{\text{ord}}, \omega_L^k(-D))$ (resp. $H^0(\overline{X}_L^{\text{ord}}(v), \omega_L^k(-D))$), where $\overline{X}(v)$ is the v -overconvergent neighborhood of the multiplicative ordinary locus $\overline{X}^{\text{ord}}$.

Andreatta, Iovita and Pilloni constructed for any weight $k \in \mathcal{W}(\mathbb{C}_p)$ and certain parameters $v, w \in \mathbb{R}_+^\times$ a Banach sheaf ω_w^k over $\overline{X}(v)$, and a natural sheaf monomorphism $\omega^k \hookrightarrow \omega_w^k$ when $k = (k_1, k_2)$ is classical (see [1]), and they describe precisely the cokernel of that monomorphism. The sheaf ω_w^k is isomorphic locally for the étale topology to the w -analytic induction of the Borel $B(\mathbb{Z}_p)$ to the Iwahoric of GL_2 with respect to the character k . Note that any character $k \in \mathcal{W}(\mathbb{C}_p)$ is locally analytic by [1, Section 2.2] and hence ω_w^k is a non-zero Banach sheaf (The sections of ω_w^k are congruent to the image of the Hodge–Tate map by [1, Proposition 4.3.1]).

The p -adic modular forms obtained by this interpolation are locally analytic overconvergent (not necessarily overconvergent), however those satisfying the slope condition of [1, Theorem 7.1.1] are overconvergent (see also [1, Proposition 2.5.1.] and [1, Proposition 7.2.1]). Note that this construction is independent of the choice of the toroidal compactification of \overline{X} (see [27, Theorem 1.6.1] and [1, Proposition 5.5.2]) and we denote the corresponding eigenvariety by \mathcal{E}_N .

B.3. Local charts of the variety \mathcal{E}_N and density of classical points of \mathcal{E}_N

Let $\chi : (\mathbb{Z}/p\mathbb{Z}^\times)^2 \rightarrow \mathbb{Q}_p^\times$ be a character, \mathcal{W}^\times the connected component of \mathcal{W} corresponding to χ , and \mathcal{E}_N^\times the union of connected components of \mathcal{E}_N given by the restriction of \mathcal{E}_N to \mathcal{W}^\times .

For $w, v \in \mathbb{R}$ let $W = \text{Spm } R$ be a small enough affinoid subdomain of \mathcal{W}^\times to ensure the existence of the Banach sheaf $\omega_w^\kappa(-D)$ of $\overline{X}(v) \times \text{Spm } R$ interpolating the Banach sheaf $\omega_w^k(-D)$ of w -analytic v -overconvergent Siegel cusp form when k varies in $\text{Spm } R$ (κ denotes here the tautological character $(\mathbb{Z}_p^\times)^2 \rightarrow R^\times$).

On the other hand, let S_κ^\dagger be the Frechet R -module of ϵ -overconvergent cuspidal Siegel families over the affinoid R and given by

$$\varinjlim_{\substack{v \rightarrow 0, \\ w \rightarrow \infty}} H^0(\overline{X}(v) \times \text{Spm } R, \omega_w^\kappa(-D)).$$

The action of the Hecke operator $\mathbb{U} = U_0.U_1$ is completely continuous on the Frechet R -module S_κ^\dagger . Let $\mathcal{T}_{W,r}$ be the image of the Hecke algebra generated over R by the image of \mathcal{H}_N in the space of endomorphisms of $S_\kappa^{\dagger, v \leq r}$, where $S_\kappa^{\dagger, \leq r}$ is the R -finite submodule of S_κ^\dagger of slope at most r for $\mathbb{U} = U_0.U_1$.⁽¹⁷⁾ It follows from the results of [4, Section II] that

$$(B.1) \quad \mathcal{E}_{N,W}^r := \text{Spm } \mathcal{T}_{W,r},$$

is an affinoid subdomain of \mathcal{E}_N and by construction $\mathcal{E}_{N,W}^r$ is finite and torsion-free over W and the $\{\mathcal{E}_{N,W}^r\}$ form an admissible covering of \mathcal{E} .

Since the ordinary locus of any toroidal compactification of the Siegel modular scheme is not an affinoid, we cannot prove that the specialization

$$H^0(\overline{X}(v) \times_{\overline{\mathbb{Q}}_p} \text{Spm } R, \omega_w^\kappa) \rightarrow H^0(\overline{X}(v), \omega_w^k)$$

is surjective and that $H^0(\overline{X}(v) \times \text{Spm } R, \omega_w^\kappa)$ is a projective R -Banach module. However, Andreatta–Iovita–Pilloni proved in [1, Proposition 8.2.3.3] a control theorem for cuspidal families and that $H^0(\overline{X}(v) \times \text{Spm } R, \omega_w^\kappa(-D))$ is a projective R -Banach module, by projecting the sheaf $\omega_w^\kappa(-D)$ to the minimal compactification of the Siegel modular scheme, and using the fact that small v -overconvergent neighborhoods of the multiplicative ordinary locus of the minimal compactification of the Siegel modular scheme are affinoid spaces, and the deep descent result [1, Proposition 8.2.2.4].

Skinner–Urban constructed in [41, Section 2] a semi-ordinary eigenvariety $\mathcal{E}_N^{|U_0|_p=1} \subset \mathcal{E}_N$ for overconvergent Siegel cusp forms of tame level $\Gamma(N)$ and genus 2 by interpolating the locally free sheaf ω^k inside a Banach sheaf ω_w^κ over the weight space \mathcal{W} using the Igusa tower. That construction is a special case of the construction given by Andreatta–Iovita and Pilloni in [1] of the eigenvariety \mathcal{E}_N , since the linearization of the Hodge–Tate map

$$\text{HT}_{H_n^D} : H_n^D \rightarrow \omega_{H_n}$$

is surjective on the multiplicative ordinary locus ($H_n \subset \overline{A}$ is the level n canonical subgroup and H_n^D is its Cartier dual), and the fact that any semi-ordinary (i.e of slope 0 for U_0) p -adic Siegel cuspforms of finite slope for U_1 overconverges to a strict neighborhood of the ordinary locus. For the

⁽¹⁷⁾Note that the action of \mathbb{U} is completely continuous on S_κ^\dagger , so we have a slope decomposition.

latter note that under the iteration of the Hecke correspondances at p , an overconvergent neighborhood of X^{ord} accumulates around the multiplicative ordinary locus X^{ord} . The correspondence U_0 improves the radius of overconvergence. Hence, the functional equation $U_0.g = U_0(g).g$ allows us to extend g to a bigger neighborhood of the multiplicative ordinary locus when $U_0(f) \neq 0$ (the function degree of [36, Theorem 3.1] increases under the iteration of U_0). Meanwhile, one can use a similar functional equation for U_1 to get classicality at the level of the sheaves when the slope satisfies the condition of [1, Proposition 7.3.1].

By construction of \mathcal{E}_N we have an algebra homomorphism $\mathcal{H}_N \rightarrow \mathcal{O}_{\mathcal{E}_N}^{\text{rig}}(\mathcal{E}_N)$, and the image lands in the subring $\mathcal{O}_{\mathcal{E}_N}^{\text{rig}}(\mathcal{E}_N)^+$ given by the global section bounded by 1 on \mathcal{E}_N . Therefore, the canonical application “system of eigenvalues” induces a correspondence between the systems of eigenvalues for Hecke operators occuring in \mathcal{H}_N of locally analytic overconvergent cuspidal Siegel eigenforms of tame level $\Gamma(N)$ and weight $k \in \mathcal{W}(\mathbb{C}_p)$ having non-zero \mathbb{U} -eigenvalue, and the set of \mathbb{C}_p -valued points of weight $k = (k_1, k_2)$ on the Siegel eigenvariety \mathcal{E}_N . Note that for any overconvergent form g corresponding to a point of \mathcal{E}_N of weights (l_1, l_2) ,

$$(B.2) \quad g | U_1 = p^{l_2-3}U_1(g).g;$$

we renormalize U_1 in the aim to have a good p -adic interpolation (see for example [41, Theorem 2.4.14]).

One has the following Lemmas proving the very Zariski density of the classical points having a crystalline representation at p in \mathcal{E}_N , which is important for applying further the results of [7, Section 4] (see the hypothesis (HT) of [7, Section 3.3.2]).

LEMMA B.3. — *Let $z \in \mathcal{E}_N$ be a classical point old at p , then there exists an affinoid neighborhood Ω of z in \mathcal{E}_N of constant slopes for U_0, U_1 and such that the old at p classical points of regular weights of Ω are very Zariski dense in it, $\kappa(\Omega)$ is an open affinoid subdomain of \mathcal{W} , and each irreducible component of Ω surjects to $\kappa(\Omega)$.*

Proof. — Note that \mathcal{E}_N is admissibly covered by $\{\mathcal{E}_{N,W}^r\}$. Hence, there exists an affinoid subdomain $\mathcal{E}_{N,W}^r$ of \mathcal{E}_N containing z and surjecting on the affinoid subdomain $W \subset \mathcal{W}$. By construction of \mathcal{E}_N , the slopes of U_0, U_1 are locally constant. Then Propositions A.1 and A.6 yields that we can shrink $\mathcal{E}_{N,W}^r$ to a smaller open affinoid subdomain Ω of \mathcal{E}_N containing z and with constant slope S_1 (resp. S_2) for the Hecke operator U_0 (resp. U_1) and such that $\kappa(\Omega)$ is an open affinoid subdomain of \mathcal{W} , and $\kappa : \Omega \rightarrow \kappa(\Omega)$ is finite

and torsion-free (so the restriction of κ to any irreducible component of Ω is surjective by Proposition A.6).

Since Ω contains the classical point z , then the points of Ω with weights satisfying the small slope conditions of [1, Theorem 7.1.1] form a Zariski dense set in Ω , because the algebraic points (l_1, l_2) of $\kappa(\Omega)$ satisfying the inequality of the small slope conditions of [1, Theorem 7.1.1] form a Zariski dense set of $\kappa(\Omega)$ (so their preimage is dense in Ω by Proposition A.6). Moreover, it follows from the criterion of classicality of overconvergent forms that the points satisfying the small slope conditions of [1, Theorem 7.1.1] are necessarily classical. Actually, Propositions A.1 and A.6, and Lemma B.2 show that classical points of Ω are very Zariski-dense in it. Finally, the assertion follows from the fact that the classical points of Ω with sufficiently regular weights satisfy the assumptions of [41, Theorem 2.4.17], and hence they are old at p . \square

COROLLARY B.4. — Let $\mathcal{E}_N^{\text{ord},1}$ be the admissible open of \mathcal{E}_N defined by

$$\mathcal{E}_N^{\text{ord},1} := \{z \in \mathcal{E}_N, |U_0(z)|_p = 1, |U_1(z)|_p = p^{-1}\},$$

$C \in \mathbb{N}_{>1}$, and Σ_C be the set of points of $\mathcal{E}_N^{\text{ord},1}$ of “algebraic weights” (k_1, k_2) satisfying $k_1 > k_2 + C \geq \text{Max}(9, C)$. Then:

- (1) The overconvergent cuspforms of Σ_C are classical and old at p .
- (2) The set Σ_C is very Zariski dense in $\mathcal{E}_N^{\text{ord},1}$.
- (3) The point π_α of $\mathcal{E}_N^{\text{ord},1}$ corresponding to π_α is an accumulation point of Σ_C .


Proof. — The points of Σ_C have slope equal to 1, Iwahoric level at p and satisfy the slope condition $1 < k_1 - k_2 + 1, k_2 \gg 0$ of the classicality criterion for overconvergent Siegel cuspforms. Hence they are necessarily classical. A direct computation shows that the points of Σ_C satisfy the assumptions of [41, Theorem 2.4.17], and hence they are necessarily old at p .

Since the algebraic weights (k_1, k_2) with $k_1 > k_2 + C \geq \text{Max}(9, C)$ are very Zariski dense in \mathcal{W} (see Lemma B.2), the assertion of (2) and (3) follows directly from the argument already used to proof Lemma B.3. \square

B.4. Siegel eigenvariety of paramodular level N

Let \mathcal{E}_Δ be the Siegel eigenvariety of tame level the paramodular group Δ . Since the classical Siegel cuspforms of level $\Delta \cap I_1$ are necessarily of level $\Gamma(N) \cap I_1$, the results of [4, II.5] yields that there exists a natural closed

immersion $\iota : \mathcal{E}_\Delta \hookrightarrow \mathcal{E}_N$ compatible with the system of Hecke eigenvalues and the weights:



Since the restricted Hecke algebra \mathcal{H}_{Np} generated over \mathbb{Z} by the Hecke operators $T_{\ell,1}, T_{\ell,2}, S_\ell$ for $\ell \nmid Np$ acts semi-simply on classical cuspidal Siegel paramodular eigenforms of cohomological weights, [4, Lemma I.9.1] implies that \mathcal{E}_Δ is reduced. Note also that \mathcal{E}_Δ is equidimensional of dimension 2.

COROLLARY B.5. — Let $\mathcal{E}_\Delta^{\text{ord},1}$ be the admissible open of \mathcal{E}_Δ defined by

$$\mathcal{E}_\Delta^{\text{ord},1} := \{z \in \mathcal{E}_\Delta, |U_0(z)|_p = 1, |U_1(z)|_p = p^{-1}\},$$

$C \in \mathbb{N}_{>1}$, and Σ_C be the set of points of $\mathcal{E}_\Delta^{\text{ord},1}$ of “algebraic weights” (k_1, k_2) satisfying $k_1 > k_2 + C \geq \text{Max}(9, C)$. Then:

- (1) The overconvergent cuspforms of Σ_C are classical and old at p .
- (2) The set Σ_C is very Zariski dense in $\mathcal{E}_\Delta^{\text{ord},1}$.
- (3) The point π_α of $\mathcal{E}_\Delta^{\text{ord},1}$ is an accumulation point of Σ_C .

Proof. — It follows immediately from Corollary B.4 and the fact that a subset of an affinoid space is a Zariski dense if and only if its intersection with any irreducible component is Zariski dense in that irreducible component (see [16, Proposition 2.2.8]). □

Appendix C. Some examples where

$$\dim H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$$

Using Nekovar’s result [34, Proposition 4.2.3] about \mathbb{L} -adic Selmer groups mentioned before Corollary 6.7 we can exhibit infinitely many examples of modular forms f of weight $k \geq 3$ such that they satisfy the condition $\dim H_{f,\text{unr}}^1(\mathbb{Q}, \rho_f(k-1)) = 1$ in Theorem 6.8. This requires finding suitable elliptic curves with ordinary reduction at p and considering their corresponding Hida family \mathcal{F} . One such example is discussed in [9, Section 9.1], where for $p = 5$ and $N = 731$ the residual Selmer group

$$H_{f,\text{unr}}^1(\mathbb{Q}, \bar{\rho}_{E,p}(1)) = H_{f,\text{unr}}^1(\mathbb{Q}, \rho_{E,p}(1) \otimes \mathbb{Q}_p/\mathbb{Z}_p)[p] = \text{Sel}_p(E)[p]$$

of the rank 1 elliptic curve E (Cremona label 731a1) is calculated to have order 5 (since the order of vanishing of $L(f, s)$ at $s = 1$ is one we know that the BSD conjecture holds). This elliptic curve has non-split reduction at

both primes dividing N and good ordinary reduction at 5, with $a_5(E) = -1$ and therefore $\alpha \neq 1$. In addition this example satisfied the condition $L_p(f_\alpha, \omega_p^{-1}, T = p) \neq 0$.

In the following assume that f is the p -ordinary stabilization of the weight two cuspform attached to a rank 1 elliptic curve E/\mathbb{Q} . Recall that \mathbb{I} is the finite flat extension of $\mathbb{Z}_p[[T]]$ generated by the Fourier coefficients of the Hida family \mathcal{F} specializing to f (\mathbb{I} is an integral domain).

Note that the cohomology groups $H_{f, \text{unr}}^i(G_{\mathbb{Q}}^{Np}, \rho_{\mathcal{F}} \otimes \chi_{\text{univ}}^{-1/2})$ of the Selmer complex are of finite type over \mathbb{I} when $i \in \{1, 2\}$ (see [34, Proposition 4.2.3]).

Let $\mathcal{P}_f \subset \mathbb{I}$ be the height one prime ideal corresponding to the system of Hecke eigenvalues of f . We have the following control theorem proved by Nekovar [34, (0.15.1.1)]

$$(C.1) \quad 0 \rightarrow H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2})) \otimes \mathbb{I}_{\mathcal{P}_f}/\mathcal{P}_f \rightarrow H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(1)) \rightarrow H_{f, \text{unr}}^2(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}) \otimes \mathbb{I}_{\mathcal{P}_f})[\mathcal{P}_f].$$

where $H_{f, \text{unr}}^2(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}) \otimes \mathbb{I}_{\mathcal{P}_f})[\mathcal{P}_f]$ means the submodule annihilated by the prime ideal \mathcal{P}_f .

Since $\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_f(1)) = 1$ Nakayama’s lemma applied to (C.1) yields that the $\mathbb{I}_{\mathcal{P}_f}$ -module $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}) \otimes \mathbb{I}_{\mathcal{P}_f})$ is monogenic. Moreover, it follows from Corollary 6.7 that $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}) \otimes \mathbb{I}_{\mathcal{P}_f})$ is a torsion-free $\mathbb{I}_{\mathcal{P}_f}$ -module, so

$$H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}) \otimes \mathbb{I}_{\mathcal{P}_f}) = H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2})) \otimes_{\mathbb{I}} \mathbb{I}_{\mathcal{P}_f}$$

is a free rank one $\mathbb{I}_{\mathcal{P}_f}$ -module. Thus there exists a principal Zariski open $D(s)$ of $\text{Spec } \mathbb{I}$ (where $s \in \mathbb{I}$) such that the localization of $H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}))$ at the non-vanishing locus $D(s)$ is a free rank one $\mathbb{I}[1/s]$ -module. On the other hand, let $\mathcal{U} \subset D(s)$ be the Zariski open defined as the complementary of the support of the \mathbb{I} -torsion part of $H_{f, \text{unr}}^2(\mathbb{Q}, \rho_{\mathcal{F}}(\chi_{\text{univ}}^{-1/2}))$. Note that the classical points of \mathcal{U} are Zariski dense, hence (C.1) yields that all the classical specialization \mathcal{F}_z of the Hida family \mathcal{F} at a point $z \in \mathcal{U}$ of weight k_z satisfy

$$\dim H_{f, \text{unr}}^1(\mathbb{Q}, \rho_{\mathcal{F}_z}(k - 1)) = 1.$$

BIBLIOGRAPHY

[1] F. ANDREATTA, A. IOVITA & V. PILLONI, “ p -adic families of Siegel modular cuspforms”, *Ann. Math.* **181** (2015), no. 2, p. 623-697.
 [2] J. ARTHUR, “Automorphic representations of $\text{GSp}(4)$ ”, in *Contributions to automorphic forms, geometry, and number theory*, Johns Hopkins University Press, 2004, p. 65-81.

- [3] J. BELLAÏCHE, “Nonsmooth classical points on eigenvarieties”, *Duke Math. J.* **145** (2008), no. 1, p. 71-90.
- [4] ———, “Eigenvarieties and p -adic L -functions”, book in preparation, <http://people.brandeis.edu/~jbellaiic/preprint/coursebook.pdf>, 2010.
- [5] ———, “Critical p -adic L -functions”, *Invent. Math.* **189** (2012), no. 1, p. 1-60.
- [6] J. BELLAÏCHE & G. CHENEVIER, “Lissité de la courbe de Hecke de GL_2 aux points Eisenstein critiques”, *J. Inst. Math. Jussieu* **5** (2006), no. 2, p. 333-349.
- [7] ———, *Families of Galois representations and Selmer groups*, Astérisque, vol. 324, Société Mathématique de France, 2009, xii+314 pages.
- [8] J. BELLAÏCHE & M. DIMITROV, “On the eigencurve at classical weight 1 points”, *Duke Math. J.* **165** (2016), no. 2, p. 245-266.
- [9] T. BERGER & K. KLOSIN, “Deformations of Saito–Kurokawa type and the paramodular conjecture”, *Am. J. Math.* **142** (2020), no. 6, p. 1821-1876.
- [10] V. G. BERKOVICH, “Étale cohomology for non-Archimedean analytic spaces”, *Publ. Math., Inst. Hautes Étud. Sci.* (1993), no. 78, p. 5-161.
- [11] A. BETINA & M. DIMITROV, “Geometry of the eigencurve at CM points and trivial zeros of Katz p -adic L -functions”, <https://arxiv.org/abs/1907.09422>, to appear in *Adv. Math.* **384** (2021).
- [12] A. BETINA, M. DIMITROV & A. POZZI, “On the failure of Gorensteinness at weight 1 Eisenstein points of the eigencurve”, <https://arxiv.org/abs/1804.00648>, to appear in *Am. J. Math.*
- [13] G. CHENEVIER, “Familles p -adiques de formes automorphes pour GL_n ”, *J. Reine Angew. Math.* **570** (2004), p. 143-217.
- [14] R. F. COLEMAN, F. Q. GOUVÊA & N. JOCHNOWITZ, “ E_2 , Θ , and overconvergence”, *Int. Math. Res. Not.* (1995), no. 1, p. 23-41.
- [15] R. F. COLEMAN & B. MAZUR, “The eigencurve”, in *Galois representations in arithmetic algebraic geometry (Durham, 1996)*, London Mathematical Society Lecture Note Series, vol. 254, Cambridge University Press, 1998, p. 1-113.
- [16] B. CONRAD, “Irreducible components of rigid spaces”, *Ann. Inst. Fourier* **49** (1999), no. 2, p. 473-541.
- [17] G. FALTINGS & C.-L. CHAI, *Degeneration of abelian varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 22, Springer, 1990, With an appendix by David Mumford, xii+316 pages.
- [18] J.-M. FONTAINE (ed.), *Périodes p -adiques*, Astérisque, vol. 223, Société Mathématique de France, 1994, Papers from the seminar held in Bures-sur-Yvette, 1988.
- [19] T. GEE & O. TAÏBI, “Arthur’s multiplicity formula for \mathbf{GSp}_4 and restriction to \mathbf{Sp}_4 ”, *J. Éc. Polytech., Math.* **6** (2019), p. 469-535.
- [20] V. HERNANDEZ, “Families of Picard modular forms and an application to the Bloch–Kato conjecture”, *Compos. Math.* **155** (2019), no. 7, p. 1327-1401.
- [21] H. HIDA, “Galois representations into $GL_2(\mathbf{Z}_p[[X]])$ attached to ordinary cusp forms”, *Invent. Math.* **85** (1986), no. 3, p. 545-613.
- [22] ———, “Control theorems of coherent sheaves on Shimura varieties of PEL type”, *J. Inst. Math. Jussieu* **1** (2002), no. 1, p. 1-76.
- [23] K. KATO, “ p -adic Hodge theory and values of zeta functions of modular forms”, in *Cohomologies p -adiques et applications arithmétiques. III*, Astérisque, vol. 295, Société Mathématique de France, 2004, p. ix, 117-290.
- [24] M. KISIN, “Overconvergent modular forms and the Fontaine-Mazur conjecture”, *Invent. Math.* **153** (2003), no. 2, p. 373-454.
- [25] ———, “Geometric deformations of modular Galois representations”, *Invent. Math.* **157** (2004), no. 2, p. 275-328.

- [26] G. LAUMON, “Fonctions zêtas des variétés de Siegel de dimension trois”, in *Formes automorphes. II. Le cas du groupe $GSp(4)$* , Astérisque, vol. 302, Société Mathématique de France, 2005, p. 1-66.
- [27] W. LÜTKEBOHMERT, “Der Satz von Remmert-Stein in der nichtarchimedischen Funktionentheorie”, *Math. Z.* **139** (1974), p. 69-84.
- [28] D. MAJUMDAR, “Geometry of the eigencurve at critical Eisenstein series of weight 2”, *J. Théor. Nombres Bordeaux* **27** (2015), no. 1, p. 183-197.
- [29] B. MAZUR, “An introduction to the deformation theory of Galois representations”, in *Modular forms and Fermat’s last theorem (Boston, MA, 1995)*, Springer, 1997, p. 243-311.
- [30] T. MIYAKE, *Modular forms*, Springer, 1989, Translated from the Japanese by Yoshitaka Maeda, x+335 pages.
- [31] M. MIYAUCHI & T. YAMAUCHI, “An explicit computation of p -stabilized vectors”, *J. Théor. Nombres Bordeaux* **26** (2014), no. 2, p. 531-558.
- [32] C. P. MOK, “Galois representations attached to automorphic forms on GL_2 over CM fields”, *Compos. Math.* **150** (2014), no. 4, p. 523-567.
- [33] J. NEKOVÁŘ, “On p -adic height pairings”, in *Séminaire de Théorie des Nombres, Paris, 1990–91*, Progress in Mathematics, vol. 108, Birkhäuser, 1993, p. 127-202.
- [34] ———, *Selmer complexes*, Astérisque, vol. 310, Société Mathématique de France, 2006, viii+559 pages.
- [35] T. OCHIAI, “Control theorem for Greenberg’s Selmer groups of Galois deformations”, *J. Number Theory* **88** (2001), no. 1, p. 59-85.
- [36] V. PILLONI, “Prolongement analytique sur les variétés de Siegel”, *Duke Math. J.* **157** (2011), no. 1, p. 167-222.
- [37] A. SAHA & R. SCHMIDT, “Yoshida lifts and simultaneous non-vanishing of dihedral twists of modular L -functions”, *J. Lond. Math. Soc.* **88** (2013), no. 1, p. 251-270.
- [38] R. SCHMIDT, “On classical Saito-Kurokawa liftings”, *J. Reine Angew. Math.* **604** (2007), p. 211-236.
- [39] ———, “Packet structure and paramodular forms”, *Trans. Am. Math. Soc.* **370** (2018), no. 5, p. 3085-3112.
- [40] ———, “Paramodular forms in CAP representations of $GSp(4)$ ”, *Acta Arith.* **194** (2020), no. 4, p. 319-340.
- [41] C. SKINNER & E. URBAN, “Sur les déformations p -adiques de certaines représentations automorphes”, *J. Inst. Math. Jussieu* **5** (2006), no. 4, p. 629-698.
- [42] ———, “The Iwasawa main conjectures for GL_2 ”, *Invent. Math.* **195** (2014), no. 1, p. 1-277.
- [43] C. M. SORENSEN, “Galois representations attached to Hilbert-Siegel modular forms”, *Doc. Math.* **15** (2010), p. 623-670.
- [44] R. TAYLOR, “On the l -adic cohomology of Siegel threefolds”, *Invent. Math.* **114** (1993), no. 2, p. 289-310.
- [45] J. TILOUINE & E. URBAN, “Several-variable p -adic families of Siegel-Hilbert cusp eigensystems and their Galois representations”, *Ann. Sci. Éc. Norm. Supér.* **32** (1999), no. 4, p. 499-574.
- [46] E. URBAN, “Selmer groups and the Eisenstein-Klingen ideal”, *Duke Math. J.* **106** (2001), no. 3, p. 485-525.
- [47] ———, “Sur les représentations p -adiques associées aux représentations cuspidales de GSp_4/\mathbb{Q} ”, in *Formes automorphes. II. Le cas du groupe $GSp(4)$* , Astérisque, vol. 302, Société Mathématique de France, 2005, p. 151-176.
- [48] R. WEISSAUER, “Four dimensional Galois representations”, in *Formes automorphes. II. Le cas du groupe $GSp(4)$* , Astérisque, vol. 302, Société Mathématique de France, 2005, p. 67-150.

- [49] T. WESTON, “Geometric Euler systems for locally isotropic motives”, *Compos. Math.* **140** (2004), no. 2, p. 317–332.

Manuscrit reçu le 15 août 2019,
révisé le 26 février 2020,
accepté le 13 mai 2020.

Tobias BERGER
School of Mathematics and Statistics
University of Sheffield
Hicks Building, Hounsfield Road
Sheffield S3 7RH (UK)
tberger@cantab.net

Adel BETINA
Faculty of Mathematics
University of Vienna
Oskar-Morgenstern-Platz 1
A-1090 Wien (Austria)
adelbetina@gmail.com