

**Promoting and maintaining changes in smoking behaviour for patients following discharge from a smokefree mental health inpatient stay: Development of a complex intervention using the Behaviour Change Wheel**

Emily Shosmith, PhD<sup>1\*</sup>, Lisa Huddleston, PhD<sup>1</sup>, Jodi Pervin, BSc<sup>1</sup>, Lion Shahab, PhD<sup>2</sup>, Peter Coventry, PhD<sup>3</sup>, Tim Coleman, MD<sup>4</sup>, Fabiana Lorencatto, PhD<sup>5</sup>, Simon Gilbody, PhD<sup>1,6</sup>, Moira Leahy<sup>7</sup>, Michelle Horspool, PhD<sup>7</sup>, Claire Paul<sup>8</sup>, Lesley Colley<sup>9</sup>, Simon Hough<sup>10</sup>, Phil Hough<sup>10</sup>, & Elena Ratschen, PhD<sup>1</sup>

<sup>1</sup> Department of Health Sciences, University of York, Heslington, York, UK

<sup>2</sup> Department of Behavioural Science and Health, University College London, London, UK

<sup>3</sup> York Environmental Sustainability Institute, University of York, UK

<sup>4</sup> Centre for Academic Primary Care, University of Nottingham, Nottingham, UK

<sup>5</sup> Centre for Behaviour Change, University College London, London, UK

<sup>6</sup> Hull York Medical School, University of York, UK

<sup>7</sup> Sheffield Health and Social Care NHS Foundation Trust, UK

<sup>8</sup> Leeds and York Partnership NHS Foundation Trust, UK

<sup>9</sup> Tees, Esk & Wear Valleys NHS Foundation Trust, UK

<sup>10</sup> Independent Peer Researcher, UK

\* Corresponding author: Emily Shosmith, email: emily.shoosmith@york.ac.uk, Department of Health Sciences, University of York, Heslington, York, YO10 5DD

© The Author(s) 2022. Published by Oxford University Press on behalf of the Society for Research on Nicotine and Tobacco.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

## **ABSTRACT**

### **Background and aims**

Evidence suggests that smokers can successfully quit, remain abstinent or reduce smoking during a smokefree mental health inpatient stay, provided behavioural/pharmacological support are offered. However, few evidence-based strategies to prevent the return to pre-hospital smoking behaviours post-discharge exist. We report the development of an intervention designed to support smoking-related behaviour change following discharge from a smokefree mental health stay.

### **Method**

We followed the Behaviour Change Wheel (BCW) intervention development process. The target behaviour was supporting patients to change their smoking behaviours following discharge from a smokefree mental health stay. Using systematic reviews, we identified the barriers/enablers, classified according to the Theoretical Domains Framework (TDF). Potential intervention functions to address key influences were identified by consulting the BCW and Behaviour Change Technique (BCT) taxonomy. Another systematic review identified effectiveness of BCTs in this context. Stakeholder consultations were conducted to prioritise/refine intervention content.

### **Results**

Barriers/enablers to supporting smoking cessation were identified within the domains of environmental context and resources (lack of staff time); knowledge (ill-informed interactions about smoking); social influences, and intentions (lack of intention to deliver support). Potential strategies to address these influences included goal setting, problem solving, feedback, social support, and information on health consequences. A strategy for operationalising these techniques into intervention components was agreed: pre-discharge evaluation sessions, personalised resource folder, tailored behavioural and text message support post-discharge, and a peer interaction group, delivered by a trained mental health worker.

### **Conclusions**

The intervention includes targeted resources to support smoking-related behaviour change in patients following discharge from a smokefree mental health setting.

## Implications

Using the BCW and TDF supported a theoretically and empirically informed process to define and develop a tailored intervention that acknowledges barriers and enablers to supporting smoking cessation in mental health settings. The result is a novel complex theory- and evidence-based intervention that will be formally tested in a randomised controlled feasibility study.

Accepted Manuscript

## INTRODUCTION

Smoking remains a leading cause of mortality and morbidity worldwide<sup>1</sup>. Smoking prevalence in the general population in England has steadily declined from 14.1% in 2019 to 13.5% in 2020<sup>2</sup>, but data indicate that prevalence remains approximately two to three times higher among people with mental health conditions in the United Kingdom<sup>3</sup>. Smokers with mental health conditions are more likely to experience greater dependence on smoking, are more likely to develop smoking-related illnesses, and long-term quit rates among this population are lower than for the general adult population<sup>4-6</sup>. Although people with mental health conditions are just as motivated to quit as those in the general population<sup>4,7</sup>, they are less likely to receive the support they require compared with smokers without mental health conditions<sup>8</sup>.

Many healthcare settings are now smokefree by law and offer smoking cessation support for smokers during their stay<sup>9</sup>. Evidence suggests that individuals can successfully remain abstinent during their inpatient stay when behavioural/pharmacological support are offered<sup>10,11</sup>. Additionally, evidence from mental health settings indicates that a period of supported abstinence may promote smoking behaviour change and motivation to quit<sup>12</sup>. However, the risk of relapse post-discharge is high, with one study reporting 76% relapsed to smoking behaviours within one day post-discharge<sup>13</sup>. Lack of support offered post-discharge and the high risk of relapse renders smoking-related resource input during the inpatient episode inefficient, as positive smoking behaviour change achieved during the inpatient stay is often lost. Identifying interventions that are effective in supporting patients around the time of discharge in maintaining or achieving positive change to their smoking behaviours has been identified as an important evidence gap in this area<sup>14</sup>.

Designing interventions to support smoking cessation would benefit from drawing on theories and frameworks from behavioural science which summarise drivers of behaviours and link these to strategies for changing behaviour<sup>15</sup>. The Behaviour Change Wheel (BCW)<sup>16</sup> and Theoretical Domains Framework (TDF) are developments in the behaviour change field that provide a systematic, theoretical basis for understanding and providing tools to change behaviour<sup>17</sup>, based on multiple models of health behaviour. The process of intervention development using the BCW and TDF is outlined in detail<sup>16</sup> and has been extensively applied to design smoking cessation interventions<sup>18-20</sup>.

### Theoretical underpinnings

The BCW was developed from an extensive review of behavioural science frameworks, designed to facilitate the development of behaviour change interventions<sup>16,21</sup>. At the centre of the BCW is a behaviour change model known as the COM-B, which postulates that behaviour change is essentially dependent on the interaction between three key

determinants: the capability (C) to perform a behaviour, the opportunity (O) and motivation (M) to perform<sup>16</sup>. The TDF is an elaboration of the COM-B model and includes 14 domains based on an integration of behavioural theories<sup>22-24</sup>. These domains prompt the consideration of a wide range of influences as they include individual-level factors, social factors, and environmental factors<sup>17</sup>.

The COM-B and TDF have been mapped to frameworks which specify different intervention strategies. The first is the BCW which specifies nine intervention types for changing behaviour (e.g., education, training)<sup>16</sup>. These intervention functions are supported by seven policy categories, representing decision types that help to support the interventions (e.g., communication/marketing, guidelines)<sup>16</sup>. The second is the more granular taxonomy of Behaviour Change Techniques (BCTs), which breaks down these broad intervention types into 93 discrete BCTs (e.g., goal setting, action planning, feedback on behaviour)<sup>24</sup>. There are published tools which pair domains from the COM-B/TDF with intervention strategies in the BCW/BCT taxonomy (BCTv1)<sup>24</sup> to specify which types of behaviour change approaches are more likely to be relevant and effective in changing different influences on behaviour<sup>16</sup>. This provides a step-wise process from moving from an initial behavioural diagnosis to identifying 'what needs to change', to selecting specific intervention components in a theory- and evidence- informed manner<sup>17</sup>.

To our knowledge, the BCW and TDF have not been applied to the context of smoking cessation following discharge from a smokefree mental health inpatient stay. The aim of this study was to apply the BCW and TDF to systematically design an evidence- and theory-based complex intervention for people with a mental health condition to support smoking-related behaviour changes following a mental health inpatient stay. This process is in line with the Medical Research Council's (MRC) guidance for developing complex interventions<sup>15</sup>.

## **METHOD**

### **Aim**

To develop an evidence- and theory-based complex intervention to support smoking-related behaviour changes for patients following discharge from a smokefree mental health inpatient stay.

### **Research design**

Intervention development involved two sequential phases: (1) synthesising findings from existing evidence and mapping these onto the BCW, and (2) using qualitative and consensus methodologies to explore acceptability and feasibility of the prototype intervention (Figure 1).

## Phase 1: Content development

The process of intervention development using the BCW and TDF is outlined in detail elsewhere<sup>16</sup>, and a detailed explanation of how the process was followed is presented in Supplementary Material 1. A summary is provided:

### *Stage one: Understanding the behaviour and identifying 'what needs to change'*

The BCW was used to understand the healthcare problem and the behaviours that could be targeted by an intervention<sup>16</sup>.

To identify what was influencing behaviours and the barriers to/enablers of change, we conducted a systematic review of the published qualitative and quantitative literature investigating smoking cessation interventions in mental health settings. We extracted and thematically synthesised barriers and enablers. These were then coded to the domains of the COM-B and TDF. The detailed methods are published elsewhere<sup>25</sup>.

### *Stage two: Identifying intervention options*

We mapped identified influences on potential intervention strategies using the aforementioned mapping tools which pair COM-B/TDF to BCW/BCTs<sup>16</sup>. This resulted in a long list of potentially relevant intervention functions. The APEASE criteria were used to guide judgements as to what functions may be the most appropriate for the intervention<sup>16</sup>. To prioritise further, and identify what interventions have been done to date in this context, we also conducted a systematic review of behavioural/pharmacological interventions that maintain abstinence following a smokefree stay and determined their effectiveness<sup>11</sup>.

We then mapped the identified intervention functions to potential policy categories that are likely to be appropriate and effective in supporting each function<sup>16</sup>. The APEASE criteria were used to help prioritise amongst these potential categories.

### *Stage three: Identifying content and implementation options*

We considered all BCTs that could be considered for any particular function, using the aforementioned mapping tools<sup>16</sup>. This resulted in a long list of potentially appropriate BCTs. The APEASE criteria were used to select those most likely to be appropriate. To further guide selection, we coded existing intervention descriptions into component BCTs using the BCTTv1 and identified which BCTs were defined as 'promising' in terms of probable

effectiveness and feasibility. We have previously reported these methods in detail<sup>11</sup>. We determined the most appropriate mode of delivery.

We selected potential intervention strategies and produced a description of a prototype intervention which was then presented to key stakeholders and an expert group of academics and clinicians (Supplementary Material 2; Table 2a).

## **Phase 2: Iterative refinement based on stakeholder and expert group input**

Consultations were conducted with stakeholders and an expert group of academics and clinicians to identify potential problems with the prototype intervention, develop solutions and assess the perceived acceptability, practicability, and ease of integration.

### *Stakeholder consultations*

Settings and participants: Consultations were conducted with four North of England NHS Trusts. Stakeholders included: (1) users of mental health services; (2) informal caregivers, and (3) mental health workers in acute adult mental health inpatient and community mental health services. The term 'mental health worker' (MHW) refers to any mental healthcare professional or healthcare associate role, and the worker did not have to be professionally qualified.

Recruitment: MHWs were invited by their service leads or through direct contact with the research team in liaison with Trusts. Patient and caregiver stakeholders were invited to volunteer through direct contact by the Trust Patient and Public Involvement (PPI) lead; by responding to advertisements in Trust newsletters.

Methods: Consultations took place remotely with individuals or groups. The prototype intervention was presented to stakeholders, and a guided discussion informed by the BCW elicited detailed feedback. Additionally, online presentation software was used during the MHW consultations. This software used 5-point Likert scales to obtain ratings for individual intervention components in relation to practicality, acceptability to staff/patients/caregivers, and the perceived ease of integration into services. The 5-point Likert scale included response anchors at either end of the scale (1 = low; 5 = high).

Analysis: Intervention component ratings were summarised using descriptive statistics. Qualitative data from the guided discussions were exported to NVivo 12 software<sup>26</sup> and analysed using thematic analysis<sup>27</sup>, employing an inductive approach, in which coding and theme development were driven by the content of the responses. One author (E.S) familiarised herself with the data and notes were made of any potential codes by identifying



recurring words or units of meaning. The same author generated initial codes from the data and organised them into meaningful groups. Codes were then organised into potential themes and all relevant coded responses were collated within the identified themes. Three authors (E.S, L.H and J.P) independently reviewed the construction of themes and relevant quotations to agree to the assignment of themes. Inductive thematic analysis was used to identify and categorise potential barriers and enablers in relation to the development and implementation of the intervention, that were subsequently mapped to the COM-B model and facilitated further intervention refinement. The findings from this process were used to refine the prototype intervention (Supplementary Material 2; Table 2b).

#### *Expert advisory group assessment of prototype intervention*

The prototype intervention was presented during expert panel consensus meetings. The expert panel involved involving academic experts in tobacco control and intervention design (n = 13), and clinicians from participating Trusts (n = 4). All experts in the advisory group had been approached to be collaborators/advisors or co-applicants for the research before it commenced, and all accepted and provided feedback. Feedback was elicited through discussion and topics related to overall perceptions, delivery pathways and potential implementation issues. During the meetings, notes of key points were taken, and written summaries of each meeting were shared with the expert panel to ensure they were accurate overviews of the discussions. Final summaries were discussed in detail to support the refinement of the intervention. The feedback from this process was used to further refine the prototype intervention (Supplementary Material 2; Table 2c).

#### *Expert advisory group and PPI group final review*

Following stakeholder consultations and expert group meetings, the revised prototype was re-presented to the expert advisory panel for final feedback. The revised prototype was also presented to members of the research programmes PPI group. This PPI group was formed following the stakeholder consultations and includes current and ex-smokers who have experiences of an inpatient mental health admission. The materials required for the intervention were developed and reviewed by our PPI group and expert panel, to ensure these materials were appropriately worded, engaging and ease to use.



## RESULTS

### Phase 1: Content development

#### *Stage one: Understanding the behaviour and identifying what needs to change*

The target behaviour of interest was defined as: adult smokers in mental health settings changing their smoking-related behaviours following discharge from a smokefree mental health inpatient stay.

The findings of the systematic review to identify the barriers and enablers to influence the behaviour of interest have been published elsewhere<sup>25</sup>. In summary, the key barriers to patients making smoking-related behaviour changes in mental health settings fell within the following TDF domains: environmental context and resources (e.g., lack of staff time); knowledge (e.g., interactions were ill-informed); social influences (e.g., smoking norms), and intentions (e.g., lack of positive intentions). Key enablers mainly fell within the domains: environmental context and resources (e.g., use of appropriate support materials) and social influences (e.g., pro-quitting social norms) (Supplementary Material 3).

#### *Stage two: Identifying intervention options*

The COM-B/TDF domains were mapped to the nine intervention functions identified within the BCW (Supplementary Material 3). The intervention functions of education, training, and enablement were included, based upon an evaluation of their affordability, practicability, effectiveness, acceptability, side-effects and safety, and equity (APEASE)<sup>16</sup>. Details of each intervention function against the APEASE criteria are provided in Supplementary Material 4; Table 4a.

The intervention functions were then mapped to the policy categories identified within the BCW (Supplementary Material 3). The policy categories of communications/marketing, guidelines, and service provision were included, based upon an evaluation of the APEASE criteria. Details of the evaluation of each policy category against the APEASE criteria are provided in Supplementary Material 4; Table 4b.

### *Stage three: Identifying content and implementation options*

The intervention functions identified were then mapped to a potential long list of BCTs using the BCTTv1<sup>16</sup> (Supplementary Material 3). The APEASE criteria and findings from the second systematic review<sup>11</sup> were used to guide the identification of the most appropriate BCTs to use within the intervention (Supplementary Material 5). The full findings of the systematic review have been published elsewhere<sup>11</sup>. In summary, nine BCTs (including 'pharmacological support', 'goal setting (behaviour)' and 'social support') were characterised as 'promising' in terms of probable effectiveness and feasibility. A review of the full BCTTv1 identified additional BCTs: 'social award', 'self-reward', 'pharmacological support', and 'restructuring the social environment', which also met the APEASE criteria.

A blended approach encompassing face-to-face and remote delivery was chosen, as the pre-discharge evaluation session and provision of resources is planned to occur on the inpatient ward pre-discharge. Post-discharge, the intervention will be delivered remotely, including behavioural support calls via telephone/video call and optional text message support. This approach was considered in terms of the evidence in relation to remote interventions and in the context of Covid-19-related restrictions. Both delivery modes met the APEASE criteria.

A prototype intervention strategy was drafted (Supplementary Material 6), based on the existing evidence and selections made throughout the BCW process.

### **Phase 2: Iterative refinement based on stakeholder and expert group input**

#### *Stakeholder consultations*

Twenty-five stakeholders participated to assess the practicality, acceptability, and ease of integration of the prototype intervention. Key stakeholders included MHWs, patients with mental health conditions and one informal caregiver (Table 1).

The prototype intervention (Supplementary Material 2; Table 2a) was well-received, and most individual components were rated highly for practicality, acceptability, and ease of integration (mean scores ranging between 4 and 4.8). The identification of a 'buddy' to provide social support post-discharge. This component had a lower score in terms of practicality, acceptability, and ease of integration compared to other components (mean scores ranged between 3.2 to 4.6) and alternative components were explored. The MHWs ratings for individual intervention components are provided in Supplementary Material 7.

Overall, four themes were identified from the guided discussion with stakeholders (Supplementary Material 8). The largest number of barriers for MHWs were classified under physical opportunity, including beliefs about limited staff time and resources. From a patient and caregiver perspective, barriers were classified under reflexive motivation, which included beliefs about MHWs not holding positive intentions to support smoking cessation.

#### Theme one: Enablers to patient engagement and intervention delivery

In relation to current support, many MHWs identified the absence of a strong link between the inpatient ward and the community. They highlighted that to provide effective support post-discharge, there needed to be improved communication between inpatient/community teams and a seamless process of transferring patient information. For many patients, starting their journey during their stay and continuing this progress into the community was perceived as an important factor in their motivation to continue engaging with smoking cessation support. Patients expressed the importance of this continuum, as opposed to inpatient support and 'separate' community support. There was a consensus that the intervention would facilitate a smoother transition from inpatient to community and would enhance patient engagement. All MHWs acknowledged that the pre-discharge assessment was critical, and it was important that the information collated in this session had to be fed into the community to avoid duplication. All stakeholders believed providing the resource pack to patients pre-discharge would 'bridge the gap' between inpatient and community support, as patient information would be in one place from inpatient to community. All stakeholders also reported pre-discharge provision was advantageous as the patients would have a plan in place pre-discharge that would continue in the community.

All stakeholders believed that setting personalised goals and the provision of a resource pack at discharge would be beneficial. All stakeholders believed the motivational and practical content included in the pack would facilitate participant engagement. Lastly, remote support post-discharge was frequently met with approval, as receiving remote behavioural support calls quickly and easily would facilitate accessibility. It was important to all stakeholders that the behavioural support was delivered by the same MHW to build a good relationship with the patient.

#### Theme two: Potential barriers to intervention delivery

MHWs frequently mentioned that capacity and resources may be limited, as staff might often lack the time or opportunity to deliver support to patients alongside their existing workload. Additionally, there was a consensus across stakeholder groups that staff were often resistant to the idea of engaging with smoking cessation support, especially when staff members smoked themselves.

### Theme three: Overall perceptions

Most MHWs agreed that the pre-discharge session designed for goal setting should be a critical component of smoking cessation support. This session is designed to tap into an individual's wishes and needs to offer patients the most appropriate and tailored support. Patients also indicated the importance of flexibility within the goal setting session as every individual would have unique goals and motivations that would require regular review.

Stakeholders believed that the resource pack was an innovative and novel idea that would be well-received by patients, as patients are provided with little written information. MHWs felt that including resources on the benefits of quitting within the pack would be advantageous. Telephone behavioural support was identified as an important, essential component by all stakeholders. Many stakeholders believed this would be well-received if the support was individualised. Lastly, the text message component was well-received by all stakeholders, with most individuals suggesting these would maintain motivation during a quit attempt.

### Theme four: Practical considerations and suggestions

The prototype intervention included a 'buddy' who could offer social support to individuals. However, most stakeholders highlighted that patients' may not be able to identify a relative/friend who is able or willing to adopt this role. All stakeholders suggested it would be beneficial to replace this component with the formation of a social support group (Supplementary Material 2; Table 2b). By offering this opportunity, stakeholders believed this would enhance the support received from others experiencing a similar situation in an informal, supportive environment. One patient suggested that this group should be organised to encourage peer-to-peer support but have a MHW present to facilitate interactions.

For the resource pack, most stakeholders suggested that the inclusion of 'success stories' which illustrated positive behaviour changes made by others in similar situations would provide motivational support. Two patients suggested additional content could be included to outline alternative activities that may offer a distraction (Supplementary Material 2; Table 2b). It was noted this information should be presented in a creative, user-friendly manner. All stakeholders were positive about the acceptability and feasibility of using motivational text messages, but with the option to opt-in/-out should they be seen as overwhelming (Supplementary Material 2; Table 2b).

## *Expert advisory group assessment of prototype intervention*

The expert advisory group perceived the intervention to be acceptable and feasible but highlighted that it would be critical to identify an appropriate person to deliver it (Supplementary Material; Table 2c). Group members acknowledged that when identifying a MHW to deliver the intervention, this should not be restricted by role, but rather those who fulfil a certain criterion. For example, experience in working with mental health and have a positive outlook on supporting people with mental health conditions to change their smoking-related behaviours. It was highlighted that there is no evidence to suggest certain job roles are better than others in terms of their effectiveness in helping individuals to positively change their smoking behaviours. The expert group indicated that it would not be advisable to exclude potential facilitators due to their job title when they may be experienced and willing to deliver the intervention effectively. A document was created in collaboration with the expert panel to provide an overview of the MHW who would deliver the intervention, outlining the role description, person specification, and training requirements.

In the systematic review, 'pros and cons' and 'framing/reframing' were identified as 'promising' in terms of likely effectiveness, but not in terms of likely feasibility<sup>11</sup>. Clinicians from the expert group discussed the importance of including these BCTs, as they appeared particularly important in a mental health inpatient context to help the individual reflect on their smokefree experience and explore their motivation to remain smokefree or positively change their smoking behaviour. During the pre-discharge evaluation, the MHW could provide motivational enhancement and strategies for managing temptations, considering the pros and cons of change. Additionally, the patient could be presented with alternative ways of thinking and counterarguments to their belief barriers and discuss behavioural self-management strategies to counter triggers.

The expert group also discussed the opt-in/-out process for the text-messaging component that was included in line with the feedback from the stakeholder consultations. The expert group acknowledged the decision to make this component optional but highlighted how text messaging support has been found to be effective for smoking cessation. Therefore, they suggested that participants should be automatically enrolled in this component unless they express a negative preference during the pre-discharge evaluation. Participants should also be able to opt-out of the component at any time by replying 'STOP' to any of the messages or by contacting the individual delivering the intervention (Supplementary Material; Table 2c).

### **The final prototype intervention**

The development process of identifying key domains through to intervention content is presented in Supplementary Material 9. The final prototype intervention consists of components that aim to support smoking-related behaviour change among patients following discharge from a smokefree mental health inpatient setting. Intervention components

include: (1) pre-discharge reflection and evaluation; (2) a personalised resource folder (My Try folder); (3) tailored behavioural support (via telephone or video call); (4) optional text-messaging support, and (5) optional opportunity for peer interaction (Supplementary Material 10). The intervention will be delivered by a trained MHW (named the 'My Try Specialist', to link with the name of the personalised resource folder). The name 'My Try' was discussed and agreed with the stakeholders, expert group, and PPI group, and was identified as appropriate as it indicated flexibility dependent on individual patient's motivations.

Supplementary Material 11 provides a detailed description of the intervention using the 12-item Template for Intervention Description and Replication (TIDieR) checklist<sup>28</sup>.

## DISCUSSION

The intervention consists of five components that together aim to support smoking-related behaviour change among patients following discharge from a smokefree mental health inpatient setting. This intervention draws on behavioural science to specifically target the barriers/enablers to supporting smoking cessation in mental health settings and draws on existing evidence. A large body of literature recognises the challenges that people with mental health conditions experience when making smoking-related behaviour changes, including low self-efficacy<sup>29,30</sup>, smoking norms<sup>25,31</sup>, widely held misconceptions about the links between mental health and smoking<sup>32,33</sup>, and a lack of support offered compared to that available to the general population<sup>34</sup>. Therefore, it is important that interventions delivered to people with mental health conditions provide flexible support tailored to individual needs and are delivered by MHWs who are experienced in assisting people with mental health conditions<sup>35,36</sup>. By examining the literature, the current prototype incorporates content that has been reported as successful for this population group<sup>11</sup>, while addressing commonly cited barriers<sup>25</sup>. Stakeholder consultations ensured that the prototype intervention met the target population's needs and preferences. This in-depth approach to intervention development increases the likelihood that it will achieve smoking-related behaviour change post-discharge. The next stage is to deliver the prototype intervention in a small-scale pilot study to test the intervention and processes for fitness of purpose. A process evaluation will highlight the need for further refinements. Subsequently, the intervention will be tested in a randomised controlled feasibility study.

The prototype intervention includes several remote components. Literature indicates that telephone support increases the chances of stopping smoking, whether the individuals are motivated to quit or not, or if they are receiving other support<sup>37</sup>. The COM-B model supports remote smoking cessation treatment as this approach may provide the opportunity to access support by removing barriers (e.g., travel and cost), particularly for those where accessing face-to-face services may be challenging<sup>38,39</sup>. Text messaging is also an established cessation modality<sup>40</sup> and has shown to promote smoking cessation in the short-term<sup>41</sup> and long-term<sup>42</sup>. The opportunity for a remote peer interaction group was selected as a



component to replace the identification of a support 'buddy'. There is strong evidence to suggest that social networks play an important role in an individual's quit attempt<sup>43,44</sup>, and we found all 'social support' BCTs were identified as 'promising' both in terms of likely effectiveness and feasibility<sup>11</sup>. Therefore, it was important to include an intervention component that aims to enhance social support, and evidence suggests remote social support can enhance smoking cessation outcomes<sup>45</sup>.

Although the remote support was well-received by the stakeholders, it is important to consider this mode of delivery may present challenges if participants have restricted access to technology and/or limited IT skills. Research has demonstrated that people with mental health conditions are at an increased risk of digital exclusion<sup>46,47</sup>. However, this was not identified as a potential barrier during our consultations, and the increasing use of remote communication to provide health services during the Covid-19 pandemic illustrates the advantages of remote counselling<sup>48</sup>.

### **Limitations**

During stakeholder consultations, we collected patient and caregiver demographics including gender and smoking status. However, we did not collect demographics such as ethnicity and socioeconomic status, which would have further helped to contextualise the findings. Secondly, informal caregivers are under-represented in the stakeholder consultations. Recruitment of informal caregivers would have provided a valuable perspective of those who have supported relatives/friends who are current or ex-smokers and have experience of using acute inpatient services. However, recruiting informal caregivers of smokers with a mental health condition is acknowledged as difficult<sup>25</sup>, and consultations were conducted at the beginning of the pandemic so recruitment was conducted remotely, restricting the opportunity to meet with potential caregivers on inpatient wards. This highlights the need for further investigation into the role of informal support networks, and the development of targeted strategies to recruit informal caregivers.

### **Conclusions**

People with mental health conditions are more likely to smoke than the general population, which contributes to widening tobacco-related health inequalities. This intervention aims to support smoking-related behaviour change in patients discharged from a smokefree mental health inpatient stay, encompassing components that have been guided by existing evidence and appropriate BCTs. It includes resources and approaches that have been developed in collaboration with stakeholders to ensure they appeal to patients with mental health conditions following a smokefree stay. Using the BCW and TDF supported a theoretically and empirically informed process to define and develop a tailored intervention that acknowledges barriers and enablers to supporting smoking cessation in mental health settings. The result is a novel complex theory- and evidence-based intervention that will be formally tested in a randomised controlled feasibility study.



**Data availability statement:** Data is available at: <https://osf.io/jdw9y/>

**Funding:** This study is funded by the National Institute for Health Research (NIHR) Programme Grants for Applied Research programme (reference: NIHR200607). Elena Ratschen and Simon Gilbody are supported by the NIHR Yorkshire and Humber Applied Research Collaboration <https://www.arc-yh.nihr.ac.uk>. Lion Shahab is supported by a programme grant from Cancer Research UK (PRCRPG-Nov21\100002) and is member of SPECTRUM, a UKPRP-funded consortium (MR/S037519/1). Peter Coventry is supported by the UK Research and Innovation Closing the Gap Network+ (ES/S004459/1) <https://www.york.ac.uk/healthsciences/closing-the-gap/> and by the National Institute for Health Research (NIHR) Yorkshire and Humber Applied Research Collaboration <https://www.arc-yh.nihr.ac.uk/>. Tim Coleman is an NIHR Senior Investigator. The views expressed are those of the authors, and not necessarily those of the NIHR or the Department of Health and Social Care.

**Conflict of interest declaration:** L.S. is a HEFCE funded member of staff at University College London. He has received honoraria for talks, an unrestricted research grant and travel expenses to attend meetings and workshops from Pfizer and an honorarium to sit on advisory panel from Johnson&Johnson, both pharmaceutical companies that make smoking cessation products. He has acted as paid reviewer for grant awarding bodies and as a paid consultant for health-care companies. Other research has been funded by the government, a community-interested company (National Centre for Smoking Cessation) and charitable sources. He has never received personal fees or research funding of any kind from alcohol, electronic cigarette, or tobacco companies. All other authors declare no conflicts of interest.

**Acknowledgements:** The authors would like to thank the stakeholders and respective Trusts involved in the consultations who contributed to recommendations for revising the prototype intervention: Sheffield Health and Social Care NHS Foundation Trust, Leeds and York Partnership NHS Foundation Trust, Tees, Esk and Wear Valley NHS Foundation Trust, and Bradford District Care NHS Foundation Trust. The authors would also like to thank the research programme's PPI group for providing feedback and assisting with the development and review of intervention materials.

## References

1. Peacock A, Leung J, Larney S, et al. Global statistics on alcohol, tobacco and illicit drug use: 2017 status report. <https://doi.org/10.1111/add.14234>. *Addiction*. 2018/10/01 2018;113(10):1905-1926. doi:<https://doi.org/10.1111/add.14234>
2. ONS. Smoking prevalence in the UK and the impact of data collection changes: 2020. Accessed 24th May, 2022. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/drugusealcoholandsmoking/>
3. Richardson S, McNeill A, Brose LS. Smoking and quitting behaviours by mental health conditions in Great Britain (1993–2014). *Addict Behav*. 2019/03/01/ 2019;90:14-19. doi:<https://doi.org/10.1016/j.addbeh.2018.10.011>
4. Richardson S, McNeill A, Brose LS. Smoking and quitting behaviours by mental health conditions in Great Britain (1993-2014). *Addict Behav*. Mar 2019;90:14-19. doi:10.1016/j.addbeh.2018.10.011
5. Glasheen C, Hedden SL, Forman-Hoffman VL, Colpe LJ. Cigarette smoking behaviors among adults with serious mental illness in a nationally representative sample. *Ann Epidemiol*. 2014/10/01/ 2014;24(10):776-780. doi:<https://doi.org/10.1016/j.annepidem.2014.07.009>
6. McClave AK, McKnight-Eily LR, Davis SP, Dube SR. Smoking Characteristics of Adults With Selected Lifetime Mental Illnesses: Results From the 2007 National Health Interview Survey. *Am J Public Health*. 2010;100(12):2464-2472. doi:10.2105/ajph.2009.188136
7. Siru R, Hulse GK, Tait RJ. Assessing motivation to quit smoking in people with mental illness: a review. *Addiction*. May 2009;104(5):719-33. doi:10.1111/j.1360-0443.2009.02545.x
8. Szatkowski L, McNeill A. The delivery of smoking cessation interventions to primary care patients with mental health problems. *Addiction*. 2013;108(8):1487-1494.
9. NICE. Stop smoking interventions and services. <https://www.nice.org.uk/guidance/ng92/chapter/Rationale-and-impact>.
10. Brose LS, Simonavicius E, McNeill A. Maintaining abstinence from smoking after a period of enforced abstinence - systematic review, meta-analysis and analysis of behaviour change techniques with a focus on mental health. *Psychol. Med*. 2018;48(4):669-678. doi:10.1017/S0033291717002021
11. Shoesmith E, Huddleston L, Lorencatto F, Shahab L, Gilbody S, Ratschen E. Supporting smoking cessation and preventing relapse following a stay in a smoke-free setting: a meta-analysis and investigation of effective behaviour change techniques. <https://doi.org/10.1111/add.15452>. *Addiction*. 2021/11/01 2021;116(11):2978-2994. doi:<https://doi.org/10.1111/add.15452>
12. Stockings EA, Bowman JA, Prochaska JJ, et al. The impact of a smoke-free psychiatric hospitalization on patient smoking outcomes: a systematic review. *Aust N Z J Psychiatry*. 2014;48(7):617-633. doi:10.1177/0004867414533835
13. Prochaska JJ, Fletcher L, Hall SE, Hall SM. Return to smoking following a smoke-free psychiatric hospitalization. Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't. *Am J Addict*. Jan-Feb 2006;15(1):15-22.
14. NICE. NICE Guidance PH48. Smoking: acute, maternity and mental health services. Accessed 24th May, 2022. <https://www.nice.org.uk/guidance/ph48>

15. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Bmj*. 2008;337
16. Michie S, Atkins L, & West R. . *The Behaviour Change Wheel: A Guide to Designing Interventions*. Silverback Publishing; 2014.
17. Richardson M, Khouja CL, Sutcliffe K, Thomas J. Using the theoretical domains framework and the behavioural change wheel in an overarching synthesis of systematic reviews. *BMJ Open*. 2019;9(6):e024950. doi:10.1136/bmjopen-2018-024950
18. Gould GS, Bar-Zeev Y, Bovill M, et al. Designing an implementation intervention with the Behaviour Change Wheel for health provider smoking cessation care for Australian Indigenous pregnant women. *Implement Sci*. 2017/09/15 2017;12(1):114. doi:10.1186/s13012-017-0645-1
19. Dugdale S, Ward J, Elison-Davies S, Davies G, Brown E. Breaking Free from Smoking: A Novel Digital Smoking Cessation Intervention for Offenders in UK Prisons. *J Smok Cessat*. 2018;13(4):189-200. doi:10.1017/jsc.2018.3
20. Passey ME, Adams C, Paul C, Atkins L, Longman JM. Improving implementation of smoking cessation guidelines in pregnancy care: development of an intervention to address system, maternity service leader and clinician factors. *Implement Sci*. 2021/11/17 2021;2(1):128. doi:10.1186/s43058-021-00235-5
21. Stanton- Fay SH, Hamilton K, Chadwick PM, et al. The DAFNE plus programme for sustained type 1 diabetes self management: Intervention development using the Behaviour Change Wheel. *Diabetic Medicine*. 2021;38(5):e14548.
22. Michie S, Johnston M, Abraham C, Lawton R, Parker D, Walker A. Making psychological theory useful for implementing evidence based practice: a consensus approach. *BMJ Qual Saf*. 2005;14(1):26-33.
23. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implement Sci*. 2011/04/23 2011;6(1):42. doi:10.1186/1748-5908-6-42
24. Michie S, Richardson M, Johnston M, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med*. 2013;46(1):81-95.
25. Huddleston L, Shoosmith E, Pervin J, Lorencatto F, Watson J, Ratschen E. A systematic review of mental health professionals, patients and carers' perceived barriers and enablers to supporting smoking cessation in mental health settings. *Nicotine Tob Res*. 2022;doi:10.1093/ntr/ntac004
26. NVivo. Version 12. 2020. www.qsrinternational.com/nvivo-qualitative-data-analysis-software/home
27. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol*. 2006/01/01 2006;3(2):77-101. doi:10.1191/1478088706qp063oa
28. Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *Bmj*. 2014;348
29. Trainor K, Leavey G. Barriers and facilitators to smoking cessation among people with severe mental illness: a critical appraisal of qualitative studies. *Nicotine Tob Res*. 2016;19(1):14-23.
30. Lubitz SF, Flitter A, Wileyto EP, et al. History and correlates of smoking cessation behaviors among smokers with serious mental illness. *Nicotine Tob Res*. 2020;22(9):1492-1499.

31. Aschbrenner KA, Naslund JA, Gill L, et al. Qualitative analysis of social network influences on quitting smoking among individuals with serious mental illness. *Journal of Mental Health*. 2019/09/03 2019;28(5):475-481. doi:10.1080/09638237.2017.1340600
32. Sheals K, Tombor I, McNeill A, Shahab L. A mixed- method systematic review and meta- analysis of mental health professionals' attitudes toward smoking and smoking cessation among people with mental illnesses. *Addiction*. 2016;111(9):1536-1553.
33. Taylor GMJ, Sawyer K, Kessler D, Munafò MR, Aveyard P, Shaw A. Views about integrating smoking cessation treatment within psychological services for patients with common mental illness: A multi-perspective qualitative study. <https://doi.org/10.1111/hex.13182>. *Health Expect*. 2021/04/01 2021;24(2):411-420. doi:<https://doi.org/10.1111/hex.13182>
34. Ratschen E, Britton J, Doody GA, Leonardi-Bee J, McNeill A. Tobacco dependence, treatment and smoke-free policies: a survey of mental health professionals' knowledge and attitudes. *Gen Hosp Psychiatry*. Nov-Dec 2009;31(6):576-82. doi:10.1016/j.genhosppsy.2009.08.003
35. Klein P, Lawn S, Tsourtos G, van Agteren J. Tailoring of a Smartphone Smoking Cessation App (Kick.it) for Serious Mental Illness Populations: Qualitative Study. *JMIR Hum Factors*. 2019;6(3):e14023-e14023. doi:10.2196/14023
36. Peckham E, Man M-S, Mitchell N, et al. Smoking Cessation Intervention for severe Mental Ill Health Trial (SCIMITAR): a pilot randomised control trial of the clinical effectiveness and cost-effectiveness of a bespoke smoking cessation service. *Health Technol Assess*. 2015/03// 2015;19(25):1-148, v-vi. doi:10.3310/hta19250
37. Hartmann-Boyce J, Hong B, Livingstone-Banks J, Wheat H, Fanshawe TR. Additional behavioural support as an adjunct to pharmacotherapy for smoking cessation. *Cochrane Database Syst Rev*. Jun 5 2019;6(6):Cd009670. doi:10.1002/14651858.CD009670.pub4
38. Matkin W, Ordóñez- Mena JM, Hartmann- Boyce J. Telephone counselling for smoking cessation. *Cochrane Database of Syst Rev*. 2019;(5)doi:10.1002/14651858.CD002850.pub4
39. Byaruhanga J, Paul CL, Wiggers J, et al. The short-term effectiveness of real-time video counselling on smoking cessation among residents in rural and remote areas: An interim analysis of a randomised trial. *J Subst Abuse Treat*. 2021/12/01/ 2021;131:108448. doi:<https://doi.org/10.1016/j.jsat.2021.108448>
40. Graham AL, Jacobs MA, Cohn AM, et al. Optimising text messaging to improve adherence to web-based smoking cessation treatment: a randomised control trial protocol. *BMJ Open*. 2016;6(3):e010687-e010687. doi:10.1136/bmjopen-2015-010687
41. Abrams LC, Ahuja M, Kodl Y, et al. Text2Quit: Results From a Pilot Test of a Personalized, Interactive Mobile Health Smoking Cessation Program. *Health Commun*. 2012/05/02 2012;17(sup1):44-53. doi:10.1080/10810730.2011.649159
42. Whittaker R, McRobbie H, Bullen C, Rodgers A, Gu Y. Mobile phone- based interventions for smoking cessation. *Cochrane Database of Syst Rev*. 2016;(4)doi:10.1002/14651858.CD006611.pub4
43. Taylor AE, Howe LD, Heron JE, Ware JJ, Hickman M, Munafò MR. Maternal smoking during pregnancy and offspring smoking initiation: assessing the role of intrauterine exposure. *Addiction*. Jun 2014;109(6):1013-21. doi:10.1111/add.12514
44. Kerr S, Woods C, Knussen C, Watson H, Hunter R. Breaking the habit: a qualitative exploration of barriers and facilitators to smoking cessation in people with enduring mental health problems. *BMC Public Health*. 2013;13(1):221-221. doi:10.1186/1471-2458-13-221
45. Goldenhersch E, Thrul J, Ungaretti J, Rosencovich N, Waitman C, Ceberio MR. Virtual Reality Smartphone-Based Intervention for Smoking Cessation: Pilot Randomized


Controlled Trial on Initial Clinical Efficacy and Adherence. *J Med Internet Res.* 2020/7/29 2020;22(7):e17571. doi:10.2196/17571

46. Spanakis P, Heron P, Walker L, et al. Use of the Internet and Digital Devices Among People With Severe Mental Ill Health During the COVID-19 Pandemic Restrictions. *Front Psychiatry.* 2021;12:732735. doi:10.3389/fpsy.2021.732735

47. Hidalgo-Mazzei D, Nikolova VL, Kitchen S, Young AH. Internet-connected devices ownership, use and interests in bipolar disorder: from desktop to mobile mental health. *Digital Psychiatry.* 2019;2(1):1-7.

48. Webster P. Virtual health care in the era of COVID-19. *Lancet.* 2020;395(10231):1180-1181. doi:10.1016/S0140-6736(20)30818-7.

Accepted Manuscript



**Figure 1.** Intervention development process



**Table 1.** Stakeholder demographics (n = 25)

<b>Mental health workers (n = 17)</b>		<b>N (%)</b>
<i>Gender</i>	Female	10 (58.8)
<i>Setting</i>	Inpatient	11 (64.7)
	Community	6 (35.3)
<i>Job role</i>	Nurse	8 (47.0)
	Healthcare assistant	3 (17.6)
	Occupational therapist	1 (5.9)
	Unit manager	1 (5.9)
	Smoking cessation advisor	2 (11.8)
	Healthy living advisor	2 (11.8)
<b>Patients (n = 7)</b>		<b>N (%)</b>
<i>Gender</i>	Female	2 (28.6)
<i>Smoking status</i>	Recently quit smoking	4 (57.1)
<i>Admission/discharge status</i>	Currently using acute inpatient service	2 (28.6)
	Community-based, and have had previous experience of using acute inpatient services	5 (71.4)
<b>Caregiver (n = 1)</b>		<b>N (%)</b>
<i>Gender</i>	Female	1 (100)
<i>Smoking status</i>	Never smoker	1 (100)

Accepted Manuscript