

Inhibition of attachment of some fouling diatoms and settlement of *Ulva lactuca* zoospores by film-forming bacterium and their extracellular products isolated from biofouled substrata in Northern Chile

Fernando Silva-Aciares*

Laboratorio de Ecología Microbiana
Departamento de Acuicultura
Facultad de Recursos del Mar
Universidad de Antofagasta
Av. Jaime Guzman s/n Casilla 170
Antofagasta, Chile
Tel: 56 55 637881
Fax: 56 55 637804
E-mail: fsaciares@uantof.cl

Carlos Riquelme

Laboratorio de Ecología Microbiana
Departamento de Acuicultura
Facultad de Recursos del Mar
Universidad de Antofagasta
Av. Jaime Guzman s/n Casilla 170
Antofagasta, Chile
Tel: 56 55 637881
Fax: 56 55 637804
E-mail: criquelme@uantof.cl

Financial support: This study was possible thanks to the financing of the project FONDEF N° D0111166 "Development of products of Microbial Origin for the Control of Biofouling in the Industry of Aquaculture" and project FONDEF N° D0411251 "Improvement biotechnology to production and application of microbial bioactive substances inhibitory of the biofouling to the marine industry".

Keywords: *Alteromonas* sp, antifouling bacteria, antifouling compounds, diatoms, marine microfouling, *Ulva* zoospores.

Abbreviations: AHL: N-acylhomoserine lactone
ASW: artificial sea water
EP: extracellular products
M9: minimal medium
SW: sea water
VNSS: vaatanen nine-salts solution

The biofouling of surfaces submerged in the marine environment includes primary colonization of the substrate by microorganisms including bacteria, microalgae, and microscopic reproductive propagules of macroorganisms such as algal zoospores. The present study reports the evaluation of the inhibitory potential of biofilms and extracellular products (EP) of the indigenous bacterium *Alteromonas* sp strain Ni1-LEM on the settlement of marine biofouling such as: (i) eight marine benthic diatoms and (ii) zoospores of the alga *Ulva lactuca*, as well as the germination of these zoospores and was compared with reference strains with proven antifouling properties, *Halomonas marina* (ATCC 25374) and *Pseudoalteromonas tunicata*. Highest antifouling activity was found for the indigenous strain. In attempts to better define the chemical nature of the

antifouling substance in the EP of the *Alteromonas* sp strain Ni1-LEM, the culture filtrates were tested for activity after heat treatment, enzymatic treatments, dialysis through semipermeable membranes, and separation into polar (aqueous) and non-polar (organic) fractions. The results suggested that the antifouling substance in the culture filtrates to be protein or peptide in nature, thermostable, hydrophilic, and equal to or greater than 3500 daltons in molecular size. Antifouling substances from bacteria may lead to the development of novel antifouling agents in the future.

Living and inert surfaces immersed in the marine environment are usually rapidly colonized by microorganisms, the first step of which includes attachment of bacteria, diatoms, fungi, and protozoans. These are often

*Corresponding author

Table 1. The effect of bacterial biofilms of *P. tunicata* (Pt), *H. marina* (Hm), *Alteromonas sp* strain Ni1-LEM and *Halomonas sp* strain Nc1-LEM on the settlement of common biofouling diatoms and the settlement and germination of the zoospores of the macroalga *U. lactuca*. Values presented are mean percentages of settlement from three replicate trials \pm one standard deviation.

Microorganisms	Treatments				
	Control	Pt	Hm	Ni1-LEM	Nc1-LEM
Diatoms (Bacillariophyta)					
<i>Nitzschia sp1</i>	95.4 \pm 1.7 ^a	52.4 \pm 5.5 ^b	59.5 \pm 7.8 ^b	41.5 \pm 3.9 ^{c*}	94.0 \pm 5.7 ^a
<i>Amphora sp</i>	100 \pm 0 ^a	43.4 \pm 4.3 ^b	52.0 \pm 0.0 ^b	36.8 \pm 7.1 ^{c*}	100 \pm 3.0 ^a
<i>Cylindrotheca closterium</i>	81.6 \pm 2.9 ^a	38.4 \pm 1.9 ^c	48.5 \pm 2.2 ^b	36.5 \pm 5.5 ^{c*}	80.2 \pm 6.3 ^a
<i>Nitzschia ovalis arnott</i>	70.9 \pm 6.8 ^a	27.9 \pm 6.3 ^c	34.2 \pm 3.4 ^b	24.4 \pm 2.4 ^{c*}	68.3 \pm 4.2 ^a
<i>Chaetoceros minutissimus</i>	71.5 \pm 2.5 ^a	25.9 \pm 4.0 ^c	36.4 \pm 1.7 ^b	22.2 \pm 3.0 ^{c*}	70.4 \pm 6.5 ^a
<i>Navícula sp1</i>	62.3 \pm 1.5 ^a	18.9 \pm 1.7 ^c	25.1 \pm 4.0 ^b	16.0 \pm 2.0 ^{c*}	65.2 \pm 3.8 ^a
<i>Navícula sp2</i>	94.1 \pm 7.8 ^a	43.4 \pm 6.8 ^c	56.2 \pm 5.1 ^b	41.2 \pm 2.1 ^{c*}	92.2 \pm 4.0 ^a
<i>Nitzschia sp2</i>	93.3 \pm 3.5 ^a	45.5 \pm 5.0 ^c	53.1 \pm 8.5 ^b	40.1 \pm 2.4 ^{c*}	94.3 \pm 3.3 ^a
Algae (Chlorophyta)					
<i>Ulva lactuca</i>	93.4 \pm 9.6 ^a	19.4 \pm 2.7 ^b	22.3 \pm 4.4 ^b	7.5 \pm 2.5 ^{c*}	94.0 \pm 8.8 ^a
	(83.5 \pm 9.3) ^a	(27.6 \pm 6.6) ^b	(31.5 \pm 3.8) ^b	(12.4 \pm 3.3) ^{c*}	(80.4 \pm 5.0) ^a

Values in () represent percentage germination of *U. lactuca* zoospores.

Different superscripted letters indicate significant differences (one way ANOVA and LSD a posteriori test with $\alpha = 0.05$ and a > b > c) of data within that row. The letter "c*" indicates the highest degree of antifouling activity.

followed by the settlement of algal spores and larvae of macro invertebrates (Dobretsov and Qian, 2002). This colonization process is termed "biofouling" and results in the modification and even deterioration of structures or surfaces submerged in the sea as a result of the undesired accumulations of micro- and macrofouling organisms, mediated by a complex mix of physical, chemical, and biological phenomena (Egan et al. 2001; Callow and Callow, 2002; Gehrke and Sand, 2003). Biofouling is a complex process which can be initiated by the adsorption of inorganic and organic compounds such as polysaccharides, proteins, and peptides within a few min of submerging a clean substrate in seawater (Egan et al. 2001). This initial step is often followed by the formation of a biofilm formed of bacteria or other microorganisms (Gehrke and Sand, 2003). The bacteria are not the only organisms which can produce initial colonization; diatoms with their sticky mucopolysaccharide secretions may also be involved in early film formation, and when they are abundant can

promote bio-corrosion of the surface (Callow and Callow, 2002). As the biofilm grows, adhesive exudates are released which help trap additional particles and microorganisms. Among these may be included algal spores, marine fungi, and protozoans, some of which may be attracted to the film by chemosensory processes (Davies and Williamson, 1995).

Bacteria can play an important role in controlling the growth of micro- and macroalgae. It has been reported that some bacteria in the genera *Flavobacterium*, *Cytophaga*, *Alteromonas*, *Pseudomonas* and *Pseudoalteromonas* and their excretion products are capable of inhibiting the growth of diatoms and microalgae which are common in harmful phytoplankton blooms (Yoshinaga et al. 1997; Kato et al. 1998; Lovejoy et al. 1998; Lee et al. 2000; Burgess et al. 2003). These bacteria are capable of acting on a broad range of phytoplanktonic components, including diatoms, dinoflagellates, and some bacteria in the genera *Vibrio*,

Acinobacter and *Flavobacterium* (Yoshinaga et al. 1997). A study carried out by Hölmstrom et al. (1996) showed the capacity for the marine bacterial strain *Pseudoalteromonas tunicata* for inhibiting growth of the common biofouling diatom *Amphora* sp. With macroalgae, effects of bacterial strains have been studied in relation to the inhibition of growth of the green alga *Enteromorpha* (Thomas and Allsopp, 1983). Inhibitory effects of *P. tunicata* on the germination of spores of the cosmopolitan green alga *Ulva lactuca* and the red alga *Polysiphonia* sp. (a common biofouling alga) have been reported by Hölmstrom et al. (1998) and Egan et al. (2001). Also, Burgess et al. (2003) reported that *Pseudomonas* sp strain no. NUDMB50-11 showed excellent inhibitory activity on the germination of the spores of *U. lactuca*.

The objective of the present study was to determine the antifouling activity of the biofilms and extracellular products (EP) of the recently isolated bacterial strain *Alteromonas* strain Ni1-LEM from a coastal substrate in northern Chile on: (i) marine benthic diatoms, and (ii) algal

zoospores from *U. lactuca*, and to compare it with reference strains with proven antifouling properties, *P. tunicata* and *Halomonas marina* (ATCC 25374). *Alteromonas* sp strain Ni1-LEM is a previously bacterium isolated at the Microbial Ecology Laboratory of the Marine Resources Faculty, Universidad de Antofagasta, Chile. The strain had shown inhibitory effects on the settlement of some marine planktonic and benthonic species in preliminary laboratory test (Ayala et al. 2006; Zapata et al. 2007).

MATERIALS AND METHODS

Bacterial strains

Evaluation was made of direct bacterial inhibition of settlement of benthic diatoms common in the microfouling community of northern Chile, as well as of the settlement and germination of spores of the cosmopolitan green macroalgae *Ulva lactuca* Linnaeus. Bacterial strains used to evaluate the inhibition as outlined above included a strain

Table 2. Effects of extracellular bacterial supernatants of strains of *P. tunicata* (Pt), *H. marina* (Hm), *Alteromonas* sp strain Ni1-LEM and *Halomonas* sp strain Nc1-LEM on the settlement of common biofouling diatoms and the settlement and germination of zoospores of the macroalga *U. lactuca*. The values presented are mean percentages of settlement in three replicate trials \pm one standard deviation.

Microorganisms	Treatments					
	Ct SW	Ct M9	Pt	Hm	Ni1-LEM	Nc1-LEM
Diatoms (Bacillariophyta)						
<i>Nitzschia sp1</i>	99.7 \pm 8.0 ^a	95.3 \pm 12.3 ^a	8.2 \pm 0.5 ^c	18.8 \pm 1.3 ^b	6.4 \pm 1.9 ^{c*}	97.2 \pm 4.6 ^a
<i>Amphora sp</i>	99.3 \pm 6.4 ^a	97.0 \pm 8.9 ^a	9.0 \pm 1.0 ^c	21.2 \pm 3.5 ^b	7.1 \pm 0.5 ^{c*}	95.2 \pm 6.2 ^a
<i>Cylindrotheca closterium</i>	100.0 \pm 5.7 ^a	97.9 \pm 19.0 ^a	12.2 \pm 2.0 ^c	8.6 \pm 5.5 ^c	8.0 \pm 2.2 ^{c*}	95.0 \pm 3.0 ^a
<i>Nitzschia ovalis arnott</i>	100.0 \pm 9.2 ^a	98.3 \pm 6.3 ^a	11.1 \pm 2.0 ^c	12.9 \pm 3.8 ^c	7.8 \pm 1.5 ^{c*}	96.6 \pm 7.0 ^a
<i>Chaetoceros minutissimus</i>	100.0 \pm 7.5 ^a	96.5 \pm 14.9 ^a	8.9 \pm 1.2 ^c	7.3 \pm 2.1 ^c	5.2 \pm 3.0 ^{c*}	95.0 \pm 10.0 ^a
<i>Navicula sp1</i>	96.6 \pm 8.9 ^a	97.3 \pm 15.0 ^a	6.2 \pm 4.0 ^c	10.3 \pm 4.0 ^c	7.0 \pm 6.5 ^{c*}	92.9 \pm 6.6 ^a
<i>Navicula sp2</i>	98.3 \pm 6.4 ^a	89.9 \pm 9.3 ^a	8.6 \pm 5.0 ^c	11.6 \pm 1.6 ^c	8.2 \pm 2.0 ^{c*}	93.6 \pm 5.0 ^a
<i>Nitzschia sp2</i>	100.0 \pm 15.5 ^a	93.9 \pm 7.0 ^a	9.2 \pm 1.5 ^c	10.4 \pm 1.4 ^c	7.7 \pm 3.3 ^{c*}	95.4 \pm 8.8 ^a
Algae (Chlorophyta)						
<i>Ulva lactuca</i>	97.2 \pm 12.7 ^a	98.3 \pm 8.6 ^a	3.3 \pm 4.4 ^c	22.8 \pm 2.5 ^b	2.0 \pm 0.3 ^{c*}	(76.1 \pm 8.2) ^a
	(80.5 \pm 9.1) ^a	(78.9 \pm 6.9) ^a	(2.3 \pm 0.8) ^c	(27.6 \pm 3.3) ^b	(1.1 \pm 0.2) ^{c*}	97.9 \pm 9.3 ^a

The values in () represent percentage of germination of zoospores of *U. lactuca*. Different superscripted letters indicate significant differences (one way ANOVA and LSD a posteriori test with $\alpha = 0.05$ and $a > b > c$) of data within that row. The letter "c*" indicates the highest degree of antifouling activity.

of *Alteromonas* sp (clone Ni1-LEM) first isolated by us in northern Chile from a natural substrate (red macroalgae; *Rhododymenia* sp), found to have anti-settlement activity against larvae of the tunicates *Ciona intestinalis* and *Pyura praeputialis* (Zapata et al. 2007). Two "reference" strains of bacteria, including *Pseudoalteromonas tunicata* and *Halomonas marina* (ATCC 25374) were tested in comparison with the Ni1-LEM strain, as these strains had been previously been reported as having activity against macrofouling (Hölmstrom et al. 1998; Maki et al. 2000). A strain of bacteria, *Halomonas* sp strain Nc1-LEM was employed as a negative "control" species as no antifouling activity has been shown for this strain.

All the strains had previously been cultured in Vaatanen Nine-Salts Solution (VNSS, Holmstrom et al. 1998) and had been stored in our microbial strain collection in 30% (v/v) glycerol-seawater at -80°C.

Diatom strains

Pieces of polystyrene 1 cm² in area which we had submerged in coastal seawater in for 14 d at 10 mt of depth to accumulate biofilm were returned to the laboratory and repeatedly rinsed in sterile seawater so as to remove loosely attached microorganisms from the surfaces. The substrates were then each immersed in a test tubes containing with 10 ml of sterile marine saline solution (SSM), and the tightly adhered microorganisms were detached from the surfaces by a 30 sec sonication of the tubes in a Cole Parmer model CPX 130 ultrasonic processor. Aliquots of 0.1 and 1.0 ml recovered from 10 samples were individually inoculated into 250 ml erlenmeyer flasks containing 100 ml of Guillard (1975) f/2 algae culture medium, and incubated for two weeks at 20°C at a light intensity of 100 $\mu\text{mol m}^{-2}\cdot\text{s}^{-1}$ with a photoperiod of 12:12 L:D. Following the incubation, the diatom species present were isolated using the methods described by Alveal et al. (1995). Eight strains of dominant diatoms were ultimately obtained from the biofilms, including *Amphora* sp; *Nitzschia sp1*; *Nitzschia sp2*; *Nitzschia ovalis* (Arnott ex Grunow in Cleve & Grunow, 1880); *Chaetoceros minutissimus* (Makarova et Proshkina-Lavrenko, 1964); *Cylindrotheca closterium* (Enrenberg, 1841); *Navícula sp1* and *Navícula sp2*. Axenic cultures of these diatoms were obtained by treating them with antibiotics to eliminate accompanying bacteria. The axenicity was verified by realizing the epifluorescent technique.

Obtaining algal zoospores

The methods of Egan et al. (2001) were used to obtain algal zoospores of *U. lactuca*. Samples of mature *U. lactuca* Linnaeus were collected from the intertidal zone on the coast at San Jorge Bay, Antofagasta, Chile, prior to sporulation as determined by microscopic examination. The thalli were cleaned of sediment and debris, and then abundantly rinsed with 0.45 μm filtered, autoclaved seawater. The algal thalli were air dried at ambient

temperature and ambient light for two hrs. Pieces of algae bearing sporangia were deposited in 200 ml beakers with 100 ml of the sterilized, filtered seawater (as above) and left exposed to light from a 60 watt (daylight) lamp.

The zoospores released were collected and deposited on one side of a 10 cm watch glass in 0.20 μm filtered, autoclaved seawater away from the light source. After 5 to 10 min, the spores which had migrated toward the brighter side of the watch glass were collected by pipetting and the procedure was repeated to assure the spores had migrated away from any associated microorganisms. The zoospores were then collected and immersed in 10 ml of the 0.20 μm filtered, autoclaved seawater, and the numbers per ml were determined using a Neubauer counting chamber.

Preparation of bacterial biofilms

The methods of Maki et al. (2000) were used to determine the effects of *Alteromonas* sp. Strain Ni1-LEM, *Pseudoalteromonas tunicata* and *Halomonas marina* (positive control) and *Halomonas* sp strain Nc1-LEM (negative control) for their comparative ability inhibit diatom microfouling. Overnight cultures of the three bacteria were grown in VNSS were centrifuged at 11000 rpm for 15 min. The supernatant was discarded and the pellet was washed and resuspended in artificial sea water (ASW), homogenized, and recentrifuged. The pellet was resuspended in 1/10 ml VNSS broth (Hölmstrom et al. 1998), and homogenized. Direct counts of the bacteria were made using the DAPI epifluorescence technique. Nuclon® multiwell polystyrene chambers (8 x 12 wells; 6 mm diameter) each received 200 μl of 1/10 VNSS per well, and were then inoculated with a bacterial suspension containing 10^8 cells·ml⁻¹, and were incubated at 20°C for 48 hrs. in one room with controlled temperature. Following incubation, the culture medium was removed from each treatment, followed by repeated rinsing with ASW to eliminate bacteria not firmly adherent to the substrate. The numbers of firmly adherent bacteria were determined by direct counts in the microscope. All the wells then received 200 μl of sterile seawater prior to making the bioassays with diatoms and algal zoospores.

Evaluation of the effects of bacterial biofilms on settlement of diatoms and algal zoospores

Once the bacterial biofilms were obtained, the multiwell chambers, containing 200 μl of 0.2 μm filtered seawater were inoculated separately with the individual diatoms at concentrations of about $3.5\cdot 10^5$ cells·ml⁻¹. Control plates were established which either did not contain bacterial films, or contained bacterial films prepared with the negative control (*Halomonas* sp strain Nc1-LEM). Experimental and control chambers were run in triplicate, for periods of 24 hrs. Following this exposure period, counts of settled diatoms were made using an Olympus

model IX50 inverted microscope at 100X. The results were expressed in percentage adherence of each diatom.

In tests for settlement and germination of *U. lactuca*, each well of a test chamber received 200 µl of zoospores suspension containing a total of about 2800 spores. The plates were incubated at ambient temperature ($24 \pm 4^\circ\text{C}$) for 2 hrs in total darkness to allow settlement of the zoospores. Following incubation, the plates were washed with ASW five times to eliminate zoospores not firmly adherent to the substrate. The plates were then examined in the inverted microscope at 100X to count the numbers of zoospores which had firmly settled. In a second group having the same treatment, incubation was carried on for three days at ambient temperature with results observed and quantified as done with the first group. Tests with the *U. lactuca* zoospores were also carried out in triplicate for each type of bacterial film tested.

$$\% \text{ germination} = \frac{\text{N}^\circ \text{ of zoospores germinated}}{\text{N}^\circ \text{ of zoospores settled}} \times 100$$

Obtaining bacterial EP in culture supernatants

EP were obtained from 1 lt cultures of the three bacterial strains tested by growing each to the stationary phase in M9 broth (minimal medium) with constant stirring at a room temperature of about 20°C . The cultures were centrifuged at 11000 rpm for 15 min at 4°C , and the supernatant was filtered twice through 0.2 µm Millipore® filters and stored at -20°C for subsequent use in the fouling inhibition bioassays. The axenic condition of the bacterial EP was verified using microscopic epifluorescence techniques.

Enzymatic treatment the EP

In order to determine if the substance active in inhibition against diatoms and algal spores was protein or peptide-like, the EP were treated with the enzymes pronase E, and carboxypeptidase G, at a final concentration of 200 µg ml⁻¹. The mixtures were incubated at 37°C for two hrs (pronase E) and at 30°C for 3 hrs (carboxypeptidase G), both at pH 7.1. Subsequently both solutions of the enzyme-treated EP were separately submitted to heating in an 80°C water bath for 20 min to denature all protein and enzymic activity. Controls were prepared for both enzymic treatments using 0.45 µm filtered, sterile seawater at the same concentrations as in the test systems with EP, as well as normal seawater and the EP without enzymic treatment in M9 medium.

Heat treatment of the EP

In order to determine if the substance inhibitory to diatoms and algal spores is affected by heating, 10 ml samples of EP were treated at 80°C for 30 min in a water bath, followed by cooling to room temperature, these samples were filtered to

0.2 µm. The treated and untreated EP were then tested to determine their comparative inhibitory effects on the diatoms and *U. lactuca* zoospores.

Obtaining a dialyzed fraction of EP

To determine the approximate molecular size of the inhibitory substance, 100 ml of the EP were dialyzed for 12 hrs at 4°C against 0.2 µm filtered sterile distilled water. Benzoinated membranes were used, having cutoffs of either 3500 or 10000 daltons (Sigma®). Each dialysate was stored under sterile conditions at -20°C until it was employed in bioassays on diatoms and *U. lactuca* zoospores.

Obtaining organic extracts of the EP

Samples of 250 ml of the EP were extracted with 100 ml of dichloromethane, separating the organic (nonpolar) phase from the aqueous (polar) phase. The solvent was removed from the organic fraction by vacuum evaporation in a Büchi rotary evaporator at 40°C . The organic extracts were redissolved in ethanol to a concentration of 9-10 mg ml⁻¹. Both the polar and non-polar fractions were used in bioassays testing for inhibitory properties on diatoms and *U. lactuca* zoospores.

Evaluating the effects of the EP on the settlement of diatoms and algal zoospores

Multiwell polystyrene chambers (as above) were used for this testing in a manner similar to that described above for the living bacterial films. In the present testing, each wells received 0.1 ml of diatom culture which was in the pre-stationary growth phase, and had been rinsed with ASW to eliminate any effects of the algae culture medium (F/2; Guillard, 1975); 0.1 ml of EP from the different treatments as described above in this section were added to the wells on the plates, giving test diatom concentrations of about $3,5 \cdot 10^5 \text{ cell mL}^{-1}$ ($2,5 \cdot 10^5 \text{ cells cm}^{-2}$). The plates were incubated at 20°C in a controlled temperature incubator at 20°C , with a 12L:12D photoperiod at $100 \mu\text{mol m}^{-2}\cdot\text{s}^{-1}$. Diatom counts were made after 24 hrs of incubation, after rinsing each chamber five times with ASW to remove any non-adherent cells, and then observed in the inverted microscope. Counts of the adherent diatoms were made for each experimental arrangement, with the results expressed as cells present per cm².

In tests with *U. lactuca* zoospores, about 2800 spores in 100 µl of sterile seawater were added to each well, followed by 100 µL of EP for each type of treatment. The preparations were incubated at a room temperature of about 20°C in total darkness for two hrs to allow zoospore settlement. Following incubation, the plates were washed with ASW five times to eliminate zoospores not firmly adherent to the substrate. The plates were then examined in the inverted microscope at 100X to count the numbers of zoospores which had firmly settled. The plates were then allowed to incubate in ambient light for 10-15 hrs, and then

observed using the inverted microscope to observe for any germination of the spores. When it was confirmed that no germination had occurred, the seawater was removed from the wells and counts were made of the attached spores. A second, identical set of preparations was run to test the

effects of the EP on germination of *U. lactuca* zoospores. In these tests, after the spores had settled and were counted, the seawater which had been removed was replaced with an equal volume of seawater containing the EP from the various treatments as described above. These systems were

Table 3. Preliminary characterization of extracellular bacterial supernatants from *P. tunicata* (Pt), *H. marina* (Hm) and *Alteromonas sp* (Ni1-LEM) on the settlement of the common biofouling diatoms *Nitzschia sp1* (Nc), *Amphora sp* (Amp), *Cylindrotheca closterium* (Cc), *Nitzschia ovalis arnott* (Fp), *Chaetoceros minutissimus* (Nm), *Navicula sp1* (Nav sp1), *Navicula sp2* (Nav sp2) and *Nitzschia sp2* (Nitz sp2) and in the settlement and germination of the spores of *U. lactuca*. The letters presented are the degree of inhibition of settlement.

Treatments	Microorganisms								
	Nc	Amp	Cc	Fp	Nm	Nav sp1	Nav sp2	Nitz sp1	<i>U. lactuca</i>
Enzymic									
Pt + Prot	A	A	A	A	A	A	A	A	A (A)
Pt + Pept	A	A	A	A	A	A	A	A	A (A)
Hm + Prot	A	A	A	A	A	A	A	A	A (A)
Hm + Pept	A	A	A	A	A	A	A	A	A (A)
Ni1-LEM + Prot	C	C	C	C	C	C	C	C	A (A)
Ni1-LEM + Pept	A	A	A	A	A	A	A	A	A (A)
Heat denatured									
Pt 80°C	A	A	A	A	A	A	A	A	C (C)
Hm 80°C	C	C	C	C	C	C	C	C	C (C)
Ni1-LEM 80°C	C	C	C	C	C	C	C	C	C (C)
Dyalizates									
Pt 3500 Da	C	C	C	C	C	C	C	C	C (C)
Pt 10000 Da	A	A	A	A	A	A	A	A	A (A)
Hm 3500 Da	B	B	B	B	C	B	C	C	C (C)
Hm 10000 Da	C	C	B	C	C	C	B	C	B (C)
Ni1 3500 Da	C	C	C	C	C	C	C	C	C (C)
Ni1 10000 Da	C	C	C	C	C	C	C	C	C (C)
Organic phase (non polar)									
Pt	A	A	A	A	A	A	A	A	A (A)
Hm	A	A	A	A	A	A	A	A	A (A)
Ni1-LEM	A	A	A	A	A	A	A	A	A (A)
Aqueous phase (polar)									
Pt	A	A	A	A	A	A	A	A	A (A)
Hm	B	B	B	B	A	A	B	A	B (B)
Ni1-LEM	A	A	A	A	A	A	A	A	A (A)

The letters in () represent the percentage germination of *U. lactuca* zoospores. Prot: Pronase E; Pept: Carboxypeptidase G

The level of inhibitory activity of treatments were categorize as non-inhibitory (A = 70-100% settlement), slightly inhibitory (B = 70-20% settlement) or strongly inhibitory (0-20% settlement).

incubated for three days at room temperature and ambient lighting conditions and then observed in the inverted microscope to determine the percentage of zoospore germination in each well. Germination was noted as positive when multiple cell clusters from each zoospore could be observed. All the experimental tests with diatoms and *U. lactuca* zoospores were carried out in triplicate.

Statistical analyses

The results obtained were submitted to a one-way analysis of variance (ANOVA, $\alpha = 0.05$) to evaluate the effect of the bacterial biofilms and their EP on the inhibition of settlement of diatoms, and on the inhibition of settlement and germination of *U. lactuca* zoospores. Determination of the factors which may have produced significant differences was carried out using LSD multiple comparison test.

RESULTS

Effect of bacterial biofilms on settlement of diatoms and algal zoospores

Results of tests on the effects of bacterial biofilms on diatom settlement, and settlement and germination of *U. lactuca* zoospores are presented in Table 1. Both the strain isolated in our laboratory (*Alteromonas* sp clon Ni1-LEM) and the reference strains *P. tunicata* significantly ($P < 0.05$) inhibited the diatoms and *U. lactuca* zoospores in comparisons with controls without biofilms, and the control prepared with films of the non-inhibitory *Halomonas* sp (clon Nc1-LEM). The highest degree of antifouling activity was noted with *Alteromonas* sp (Clon Ni1-LEM), in comparison with the reference strains.

Effects of EP on the settlement of diatoms and algal zoospores

Results obtained on the effects of EP on diatom settlement after 24 hrs incubations and on settlement and germination of *U. lactuca* zoospores are presented in Table 2. Similarly to the results obtained with the bacterial biofilms, *Alteromonas* sp clon Ni1-LEM and the reference strain *P. tunicata* significantly ($P < 0.05$) inhibited the diatoms and *U. lactuca* zoospores in comparisons with controls done using sea water (SW), M9 and the negative control preconditioned with the biofilm produced by the inactive bacterium *Halomonas* sp (clon Nc1-LEM). Again, the highest degree of antifouling activity was noted with *Alteromonas* sp (Clon Ni1-LEM).

Enzymatic treatments of EP, and their effects on inhibition of diatoms and zoospores of *U. lactuca*

The results obtained on the effects of the two enzyme treatments of EP on their capacity for fouling inhibition of diatoms and algal spores are presented in Table 3. These results showed that both the enzymes used suppressed the

inhibitory activity of the reference strains *P. tunicata* and *H. marina* showing high values for settlement of the diatoms, and settlement and germination of the *U. lactuca* zoospores compared with results from untreated supernatants, suggesting that inhibitory proteins or peptides had been denatured by the enzymic treatments. Only the enzyme carboxypeptidase G affected the inhibitory activity of the *Alteromonas* sp clon Ni1-LEM supernatant on the diatoms in this study, suggesting that the inhibitor was peptide-like. In contrast, since the settlement and germination of *U. lactuca* spores was affected by both enzyme-treated supernates, compounds inhibitory to *Ulva* could have been of both protein and/or peptide nature.

Heat treatment of EP and their inhibitory effects on benthic diatoms and *U. lactuca* zoospores

The results obtained after exposure of the diatoms and algal zoospores to the 80°C heat-treated bacterial supernatants (Table 3) indicated that the compound active in fouling inhibition was thermoresistant, since the results after the heat treatment were the same with the EP as those obtained without the heat treatment (Table 2).

Effects of dialyzed fractions of EP on diatoms and *U. lactuca* zoospores

The results obtained by separation by molecular size of EP using dialysis at 3500 and 10000 daltons are shown in Table 3 and reveal that for *P. tunicata* the inhibitory fraction is found in the supernatant dialyzed at 3500 daltons, having inhibitory effects on the settlement of the eight diatom species, and on the germination of *U. lactuca*. With extracellular supernatants from *H. marina* and *Alteromonas* sp (clon Ni1-LEM), the results show the inhibitory effects of the 3500 and 10000 dalton fractions on the study organisms (Table 3). These results suggest that the active compound from *P. tunicata* had a size equal to or greater than 3500 and less than 10000 daltons, and greater or equal to 3500 daltons for *H. marina* and *Alteromonas* sp (clon Ni1-LEM).

Effects of organic extracts and the aqueous phase of EP on the settlement of diatoms and zoospores of *U. lactuca*.

The results presented in Table 3 show that only the aqueous phase of the bacterial supernatants of the three bacterial strains tested showed inhibition of settlement of the eight diatom species and settlement and germination of *U. lactuca* zoospores. This result suggested that the active component was of a polar compound.

DISCUSSION

Table 1 and Table 2 shows the significant effects of bacterial biofilms and their EP of "reference" strain of bacteria *Pseudoalteromonas tunicate* and *Halomonas marina* and native bacterium *Alteromonas* sp strain Ni1-LEM, on inhibition of settlement of common fouling

diatoms and settlement and germination of *U. lactuca* zoospores, with the highest activity attributable to the *Alteromonas* strain Ni1-LEM. Various authors have indicated that bacterial biofilms were able to inhibit the settlement of benthic diatoms (Gawne et al. 1998; Hölmstrom et al. 2000) and settlement and subsequent germination of algal zoospores (Egan et al. 2001; Patel et al. 2003). On the other hand, the anti-diatom effects of the bacterium HYK0203-SK02 *Pseudomonas putida* were shown to be effective against the pelagic diatoms *Stephanodiscus hantzschii* and the cyanobacterium *Microcystis aeruginosa* (Kang et al. 2005). Most bacterial isolates from the alga *Ulva reticulata* inhibited the growth of the diatoms *Nitzschia paleacea* but did not cause its mortality in a study by Dobretsov and Qian (2002). These same authors attribute the effects of biofilms on surface fouling to their physical and chemical properties. These moderating factors include the strains of bacteria and diatoms involved; adhesive adaptations of the microorganisms including flagellae, pili, peduncles, feet or rafia, and the sizes and shapes of microorganisms involved (Gawne et al. 1998; Wetherbee et al. 1998; Egan et al. 2001; Pasmore and Costerton, 2003). Also of importance is the capacity of the microorganisms for the production of extracellular adhesive substances such as polysaccharides (Wustman et al. 1997) glycoproteins, and polyglycans (Lind et al. 1997). One of the most extensively studied bacteria with antifouling activity has been *Pseudoalteromonas tunicata* isolated directly from the surface of the tunicate *Ciona intestinalis*. This bacterium was shown to be able to inhibit the settlement and growth of a large number of organisms including invertebrate larvae, bacteria, diatoms, fungi, and macroalgal zoospores (Hölmstrom and Kjelleberg, 1999; Egan et al. 2000; Egan et al. 2001). Studies by Maki et al (1988) using 18 different bacterial strains, evaluated their effect on settlement of cirripeds, finding seven inhibitory species. The most active inhibitory species was *Deleya marina* (presently known as *Halomonas marina*). Subsequent studies with *H. marina* showed that this bacterium could attach to different substrates on which the cirriped larvae then showed variable settlement responses, suggesting a complex interaction between the substrate, the bacterial strain, and the larva (Maki et al. 2000). There are, however, no reports in the literature on the effects of *H. marina* on the settlement of diatoms or settlement and germination of *U. lactuca* zoospores. Moreover, previous studies have shown that biofilm of our native bacterium *Alteromonas* sp strain Ni1-LEM and their EP inhibits the byssal formation in the mussel *Semimitylus algosus* (Ayala et al. 2006) and inhibits the settlement of tunicate larvae *Ciona intestinalis* and *Pyura praeputialis* (Zapata et al. 2007). This bacteria has an extended stationary growth phase, during which it produces metabolites actively inhibitory to fouling organisms (data no show). This results are in agreement with Egan et al. (2001) and Lau and Qian, (2001), who indicated that *P. tunicata* has an extended stationary phase, in which it produces at least five extracellular components that inhibited the settlement of a broad range of fouling

organisms. Stationary-phase supernatant was used because within the marine environment most bacteria spend long periods of time in stationary phase and commonly produce active secondary metabolites in this stage.

In the present study, the assays of supernatants treated with pronase E, carboxypeptidase G, organic extracts, aqueous phase extract, and heat treated extracts (Table 3) suggested that the actively inhibitory products of *Alteromonas* sp strain Ni1-LEM which were able to inhibit the attachment of diatoms and settlement and germination of *U. lactuca* zoospores were proteins or thermostable peptides of low molecular weight and with polar characteristics. This agrees with Holmström and Kjelleberg (1999) and Egan et al. (2001) that reported that polar, heat stable, low molecular weight peptides or proteins inhibited the settlement of bacteria, fungi, microalgae, algal zoospores, and invertebrates. Later, results from high performance liquid chromatography (HPLC) carried out on active extracellular fractions of our bacterium showed that the protein or peptide involved as an active inhibitory component against marine diatoms (data no shown) and tunicate larvae of *C. intestinalis* and *P. praeputialis* (Zapata et al. 2007), was not a hydrophobic protein, but was polar, and thus was easily dispersed through the marine environment. In contrast to our results in *Alteromonas* sp strain Ni1-LEM, Egan et al (2001) have indicated that the anti-algal activities of *P. tunicata* produces an extracellular component that is heat sensitive. It is feasible that low molecular weight proteins and peptides are less affected by heat. Examples of such peptides are the heat stable enterotoxins produced by *Escherichia coli* (Robins-Browne, 1994). There is little published information on the nature of the inhibitory activity of *Alteromonas* species in the establishment of fouling communities. For instance, Kon-ya et al. (1995) identified the ubiquinone-8 molecule as responsible for the anti-larval activity shown by *Alteromonas* sp. KK10304 isolated from a marine sponge. 2-n-pentyl-4-quinolinol produced by the marine bacterium *Alteromonas* sp. inhibited the growth of plankton diatoms (Long et al. 2003).

The EP of our bacterium functioned directly as chemosensory inhibitory signals which were easily diffused throughout the aqueous medium, and could be internal components in the transmission of biochemical signals. Decho et al. (1998) suggested that these compounds represented a logical option for a functional signaling molecule in the marine environment due to its solubility in water and ease of synthesis since the biochemical machinery needed for producing specialized proteins and peptides are readily available within most organisms, and that a variety of signals could be produced depending on the lengths and peptide sequences produced. On the other hand, Tait et al. (2005) have demonstrated that bacterial N-acylhomoserine lactone (AHL) (Family of quorum sensing signalling molecules) molecules are required before there is attraction of zoospores to bacterial biofilm. Zoospores are attracted to biofilms of the wild type *Vibrio anguillarum*

but not to mutants that do not produce AHLs. They have demonstrated that attraction is disrupted with recombinant *V. anguillarum* producing AiiA (autoinducer inactivation) proteins. This protein, discovered by Dong et al (2000) in an *Bacillus* isolate enzymatically inactivates AHLs by hydrolysing the homoserine lactone ring. Presumably in case of our bacterium, the protein or peptide of the EP would be the cause of the loss of AHL synthesis and as consequence, the inhibition of the settlement of diatoms and *U. lactuca* zoospores. Certainly bacterial antifouling activity in general deserves further studies.

At the moment, the mechanism by which *Alteromonas* sp strain Ni1-LEM inhibits the settlement of components of biofouling, are not known. Nevertheless, Lovejoy et al. (1998), suggested that bacteria inhibitory to algal growth acted by means of two mechanisms, one direct and one indirect; an example of the first mechanism is that in *Cytophaga* sp. (strain J18/M01) which induces lysis in diatom cultures beginning after two hrs of exposure to living bacteria, but not by EP of the same bacterial strain (Imai et al. 1993). Indirect attacks effect inhibition of algae through emission from the bacteria of chemical products or extracellular exudates. Lovejoy et al. (1998) evaluated the effects of strains of *Pseudoalteromonas* sp isolated from the Huon Estuary, Tasmania (Australia) on the growth of the red-tide dinoflagellates *Gymnodinium catenarum*, *Chatonella marina* and *Heterosigma akashiwo*, finding that the presence of the marine bacteria produced adverse effects on the three dinoflagellates. The deleterious effects were related to the production of high molecular weight extracellular substances by the bacteria. In addition, the bacterium, *Pseudoalteromonas* sp A28 was demonstrated to lyse marine algae via the production of extracellular proteases (Lee et al. 2000). Egan et al. (2001) reported that the antifouling substance as cited in the preceding affected the cell membrane of zoospores of *U. lactuca*, inhibiting their settlement and germination. This was attributed to the fact that the plasma of the membrane was exposed prior to settlement; although once the zoospore settled it formed a protective cell wall. A second mechanism proposed to explain the inhibitory effect of the active component produced by the bacteria was that the component functioned as a negative biochemical signal for settlement and germination, preventing their attachment and ultimately resulting in their death (Egan et al. 2001).

The preceding discussion implies the feasibility of developing bacterial biofilms on natural marine substrates, and that the EP of these biofilms could function in inhibition of settlement of microalgae and macroalgae components of fouling communities. The results shown in this study are important for future research into the biology, ecology, microbiology and marine botany which attempts to elucidate biological interactions which occur among different organisms associated with surfaces in the marine environment, and with the identification of natural

metabolites which could in the future lead to the development of applied methods for the mitigation of economically important marine fouling.

REFERENCES

- ALVEAL, K.; FERRARIO, M.; OLIVEIRA, E. and SAR, E. *Manual de Métodos Ficológicos*. Editorial Anibal Pinto S.A., Concepción, Chile, 1995, p. 863.
- AYALA, Cristina; CLARKE, Marcela and RIQUELME, Carlos. Inhibition of byssal formation in *Semimytilus algosus* (Gould, 1850) by a film-forming bacterium isolated from biofouled substrata in northern Chile. *Biofouling*, 2006, vol. 22, no. 1, p. 61-68.
- BURGESS, J. Grant; BOYD, Kenneth G.; ARMSTRONG, Evelyn; JIANG, Zhong; YAN, Liming; BERGGREN, Matz; MAY, Ulrika; PISACANE, Tony; GRANMO, Ake and ADAMS, David R. The development of a marine natural product-based antifouling paint. *Biofouling*, 2003, vol. 19, no. 1, p. 197-205.
- CALLOW, Maureen E. and CALLOW, James A. Marine biofouling: a sticky problem. *Biologist*, 2002, vol. 49, no. 1, p. 1-5.
- DAVIES, A. and WILLIAMSON, P. Marine Biofouling: A sticky problem. NERC News, April 1995, Marine BioFoulNet.
- DECHO, Alan W.; BROWNE, Kenneth and ZIMMERFAUST, Richard K. Chemical cues: why basic peptides are signal molecules in marine environments. *Limnology and Oceanography*, November 1998, vol. 43, no. 7, p. 1410-1417.
- DOBRETISOV, Sergey V. and QIAN, Pei-Yuan. Effect of bacteria associated with the green alga *Ulva reticulata* on marine micro-and macrofouling. *Biofouling*, January 2002, vol. 18, no. 3, p. 217-228.
- DONG, Yi-Hu; XU, Jin-Ling; LI, Xian-Zhen and ZHANG, Lian-Hui. AiiA, an enzyme that inactivates the acylhomoserine lactone quorum sensing signal and attenuates the virulence of *Erwinia carotovora*. *Proceedings of the National Academy of Sciences of the United States of America*, March 2000, vol. 97, no. 7, p. 3526-3531.
- EGAN, Suhelen; THOMAS, Torsten; HOLMSTRÖM, Carola and KJELLEBERG, Staffan. Phylogenetic relationship and antifouling activity of bacterial epiphytes from the marine algae *Ulva lactuca*. *Environmental Microbiology*, June 2000, vol. 2, no. 3, p. 343-347.
- EGAN, Suhelen; JAMES, Sally; HOLMSTRÖM, Carola and KJELLEBERG, Staffan. Inhibition of algal spore germination by the marine bacterium *Pseudoalteromonas*

tunicata. *FEMS Microbiology Ecology*, March 2001, vol. 35, no. 1, p. 67-73.

GAWNE, Benjamin; WANG, Yan; HOAGLAND, Kyle D. and GRETZ, Michael R. Role of bacterial and bacterial exopolymer in the attachment of *Achnanthes longipes* (bacillariophyceae). *Biofouling*, 1998, vol. 13, no. 2, p. 137-156.

GEHRKE, T. and SAND, W. Interactions between microorganisms and physicochemical factors cause MIC of steel pilings in harbours (ALWC). *NACE International Corrosion*, 2003, p. 1-8.

GUILLARD, R. *Culture of phytoplankton for feeding marine invertebrates*. Plenum Press, New York, USA, 1975. p. 29.

HOLMSTROM, Carola; JAMES, Sally; EGAN, Suhelen and KJELLEBERG, Staffan. Inhibition of common fouling organisms by marine bacterial isolates with special reference to the role of pigmented bacteria. *Biofouling*, 1996, vol. 10, no. 1-3, p. 251-259.

HOLMSTROM, Carola; JAMES, Sally; NEILAN, Brett A.; WHITE, David C. and KJELLEBERG, Staffan. *Pseudoalteromonas tunicata* sp nov., a bacterium that produces antifouling agents. *International Journal of Systematic Bacteriology*, October 1998, vol. 48, no. 4, p. 1205-1212.

HOLMSTROM, Carola and KJELLEBERG, Staffan. Marine *Pseudoalteromonas* species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiology Ecology*, December 1999, vol. 30, no. 4, p. 285-293.

HOLMSTROM, C.; STEINBERG, P.; CHRISTOV, V. and KJELLEBERG, S. Bacteria immobilized in hydrogels: a novel concept to prevent development of biofouling communities. *Biofouling*, 2000, vol. 15, no. 1-3, p. 109-117.

IMAI, I.; ISHIDA, Y. and HATA, Y. Killing of marine phytoplankton by a gliding bacterium *Cytophaga* sp., isolated from the coastal sea of Japan. *Marine Biology*, August 1993, vol. 116, no. 4, p. 527-532.

KANG, Y.-H.; KIM, J.-D.; KIM, B.-H.; KONG, D.-S. and HAN, M.-S. Isolation and characterization of a bio-agent antagonistic to the diatom, *Stephanodiscus hantzschii*. *Journal of Applied Microbiology*, May 2005, vol. 98, no. 5, p. 1030-1038.

KATO, Junichi; AMIE, Junya; MURATA, Yoshinori; KURODA, Akio; MITSUTANI, Atsushi and OHTAKE, Hisao. Development of a genetic transformation system for an alga-lysing bacterium. *Applied and Environmental Microbiology*, June 1998, vol. 64, no. 6, p. 2061-2064.

KON-YA, K.; SHIMIDZU, N.; OTAKI, N.; YOKOYAMA, A.; ADACHI, K. and MIKI, W. Inhibitory effect of bacterial ubiquinones on the settling of barnacles, *Balanus amphitrite*. *Cellular and Molecular Life Sciences*, February 1995, vol. 51, no. 2, p. 153-155.

LAU, S.C.K. and QIAN, P.-Y. Larval settlement in the serpulid polychaete *Hydroides elegans* (Haswell) in response to bacterial film: an investigation of the nature of putative larval settlement cue. *Marine Biology*, February 2001, vol. 138, no. 2, p. 321-328.

LEE, Sun-Og; KATO, Junichi; TAKIGUCHI, Noboru; KURODA, Akio; IKEDA, Tsukasa; MITSUTANI, Atsushi and OHTAKE, Hisao. Involvement of an extracellular protease in algicidal activity of the marine bacterium *Pseudoalteromonas* sp strain A28. *Applied and Environmental Microbiology*, October 2000, vol. 66, no. 10, p. 4334-4339.

LIND, Jan; HEIMANN, Kirsten; MILLER, Elizabeth; VAN VLIET, Catherine; HOOGENRAAD, Nicholas J. and WETHERBEE, Richard. Substratum adhesion and gliding in a diatom are mediated by extracellular proteoglycans. *Planta*, September 1997, vol. 203, no. 2, p. 213-221.

LONG, Richard A.; QURESHI, Asfia; FAULKNER, D. John and AZAM, Farooq. 2-n-pentyl-4-quinolinol produced by a marine *Alteromonas* sp. and its potential ecological and biogeochemical roles. *Applied and Environmental Microbiology*, January 2003, vol. 69, no. 1, p. 568-576.

LOVEJOY, Connie; BOWMAN, John and HALLEGRAEF, Gustaaf. Algicidal effects of a novel marine *Pseudoalteromonas* isolate (class *Proteobacteria*, gamma sub-division) on harmful algal bloom species of the genera *Chattonella*, *Gymnodinium*, and *Heterosigma*. *Applied and Environmental Microbiology*, August 1998, vol. 64, no. 8, p. 2806-2813.

MAKI, J.S.; RITTSCHOF, D.; COSTLOW, J.D. and MITCHELL, R. Inhibition of attachment of larval barnacles, *Balanus amphitrite*, by bacterial surface films. *Marine Biology*, February 1988, vol. 97, no. 2, p. 199-206.

MAKI, J.S.; DING, L.; STOKES, J.; KAVOURAS, J.H. and RITTSCHOF, D. Substratum/ bacterial interaction and larval attachment: Films an exopolysaccharides of *Halomonas marina* (ATCC 25374) and their effects on barnacle cyprid larvae, *Balanus amphitrite* Darwin. *Biofouling*, 2000, vol. 16, no. 2-4, p. 159-170.

PASMORE, Mark and COSTERTON, J. William. Biofilms, bacterial signaling, and their ties to marine biology. *Journal of Industrial Microbiology and Biotechnology*, July 2003, vol. 30, no. 7, p. 407-413.

PATEL, Pratixa; CALLOW, Maureen E.; JOINT, Ian and CALLOW, James A. Specificity in the settlement-

modifying response of bacterial biofilms towards zoospores of the marine alga *Enteromorpha*. *Environmental Microbiology*, May 2003, vol. 5, no. 5, p. 338-349.

ROBINS-BROWNE, R. *Escherichia coli* strains that cause diarrhoea: models of bacterial pathogenesis. In: GILBERT, G.L. ed. *Recent Advances in Microbiology*. Melbourne, The Australian Society for Microbiology Inc., 1994, p. 292-375.

TAIT, Karen; JOINT, Ian; DAIKYN, Mavis; MILTON, Debra L.; WILLIAMS, Paul and CAMARA, Miguel. Disruption of quorum sensing in seawater abolishes attraction of zoospores of the green alga *Ulva* to bacterial biofilms. *Environmental Microbiology*, February 2005, vol. 7, no. 2, p. 229-240.

THOMAS, R.W.S.P. and ALLSOPP, D. The effects of certain periphytic marine bacteria upon the settlement and growth of *Enteromorpha*, a fouling alga. *Biodeterioration*, 1983, vol. 5, p. 348-357.

WETHERBEE, Richard; LIND, Jan L.; BURKE, Jo and QUATRANO, Ralph S. The first kiss: establishment and control of initial adhesion by raphids diatoms - Minireview. *Journal of Phycology*, February 1998, vol. 34, no. 1, p. 9-15.

WUSTMAN, Brandon A.; GRETZ, Michael R. and HOAGLAND, Kyle D. Extracellular matrix assembly in diatoms (Bacillariophyceae): I. A model of adhesives based on chemical characterization and localization of polysaccharides from the marine diatom *Achnantes longipes* and other diatoms. *Plant Physiology*, April 1997, vol. 113, no. 4, p. 1059-1069.

YOSHINAGA, I.; KAWAI, T. and ISHIDA, Y. Analysis of algicidal ranges of the bacteria killing the marine dinoflagellate *Gymnodinium mikimotoi* isolated from Tanabe Bay, Wakayama Pref., Japan. *Fisheries Science*, February 1997, vol. 63, no. 1, p. 94-98.

ZAPATA, Manuel; SILVA, Fernando; LUZA, Yery; WILKENS, Marcela and RIQUELME, Carlos. The inhibitory effect of biofilms produced by wild bacterial isolates to the larval settlement of the fouling ascidia *Ciona intestinalis* and *Pyura praeputialis*. *Electronic Journal of Biotechnology* [online]. 15 January 2007, vol. 10, no. 1. Available from Internet: <http://www.ejbiotechnology.info/content/vol10/issue1/full/6/index.html>. ISSN 0717-3458.