

A microbial community analysis of the octocoral *Eunicea fusca*

Angela Duque-Alarcón¹ · Lory Z. Santiago-Vázquez^{1,2} · Russell G. Kerr^{1,3} ✉

¹ Florida Atlantic University, Department of Chemistry and Biochemistry, Center of Excellence in Biomedical and Marine Biotechnology, Boca Raton, FL, USA

² University of Houston-Clear Lake, Biotechnology and Biology, Houston, TX, USA

³ University of Prince Edward Island, Departments of Chemistry and Biomedical Sciences, Charlottetown, PE, Canada

✉ Corresponding author: rkerr@upeu.ca

Invited Article

Published online: September 15, 2012

© 2012 by Pontificia Universidad Católica de Valparaíso, Chile

Abstract While there is a significant and growing body of knowledge describing the microbial communities of marine invertebrates such as sponges, there are very few such studies focused on octocorals. The octocoral *Eunicea fusca* is common on reefs in various regions of the Caribbean and has been the subject of natural product investigations. As part of an effort to describe the microbial community associated with octocorals, a culture-independent analysis of the bacterial community of *E. fusca* was conducted. Specifically, a 16S rDNA clone library analysis was performed to provide baseline data. A total of 40 bacteria members from 11 groups were found. In general, Proteobacteria were the dominant group with a total of 24 species and α -Proteobacteria represented the highest percentage of bacteria associated with *E. fusca* (27.5%). Other prominent groups observed were Acidobacteria, Actinobacteria, Cyanobacteria, Planctomycetes, δ -Proteobacteria, Lentisphaerae and Nitrospirae. This is the first analysis of bacterial populations associated with the gorgonian *E. fusca*.

Keywords: 16S rDNA, bacteria, culture-independent analysis, *Eunicea fusca*, microbial community, octocoral

INTRODUCTION

The octocoral *Eunicea fusca* (Subclass Octocorallia, Order Gorgonacea, Family Plexauridae) is the source of anti-inflammatory compounds such as fuscicol and the fuscocides A and B. These act as selective inhibitors of leukotriene synthesis (Shin and Fenical, 1991; Jacobson and Jacobs, 1992a; Jacobson and Jacobs, 1992b). As exemplified by *E. fusca*, a variety of structurally unique, bioactive compounds have been isolated from octocorals (Faulkner, 2002). It has been suggested that bacteria living in symbiosis with these organisms have an important role in the production of these bioactive compounds (Radjasa and Sabdono, 2009). For instance, a *Vibrio* sp. isolated from the surface of the soft coral *Sinularia polydactyla* is the producer of Aqabamycins A-G, novel metabolites with antibacterial and cytotoxic activities (Al-Zereini et al. 2010). Even though octocorals are well known for their secondary metabolites, their associated microbial communities are relatively understudied in comparison to other well-studied marine invertebrates such as sponges and scleractinian corals (Enticknap et al. 2006; Littman et al. 2010).

In addition to their possible involvement in secondary metabolite production, symbiotic bacteria may also contribute to the coral's physiology and health (Rosenberg et al. 2007; Shnit-Orland and Kushmaro, 2009). Since corals have no adaptive immune system, bacteria associated with healthy corals are believed to play an important role in innate immunity (Reshef et al. 2006). For instance, it has been hypothesized that the microbial community of the mucopolysaccharide surface layer of *Gorgonia ventalina* plays a role in protection against disease by occupying niches that otherwise might be inhabited by pathogens, and also by producing antimicrobial compounds (Gil-Agudelo et al. 2006).

Bacterial biofilms have been implicated in promoting larvae metamorphosis of the scleractinian corals (Webster et al. 2004). It is possible that some of these bacteria can enter into a symbiotic relationship with the developing coral. Some of the most exciting and recent studies are examining the relationship between coral disease and bacterial population changes. For instance, microbial communities associated with *Acropora millepora* were shown to change during bleaching events shifting to *Vibrio*-dominated communities just prior to visual bleaching signs (Bourne et al. 2008). Changes in bacterial populations associated with bleached or azooxanthellate corals (Koren and Rosenberg, 2006) and with diseased corals have also been reported (Sunagawa et al. 2009). Therefore, an understanding of microbial communities in the relatively unstudied octocorals is of great importance.

The small number of studies that have examined the microbial assemblages of soft corals have reported a predominance of γ -Proteobacteria (Brück et al. 2007; Santiago-Vázquez et al. 2007). Studies that have examined the culturable bacteria of soft corals have also observed an abundance of γ -Proteobacteria followed closely by α -Proteobacteria (Ivanova et al. 2005) in addition to a minor presence of Bacteroidetes in the soft coral *Paragorgia arborea* (Nedashkovskaya et al. 2005). Interestingly, an examination of bacteria associated with *Plexaura fusifera* showed that unbleached corals were dominated by low G+C Firmicutes whereas bleached corals were dominated by γ -Proteobacteria demonstrating the specificity of these relationships (Frenz-Ross et al. 2008). While little is known about bacterial communities in octocorals, scleractinian coral-bacteria associations have been found to be specific (Rohwer et al. 2002; Lema et al. 2012).

The goal of this study is to examine the bacterial communities closely associated with *E. fusca* by culture-independent techniques (CIT; 16S rDNA clone library) to provide a baseline for other studies. CIT approaches are known to provide a broader analysis of bacterial populations when compared to culture dependent studies. It has been estimated that only 0.1% to 1% of microorganisms can be cultured in the laboratory (Jannasch and Jones, 1959). The DNA of the bacteria was amplified by universal eukaryotic 16S rDNA primers and a clone library was constructed. Restriction fragment length polymorphism (RFLP) analysis was used as a dereplication tool. Subsequently, PCR amplicons of interest were sequenced and analyzed. In accordance with previous observations from other soft corals, it was shown that γ -Proteobacteria were the most prominent bacteria observed. A study of the microbial ecology of this coral will contribute to the identification of potential producers of secondary metabolites and will provide a preliminary baseline of the bacterial community of the coral in its healthy state.

MATERIALS AND METHODS

Supplies

All chemicals were obtained from Sigma Chemical Co. (St. Louis, MO, USA). General laboratory supplies were obtained from Fisher Scientific (Suwanee, GA, USA). Pre-formulated bacterial broths and agars were supplied by Difco Laboratories (Detroit, MI, USA).

Coral sample collection

Eunicea fusca was collected at Hillsboro Ledge, Deerfield Beach, Florida, USA, on April 2005 by SCUBA diving at depths of approximately 10 m. Coral colonies were handled with gloves and stored in individual plastic bags. Individual colonies were rinsed several times with sterile seawater, immediately flash frozen in liquid nitrogen, and stored at -80°C until DNA extraction was performed.

DNA extraction

Genomic DNA (gDNA) was extracted from 100 mg of flash frozen *E. fusca* using the DNeasy Plant Mini Kit (Qiagen, Valencia, CA, USA). *E. fusca* was ground with liquid nitrogen using a sterile mortar and pestle. The coral slurry was transferred to a sterile 1.5 mL centrifuge tube and DNA was extracted according to manufacturer's instructions. Samples were eluted 2X in a total of 60 μL of ddH₂O (double distilled water). Extracted DNA was analyzed by gel electrophoresis. Samples were loaded, along with the appropriate molecular weight markers, on a 1.5% (w/v) agarose gel, 1X TBE (Tris-Borate EDTA) buffer, pH 8.3 and 1 $\mu\text{g}/\mu\text{L}$ ethidium bromide. Electrophoresis was performed at 100V for 60 min. DNA

bands were visualized in a Typhoon 9410 Imager (GE Healthcare, Piscataway, NJ, USA). DNA concentration was quantified by reading the absorbance at 260 nm on a SmartSpec 3000 spectrophotometer (Bio-Rad, Hercules, CA, USA).

Amplification and purification of 16S rDNA

Purified DNA from coral tissue was diluted to 1:10, 1:100, and 1:1000 of the original concentration with ddH₂O. These dilutions were used for 16S rDNA amplification with universal eubacterial primers FC27 (5'-AGAGTTTGATCCTGGCTCAG-3') and RC1492 (5'TACGGYTACCTTGTTACGACTT3'; Mincer et al. 2004). Each 50 µL PCR reaction consisted of 1 µL template DNA at the different concentrations, 25 µL Go Taq Green Master Mix (Promega, Valencia, CA, USA), 1 µL of each primer, and 22 µL of ddH₂O. PCR cycling conditions were as follows: initial denaturation at 95°C for 5 min, followed by 35 cycles of 95°C for 30 sec, 55°C for 30 sec, 72°C for 1.5 min. A final extension of 7 min at 72°C was added. PCR products were confirmed by gel electrophoresis. PCR amplicons of the anticipated size (~1.5kb) were selected for cloning. The PCR products were eluted from the gel with the MinElute Gel Extraction Kit (Qiagen) following manufacturer's instructions. Samples were eluted twice in a total of approximately 10 µL. Two µL of the eluted product was loaded on a 1.2% agarose gel alongside a quantifying ladder 1 kb Plus DNA Ladder E-Gel® DNA markers (Invitrogen, Carlsbad, CA) to estimate quantity of the eluted PCR amplicon in preparation for cloning.

16S rDNA cloning

The 16S rDNA fragments were cloned using the TOPO TA Cloning Kit for Sequencing according to the manufacturer's instructions (Invitrogen, Life Technologies, Carlsbad, CA, USA). A 100 µL aliquot from each transformation was plated onto pre-warmed selective LB plates containing Ampicillin (50 mg/mL) and incubated over night at 37°C. A minimum of 96 clones from each cloning plate were randomly selected and PCR-amplified using M13F (GTAAAACGACGGCCAG) and M13R (GTAAAACGACGGCCAGT) primers following manufacturer's cycling conditions. PCR products were separated by 1.5% (w/v) agarose gel electrophoresis and visualized on a Typhoon. Clones with the correct amplicon showed a single band at approximately 1700 bp. The number of clones ultimately picked was determined by the subsequent diversity observed by RFLP.

Restriction Fragment Length Polymorphism (RFLP)

The 16S rDNA PCR clones were digested with the restriction enzyme *HhaI* (Promega, Madison, WI, USA). The restriction digest reaction contained 5 µL PCR product, 0.1 µL *HhaI* (2 U/µL), 1 µL reaction buffer, 0.1 µL BSA, and 3.8 µL sterile ddH₂O and was incubated at 37°C for 1 hr and at 65°C for 10 min. The resulting fragments were loaded on a 1.2% agarose gel and analyzed by electrophoresis as detailed above. Restriction patterns and OTUs (operational taxonomic units) were compared and 2-3 clones that displayed different patterns were chosen for sequencing analysis.

DNA sequencing and analyses

Clones with unique RFLP patterns were sequenced using the Big Dye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Forster City, CA, USA) on an ABI 3100 automated sequencer (Applied Biosystems) according to the manufacturer's instructions. Alternatively, sequences were sent to BioAnalytical Services Laboratory, Center of Marine Biotechnology, University of Maryland Biotechnology Institute, Baltimore, MD, USA.

Sequences were viewed and edited using Chromas Lite 2000 (Technelysium, Tewantin, Australia; www.technelysium.com.au/chromas.html) and EditSeq (Lasergene, DNASTAR, Madison, WI, USA). Contiguous (contigs) sequences were built by Seqman (Lasergene).

16S rDNA sequence comparison

The 16S rDNA sequences were subjected to Basic Local Alignment Search Tool (BLAST) analysis at the National Center for Biotechnology Information (NCBI) database (www.ncbi.nlm.nih.gov). The

nucleotide sequences of the 16S rDNA gene sequences reported are available in GenBank under accession numbers EF657844 to EF657883.

Phylogenetic analysis

The 16S rDNA sequences of the clones were aligned and a using MegAlign (Lasergene) and the ClustalW algorithm. Subsequently, a phylogenetic tree of the 16S rDNA sequences was constructed using Phylip and the neighbour-joining method. Each reconstructed group was statistically evaluated by bootstrapping with a 1000 replicates (Nei and Kumar, 2000).

RESULTS AND DISCUSSION

This study analyzed the microbial community of bacteria associated with *Eunicea fusca*. There is evidence demonstrating that bacteria may be the source of biologically active natural products initially isolated from octocorals and in other marine organisms (Unson et al. 1994; Radjasa and Sabdono, 2009; Al-Zereini et al. 2010; Donia et al. 2011) and thus it has become a priority to gain a detailed understanding of the microbial communities in such systems. This knowledge may lead to an efficient and sustainable supply of these compounds that does not rely on harvesting coral tissue from the environment. In addition, the physiological role of these bacteria in the octocoral host is not understood.

A 16S rDNA clone library was constructed from an *E. fusca* genomic DNA sample. The resulting transformants were screened by RFLP analysis with *HhaI* as the restriction endonucleases that allowed for the clustering of similar clones. This analysis analyzed a total of 310 clones with the presence of 40 unique OTUs. Representatives of these clones were chosen for sequence analysis.

The taxonomic relatedness of the bacteria associated with *E. fusca* clones was deduced by the construction of a phylogenetic tree inferred from comparative sequence analyses of the 16S rDNA sequence from each clone (Figure 1 and Table 1). Based on this comparison, eleven distinct groups were observed with representatives from the sub-divisions α -Proteobacteria (27.5%), β -Proteobacteria (15.0%), γ -Proteobacteria (15.0%), Planctomycetes (12.5%), CFB (12.5%), Lentisphaerae (5.0%), Actinobacteria (2.5%), δ -Proteobacteria (2.5%), Acidobacteria (2.5%), Nitrospirae (2.5%), and Cyanobacteria (2.5%). Unpublished results from our group analyzing *E. fusca*'s culturable bacteria showed the collection of isolates was dominated by α -Proteobacteria (77%) followed by γ -Proteobacteria (13%), β -Proteobacteria (7%), and the division Cytophaga-Flexibacter-Bacteroidetes (CFB; 3%). In both cases bacterial populations were dominated by Proteobacteria in agreement with previous reports for other octocorals (Harder et al. 2003; Ivanova et al. 2005; Brück et al. 2007; Santiago-Vázquez et al. 2007; Webster and Bourne, 2007).

A survey of the identified taxa revealed candidates that hold promise as potential producers of bioactive secondary metabolites. These include the γ -Proteobacteria EF0605 (Accession no. EF657879). Evidence suggests that the related marine bacterium *Pseudomonas rhizosphaerae* may be the producer of potent antibacterial and antilarval secondary metabolites (Qi et al. 2009). Similarly, *Pseudomonas* spp. isolated from marine sources were found to be the source of potent antimicrobial compounds (Needham et al. 1994; Charyulu et al. 2009). A second candidate is the γ -Proteobacteria EF1505 (EF657861). This bacterium is related to *Candidatus Endobugula sertula*, the endosymbiont bacterium of the bryozoan *Bugula neritina* and also the known source of the bryostatins, a family of macrocyclic lactones with anticancer activity (Davidson et al. 2001). Finally, EF0705 (EF657868) and EF3305 (EF657846) showed a close phylogenetic association to actinobacteria. Marine actinomycetes are well known prolific sources of secondary metabolites (Lam, 2006).

Coral bacteria are believed to play an important role in the host's health. Corals are known to have symbiotic relationships with nitrogen-fixing bacteria (diazotrophs; Lema et al. 2012) therefore bacteria were examined for their potential involvement in the nitrogen cycle. A group of five α -Proteobacteria, EF2205 (EF657848), EF2405 (EF657850), EF2505 (EF657870), EF2605 (EF657871), and EF3905 (EF657852) were found to be related to *Pseudovibrio denitrificans* (AY486423). This coastal seawater organism is a facultative anaerobe capable of denitrification (Shieh et al. 2004). This process may allow bacteria to contribute to the health of the host by removing excess nitrate, and therefore controlling the population of the symbiotic dinoflagellate *Symbiodinium*. Excess *Symbiodinium* and the

subsequent production of harmful reactive oxygen species (ROS) are contributing factors to coral bleaching (Downs et al. 2002; Merle et al. 2007). Another isolate potentially involved in the Nitrogen cycle of *E. fusca* is the Nitrospirae EF0705 (EF657868). This bacterium is related to *Nitrospira marina*, a chemolithotrophic nitrite-oxidizing bacterium (Watson et al. 1986). In addition to nitrogen, bacteria involved in the carbon cycle are known to be associated with corals (Wegley et al. 2007). The species EF0505 (EF657862), part of the CFB group, is closely related to CFB found in coastal ocean bacterioplankton known to express transporter genes in response to DOC and therefore contribute to carbon turnover (Poretsky et al. 2010). Excess dissolved organic carbon (DOC) is detrimental to the health of corals (Kline et al. 2006).




Fig. 1 Phylogenetic tree showing the phylogenetic relationships between *Eunicea fusca*'s associated bacteria as determined by 16S rDNA sequence analysis. Numbers at nodes are percentages indicating levels of bootstrap support, based on neighbour-joining analysis of 1000 re-sampled data sets. Reference sequences derived from GenBank entries are written with their corresponding accession numbers.

Our findings concur with data reported for other octocorals (Brück et al. 2007; Santiago-Vázquez et al. 2007) and suggest that a more in depth assessment using other techniques such as fluorescent *in situ* hybridization (FISH) and next generation sequencing is warranted. Such tools can be used to examine how these associations vary with geographical location and health state of *E. fusca*.

It can be concluded from this initial study that *E. fusca* hosts a diverse and complex assemblage of coral-associated bacteria dominated by Proteobacteria, many of which are only distantly related to previously described bacteria. This is the first study of the microbial community associated with the coral *E. fusca* which provides a baseline for further studies in the areas of coral health and natural products.

Table 1. *Eunicea fusca's* bacterial associates.

Phylogenetic Association	Clone	Accession no.	Closest GenBank Homologue and Accession no.	% Similarity	
α-Proteobacteria	EF2105	EF657847	Uncultured Rhizobiales bacterium AB294974	95	
	EF2205	EF657848	Uncultured Alpha Proteobacterium clone Cobs2TisF11 EU246824	97	
	EF2005	EF657849	Uncultured Alpha Proteobacterium AJ581348	95	
	EF2405	EF657850	Uncultured Alpha Proteobacterium clone FII-AN025 JQ579966	93	
	EF3405	EF657851	Uncultured Rhizobiales bacterium clone B255_A3 EF092222	93	
	EF3905	EF657852	Uncultured Alpha Proteobacterium clone BBD_217_35 DQ446160	99	
	EF4105	EF657853	Uncultured bacterium clone LX E63 JF514279	96	
	EF2505	EF657870	Uncultured Rhodobacteraceae bacterium clone MD3.59 FJ425647	95	
	EF2605	EF657871	Uncultured Rhodobacteraceae bacterium clone MD3.59 FJ425647	95	
	EF2705	EF657881	Uncultured Alpha Proteobacterium EU544719	88	
	EF1205	EF657882	Uncultured Alpha Proteobacterium clone LC1-25 DQ289899	94	
	β-Proteobacteria	EF0905	EF657857	Uncultured bacterium clone Mann16S_H10 FJ952693	91
		EF0105	EF657858	Uncultured bacterium clone Mann16S_H10 FJ952693	92
		EF1105	EF657859	Uncultured bacterium clone Mann16S_H10 FJ952693	94
EF4005		EF657860	Uncultured bacterium clone transformant11 GU799620	92	
EF3205		EF657877	Uncultured bacterium clone GU799620	92	
EF3605		EF657878	Uncultured bacterium clone Mann16S_H10 FJ952693	93	
γ-Proteobacteria	EF0305	EF657844	Uncultured <i>Spongiobacter</i> sp. clone EC79 DQ889911	96	
	EF1505	EF657861	<i>Endobugula sertula</i> strain BnSP AF006606	91	
	EF1805	EF657865	Uncultured Gamma Proteobacterium clone XE2E10 JN596612	92	
	EF0605	EF657879	Uncultured <i>Pseudomonas</i> sp. clone VHS-B4-54 DQ395002	91	
	EF1005	EF657880	Uncultured bacterium clone QSW4 JQ347469	96	
	EF1305	EF657883	Uncultured Gamma Proteobacterium clone RII-OX016 JQ580103	88	
δ-Proteobacteria	EF2905	EF657863	Uncultured Delta Proteobacterium clone BME87 DQ917833	93	
CFB group	EF0205	EF657854	Uncultured bacterium clone 5S54 JF272214	97	
	EF1605	EF657855	Uncultured Bacteroidetes bacterium clone FII-TR116 JQ579911	95	
	EF3705	EF657856	Uncultured Bacteroidetes bacterium clone d149 GQ850585	93	
	EF0505	EF657862	Uncultured <i>Flavobacterium</i> sp. FJ745113	94	
	EF1705	EF657864	Uncultured bacterium clone SHFH609 FJ203538	93	
	Planctomycetes	EF0805	EF657872	Uncultured Planctomycete clone XE2D09 JN596618	96
EF3105		EF657873	Uncultured Planctomycete clone XE2D09 JN596618	98	
EF2805		EF657874	Uncultured Planctomycete clone RII-OX078 JQ580165	93	
EF2305		EF657875	Uncultured Planctomycete clone RII-AN084 JQ580463	92	
EF3805		EF657876	Uncultured Planctomycete clone RII-OX078 JQ580165	94	
Lentisphaerae		EF1905	EF657866	Uncultured Lentisphaerae bacterium clone PRTBB8480 HM798920	98
	EF0405	EF657867	Uncultured Lentisphaerae bacterium clone PRTBB8480 HM798920	98	
Acidobacteria	EF3505	EF657845	Uncultured Acidobacteria bacterium clone BME6 DQ917814	92	
Actinobacteria	EF3305	EF657846	Uncultured Actinobacterium clone II10D FJ205382	95	
Nitrospirae	EF0705	EF657868	<i>Nitrospira</i> sp. enrichment culture clone Aa01 EU055608	95	
Cyanobacteria	EF1405	EF657869	<i>Oscillatoria rosea</i> IAM M-220 AB003164	95	

ACKNOWLEDGMENTS

The authors acknowledge the crew of the MV Diversity (Florida, USA) for assistance with the collections, and Julie J. Enticknap for her assistance with phylogenetic data analysis.

Financial support: The authors gratefully acknowledge financial support from the National Science Foundation under a grant awarded to L.Z. Santiago-Vázquez (awards #0310283 and 0514500) and the Center of Excellence in Biomedical and Marine Biotechnology, Florida Atlantic University. R.G. Kerr acknowledges support from the Natural Sciences and Engineering Council of Canada (NSERC), the Canada Research Chair Program and the Jeanne and Jean-Louis Lévesque Foundation.

REFERENCES

- AL-ZEREINI, W.; FOTSO FONDJA YAO, C.B.; LAATSCH, H. and ANKE, H. (2010). Aqabamycins A-G: Novel nitro maleimides from a marine *Vibrio* species. I. Taxonomy, fermentation, isolation and biological activities. *The Journal of Antibiotics*, vol. 63, no. 6, p. 297-301.
- BOURNE, D.; IIDA, Y.; UTHICKE, S. and SMITH-KEUNE, C. (2008). Changes in coral-associated microbial communities during a bleaching event. *The ISME Journal*, vol. 2, no. 4, p. 350-363. [\[CrossRef\]](#)
- BRÜCK, T.B.; BRÜCK, W.M.; SANTIAGO-VÁZQUEZ, L.Z.; MCCARTHY, P.J. and KERR, R.G. (2007). Diversity of the bacterial communities associated with the azooxanthellate deep water octocorals *Leptogorgia minimata*, *Iciligorgia schrammi*, and *Swiftia exertia*. *Marine Biotechnology*, vol. 9, no. 5, p. 561-576. [\[CrossRef\]](#)
- CHARYULU, E.M.; SEKARAN, G.; RAJAKUMAR, G.S. and GNANAMANI, A. (2009). Antimicrobial activity of secondary metabolite from marine isolate, *Pseudomonas* sp. against Gram positive and negative bacteria including MRSA. *Indian Journal of Experimental Biology*, vol. 47, no. 12, p. 964-968.
- DAVIDSON, S.; ALLEN, S.; LIM, G.; ANDERSON, C. and HAYGOOD, M. (2001). Evidence for the biosynthesis of bryostatins by the bacterial symbiont "*Candidatus* Endobugula sertula" of the bryozoan *Bugula neritina*. *Applied and Environmental Microbiology*, vol. 67, no. 10, p. 4531-4537. [\[CrossRef\]](#)
- DONIA, M.S.; FRICKE, W.F.; PARTENSKY, F.; COX, J.; ELSHAHAWI, S.I.; WHITE, J.R.; PHILLIPPY, A.M.; SCHATZ, M.C.; PIEL, J.; HAYGOOD, M.G.; RAVEL, J. and SCHMIDT, E.W. (2011). Complex microbiome underlying secondary and primary metabolism in the tunicate-*Prochloron* symbiosis. *Proceedings of the National Academy of Sciences of the United States of America*, vol. 108, no. 51, p. E1423-E1432. [\[CrossRef\]](#)
- DOWNS, C.; FAUTH, J.E.; HALAS, J.C.; DUSTAN, P.; BEMISS, J. and WOODLEY, C.M. (2002). Oxidative stress and seasonal coral bleaching. *Free Radical Biology and Medicine*, vol. 33, no. 4, p. 533-543. [\[CrossRef\]](#)
- ENTICKNAP, J.J.; KELLY, M.; PERAUD, O. and HILL, R.T. (2006). Characterization of a culturable alphaproteobacterial symbiont common to many marine sponges and evidence for vertical transmission via sponge larvae. *Applied and Environmental Microbiology*, vol. 72, no. 5, p. 3724-3732. [\[CrossRef\]](#)
- FAULKNER, D.J. (2002). Marine natural products. *Natural Product Reports*, vol. 19, no. 1, p. 1-49. [\[CrossRef\]](#)
- FRENZ-ROSS, J.L.; ENTICKNAP, J.J. and KERR, R.G. (2008). The effect of bleaching on the terpene chemistry of *Plexaurella fusifera*: Evidence that zooxanthellae are not responsible for sesquiterpene production. *Marine Biotechnology*, vol. 10, no. 5, p. 572-578. [\[CrossRef\]](#)
- GIL-AGUDELO, D.L.; MYERS, C.; SMITH, G.W. and KIM, K. (2006). Changes in the microbial communities associated with *Gorgonia ventalina* during aspergillosis infection. *Diseases of Aquatic Organisms*, vol. 69, no. 1, p. 89-94. [\[CrossRef\]](#)
- HARDER, T.; LAU, S.C.K.; DOBRETISOV, S.; FANG, T.K. and QIAN, P.-Y. (2003). A distinctive epibiotic bacterial community on the soft coral *Dendronephthya* sp. and antibacterial activity of coral tissue extracts suggest a chemical mechanism against bacterial epibiosis. *FEMS Microbiology Ecology*, vol. 43, no. 3, p. 337-347. [\[CrossRef\]](#)
- IVANOVA, E.P.; BOWMAN, J.P.; LYSENKO, A.M.; ZHUKOVA, N.V.; GORSHKOVA, N.M.; KUZNETSOVA, T.A.; KALINOVSKAYA, N.I.; SHEVCHENKO, L.S. and MIKHAILOV, V.V. (2005). *Erythrobacter vulgaris* sp. nov., a novel organism isolated from the marine invertebrates. *Systematic and Applied Microbiology*, vol. 28, no. 2, p. 123-130. [\[CrossRef\]](#)
- JACOBSON, P.B. and JACOBS, R.S. (1992a). Fuscoidin: An anti-inflammatory marine natural product which selectively inhibits 5-lipoxygenase. Part I: Physiological and biochemical studies in murine inflammatory models. *Journal of Pharmacology and Experimental Therapeutics*, vol. 262, no. 2, p. 866-873.
- JACOBSON, P.B. and JACOBS, R.S. (1992b). Fuscoidin: An anti-inflammatory marine natural product which selectively inhibits 5-lipoxygenase. Part II: Biochemical studies in the human neutrophil. *Journal of Pharmacology and Experimental Therapeutics*, vol. 262, no. 2, p. 874-882.
- JANNASCH, H.W. and JONES, G.E. (1959). Bacterial populations in sea water as determined by different methods of enumeration. *Limnology and Oceanography*, vol. 4, no. 2, p. 128-139. [\[CrossRef\]](#)
- KLINE, D.I.; KUNTZ, N.M.; BREITBART, M.; KNOWLTON, N. and ROHWER, F. (2006). Role of elevated organic carbon levels and microbial activity in coral mortality. *Marine Ecology Progress Series*, vol. 314, p. 119-125. [\[CrossRef\]](#)

- KOREN, O. and ROSENBERG, E. (2006). Bacteria associated with mucus and tissues of the coral *Oculina patagonica* in summer and winter. *Applied and Environmental Microbiology*, vol. 72, no. 8, p. 5254-5259. [\[CrossRef\]](#)
- LAM, K.S. (2006). Discovery of novel metabolites from marine actinomycetes. *Current Opinion in Microbiology*, vol. 9, no. 3, p. 245-251. [\[CrossRef\]](#)
- LEMA, K.A.; WILLIS, B.L. and BOURNE, D.G. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Applied and Environmental Microbiology*, vol. 78, no. 9, p. 3136-3144. [\[CrossRef\]](#)
- LITTMAN, R.A.; BOURNE, D.G. and WILLIS, B.L. (2010). Responses of coral-associated bacterial communities to heat stress differ with *Symbiodinium* type on the same coral host. *Molecular Ecology*, vol. 19, no. 9, p. 1978-1990. [\[CrossRef\]](#)
- MERLE, P.L.; SABOURAULT, C.; RICHIER, S.; ALLEMAND, D. and FURLA, P. (2007). Catalase characterization and implication in bleaching of a symbiotic sea anemone. *Free Radical Biology and Medicine*, vol. 42, no. 2, p. 236-246. [\[CrossRef\]](#)
- MINCER, T.J.; SPYERE, A.; JENSEN, P.R. and FENICAL, W. (2004). Phylogenetic analyses and diterpenoid production by marine bacteria of the genus *Saprospira*. *Current Microbiology*, vol. 49, no. 4, p. 300-307. [\[CrossRef\]](#)
- NEDASHKOVSKAYA, O.I.; KIM, S.B.; LYSENKO, A.M.; FROLOVA, G.M.; MIKHAILOV, V.V. and BAE, K.S. (2005). *Bizionia paragorgiae* gen. nov., sp. nov., a novel member of the family *Flavobacteriaceae* isolated from the soft coral *Paragorgia arborea*. *International Journal of Systematic and Evolutionary Microbiology*, vol. 55, no. 1, p. 375-378. [\[CrossRef\]](#)
- NEEDHAM, J.; KELLY, M.T.; ISHIGE, M. and ANDERSEN, R.J. (1994). Andrimid and moiramides A-C, metabolites produced in culture by a marine isolate of the bacterium *Pseudomonas fluorescens*: Structure elucidation and biosynthesis. *The Journal of Organic Chemistry*, vol. 59, no. 8, p. 2058-2063. [\[CrossRef\]](#)
- NEI, M. and KUMAR, S. (2000). *Molecular Evolution and Phylogenetics*. Oxford University Press, New York. 333 p. ISBN 978-0-195-13585-5.
- PORETSKY, R.S.; SUN, S.; MOU, X. and MORAN, M.A. (2010). Transporter genes expressed by coastal bacterioplankton in response to dissolved organic carbon. *Environmental Microbiology*, vol. 12, no. 3, p. 616-627. [\[CrossRef\]](#)
- QI, S.H.; XU, Y.; GAO, J.; QIAN, P.Y. and ZHANG, S. (2009). Antibacterial and antilarval compounds from marine bacterium *Pseudomonas rhizosphaerae*. *Annals of Microbiology*, vol. 59, no. 2, p. 229-233. [\[CrossRef\]](#)
- RADJASA, O.K. and SABDONO, A. (2009). Bacterial symbionts of reef's invertebrates as a sustainable source of marine natural products. *Current Research in Bacteriology*, vol. 2, no. 1, p. 7-13. [\[CrossRef\]](#)
- RESHEF, L.; KOREN, O.; LOYA, Y.; ZILBER-ROSENBERG, I. and ROSENBERG, E. (2006). The coral probiotic hypothesis. *Environmental Microbiology*, vol. 8, no. 12, p. 2068-2073. [\[CrossRef\]](#)
- ROHWER, F.; SEGURITAN, V.; AZAM, F. and KNOWLTON, N. (2002). Diversity and distribution of coral-associated bacteria. *Marine Ecology Progress Series*, vol. 243, p. 1-10. [\[CrossRef\]](#)
- ROSENBERG, E.; KOREN, O.; RESHEF, L.; EFRONY, R. and ZILBER-ROSENBERG, I. (2007). The role of microorganisms in coral health, disease and evolution. *Nature Reviews Microbiology*, vol. 5, no. 5, p. 355-362. [\[CrossRef\]](#)
- SANTIAGO-VÁZQUEZ, L.Z.; BRÜCK, T.B.; BRÜCK, W.M.; DUQUE-ALARCÓN, A.P.; MCCARTHY, P.J. and KERR, R.G. (2007). The diversity of the bacterial communities associated with the azooxanthellate hexacoral *Cirripathes lutkeni*. *The ISME Journal*, vol. 1, no. 7, p. 654-659. [\[CrossRef\]](#)
- SHIEH, W.Y.; LIN, Y.-T. and JEAN, W.D. (2004). *Pseudovibrio denitrificans* gen. nov., sp. nov., a marine, facultatively anaerobic, fermentative bacterium capable of denitrification. *International Journal of Systematic and Evolutionary Microbiology*, vol. 54, no. 6, p. 2307-2312. [\[CrossRef\]](#)
- SHIN, J. and FENICAL, W. (1991). Fuscosides A-D: Anti-inflammatory diterpenoid glycosides of new structural classes from the Caribbean gorgonian *Eunicea fusca*. *The Journal of Organic Chemistry*, vol. 56, no. 9, p. 3153-3158. [\[CrossRef\]](#)
- SHNIT-ORLAND, M. and KUSHMARO, A. (2009). Coral mucus-associated bacteria: A possible first line of defense. *FEMS Microbiology Ecology*, vol. 67, no. 3, p. 371-380. [\[CrossRef\]](#)
- SUNAGAWA, S.; WILSON, E.C.; THALER, M.; SMITH, M.L.; CARUSO, C.; PRINGLE, J.R.; WEIS, V.M.; MEDINA, M. and SCHWARZ, J.A. (2009). Generation and analysis of transcriptomic resources for a model system on the rise: The sea anemone *Aiptasia pallida* and its dinoflagellate endosymbiont. *BMC Genomics*, vol. 10, no. 258. [\[CrossRef\]](#)
- UNSON, M.; HOLLAND, N. and FAULKNER, D. (1994). A brominated secondary metabolite synthesized by the cyanobacterial symbiont of a marine sponge and accumulation of the crystalline metabolite in the sponge tissue. *Marine Biology*, vol. 119, no. 1, p. 1-11. [\[CrossRef\]](#)
- WATSON, S.W.; BOCK, E.; VALOIS, F.W.; WATERBURY, J.B. and SCHLOSSER, U. (1986). *Nitrospira marina* gen. nov. sp. nov.: A chemolithotrophic nitrite-oxidizing bacterium. *Archives of Microbiology*, vol. 144, no. 1, p. 1-7. [\[CrossRef\]](#)
- WEBSTER, N.S.; SMITH, L.D.; HEYWARD, A.J.; WATTS, J.E.M.; WEBB, R.I.; BLACKALL, L.L. and NEGRI, A.P. (2004). Metamorphosis of a scleractinian coral in response to microbial biofilms. *Applied and Environmental Microbiology*, vol. 70, no. 2, p. 1213-1221. [\[CrossRef\]](#)
- WEBSTER, N.S. and BOURNE, D. (2007). Bacterial community structure associated with the Antarctic soft coral, *Alcyonium antarcticum*. *FEMS Microbiology Ecology*, vol. 59, no. 1, p. 81-94. [\[CrossRef\]](#)
- WEGLEY, L.; EDWARDS, R.; RODRIGUEZ-BRITO, B.; LIU, H. and ROHWER, F. (2007). Metagenomic analysis of the microbial community associated with the coral *Porites astreoides*. *Environmental Microbiology*, vol. 9, no. 11, p. 2707-2719. [\[CrossRef\]](#)

How to reference this article:

DUQUE-ALARCÓN, A.; SANTIAGO-VÁZQUEZ, L.Z. and KERR, R.G. (2012). A microbial community analysis of the octocoral *Eunicea fusca*. *Electronic Journal of Biotechnology*, vol. 15, no. 5. <http://dx.doi.org/10.2225/vol15-issue5-fulltext-11>