

Relationship of left ventricular and atrial dimensions with moderate to severe left ventricular diastolic dysfunction (grade II and above)

Muhammad Hamza, Mishal Fatima, Muhammad Masood, Hafiz Umar Masood, Ghazal Tasleem, Hassaan Ahmed, Maha Nadir, Zubair Satti

Rawalpindi Medical College, Cardiology.

Abstract:

Introduction: Left ventricular diastolic dysfunction (DD) is an entity in which the ventricle fails to fill up properly due to impaired ventricular relaxation and/or decreased compliance. The diagnosis of diastolic dysfunction is based on a variety of parameters in doppler echocardiography. However, some parameters like interventricular septal thickness in diastole (IVSd), posterior wall thickness in diastole (PWd), left ventricular internal end diastolic and systolic diameters (LVIDD and LVISD) along with left atrial diameters (LAD) have yet to be evaluated for the diagnostic workup of DD.

Methods: A case control study was done in the cardiology department from patient records from 2016 to 2018. Patients were diagnosed as diastolic dysfunction grade II and above by doppler echocardiography. IVSd, PWd, LVIDD, LAD, LVISD were obtained through 2-D echocardiography.

Results: Patients with DD had greater LAD, IVSd and PWd and decreased LVIDD and LVISD as compared to control group. Overall, IVSd was the most significant predictor (OR 1.52 95%CI 1.35-1.71) of DD followed by PWd and LAD. Similarly, LAD, IVSd and PWd had higher sensitivity and specificity than LVIDD and LVISD.

Conclusion: IVSd, LAD and PWd showed significant performance in the diagnosis of diastolic dysfunction and hence can be used as a screening and diagnostic tool in diastolic dysfunction of grade II and above.

Keywords: Heart failure; diastolic; echocardiography; left ventricle.

DOI: <https://dx.doi.org/10.4314/ahs.v20i4.27>

Cite as: Hamza M, Fatima M, Masood M, Masood HU, Tasleem G, Ahmed H, et al. Relationship of left ventricular and atrial dimensions with moderate to severe left ventricular diastolic dysfunction (grade II and above). *Afri Health Sci.* 2020;20(4):1749-53. <https://dx.doi.org/10.4314/ahs.v20i4.27>

Introduction

Left ventricular diastolic dysfunction (DD) is a clinical entity in which the ventricle fails to fill up properly due to impaired ventricular relaxation and/or decreased compliance of the ventricles. The most common cause of this is concentric hypertrophy due to arterial hypertension, senility and aortic stenosis but other conditions like restrictive cardiopathy and constrictive pericarditis can also cause diastolic dysfunction. This adaptive response is mainly due to sustained hypertension, aortic stenosis and advanced age. Diastolic dysfunction has similar symptoms as systolic dysfunction like dyspnea, exercise intolerance and edema but it has preserved or slightly decreased ejection fraction¹. DD is graded into 3 types from mild (grade I), moderate (grade II) and severe (grade III)² The diagnosis of diastolic dysfunction

is based on doppler echocardiography and E/A ratio (E is the velocity of blood during early phase of diastole and A is the velocity during atrial phase) is the main diagnostic criteria to grade diastolic dysfunction. The left ventricular geometric indices like interventricular septal thickness in diastole (IVSd), posterior wall thickness in diastole (PWd), left ventricular internal end diastolic and systolic diameters (LVIDD and LVISD) along with left atrial diameters (LAD) have mostly remained in the realm of theoretical description of diastolic dysfunction. Literature review shows that these indices remain neglected in the clinical screening or diagnosis of diastolic dysfunction^{3,4}.

DD is quite a common disease with an overall 25-30% prevalence in general population and its prevalence increases with age. It is correlated with significant cardiac morbidity and mortality if remained undiagnosed and untreated.⁵ This problem is made even more challenging due to the relative ambiguity and limitations in its diagnostic indices (namely pseudo normalization of E/A in grade II disease)^{1,2,5}. Therefore, multi modal diagnostic criteria involving many indices are encouraged for diagnostic work up of DD.²

Corresponding author:

Muhammad Hamza,
Rawalpindi Medical College, Cardiology.
Email: goodoldhamza@gmail.com

Left ventricular geometrical indices have been mainly confined to theoretical concept of DD and little work has been done on their application in clinical diagnosis. These indices can easily be assessed using echocardiography and can provide additional tools in the multi modal diagnostic arsenal available for DD.

Many studies have shown the values of left ventricular geometrical indices in DD. IVSd, PWD, LAD and ejection fraction (EF) increases while LVIDD and LVISD decreases in DD^{6,4}. However, there is a relative dearth of literature that evaluates the diagnostic abilities of these indices. Recently, a novel parameter LAD/LVIDD was proposed as a screening test for DD of grade II and above⁶.

Thus, this study aims to evaluate left ventricular geometrical indices in the screening and diagnosis of DD grade II and above.

Materials and Methods

A case control study was done in the cardiology department from patient records from 2016 to 2018. Left Atrial Diameters (LAD) were obtained on 2-Dimensional (2-D) Echocardiography in apical views (four chamber view) during end systole. Interventricular septal and posterior wall thicknesses at end-diastole along with left ventricular internal diameters at both end diastole and end systole were obtained on 2-Dimensional (2-D) Echocardiography in parasternal long axis view (4 chamber view).

In patients with normal EF without any evidence of myocardial disease, diastolic dysfunction was diagnosed on 2D and Doppler Echocardiography when all three of these variables were abnormal (Annular septal e' velocity less than 7cm/sec or lateral e' velocity less than 10cm/sec, E/e' ratios greater than 14, LA volume index greater than 34mL/m²) as per American society of Echocardiography guidelines². Grade 2 DD was then categorized when E/A ratios were either less than or equal to 0.8 with Mitral E velocities greater than 50 cm/sec or when E/A ratios were between 0.8 and 2.2.

In patients with reduced EF (less than 50%) or patients with preserved EF (greater or equal to 50%) and myocardial disease (e.g. Coronary Artery Disease) diastolic dysfunction grade II was diagnosed when E/A ratios

were either less than or equal to 0.8 with Mitral E velocities greater than 50 cm/sec or when E/A ratios were between 0.8 and 2 along with these two abnormal variables (E/e' greater than 14 and LA volume index greater than 34mL/m²). Patients having E/A ratio greater or equal to 2 were categorized as Diastolic Dysfunction grade III. Tricuspid regurgitation (TR) velocities were not acquired for the assessment of diastolic dysfunction and therefore an approach consisting of only E/e', E/A, LA volume index and mitral annular velocities was utilized as per American Society of Echocardiographic guidelines 2016 when TR velocities are not available²

178 cases were then randomly selected and compared with 191 controls after careful matching for age, sex, hypertension, diabetes. Patients with rhythm abnormalities and valvular abnormalities were excluded (to facilitate the diagnosis of diastolic dysfunction and to minimize the effect on atrial and ventricular dimensions). Patients with grade I diastolic dysfunction are asymptomatic with a relatively benign course that does not progress in severity. It was excluded from this study due to its negligible effect on ventricular and atrial geometry⁷. Left ventricular geometrical characteristics are size, weight, thickness and volume of the left ventricles that can be obtained using echocardiography. Diastolic dysfunction (DD) is defined as symptoms of systolic dysfunction with preserved or slightly reduced ejection fraction with impaired left ventricular relaxation and decreased compliance.

IVSd, PWD, LAD, LVIDD, LVISD and EF were compared between cases and controls using independent t-test. These indices were correlated with diastolic dysfunction using Spearman correlation. Binary logistic regression (using the above indices) was used to predict diastolic dysfunction. A Receiver Operating Characteristic (ROC) curve was used to check sensitivity and specificity of the above indices. IBM Statistical package for social sciences (SPSS) version 23 was used. A p-value of less than 0.05 was considered to be significant.

Results

The patients with diastolic dysfunction were older and had greater LAD, IVSd and PWD as compared to control group. LVIDD and LVISD were significantly decreased in the diseased group (Table 1).

Table 1: Left atrial and ventricular characteristics of the study population

Character	Control group (n=191)	Disease group with DD of grade II and above (n=178)	Spearman Correlation R	P-value
Gender	M= 81, F=110	M=73, F=105	---	0.8
Age	49.5 ± 18.7	54.8 ± 16	0.20	<0.001
LAD (mm)	34.9 ± 7.4	44.6 ± 7	0.50	<0.001
LVIDD (mm)	48.8 ± 7.9	43.4 ± 5.2	-0.30	<0.001
LVISD (mm)	33.4 ± 8.1	26.3 ± 3.9	-0.40	<0.001
IVSd (mm)	9.3 ± 3.0	11.4 ± 2.1	0.46	<0.001
PWd (mm)	9.6 ± 3.1	10.6 ± 1.6	0.30	<0.001
EF (%)	49.1 ± 13.3	56.1 ± 7.3	0.30	<0.001

Values are represented as mean ± SD LAD= Left atrial Diameter, LVIDD= Left ventricular internal end diastolic diameter, LVISD= Left Ventricular internal end systolic diameter, IVSd = Interventricular septal diameter, PWd= Posterior Wall Diameter, EF= Ejection Fraction

Overall, IVSD was most predictive of DD (OR 1.52, 95% CI 1.35-1.71) as compared to other variables (Table 2).

Table 2: Binary logistic Regression for diastolic dysfunction

Variable	Regression Coefficients (B)	Odds Ratio (ExpB)	95 % Confidence Interval	P-value
IVSd	0.421	1.52	1.35 - 1.71	0.009
PWd	0.185	1.20	1.09 - 1.32	0.012
LAD	0.164	1.18	1.13 - 1.2	<0.001
LVISD	-0.154	0.85	0.82 - 0.89	<0.001
EF	0.064	1.06	1.04 - 1.09	<0.001

LAD= Left atrial Diameter, LVIDD= Left ventricular internal end diastolic diameter, LVISD= Left Ventricular internal end systolic diameter, IVSd = Interventricular septal diameter, PWd= Posterior Wall Diameter, EF= Ejection Fraction

LAD, IVSd and PWd had higher specificity and sensitivity on ROC curve as compared to LVISD and LVIDD. A new parameter combining LAD, IVSd and PWd was 82% sensitive and 78% specific (Table 3).

Table 3: Receiver operating characteristic Curve for diastolic dysfunction

Variable	Area under the curve	P-value	Selected cut off value	Sensitivity	Specificity
LAD	0.82	<0.001	40.5mm	74.2%	79%
IVSd	0.76	<0.001	10.8mm	72%	76%
PWd	0.72	<0.001	10.5mm	71.3%	72%
LVISD	0.25	<0.001	28.5mm	32%	31.9%
LVIDD	0.26	<0.001	44.5mm	37.1%	34%
LAD+IVSd+PWd	0.84	<0.001	58.9 mm	82.6%	74%

LAD= Left atrial Diameter, LVIDD= Left ventricular internal end diastolic diameter, LVISD= Left Ventricular internal end systolic diameter, IVSd = Interventricular septal diameter, PWd= Posterior Wall Diameter, EF= Ejection Fraction

Discussion

The ambiguity surrounding diastolic dysfunction stems from its resemblance to systolic dysfunction in clinical presentation and investigations like ECG or chest radiography. Diastolic dysfunction has a chronic long standing course which is asymptomatic in initial stages but can cause significant complications like ischemic heart disease, heart failure or atrial fibrillation in more severe cases. Diastolic dysfunction is common in long standing hypertension, advanced age and diabetes, all of which are also independent risk factors for ischemic heart disease⁸. Therefore, early diagnosis of diastolic dysfunction in these clinical settings can help improve outcomes⁹ by aggressive treatment and management (diuretics, calcium channel blockers, beta blocker (especially nebivolol), angiotensin receptor blocker^{10,11}). Screening of diastolic dysfunction has also led to improvement in better control of underlying cause (diabetes, hypertension or aortic stenosis) and thus improving outcomes and preventing dire complications¹².

The diagnosis of diastolic dysfunction is similarly mired in ambiguity and multiple modalities involving various investigations are used in its confirmation. 2-D doppler echocardiography is routinely used to evaluate DD due to its ease of use and cost. Multiple doppler indices like E/A ratio or E/e' ratio are used for this purpose¹³ but these indices also have many limitations as well^{2,14}. Therefore, new echocardiographic parameters are continuously investigated to be used alongside Doppler indices to enhance diagnostic fidelity⁶. Ventricular and atrial dimensions can therefore provide a new avenue to augment current diagnostic criteria. IVSd, PWd, LAD, LVIDD and LVISD have been shown to be significantly associated with diastolic dysfunction in our study as well as in other studies^{7,15}. LAD, IVSd and PWd were the most promising dimensions in our study. LAD was 74% sensitive and 79% specific for grade II DD and above. A slightly less sensitivity and specificity for LAD was calculated by another researcher (69% sensitive and 77% specific for grade II and above)¹⁶ whereas others have demonstrated a slightly better specificity and sensitivity of LAD than the present study¹⁷. This demonstrates a promising role of LAD as an adjunct to doppler indices for the evaluation of DD. Similarly ventricular dimensions have shown moderate to poor correlation with diastolic dysfunction in this study. The best association was seen with IVSd with $r=0.457$, AUC 0.762, 72% sensitive and 76% specific. IVSd has been shown to be significantly increased in DD in other studies^{4,7},

however no study could be found that has evaluated its efficacy in the diagnosis of DD. LVIDD and LVISD is negatively correlated with DD in our study but it is not highly specific or sensitive. Other studies have shown a positive correlation of DD with LVIDD and LVISD^{7,17,18} which suggests mixed systolic and diastolic dysfunction among patients in these studies. Others have shown a negative correlation⁶. PWd is also significantly associated with DD and is specific and sensitive in the diagnosis of DD as shown by other studies^{7,17}.

A combination of IVSd, PWd and LAD was 82% sensitive and 74% specific on ROC curve which is higher than each of the dimension alone. This shows that a combined approach utilizing all these dimensions would be helpful in making a correct diagnosis. However, there still is a need to compare and combine these parameters with current doppler indices. This would help ascertain real life benefit of these dimensions in the diagnosis of DD.

Conclusion

IVSd, LAD and PWd showed significant performance in the diagnosis of diastolic dysfunction and hence can be used as a screening and diagnostic tool in diastolic dysfunction of grade II and above.

References

1. Gutierrez C, Blanchard DG. Diastolic heart failure: challenges of diagnosis and treatment. *Am Fam Physician*. 2004 Jun;69(11):2609–16.
2. Nagueh SF, Smiseth OA, Appleton CP, Byrd BF, Dokainish H, Edvardsen T, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr Internet*. 2016;29(4):277–314. Available from: <http://dx.doi.org/10.1016/j.echo.2016.01.011>
3. Osranek M, Seward JB, Buschenreithner B, Bergler-Klein J, Heger M, Kklar U, et al. Diastolic function assessment in clinical practice: the value of 2-dimensional echocardiography. *Am Heart J*. 2007 Jul;154(1):130–6.
4. Adamu GU, Katibi AI, Opadijo GO, Omotoso ABO, Araoye AM. Prevalence of left ventricular diastolic dysfunction in newly diagnosed Nigerians with systemic hypertension: a pulsed wave Doppler echocardiographic study. *Afr Health Sci*. 2010 Jun;10(2):177–82.
5. Naylor M, Cooper LL, Enserro DM, Xanthakis V, Larson MG, Benjamin EJ, et al. Left Ventricular Diastolic Dysfunction in the Community: Impact of Diag-

- nostic Criteria on the Burden, Correlates, and Prognosis. *J Am Heart Assoc.* 2018 Jun 1;7(11):e008291.
6. Ul Haq I, Saeed M, Hamza M, Ali AS, Nadir M. Left atrial to left ventricular end diastolic diameter ratio in left ventricular diastolic dysfunction. *Pakistan J Physiol.* 2017;13(1):15–8.
 7. Cacciapuoti F, Scognamiglio A, Paoli VD, Romano C, Cacciapuoti F. Left Atrial Volume Index as Indicator of Left Ventricular Diastolic Dysfunction: Comparison between Left Atrial Volume Index and Tissue Myocardial Performance Index. *J Cardiovasc Ultrasound.* 2012;20(1):25–9.
 8. Jeong E-M, Dudley Jr SC. Diastolic dysfunction. *Circ J.* 2015;79(3):470–7.
 9. Ghany R, Palacio A, Chen G, Dawkins E, Ghany A, Forbes E, et al. A screening echocardiogram to identify diastolic dysfunction leads to better outcomes. *Echocardiography.* 2017 Aug;34(8):1152–8.
 10. Flather MD, Shibata MC, Coats AJS, Van Veldhuisen DJ, Parkhomenko A, Borbola J, et al. Randomized trial to determine the effect of nebivolol on mortality and cardiovascular hospital admission in elderly patients with heart failure (SENIORS). *Eur Heart J.* 2005 Feb;26(3):215–25.
 11. Aziz F, Tk L, Enweluzo C. Diastolic Heart Failure : A Concise Review. *J Clin Med Res.* 2013;5(5):327–34.
 12. Romano S, Di Mauro M, Fratini S, Guarracini L, Guarracini F, Poccia G, et al. Early diagnosis of left ventricular diastolic dysfunction in diabetic patients: a possible role for natriuretic peptides. *Cardiovasc Diabetol.* 2010;9:89.
 13. Dokainish H. Left ventricular diastolic function and dysfunction: Central role of echocardiography. *Glob Cardiol Sci Pract.* 2015;2015:3.
 14. Cahill JM, Horan M, Quigley P, Maurer B, McDonald K. Doppler-echocardiographic indices of diastolic function in heart failure admissions with preserved left ventricular systolic function. *Eur J Heart Fail.* 2002;4(4):473–8.
 15. van Dalen BM, Strachinaru M, van der Swaluw J, Geleijnse ML. A simple, fast and reproducible echocardiographic approach to grade left ventricular diastolic function. *Int J Cardiovasc Imaging.* 2016;32(5):743–52.
 16. Pritchett AM, Mahoney DW, Jacobsen SJ, Rodeheffer RJ, Karon BL, Redfield MM. Diastolic dysfunction and left atrial volume: A population-based study. *J Am Coll Cardiol.* 2005;45(1):87–92.
 17. El Aouar LMM, Meyerfreud D, Magalhães P, Rodrigues SL, Baldo MP, Brasil Y, et al. Relationship between left atrial volume and diastolic dysfunction in 500 Brazilian patients. *Arq Bras Cardiol.* 2013;101(1):52–8.
 18. Germing A, Gotzmann M, Schikowski T, Vierkötter A, Ranft U, Mügge A, et al. Diastolic dysfunction without abnormalities in left atrial and left ventricular geometry does not affect quality of life in elderly women. *Exp Clin Cardiol.* 2011;16(2):2010–2.