



TITLE:

SHELAH-STRONG TYPE AND ALGEBRAIC CLOSURE OVER A HYPERIMAGINARY (Model theoretic aspects of the notion of independence and dimension)

AUTHOR(S):

LEE, HYOOYON

CITATION:

LEE, HYOOYON. SHELAH-STRONG TYPE AND ALGEBRAIC CLOSURE OVER A HYPERIMAGINARY (Model theoretic aspects of the notion of independence and dimension). 数理解析研究所講究録 2022, 2218: 93-99

ISSUE DATE:

2022-05

URL:

<http://hdl.handle.net/2433/277123>

RIGHT:

SHELAH-STRONG TYPE AND ALGEBRAIC CLOSURE OVER A HYPERIMAGINARY

HYOYOON LEE

DEPARTMENT OF MATHEMATICS, YONSEI UNIVERSITY

ABSTRACT. We characterize Shelah-strong type over a hyperimaginary with the algebraic closure of a hyperimaginary. Also, we present and take a careful look at an example that witnesses $\text{acl}^{\text{eq}}(e)$ is not interdefinable with $\text{acl}(e)$ where e is a hyperimaginary.

Fix a first order language \mathcal{L} , complete theory T and monster model \mathcal{M} . **Throughout, fix a hyperimaginary $e = a_E$ where a is a (possibly infinite) real tuple and E is an \emptyset -type-definable equivalence relation on $\mathcal{M}^{|a|}$.**

Most of the facts and remarks whose proofs are omitted can be found in the author's dissertation [6].

Fact 1.

- (1) A real tuple b is simply $b/(\bigwedge_{i<\alpha} x_i = y_i)$ where $b = (b_i)_{i<\alpha}$, hence can be seen as (that is, interdefinable with) a hyperimaginary; an imaginary tuple $(b_i/F_i)_{i<\alpha}$ is $(b_i)_{i<\alpha}/(\bigwedge_{i<\alpha} F_i(x_i, y_i))$ where all x_i, y_i 's are disjoint, hence is a hyperimaginary as well. **In this regard, considering over a set of real elements or a set of imaginaries can be safely replaced by considering over a single hyperimaginary.**
- (2) In the same manner as above, a sequence of hyperimaginaries can be regarded as a single hyperimaginary: A tuple of hyperimaginaries $(b_i/F_i)_{i<\alpha}$ is interdefinable with $(b_i)_{i<\alpha}/(\bigwedge_{i<\alpha} F_i(x_i, y_i))$ where all x_i, y_i 's are disjoint.

Definition 2.

- (1) For any hyperimaginary e' , we denote $e' \in \text{dcl}(e)$ and say e' is *definable over e* if $f(e') = e'$ for all $f \in \text{Aut}_e(\mathcal{M})$.
- (2) For any hyperimaginary e' , we denote $e' \in \text{bdd}(e)$ and say e' is *bounded over e* if $\{f(e') : f \in \text{Aut}_e(\mathcal{M})\}$ is bounded.

Remark 3. In Definition 2, $e' \in \text{dcl}(e)$ and $e' \in \text{bdd}(e)$ are independent of the choice of a monster model \mathcal{M} .

Proof. It is easy, but anyway we prove it. Let $\mathcal{M} \prec \mathcal{M}'$ be monster models of T . Suppose that there are only κ -many automorphic images of e' in \mathcal{M} , whereas there are at least κ^+ images in \mathcal{M}' . Say $e' = b_F$ where b is a real tuple and F is an \emptyset -type-definable equivalence relation. Let $(b_i/F)_{i<\kappa^+}$ be an enumeration of automorphic images of b_F in \mathcal{M}' . Since there is $(b'_i)_{i<\kappa^+} \equiv_a (b_i)_{i<\kappa^+}$ where each $b'_i \in \mathcal{M}$, there are at least κ^+ -many conjugates of b_F in \mathcal{M} (recall $e = a/E$), a contradiction. \square

Fact 4.

- (1) A hyperimaginary b_F is called *countable* if $|b|$ is countable. It's not so difficult to prove that any hyperimaginary is interdefinable with a sequence of countable hyperimaginaries (see, for example [5, Lemma 4.1.3]).

- (2) From now on, definable closure of \mathbf{e} , $\text{dcl}(\mathbf{e})$ will be seen as an actual (small) set, the set of all countable hyperimaginaries which are definable over \mathbf{e} : In this way, $\mathbf{e}' \in \text{dcl}(\mathbf{e})$ now means that there is a sequence of countable hyperimaginaries that is interdefinable with \mathbf{e}' and fixed by any $f \in \text{Aut}_{\mathbf{e}}(\mathcal{M})$. Also note that $f \in \text{Aut}_{\text{dcl}(\mathbf{e})}(\mathcal{M})$ if and only if f fixes all hyperimaginaries that are definable over \mathbf{e} . As pointed out in Fact 1(2), $\text{dcl}(\mathbf{e})$ also can be seen as a single hyperimaginary.
- (3) Likewise, the bounded closure of \mathbf{e} , $\text{bdd}(\mathbf{e})$ is the set of all countable hyperimaginaries which are bounded over \mathbf{e} . In the same way as above, $\mathbf{e}' \in \text{bdd}(\mathbf{e})$ means that there is a sequence of countable hyperimaginaries that is interdefinable with \mathbf{e}' , and the number of \mathbf{e} -automorphic images of it is bounded. Again, $f \in \text{Aut}_{\text{bdd}(\mathbf{e})}(\mathcal{M})$ is equivalent to saying that f fixes all hyperimaginaries that are bounded over \mathbf{e} .

Remark & Definition 5.

- (1) For a hyperimaginary \mathbf{e}' , denote $\mathbf{e}' \in \text{acl}(\mathbf{e})$ and say \mathbf{e}' is *algebraic over \mathbf{e}* if $\{f(\mathbf{e}') : f \in \text{Aut}_{\mathbf{e}}(\mathcal{M})\}$ is finite. As in Remark 3, this definition is independent of the choice of a monster model.
- (2) As in Fact 4, the *algebraic closure* of \mathbf{e} , $\text{acl}(\mathbf{e})$ can be regarded as a bounded set of countable hyperimaginaries, which is interdefinable with a single hyperimaginary $b_F \in \text{bdd}(\mathbf{e})$ (but possibly $b_F \notin \text{acl}(\mathbf{e})$).
- (3) Note that given $d_i/L_i \in \text{acl}(\mathbf{e})$ ($i \leq n$), as pointed out in Fact 1, $(d_0/L_0, \dots, d_n/L_n)$ is interdefinable with a single $d_L \in \text{acl}(\mathbf{e})$. Hence by compactness, for any hyperimaginaries b_F and c_F ,

$$b_F \equiv_{\text{acl}(\mathbf{e})} c_F \text{ if and only if } b_F \equiv_{d_L} c_F \text{ for any } d_L \in \text{acl}(\mathbf{e}).$$

Definition 6.

- (1) $\text{Aut}_{\mathbf{e}}(\mathcal{M}) = \{f \in \text{Aut}(\mathcal{M}) : f(\mathbf{e}) = \mathbf{e}\}$ (f may permute the elements of \mathbf{e}).
- (2) $\text{Autf}_{\mathbf{e}}(\mathcal{M})$ is a subgroup of $\text{Aut}_{\mathbf{e}}(\mathcal{M})$ generated by

$$\{f \in \text{Aut}_{\mathbf{e}}(\mathcal{M}) : f \in \text{Aut}_M(\mathcal{M}) \text{ for some } M \models T \text{ such that } \mathbf{e} \in \text{dcl}(M)\}.$$

It can be easily seen that $\text{Autf}_{\mathbf{e}}(\mathcal{M})$ is a normal subgroup of $\text{Aut}_{\mathbf{e}}(\mathcal{M})$.

- (3) The *Lascar group over of T \mathbf{e}* is the quotient group

$$\text{Gal}_{\mathbf{L}}(T, \mathbf{e}) = \text{Aut}_{\mathbf{e}}(\mathcal{M}) / \text{Autf}_{\mathbf{e}}(\mathcal{M}).$$

Remark 7.

- (1) Up to isomorphism, $\text{Gal}_{\mathbf{L}}(T, \mathbf{e})$ is independent of the choice of a monster model \mathcal{M} .
- (2) There are well-defined maps μ and ν such that:

$$\begin{aligned} \text{Aut}_{\mathbf{e}}(\mathcal{M}) &\xrightarrow{\mu} S_M(M) \xrightarrow{\nu} \text{Gal}_{\mathbf{L}}(T, \mathbf{e}) \\ f &\mapsto \text{tp}(f(M)/M) \mapsto \bar{f} = \pi(f) \end{aligned}$$

where M is a small model of T such that $\mathbf{e} \in \text{dcl}(M)$, and $\pi : \text{Aut}_{\mathbf{e}}(\mathcal{M}) \rightarrow \text{Gal}_{\mathbf{L}}(T, \mathbf{e})$ is the canonical projection.

The topology of $\text{Gal}_{\mathbf{L}}(T, \mathbf{e})$ is given by the topology induced by the quotient map ν , and it is independent of the choice of M .

Fact 8.

- (1) $\text{Gal}_{\mathbf{L}}(T, \mathbf{e})$ is a topological group.

- (2) Let $H \leq \text{Aut}_e(\mathcal{M})$ and let $H' = \pi(H) \leq \text{Gal}_L(T, \mathbf{e})$. Then H' is closed in $\text{Gal}_L(T, \mathbf{e})$ and $H = \pi^{-1}(H')$, if and only if $H = \text{Aut}_{e'}(\mathcal{M})$ for some hyperimaginary $e' \in \text{bdd}(\mathbf{e})$.
- (3) Let $H' \leq \text{Gal}_L(T, \mathbf{e})$ be closed and F be an \emptyset -type-definable equivalence relation. Then for $H = \pi^{-1}(H')$, $x_F \equiv_e^H y_F$ is equivalent to $x_F \equiv_{e'} y_F$ for some hyperimaginary $e' \in \text{bdd}(\mathbf{e})$, and hence $x_F \equiv_e^H y_F$ is an e' -invariant type-definable bounded equivalence relation. Especially, if $H' \trianglelefteq \text{Gal}_L(T, \mathbf{e})$, then $x_F \equiv_e^H y_F$ is e -invariant.

Definition 9.

- (1) $\text{Gal}_L^0(T, \mathbf{e})$ denotes the connected component of the identity in $\text{Gal}_L(T, \mathbf{e})$.
- (2) $\text{Aut}_s(\mathcal{M}, \mathbf{e}) := \pi^{-1}(\text{Gal}_L^0(T, \mathbf{e}))$.
- (3) Two hyperimaginaries b_F and c_F are said to have the same *Shelah-strong type* if there is $f \in \text{Aut}_s(\mathcal{M}, \mathbf{e})$ such that $f(b_F) = c_F$, denoted by $b_F \equiv_e^s c_F$.

Remark 10. Note that $\text{Gal}_L^0(T, \mathbf{e})$ is a normal closed subgroup of $\text{Gal}_L(T, \mathbf{e})$ ([4]) and \equiv_e^s is the orbit equivalence relation $\equiv_e^{\text{Aut}_s(\mathcal{M}, \mathbf{e})}$, thus \equiv_e^s is type-definable over \mathbf{e} by Fact 8(3). We denote

$$\text{Gal}_s(T, \mathbf{e}) := \text{Gal}_L(T, \mathbf{e}) / \text{Gal}_L^0(T, \mathbf{e}) \cong \text{Aut}_e(\mathcal{M}) / \text{Aut}_s(\mathcal{M}, \mathbf{e}).$$

Thus $\text{Gal}_s(T, \mathbf{e})$ is a profinite (i.e. compact and totally disconnected) topological group. $\text{Gal}_L^0(T, \mathbf{e})$ is the intersection of all closed (normal) subgroups of finite indices in $\text{Gal}_L(T, \mathbf{e})$, since such an intersection is the identity for a profinite group ([4]).

Proposition 11.

- (1) $\text{Aut}_s(\mathcal{M}, \mathbf{e}) = \text{Aut}_{\text{acl}(\mathbf{e})}(\mathcal{M})$.
- (2) Let b_F, c_F be hyperimaginaries. The following are equivalent.
- $b_F \equiv_e^s c_F$.
 - $b_F \equiv_{\text{acl}(\mathbf{e})} c_F$.

Proof. (1). We claim first that

$$\text{Gal}_L^0(T, \mathbf{e}) = \bigcap \{ \pi(\text{Aut}_{d_L e}(\mathcal{M})) : d_L \in \text{acl}(\mathbf{e}) \}.$$

Let $d_L \in \text{acl}(\mathbf{e})$ where d_L is a hyperimaginary. Say $d_L^0 (= d_L), \dots, d_L^n$ are all the conjugates of d_L over \mathbf{e} . Then any $f \in \text{Aut}_e(\mathcal{M})$ permutes the set $\{d_L^0, \dots, d_L^n\}$. Hence it follows that $\text{Aut}_{d_L e}(\mathcal{M})$ has a finite index in $\text{Aut}_e(\mathcal{M})$. Thus (due to Fact 8(2)) $\pi(\text{Aut}_{d_L e}(\mathcal{M}))$ is a closed subgroup of finite index in $\text{Gal}_L(T, \mathbf{e})$. Then as in Remark 10, we have $\text{Gal}_L^0(T, \mathbf{e}) \leq \pi(\text{Aut}_{d_L e}(\mathcal{M}))$.

Conversely, given a normal closed subgroup $H' \leq \text{Gal}_L(T, \mathbf{e})$ of finite index and $H := \pi^{-1}(H')$, Fact 8(2) says $H' = \pi(\text{Aut}_{b_F e}(\mathcal{M}))$ for some $b_F \in \text{bdd}(\mathbf{e})$. But since H' is of finite index, the same holds for $H = \text{Aut}_{b_F e}(\mathcal{M})$ in $\text{Aut}_e(\mathcal{M})$, and we must have $b_F \in \text{acl}(\mathbf{e})$. Thus the claim follows from Remark 10.

Therefore

$$\begin{aligned} \text{Aut}_s(\mathcal{M}, \mathbf{e}) &= \pi^{-1}(\text{Gal}_L^0(T, \mathbf{e})) = \pi^{-1}\left(\bigcap \{ \pi(\text{Aut}_{d_L e}(\mathcal{M})) : d_L \in \text{acl}(\mathbf{e}) \}\right) \\ &= \bigcap \{ \text{Aut}_{d_L e}(\mathcal{M}) : d_L \in \text{acl}(\mathbf{e}) \} = \text{Aut}_{\text{acl}(\mathbf{e})}(\mathcal{M}), \end{aligned}$$

where the last equality follows by Remark & Definition 5(3).

- (2) follows from (1). □

Recall that $\text{acl}^{\text{eq}}(\mathbf{e}) := \{\mathbf{e}\} \cup (\text{acl}(\mathbf{e}) \cap \mathcal{M}^{\text{eq}})$ is the *eq-algebraic closure* of \mathbf{e} , where as usual \mathcal{M}^{eq} is the set of all imaginary elements (equivalence classes of \emptyset -definable equivalence relations) of \mathcal{M} . Good summary of basic facts concerning imaginary elements can be found in [1, Chapter 1]. The following remark is proved using the proof of [9, Theorem 21].

Remark 12. For any small set A of imaginaries, $\text{acl}^{\text{eq}}(A) (= \text{acl}(A) \cap \mathcal{M}^{\text{eq}})$ is interdefinable with $\text{acl}(A)$.

Proof. Recall that $\text{Gal}_{\mathbb{L}}^0(T, A)$ is the intersection of all closed (normal) subgroups of finite indices in $\text{Gal}_{\mathbb{L}}(T, A)$ (Remark 10). Let H' be a closed subgroup of finite index in $\text{Gal}_{\mathbb{L}}(T, A)$. It suffices to show that $H' = \pi(\text{Aut}_{\mathbf{b}A}(\mathcal{M}))$ for some $\mathbf{b} \in \text{acl}^{\text{eq}}(A)$; by Fact 8(2), we have

$$\begin{aligned} \text{Gal}_{\mathbb{L}}^0(T, \mathbf{e}) &= \bigcap \{H' : H' \text{ is a closed subgroup of finite index in } \text{Gal}_{\mathbb{L}}(T, A)\} \\ &\subseteq \bigcap \{\pi(\text{Aut}_{d_L A}(\mathcal{M})) : d_L \in \text{acl}^{\text{eq}}(A)\}; \end{aligned}$$

thus if we show that $H' = \pi(\text{Aut}_{\mathbf{b}A}(\mathcal{M}))$ for some $\mathbf{b} \in \text{acl}^{\text{eq}}(A)$, then $\text{Gal}_{\mathbb{L}}^0(T, A) = \bigcap \{\pi(\text{Aut}_{d_L A}(\mathcal{M})) : d_L \in \text{acl}^{\text{eq}}(A)\}$. Taking π^{-1} , we get $\text{Aut}_{\text{acl}(A)}(\mathcal{M}) = \text{Aut}_{\text{acl}^{\text{eq}}(A)}(\mathcal{M})$ (by a similar manner as in the last lines of the proof of Proposition 11(1)).

Since H is closed in $\text{Gal}_{\mathbb{L}}(T, A)$, by Fact 8(3), $H = \pi(\text{Aut}_{c_F A}(\mathcal{M}))$ for some hyperimaginary $c_F \in \text{bdd}(A)$. But H has finite index in $\text{Gal}_{\mathbb{L}}(T, A)$, hence (by Fact 8(2),) $c_F \in \text{acl}(A)$. Say $\{c_F = c_0/F, \dots, c_{n-1}/F\}$ is the set of all A -conjugates of c_F .

We may assume that F is closed under conjunction and all formulas in F are symmetric and reflexive. Note that by compactness, there is $\delta \in F$ such that for all $i < j < n$,

$$c_i c_j \neq \exists z_0 z_1 z_2 (\delta(x, z_0) \wedge \delta(z_0, z_1) \wedge \delta(z_1, z_2) \wedge \delta(z_2, y)).$$

Let $\delta^4(x, y) \equiv \exists z_0 z_1 z_2 (\delta(x, z_0) \wedge \delta(z_0, z_1) \wedge \delta(z_1, z_2) \wedge \delta(z_2, y))$, and define $\delta^m(x, y)$ similarly for $m < \omega$. Note that in particular, $\delta(c_i, \mathcal{M})$'s are pairwise disjoint.

Let d be any realization of $\text{tp}(c_0/A)$. Then $d \models \bigvee_{i < n} F(x, c_i)$, thus $d \models \bigvee_{i < n} \delta(x, c_i)$, implying that there is $\varphi(x) \in \text{tp}(c_0/A)$ such that $\varphi(x) \models \bigvee_{i < n} \delta(x, c_i)$, that is, $\varphi(\mathcal{M})$ can be partitioned as $\{\varphi(\mathcal{M}) \cap \delta(c_i, \mathcal{M}) : i < n\}$. Note that we can say $\varphi(x)$ is A -invariant; this is possible because A is a set of imaginaries, not a hyperimaginary.

Claim 1. For any $a', a'' \models \varphi(x)$,

$$a' a'' \models \delta^2(x, y) \text{ if and only if } a', a'' \in \varphi(\mathcal{M}) \cap \delta(c_i, \mathcal{M}) \text{ for some } i < n.$$

Proof. Assume $\models \delta^2(a', a'')$, hence there is some a^* such that $\models \delta(a', a^*) \wedge \delta(a^*, a'')$. Suppose a' and a'' belong to different components for a contradiction. Then

$$\models \delta(c_i, a') \wedge \delta(a', a^*) \wedge \delta(a^*, a'') \wedge \delta(a'', c_j)$$

for some $i \neq j < n$, implying $c_i c_j \models \delta^4(x, y)$, a contradiction.

For the converse, suppose $a', a'' \in \varphi(\mathcal{M}) \cap \delta(c_i, \mathcal{M})$ for some $i < n$. Then $\models \delta(a', c_i) \wedge \delta(c_i, a'')$. \square

Now define

$$L(x, y) \equiv (\neg\varphi(x) \wedge \neg\varphi(y)) \vee (\varphi(x) \wedge \varphi(y) \wedge \delta^2(x, y)).$$

Since $\varphi(x)$ is A -invariant, L is an A -definable equivalence relation with finitely many classes, $\neg\varphi(\mathcal{M}), \varphi(\mathcal{M}) \cap \delta(c_0, \mathcal{M}), \dots, \varphi(\mathcal{M}) \cap \delta(c_{n-1}, \mathcal{M})$. Note that some imaginary $\mathbf{b} (\in \text{acl}(A))$ is interdefinable with c/L ([1, Lemma 1.10]).

Claim 2. c/F and \mathbf{b} (or equivalently, c/L) are interdefinable over A .

Proof. Let $f \in \text{Aut}_A(\mathcal{M})$. Then

$$\begin{aligned} f(c/F) = c/F \text{ iff } F(f(c), c) \text{ holds iff } \models \delta^2(f(c), c) \\ \text{iff } L(f(c), c) \text{ holds iff } f(c/L) = c/L, \end{aligned}$$

where the second logical equivalence follows since: Otherwise, $\models \delta^2(f(c), c)$ but $F(c_i, f(c))$ and $F(c, c_j)$ hold for some $i \neq j < n$. But then we have $\models \delta^4(c_i, c_j)$, a contradiction. \square

By Claim 2, $H = \pi(\text{Aut}_{cFA}(\mathcal{M})) = \pi(\text{Aut}_{\mathbf{b}A}(\mathcal{M}))$ where $\mathbf{b} \in \text{acl}^{\text{eq}}(A)$. \square

However, contrary to [5, Corollary 5.1.15], in general $\text{acl}(e)$ and $\text{acl}^{\text{eq}}(e)$ need not be interdefinable; the error occurred there due to the incorrect proof of [5, 5.1.14(1) \Rightarrow (2)]. An example presented in [3] for another purpose supplies a counterexample. Consider the following 2-sorted model:

$M = ((M_1, S_1, \{g_{1/n}^1 : n \geq 1\}), (M_2, S_2, \{g_{1/n}^2 : n \geq 1\}), \delta)$ where

- (1) M_1 and M_2 are unit circles centered at origins of two disjoint (real) planes.
- (2) S_i is a ternary relation on M_i , defined by $S_i(b, c, d)$ holds if and only if b, c and d are in clockwise-order.
- (3) $g_{1/n}^i$ is a unary function on M_i such that $g_{1/n}^i(b) = \text{rotation of } b \text{ by } 2\pi/n\text{-radians clockwise.}$
- (4) $\delta : M_1 \rightarrow M_2$ is the double covering, i.e. $\delta(\cos t, \sin t) = (\cos 2t, \sin 2t)$.
- (5) Let \mathcal{M} be a monster model of $\text{Th}(M)$ and $\mathcal{M}_1, \mathcal{M}_2$ be the two sorts of \mathcal{M} .

In [2, Theorems 5.8 and 5.9], it is shown that each $\text{Th}(\mathcal{M}_i)$ has weak elimination of imaginaries (that is, for any imaginary element c , there is a finite real tuple b such that $c \in \text{dcl}(b)$ and $b \in \text{acl}(c)$), using the B. Poizat's notion of weak elimination of imaginaries ([7, Chapter 16.5]). The following fact is a folklore, whose explicit proof was observed in RIMS model theory workshop by I. Yoneda ([8]).

Fact 13. *A (complete) theory T has weak elimination of imaginaries if and only if every definable set has a smallest algebraically closed set over which it is definable.*

Remark & Definition 14.

- (1) For each element b of sort $i = 1, 2$, $g_r^i(b)$ means $(g_{1/n}^i)^m(b)$ where r is a rational number m/n .
- (2) For each element b of sort 2, $\delta^{-1}(b) = \{c_0, c_1\}$, the δ -preimage of b .
- (3) For a set of elements $B = B_1 \cup B_2$ of \mathcal{M} where each element of B_i is of sort i ,

$$\begin{aligned} \text{cl}(B) = \{g_r^1(b) : r \in \mathbb{Q}, b \in B_1\} \cup \{\delta(g_r^1(b)) : r \in \mathbb{Q}, b \in B_1\} \\ \cup \{g_r^2(b) : r \in \mathbb{Q}, b \in B_2\} \cup \bigcup_{r \in \mathbb{Q}, b \in B_2} \delta^{-1}(g_r^2(b)). \end{aligned}$$

- (4) Note that in the above item, the substructure generated by B is formed by omitting the last union: $\bigcup_{r \in \mathbb{Q}, b \in B_2} \delta^{-1}(g_r^2(b))$.

Lemma 15. *Let $B = \{b_0, \dots, b_{n-1}\}$ be a subset of \mathcal{M} . Then*

$$\text{acl}(B) = \text{cl}(B).$$

Proof. Say $B = \{b_0, \dots, b_{m-1}, b_m, \dots, b_{n-1}\}$ where b_0, \dots, b_{m-1} are of sort 1 and the others are of 2. Choose any element b of sort 1. If

$$b \notin \{g_r^1(b_i) : r \in \mathbb{Q}, i < m\} \cup \bigcup_{r \in \mathbb{Q}, m \leq i < n} \delta^{-1}(g_r^2(b_i)),$$

then $b \notin \text{acl}(B)$ since there are infinitely many elements which are infinitesimally close to b and there is an B -automorphism mapping b to each such element.

Likewise, for an element b of sort 2, if

$$b \notin \{g_r^2(\delta(b_i)) : r \in \mathbb{Q}, i < m\} \cup \{g_r^2(b_i) : r \in \mathbb{Q}, m \leq i < n\},$$

then $b \notin \text{acl}(B)$. Thus $\text{acl}(B) \subseteq \text{cl}(B)$.

For the converse, it is easy to observe that

$$\begin{aligned} & \{g_r^1(b_i) : r \in \mathbb{Q}, i < m\} \cup \{g_r^2(\delta(b_i)) : r \in \mathbb{Q}, i < m\} \\ & \cup \{g_r^2(\delta(b_i)) : r \in \mathbb{Q}, m \leq i < n\} \subseteq \text{dcl}(B) \text{ and} \\ & \bigcup_{r \in \mathbb{Q}, m \leq i < n} \delta^{-1}(g_r^2(b_i)) \subseteq \text{acl}(B) \end{aligned}$$

since each $b \in \bigcup_{r \in \mathbb{Q}, m \leq i < n} \delta^{-1}(g_r^2(b_i))$ has at most two B -automorphic images (has only one B -automorphic image if $m \neq 0$). \square

Proposition 16. *Th(\mathcal{M}) has weak elimination of imaginaries.*

Proof. Let $\varphi(x, y_0, \dots, y_{n-1}) \in \mathcal{L}$ and $B = \{b_0, \dots, b_{n-1}\} = \{b_0, \dots, b_{m-1}\} \cup \{b_m, \dots, b_{n-1}\}$ where b_0, \dots, b_{m-1} are of sort 1 and the others are of 2. According to Fact 13, it suffices to show that there is a smallest algebraically closed set over which $\varphi(\mathcal{M}, B) \equiv \varphi(\mathcal{M}, b_0, \dots, b_{n-1})$ is definable.

Since there is some c_i such that $\delta(c_i) = b_i$ for each $i \in \{m, \dots, n-1\}$, we may assume that every element of B is of sort 1. Choose $D = \{d_0, \dots, d_{k-1}\} \subseteq B$ such that $\{g_r^1(d_i) : r \in \mathbb{Q}, i < k\} = \{g_r^1(b_i) : r \in \mathbb{Q}, i < n\}$ and $d_i \notin \text{cl}(D) \setminus \{d_i\}$ for each $i < k$. Then $\varphi(\mathcal{M}, B)$ is definable over D and there is some minimal subset D' of D such that $\varphi(\mathcal{M}, B)$ is definable over $\text{acl}(D')$ by Lemma 15. \square

Now for $i = 1, 2$, we let $E_i(x, y)$ if and only if x and y in \mathcal{M}_i are infinitesimally close, i.e.

$$E_i(x, y) := \bigwedge_{1 < n} (S_i(x, y, g_{1/n}^i(x)) \vee S_i(y, x, g_{1/n}^i(y))),$$

which is an \emptyset -type-definable equivalence relation. Let $b \in \mathcal{M}_2$, $c, c' \in \mathcal{M}_1$ where $\delta(c) = \delta(c') = b$. Note that c, c' are antipodal to each other and $c/E_1, c'/E_1$ are conjugates over b/E_2 , hence $c/E_1, c'/E_1 \in \text{acl}(b/E_2)$.

Theorem 17. *$\text{acl}(b/E_2)$ and $\text{acl}^{\text{eq}}(b/E_2)$ are not interdefinable.*

Proof. We prove following Claim and then conclude.

Claim. $\text{acl}^{\text{eq}}(b/E_2)$ is interdefinable with b/E_2 .

Proof. To lead a contradiction, suppose that there are distinct imaginaries $d_1, d_2 \in \text{acl}^{\text{eq}}(b/E_2)$ such that $d_1 \equiv_{b/E_2} d_2$. Weak elimination of imaginaries of $\text{Th}(\mathcal{M})$ (Proposition 16) implies that $\text{acl}^{\text{eq}}(d_1, d_2)$ and $D := \{d \in \mathcal{M} : d \in \text{acl}^{\text{eq}}(d_1, d_2)\}$ are interdefinable (*). In particular, $D \subseteq \text{acl}^{\text{eq}}(b/E_2) \cap \mathcal{M}$. However, for any infinitesimally close $d, d' \in \mathcal{M}_i$ ($i = 1, 2$), there is $f \in \text{Aut}_{b/E_2}(\mathcal{M})$ sending d to d' . Hence indeed $D = \emptyset$, which contradicts (*) (because $d_1 \equiv_{b/E_2} d_2$ and $d_1 \neq d_2 \in \text{acl}^{\text{eq}}(d_1, d_2)$). \square

Now $c/E_1, c'/E_1 \in \text{acl}(b/E_2) \setminus \text{dcl}(b/E_2) = \text{acl}(b/E_2) \setminus \text{dcl}(\text{acl}^{\text{eq}}(b/E_2))$.

□

REFERENCES

- [1] Enrique Casanovas, Simple theories and hyperimaginaries (Lecture Notes in Logic), Cambridge University Press, 2011.
- [2] Jan Dobrowolski, Byunghan Kim and Junguk Lee, The Lascar groups and the first homology groups in model theory, *Annals of Pure and Applied Logic* **168**, (2017), 2129–2151.
- [3] Jan Dobrowolski, Byunghan Kim, Alexei Kolesnikov and Junguk Lee, The relativized Lascar groups, type-amalgamation, and algebraicity, *Journal of Symbolic Logic*, **86**, (2021), 531–557.
- [4] Karl H. Hofmann and Sidney A. Morris, The structure of compact groups, De Gruyter, 2013.
- [5] Byunghan Kim, Simplicity theory, Oxford University Press, 2014.
- [6] Hyoyoon Lee, Quotient groups of the Lascar group and strong types in the context of hyperimaginaries, PhD thesis, Yonsei University, in preparation.
- [7] Bruno Poizat, A Course in Model Theory: An Introduction to Contemporary Mathematical Logic, Springer, 2000.
- [8] Ikuo Yoneda, On two definitions of weak elimination of imaginaries, RIMS Model Theory Workshop 2021, December 2021.
- [9] Martin Ziegler, Introduction to the Lascar group, Tits buildings and the model theory of groups, London Math. Lecture Note Series, **291**, Cambridge University Press, (2002), 279–298.

DEPARTMENT OF MATHEMATICS
 YONSEI UNIVERSITY
 50 YONSEI-RO SEODAEMUN-GU
 SEOUL 03722
 SOUTH KOREA

Email address: `alternative@yonsei.ac.kr`