

Review

# Halophyte Plants as Potential Sources of Anticancer Agents: A Comprehensive Review

Luísa Custodio <sup>1,\*</sup>, Pedro Garcia-Caparros <sup>2</sup>, Catarina Guerreiro Pereira <sup>1</sup> and Pedro Castelo-Branco <sup>3,4,5</sup>

<sup>1</sup> Centre of Marine Sciences, Faculty of Sciences and Technology, Campus of Gambelas, University of Algarve, 8005-139 Faro, Portugal

<sup>2</sup> Agronomy Department of Superior School Engineering, University of Almeria, 04120 Almería, Spain

<sup>3</sup> Faculty of Medicine and Biomedical Sciences (FMCB), Campus de Gambelas, University of Algarve, 8005-139 Faro, Portugal

<sup>4</sup> Algarve Biomedical Center Research Institute (ABC-RI), 8005-139 Faro, Portugal

<sup>5</sup> Champalimaud Research Program, Champalimaud Center for the Unknown, 1400-038 Lisbon, Portugal

\* Correspondence: [lcustodio@ualg.pt](mailto:lcustodio@ualg.pt)

**Abstract:** Salt-tolerant plants (halophytes) are widely distributed worldwide in several environments such as coastal salt marshes, sand dunes, and inland deserts. To cope with the harsh conditions that characterize those habitats, which include high salinity and radiation levels, such plants have developed morphological and physiological traits, the latter including the synthesis and accumulation of important secondary metabolites such as alkaloids and polyphenols. While essential in maintaining plant homeostasis, these compounds are highly valued in the medical field for the treatment of several human diseases, including cancer. Cancer is one of the most life-threatening disorders worldwide, which accentuates the need to improve current cancer therapies and minimize potential adverse secondary side-effects. In this context, the pharmacological evaluation of natural compounds has attracted growing interest since nature has already provided some important anti-cancer drugs. This review compiles, for the first time, research regarding the anticancer activity of halophytes from different families, including, whenever possible, the bioactive molecules involved in such therapeutical properties along with possible mechanisms of action. The introduction section provides some pertinent information regarding cancer and a summary of the most important characteristics of halophytes. The next section gives information regarding the in vitro and in vivo cytotoxic properties of several halophyte species, grouped by families, including contents in bioactive metabolites and proposed modes of action, if possible. Lastly, the conclusion presents the most relevant metabolites and/or promising species and extracts that could be further explored in anticancer drug research.

**Keywords:** cancer; natural products; cytotoxic; salt-tolerant plants; antitumoral drugs

**Citation:** Custodio, L.; Garcia-Caparros, P.; Pereira, C.G.; Castelo-Branco, P. Halophyte Plants as Potential Sources of Anticancer Agents: A Comprehensive Review. *Pharmaceutics* **2022**, *14*, 2406. <https://doi.org/10.3390/pharmaceutics14112406>

Academic Editors: Noelia Duarte and Ana Paula Francisco

Received: 13 July 2022

Accepted: 4 November 2022

Published: 8 November 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Cancer is a group of diseases characterized by malignant neoplasms arising from the abnormal and uncontrolled cell proliferation that invades and destroys the surrounding tissue and, if not controlled, can result in death [1,2]. With an established complex link between cancer and aging and the increased risk factors of an unhealthy lifestyle, cancer is undoubtedly a major health problem in developed and developing countries [3–7]. It is estimated that, by 2040, the number of new cancer cases diagnosed per year will be higher than 27.5 million worldwide, with 16.3 million cancer deaths due to population growth and aging [8]. Globally, lung and breast cancers are the most frequently diagnosed and are the leading causes of cancer-related death in men and women, respectively [9]. Significant advances have been made in cancer research in the last 20 years concerning its general biology, prevention, and treatment [10]. Currently, chemotherapy is one of the

most common cancer therapeutic approaches, although secondary effects are known and include a negative impact on the immune system, and the effectiveness of this therapeutic approach can be limited by drug resistance [11,12]. Therefore, new anticancer drugs and therapies that concomitantly minimize its harmful side-effects are still in urgent demand.

In the war against cancer, the role of natural compounds has become of crucial interest, and the scientific community has placed great effort in identifying novel sources of antitumoral molecules among medicinal plants and, more recently, marine organisms [10,13]. The pharmacological evaluation of natural compounds has rendered some important anticancer drugs such as paclitaxel and its derivatives from the bark of the Pacific yew tree (*Taxus brevifolia* Nutt, family: Taxaceae), or vincristine and vinblastine from Madagascar periwinkle (*Catharanthus roseus* (L.) G. Don, family: Apocynaceae) [14]. However, in addition to glycophyte plants and some marine species, other organisms including halophytes contain metabolites with potential use as anticancer drugs.

Although the definition of halophytes has yet to reach consensus, some authors define them as plants completing their life cycle in environments with a salt concentration of around 200 mM of sodium chloride (NaCl) (20 dS m<sup>-1</sup>) or more [15–17]. They represent nearly 1% of the global flora diversity and inhabit different ecosystems such as beaches, rocky shores, saltmarshes, estuaries, and inland deserts [17–19]. Such areas are characterized by stressful abiotic conditions, (e.g., high salinity, radiation levels, and drought), which trigger the production and accumulation of reactive oxygen species (ROS) in the plants, leading to cellular and tissue damage, metabolic disorders, and senescence. Halophytes are equipped with strong antioxidant defense systems to counteract the negative effects of ROS, including antioxidant enzymatic mechanisms and the synthesis of secondary metabolites, such as phenolic compounds, saponins, and alkaloids [18,20]. These molecules have a crucial protective role in the plant and exhibit relevant bioactivities, including antioxidant, anti-inflammatory, and antitumoral, which are linked to beneficial therapeutic properties and could help explain the use of some halophytes in traditional medicine and as food [18,21,22].

There are several studies focused on the anticancer properties of a high number of halophytic species, but this information is scattered in the literature. Therefore, our aim was to compile and summarize existing information regarding the anticancer activity of halophytes, the molecules involved in such properties, and the possible mechanisms of action (Tables 1–4). The Web of Science database and Google Scholar (as a search engine) were consulted to retrieve the most updated articles on the topic under investigation. The keywords “salt-tolerant plants” and “halophytes” were used alone or in combination with, for example, “anticarcinogenic”, “antitumoral”, or “anticancer”. Only English articles having full text were considered. The contribution was distributed according to the different families and species, which were previously checked in the eHALOPH database [23]. The correspondence between the name of the cell lines referred in the text and the type of cancer can be found in Table 1, together with a list of halophyte species, organs and extracts tested, cell lines/types of cancer assessed, and obtained half-maximal inhibitory concentrations (IC<sub>50</sub>) values and proposed mechanisms of action, when provided by the authors. However, at the end of the study, a summary is presented referring only to the results obeying the criteria of cytotoxic activity for crude extracts (IC<sub>50</sub> < 30 µg/mL; Table 3) and isolated compounds (IC<sub>50</sub> ≤ 10 µM; Table 4) considered relevant to proceed for antitumoral applications, as established by the American National Cancer Institute (NCI) [24] and the National Institutes of Health (NIH) [25], respectively.

## 2. Anticancer Activity of Halophytes

Halophytes are a valuable source of bioactive molecules for the prevention and treatment of human chronic diseases, including cancer [26]. In this review, information is presented by family, namely, *Acanthaceae*, *Aizoaceae*, *Amaranthaceae*, *Apiaceae*, *Asteraceae*, *Brassicaceae*, *Convolvulaceae*, *Cymodoceaceae*, *Cyperaceae*, *Fabaceae*, *Juncaceae*, *Malvaceae*, *Myrtaceae*, *Plantaginaceae*, *Plumbaginaceae*, *Poaceae*, *Portulacaceae*, *Rhizophoraceae*, *Solanaceae*,

and *Tamaricaceae*. Data regarding the anticancer activity of different halophyte species, including organs and extracts tested, cell lines/types of cancer assessed, and obtained IC<sub>50</sub> values, are summarized in Table 1. Table 2 contains detected molecules in the active extracts from halophyte species with anticancer properties, while Table 3 summarizes information on extracts/fractions of selected species that could be further explored in the search for new drug-leads, as established by the American National Cancer Institute (NCI) [24], including cytotoxic activity and chemical composition of the extracts (when possible), cell lines tested, obtained IC<sub>50</sub> values, and proposed mechanisms of action. Lastly, data related to the cytotoxic activity of selected isolated compounds, according to the criteria of the National Institutes of Health (NIH) [25], are presented on Table 4.

### 2.1. *Acanthaceae* Family

*Acanthaceae* currently comprises 192 genera and 5504 species growing in Indonesia, Malaysia, Africa, Brazil, and Central America [27,28]. The presence of bioactive metabolites such as alkaloids, phenols, terpenoids, tannins, quinones, cardiac glycosides, saponins, carbohydrates, and flavonoids, highlights the medicinal relevance of this family in the treatment of several diseases, including cancer [29]. This review focuses on the anticancer activity of the halophytic genera *Acanthus* and *Avicennia*.

*Acanthus ilicifolius* L. (holly leaved acanthus) is a mangrove shrub rich in triterpenes, alkaloids, and flavonoids, traditionally used in Chinese and Indian medicine against asthma, headache, and skin diseases [30]. An ethanolic leaf extract from *A. ilicifolius* significantly reduced the viability of HepG2 cells in a dose-dependent manner (92% of cytotoxicity at 100 µg/mL) via apoptosis induction potentially associated with DNA damage [31] (Table 1). A water root extract from *A. ilicifolius* reduced the viability of HepG2 cells, as observed in the 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide (MTT) assay (IC<sub>50</sub> value of 39.76 µg/mL after 48 h of incubation) [32], via apoptosis induction related to DNA damage (Table 1). Ethanol extracts from leaves and roots of holly leaved acanthus were also able to reduce the viability of MCF7 cells (breast carcinoma), with IC<sub>50</sub> values of 24.22 and 29.20 µg/mL for leaf and root extracts, and of PA1 cells (ovarian carcinoma), with IC<sub>50</sub> values of 15.74 µg/mL and 20.00 µg/mL for leaf and root, respectively (Table 1) [33]. The chemical profile of the active extracts was not established. *Acanthus ebracteatus* Vahl. (holly mangrove), is traditionally used in Thai medicine as an anticancer agent [34]. A protein hydrolysate (<3 kDa) from the aerial parts of *A. ebracteatus*, reduced the viability of A431 cells, with an IC<sub>50</sub> value of 425.9 ng protein/mL after 24 h of incubation (Table 1). Partially purified peptides were obtained from this hydrolysate and resulted in the synergistic effects against cell viability of A431 cells, via apoptosis induction [34].

*Avicennia* comprises mangroves distributed in the Indo-Western Pacific and Atlantic eastern Pacific areas. Some *Avicennia* species are effective against cancer due to the presence of different classes of molecules, including alkaloids, phenols, flavonoids, tannins, iridoid glucosides, and terpenoids [35]. A methanol leaf extract from *Avicennia alba* Blume significantly reduced the viability of MCF7 and HeLa cells, evaluated by the MTT assay, with IC<sub>50</sub> values of 57.02 and 44.30 µg/mL, respectively, after 48 h of incubation [36] (Table 1). After treatment with the leaf extract of *A. alba* both cell lines exhibited morphological modifications, i.e., a reduction in cell size and cell detachment. The extract was analyzed using gas chromatography–mass spectrometry (GC–MS) allowing for the identification of 12 compounds, the most abundant ones being catechol borane (11%), neophytadiene (17%), and hexadecanoic acid (29%) (Table 2), but their cytotoxicity toward the tested cell lines was not established [36]. Moreover, since GC–MS mainly allows for the detection of lipophilic molecules, one cannot exclude the possibility that hydrophilic compounds with cytotoxic properties may be present in the methanol leaf extract but were not detected with the chromatographic system used. A chloroform/methanol extract from leaves of *A. alba* reduced the viability of WiDr cells (IC<sub>50</sub>

= 173.78  $\mu\text{g/mL}$  after 24 h of incubation) using doxorubicin as the positive control and induced cell arrest in the G0–G1 phase, resulting in apoptosis induction [37] (Table 1).

*Avicennia marina* (Forssk.) Vierh, one of the most abundant and common mangrove species, is used in traditional medicine for the treatment of different diseases, including rheumatism and smallpox, and it contains different molecules with therapeutic properties, such as luteolin 7-O-methylether and chrysoeriol 7-O-glucoside [38]. An ethyl acetate extract of leaves and stems of *A. marina* reduced the viability of MCF7 cells (MTT assay), with values of cell growth inhibition of 65% and 75% after treatment for 48 h with the extract at 100 and 200  $\mu\text{g/mL}$ , respectively [39] (Table 1). When applied at the concentration of 200  $\mu\text{g/mL}$  the extract induced apoptosis in a concentration-dependent manner through ROS production and disruption of the mitochondrial membrane potential ( $\Delta\psi\text{m}$ ); however, no PARP-1 cleavage was detected, and a significant decrease in caspase-7 protein levels was observed. A significant level of autophagy was also detected at the same concentration [39]. A methanol extract of the same species reduced the viability of HeLa cells in a concentration-dependent manner (tested concentrations: 50, 100, 250, 500, and 1000  $\mu\text{g/mL}$ ) with an  $\text{IC}_{50}$  value of 107  $\mu\text{g/mL}$  [40] (Table 1). The chemical profile of the extracts was not established by the authors and, therefore, the bioactive compounds were not identified. In another study targeting the same species [38], water, ethanol, methanol, and ethyl acetate extracts from leaves and seeds were tested toward different human cancer cell lines, namely, breast (AU565, MDA-MB-231, and BT483), liver (HepG2 and Huh7), and NIH3T3 cells. The ethyl acetate extract from leaves had the highest cytotoxicity, especially toward cancer cells, which was attributed by the authors to its highest phenol content. The ethyl acetate extract suppressed xenograft MDA-MB-231 tumor growth in nude mice, and induced apoptosis in cancer cells through apoptosis inhibition and inhibition of cell migration. This extract was further fractionated by column chromatography, and resulting fractions were retested for anticancer properties. The most active fractions, F2-5, F3-2-9, and F3-2-10, had  $\text{IC}_{50}$  values lower than 20  $\mu\text{g/mL}$ , and they were profiled by  $^1\text{H-NMR}$  and  $^{13}\text{C-NMR}$ , allowing for the identification of the flavonoids luteolin and quercetin standards, and avicennones D and E. The antiproliferative activity of the flavonoids was confirmed [38]. Additional studies about the anticarcinogenic activity of other halophytic species of the *Acanthaceae* family can be found in Table 1.

## 2.2. Aizoaceae Family

*Aizoaceae* contains mostly succulent plants with 1910 species currently recognized, distributed across 125 genera [41,42]. The *Mesembryanthemum* genus, for example, is used in traditional medicine to treat several diseases such as hepatic conditions and diabetes, and it contains several bioactive compounds, including flavonoids and catechins [43]. An ethanol crude extract from leaves of *Mesembryanthemum crystallinum* L. (ice plant), and obtained hexane, ethyl acetate, butanol, and water fractions were tested for their effect on the growth of HCT116 cells [44] (Table 1). The application of the ethyl acetate and butanol fractions resulted in a dose-dependent inhibition of cell growth (at 125, 250, and 500  $\mu\text{g/mL}$ , applied for 24 and 48 h), as observed in the MTT assay, a reduction in the levels of intracellular ROS (at 250 and 500  $\mu\text{g/mL}$ , using 2',7'-dichlorofluorescein-diacetate), and apoptosis induction (at 250 and 500  $\mu\text{g/mL}$ , during 48 h). Treatment with the butanol fraction also resulted in cell-cycle arrest in the G2/M phase [44]. Samples were evaluated for total phenolic content (TPC), using the Folin–Ciocalteu (F–C) assay, and it was observed that the butanol fraction had the highest TPC (5.4 mg gallic acid equivalent/g) [44], suggesting that phenolics may be related to the detected cytotoxicity. However, no attempt was made by the authors to identify the active molecules present in this sample.

The application of ethanol, methanol, acetone, hexane, and diethyl-ether extracts from whole plants of *Sesuvium portulacastrum* L. (sea purslane) decreased the viability of MDA-MB-231, IMR32, and HCT116 cell lines, as assessed by MTT after 24 h of incubation, using doxorubicin as the standard [45]. The hexane extract had the highest cytotoxic

activity with IC<sub>50</sub> values of 942.07, 703.40, and 407.87 µg/mL for MDA-MDB-231, IMR32, and HCT116 cell lines (Table 1). Despite the considerably high IC<sub>50</sub> values obtained, the possible mechanisms of action of the extracts were evaluated using phase contrast-inverted microscope observation and propidium iodide staining, which revealed the presence of typical features of apoptotic cells, including nuclear condensation, cell shrinkage, and the presence of apoptotic bodies [45] (Table 1). No attempt was made by the authors to unravel the chemical composition of the extracts.

### 2.3. Amaranthaceae Family

*Amaranthaceae* currently comprises 2398 species grouped in 182 genera [46], which can be found in a wide range of habitats from arid and semiarid regions to saline environments and subtropical areas; it contains flavonoids, phenolic acids, terpenes, and triterpene saponins, and it displays important pharmacological properties such as anticarcinogenic [47,48].

*Arthrocnemum* sp. is a small genus of succulent halophytes present in coastal marshes of South Africa, Asia, Mediterranean, Europe, and Northern America, and it contains several compounds such as phenols, flavonoids, and tannins with described therapeutic properties, including antibiotic, hypoglycemic, and antitumoral [49,50]. The application of an 80% methanol extract from shoots of *Arthrocnemum indicum* (Willd.) Moq. significantly reduced the viability of Caco-2 cells, as observed by the MTT assay [50] (Table 1). Fluorescence microscopy observation through DAPI (4,6-diamidino-2-phenylindole) staining showed that cells exhibited a decline of DNA synthesis, while flow cytometry allowed for the observation of cell-cycle arrest at the G2/M phase after 72 h of exposure to the extract at the concentration of 100 µg/mL [51] (Table 1). The extract was profiled by liquid chromatography/electrospray ionization time-of-flight mass spectrometry (LC/ESI-TOF-MS), and high levels of phenolic compounds were detected such as gallic acid, cyanidin, chrysoeriol, quercetin, catechol, syringic acid, and luteolin (Table 2), although their cytotoxicity toward Caco-2 cells was not established [51]. However, it is known that some of the detected compounds exhibit antitumoral properties. For example, chrysoeriol was identified as a selective inhibitor of the PI3K–AKT–mTOR pathway and, therefore, linked with cell cycle regulatory effects; it also reduced the proliferation of human multiple myeloma cells (RPMI 8226 and KM3, with IC<sub>50</sub> values of 26 and 35 µmol/L at 48 h) but not of peripheral blood mononuclear cells (PBMNCs) [52].

Several *Atriplex* species are halophytic [53]. An ethanol extract from leaves of *Atriplex halimus* L. (sea orache) significantly reduced the viability of HepG2 cells (IC<sub>50</sub> = 54.86 µg/mL after 24 h of incubation) (Table 1) via apoptosis induction linked to the expression of TP53, BCL2, and BAX genes [54] (Table 1). No information was provided by the authors regarding the chemical components of such extract, but it is known that *A. halimus* is characterized by the presence of different bioactive compounds, including syringetin derivatives and the flavonol glycosides atriplexoside A [3'-O-methylquercetin-4'-O-β-apiofuranoside-3-O-(6''-O-α-rhamnopyranosyl-β-glucopyranoside)] and atriplexoside B [3'-O-methylquercetin-4'-O-(5''-O-β-xylopyranosyl-β-apiofuranoside)-3-O-(6''-O-α-rhamnopyranosyl-β-glucopyranoside)], which may be related to the detected cytotoxic activity [55–60].

*Chenopodium* contains weedy herbs native to Asia, Europe, and America, rich in phenolics, saponins, and triterpenoids, conferring them therapeutic properties such as laxative, analgesic, and anticarcinogenic [61]. The edible species *Chenopodium album* L. (lamb's quarters) is valued in traditional medicine for its anticancer properties [62]. A petroleum ether extract of branches and leaves from *C. album* reduced the viability of A549 cells in a dose-dependent manner, showing an IC<sub>50</sub> of 33.31 µg/mL using gemcitabine as a positive control, which was linked to cell-cycle arrest at the G1 phase [63] (Table 1). Essential oil from whole plants of *Chenopodium ambrosioides* L. (Mexican tea) decreased the viability of MCF7 cells by inducing DNA fragmentation (IC<sub>50</sub> values of 18.75, 9.45, and

10.50 µg/mL at 6, 24, and 48 h, respectively) when compared to the control treatment [64] (Table 1). In another study, an essential oil of the same species also displayed a significant cytotoxic activity toward RAJI cells, with an IC<sub>50</sub> value of 1.0 µg/mL, which was probably related to the high levels of ascaridole, detected using GC with flame ionization detection (GC-FID), GC-MS, and proton nuclear magnetic resonance (1H-NMR) [65] (Tables 1 and 2). Ascaridol is a monoterpene with a strong in vitro capacity to decrease the growth of different tumor cell lines and is, therefore, considered as a strong candidate for the treatment of cancer [65]. *Chenopodium quinoa* Willd is a pseudocereal originated from the Andes of South America with recognized nutritional and functional properties that rendered the species the status of “functional food” [66]. Polysaccharides were extracted from quinoa seeds with petroleum ether and purified by column chromatography, revealing mainly galacturonic acid and glucose monosaccharides. This polysaccharide fraction was applied toward cancer (SMMC 7721 and MCF7) and “normal” (L02 and MCF 10A) cell lines, for 24 and 48 h, allowing for IC<sub>50</sub> values ranging from 53.4 to 121.4 µg/mL, without affecting “normal” cell viability [66] (Table 1).

*Salicornia* species (glasswort, sea asparagus, or samphire) are edible succulent halophytes highly appreciated in gourmet cuisine due to their organoleptic properties [67]. The ethyl acetate and methanol leaf extracts from *Salicornia europaea* L. reduced the viability of MCF7 cells with IC<sub>50</sub> values of 97.9 and 117.1 µg/mL for the ethyl acetate and methanol extracts, respectively, after 24 h of incubation [68]. A qualitative analysis of the extracts identified tannins and saponins, which may be associated with the in vitro antitumor capacity. Moreover, a GC-MS analysis showed the presence of 32 and 29 compounds in corresponding ethyl acetate and methanol extracts, respectively, which were not assessed individually against the cancer cells [68]. Tannins are phenolic compounds with a high molecular weight, while saponins contain a steroidal or triterpenoid aglycone and one or more sugar chains [69]. Both groups of compounds exhibit antitumoral properties; for example, saponins generally decrease cellular proliferation by binding to cholesterol structures on cell membranes, forming pores and holes by binding to specific receptors followed by the induction of apoptosis, while tannins can hamper cancer cell proliferation through antioxidant processes and apoptosis induction [70,71]. However, the presented results must be analyzed with caution, since the phytochemical analysis reported by the authors only targets lipophilic compounds, without unraveling the possible cytotoxic molecules present in the methanol (polar) extract.

*Suaeda* comprises around 80 to 100 succulent species distributed in semideserts, deserts, and seashores. These species are usually edible and contain several bioactive components, such as phenolics, flavonoids, and terpenoids, conferring them with bioactive properties, including anticarcinogenic [72,73]. A hexane extract from leaves of *Suaeda ruticose* (L.) Forssk. (shrubby seablight) reduced the viability of HCT116, HepG2, and MCF7 cell lines, being more effective toward HCT116 cells (IC<sub>50</sub> = 17.2 µg/mL) [74] (Tables 1 and 4). The cytotoxicity of the extract was tested using the sulforodamine B (SRB) assays for 72 h using doxorubicin as a positive control with IC<sub>50</sub> values of approximately 0.5 µg/mL. This extract caused cell-cycle arrest at the G<sub>0</sub>-G<sub>1</sub> phase and induced apoptosis, especially in HCT116 cells that exhibited chromatin condensation and membrane blebbing (Table 4). The extract was profiled using liquid chromatography coupled to electrospray ionization quadrupole time-of-flight mass spectrometry (LC-ESI-QTOF-MS/MS) and several molecules were identified, including monoterpenes (dihydrojasmane, jasmolone, terpinene-4-ol), diterpenes (pimaric acid, steviol, and momilactone B), and phenolics (quercinol, zingerone, zingerol, and neovaflan) (Table 4), but its contribution to the detected cytotoxicity was not directly established [74]. However, there are several reports on the antitumoral properties of phenolic compounds [56–59]. Specifically, zingerone reduced the viability of HCT116 cells through ROS-mediated apoptosis in colon cancer cells [75]. Monoterpenes and diterpenes have reported cytotoxic properties [76,77]. For instance, dihydrojasmane was identified as one of the

active compounds toward HeLa and NIH/3T3 cell lines, in a methanol extract from *Rumex hastatus* D. Don [78], while steviol significantly reduced the viability of MCF7 cells via apoptosis induction [79]. A dichloromethane extract from the shoots of *S. fruticosus* significantly reduced the viability of A549 (IC<sub>50</sub> = 49 µg/mL), DLD1 (IC<sub>50</sub> = 10 µg/mL), Caco-2 (IC<sub>50</sub> = 140 µg/mL), and HT-29 cells (IC<sub>50</sub> = 12 µg/mL), which was assessed using the resazurin reduction test after 48 h of incubation (Tables 1 and 4). Nevertheless, the bioactive compounds present in the chemical profile of the extracts were not identified [80]. Additional studies on the anticarcinogenic activity of other halophytic species of this family are summarized in Table 1.

#### 2.4. Apiaceae Family

*Apiaceae* currently comprises 3916 species distributed across 457 genera dispersed worldwide [81], and it is characterized by the presence of polyacetylenes displaying important bioactivities, including anticarcinogenic [82]. In this family, there are two genera of halophytic species, *Eryngium* sp. and *Crithmum* sp., with described cytotoxic properties against human cancer cells.

*Eryngium* includes approximately 250 species present in Eurasia, North Africa, North and South America, and Australia. *Eryngium* species are frequently used as ornamentals or as a food source and are valued in traditional medicine as a diuretic, to treat diarrhea, headaches, and digestive problems. These properties are potentially associated with the presence of terpenoids, triterpenoids, polyacetylenes, flavonoids, and coumarins [83]. The cytotoxic effects of aqueous extracts from shoots and roots of *Eryngium maritimum* L. were assessed against human cancer lines (HepG2, HEP-2, and U138MG) and on a “normal” cell line (Vero), using the MTT assay after 24 h of incubation [70] (Table 1). The root extracts were overall more cytotoxic, and the IC<sub>50</sub> values ranged from 30.3 to 50 µg/mL. However, a strong reduction in cellular viability was also observed in the “normal” cell line, which suggest no selectivity toward cancer cells [70] (Table 1). Although the phytochemical profile of the extracts was not established by the authors, other reports identified several cytotoxic molecules in *E. maritimum*, including saponins, which create pores and perforations on the cellular membranes by binding to specific receptors, followed by apoptosis induction by, for example, stimulation of the cytochrome c-caspase 9-caspase 3-pathways [70,71].

*Crithmum maritimum* L. (sea fennel) is traditionally used as a cooking ingredient and in folk medicine for its stimulating and diuretic effects [84]. An ethyl acetate extract from the whole sea fennel plant reduced the proliferation of Huh7 and HepG2 cell lines, by interfering with the cell cycle, specifically on the shift of phases with increasing number of cells in the G<sub>0</sub>/G<sub>1</sub> phase after 24 h of incubation and in the G<sub>2</sub>/M phase after 48 h [85]. The extract induced an increase in necrotic and apoptotic cancer cells, assessed by cytofluorimetric analysis although the values of IC<sub>50</sub>, but the chemical profile of the extract was not assessed [85] (Table 1).

#### 2.5. Asteraceae Family

*Asteraceae* currently contains 1733 genera and 35,988 species [86], including several used as ornamentals (e.g., *Calendula* and *Chrysanthemum*) or for food (e.g., *Helianthus annuus*) and medicinal (e.g., *Artemisia* sp. and *Echinacea* sp.) purposes [87]. *Achillea millefolium* L. (yarrow) is an aromatic perennial herb with traditional medicinal uses such as wound healing and anti-inflammatory activity, and it contains different classes of bioactive compounds, including flavonoids and terpenoids [88]. A methanol extract from yarrow shoots was applied toward prostate cancer (DU145) and “normal” skin (HFFF2) cells, at different concentrations (20, 100, 500, 1000, and 2000 µg/mL), alone or in combination with bleomycin, an anticancer agent. The yarrow extract was not toxic toward “normal” cells, but had significantly enhanced cytotoxicity induced by bleomycin showing 60% and 49% survival rate at doses of 1000 and 2000 µg/mL, respectively, which may indicate that this extract contains molecules able to improve the effectiveness of

bleomycin, while minimizing negative side-effects caused by toxicity toward “normal” cells. [88] (Table 1). The extract was not profiled for chemical components, but there are reports of the presence of several cytotoxic compounds in yarrow, such as achillinin A (guaianolide) and casticin (flavonoid) [88,89]. In another work, petroleum ether, ethyl acetate, methanol, and water extracts were prepared from aerial organs of yarrow and tested for toxicity, using the MTT assay, on human cancer cell lines [90] (Table 1). The strongest cytotoxic effect was observed after application of the ethyl acetate extract on HeLa ( $IC_{50} = 0.58 \mu\text{g/mL}$ ) and K562 cells ( $IC_{50} = 0.73 \mu\text{g/mL}$ ), followed by the water (MCF-7,  $IC_{50} = 0.87 \mu\text{g/mL}$ ) and the petroleum ether extract (K562,  $IC_{50} = 0.87 \mu\text{g/mL}$ ) (Tables 1 and 4) [90]. The cytotoxic activity of these extracts may be associated with its contents in phenolic acids, such as apigenin and chlorogenic, *p*-coumaric, and rosmarinic acids (Table 4), which showed the capacity to block oncogenic pathways due to the activation of caspases [91]. A methanol extract from aerial parts of *Limbarda crithmoides* (L.) Dumort (commonly known as *Inula crithmoides* L.) was able to decrease the viability of acute myeloid leukemia cells (OCI-AML3) when applied at 100 and 200  $\mu\text{g/mL}$ , for 24 h [92] (Table 1). This extract was submitted to a solvent–solvent partitioning, affording *n*-hexane, dichloromethane, and aqueous methanol-soluble fractions. The hexane and dichloromethane fractions exhibited a strong cytotoxicity toward OCI-AML3 cells at concentrations of 15 or 10  $\mu\text{g/mL}$ , which was ascribed to an increase in apoptotic cells, especially in the G0/G1 phase by the mitochondria-dependent pathway. The hexane extract was then further fractionated, leading to the isolation of two molecules, which were identified by NMR as the thymol derivatives 10-acetoxy-8,9-epoxythymol tiglate and 10-acetoxy-9Z-chloro-8,9-dehydrothymol, with the latter being the most active, causing a decrease in cell viability at 1.25  $\mu\text{g/mL}$  associated with apoptosis induction [92].

#### 2.6. Brassicaceae Family

*Brassicaceae* is one of the largest dicotyledon family of flowering plants, including model species and commercial crops, with 341 genera and 3921 species recognized at the moment [93,94], but few studies have described the potential anticarcinogenic properties of halophytes belonging to this family. One example is the annual halophyte *Cakile maritima* Scop. (sea rocket), which is confined to maritime strandlines of sand and has agronomic (oilseed and phytoremediation) and medicinal (diuretic, antiscorbutic, and purgative) properties [95]. Hexane, ethyl acetate, and methanol fractions were obtained from a methanol extract prepared from aerial organs of sea rocket and tested for antiproliferative properties on Caco-2 and HeLa carcinoma cells [96] (Table 1). The hexane fraction significantly reduced the viability of Caco-2 and HeLa cells, with  $IC_{50}$  values of 12 and 126  $\mu\text{g/mL}$  after 24 h of incubation; respectively. Cisplatin was used as the positive control with values of 69  $\mu\text{g/mL}$  (Caco-2) and 85  $\mu\text{g/mL}$  (HeLa). The extract was profiled using GC–MS, and the major molecules identified were, by area, 2-hydroxy-1,8-cineole, decane, and limonene, which may contribute to the detected antiproliferative activity [96] (Table 4). For instance, limonene has been reported with anticarcinogenic activity in HepG2 cells due to apoptosis induction [97].

#### 2.7. Convolvulaceae Family

*Convolvulaceae* currently contains 50 genera and 1952 species widely distributed in tropical and temperate regions of the world [98,99]. *Calystegia soldanella* (L.) R. Br. ex Roem. & Schult. (shore bindweed) is a perennial edible herb commonly found in coastal sand dunes and foredunes of South Korea, East Asia, Europe, and the Pacific. Shore bindweed is traditionally used for the treatment of, for example, rheumatic arthritis and scurvy, and it displays relevant biological properties, including anticancer [100]. An 85% aqueous fraction from combined methylene chloride and methanol crude extracts obtained from whole plants of shore bindweed significantly decreased HepG2 cellular viability after 24 h of incubation, in a concentration-dependent manner, via cell-cycle arrest at the G0–G1 and S phases and apoptosis induction [100]. In another study, a



methanol crude extract from whole plants of the same species exhibited a potent cytotoxic activity toward A549 cells (human lung cancer) and Col2 cells (human colon cancer), with  $IC_{50}$  values of 8.0  $\mu\text{g/mL}$  and 27.4  $\mu\text{g/mL}$ , respectively [101]. A hydroalcoholic extract from shoots of *Cressa cretica* L. decreased the viability of HepG2 cells ( $IC_{50}$  value = 2300  $\mu\text{g/mL}$  after 72 h of incubation), by increasing the expression of the proapoptotic protein BAX in detriment to antiapoptotic proteins (BCL2) [102].

### 2.8. Cymodoceaceae Family

This family presently contains six genera and 17 species of seagrasses [103,104]. A water leaf extract from *Cymodocea rotundata* Ehrenb Hempr. ex Aschers. was supplemented with silver nitrate ( $\text{AgNO}_3$ , 1 M), to produce silver nanoparticles (AgNPs). AgNPs have several medical applications, including coating of medical devices and wound dressings, and they exhibit cytotoxic activity [105]. The obtained AgNPs exhibited high cytotoxicity toward MG63 cells with an  $IC_{50}$  value of 25.31  $\mu\text{g/mL}$  after 48 h of incubation [105]. AgNPs were also produced by combining a water leaf extract of *Cymodocea serrulata* (R. Br.) Aschers. & Magnus and  $\text{AgNO}_3$  (1 M), and they reduced the viability of A549 cells in a direct dose–response manner ( $IC_{50}$  = 100  $\mu\text{g/mL}$  after 24 h of incubation). The higher cytotoxicity of the AgNPs was ascribed to the easy permeability to the cellular barriers and their high affinity to biological macromolecules, as well as their capacity to release ROS [106]. No attempt was made to identify the active molecules present in those samples [105,106]. A hydroethanolic extract from shoots of the same species inhibited the proliferation of HepG2 cells ( $IC_{50}$  value of 82.92  $\mu\text{g/mL}$  after 24 h of incubation with camptothecin as a positive control with an  $IC_{50}$  value of 8  $\mu\text{g/mL}$ ) [107]. The extract had a high concentration of tannins, flavonoids, and terpenoids, with described antitumoral properties [108], but no attempt was made by the authors to identify the possible bioactive molecules [107].

### 2.9. Cyperaceae Family

*Cyperaceae* currently comprises 92 genera and 5888 species with a cosmopolitan distribution [109,110]. *Cyperus rotundus* L. (nut grass) is a perennial halophyte species traditionally used in the treatment of several pathologies, such as stomach disorders [111]. Methanol, ethanol, and water extracts from nut grass rhizomes were tested toward MDA-MB-231 cells, for 24 h [112] (Table 1). The highest reduction in cell viability was obtained with the ethanol extract, with an  $IC_{50}$  value of 225  $\mu\text{g/mL}$ , through apoptosis induction via upregulation of the death receptor 4 (DR4), DR5, and proapoptotic BAX, and downregulation of antiapoptotic survivin and BCL2 [112]. The active extract was not chemically profiled, but it is known that the rhizomes of nut grass are rich in several bioactive molecules, including flavonoids, tannins, and sesquiterpenes [113], with reported cytotoxic activity against cancer cell lines.

### 2.10. Fabaceae Family

This family currently comprises 778 genera and 22,356 species [114] and includes five genera of halophytic species, namely, *Alhagi*, *Glycyrrhiza*, *Melilotus*, *Prosopis*, and *Sesbania*, with described anticarcinogenic properties [115].

The genus *Alhagi* is distributed throughout Asia, Australia, and Europe and used traditionally for the treatment of, for example, gastroenteritis, ulcers, and rheumatoid arthritis [116]. *Alhagi maurorum* Medik (camelthorn) is used for its anti-inflammatory properties, which are ascribed to the presence of the triterpenoid lupeol [117]. In fact, lupeol was isolated from a methanol extract of camelthorn aerial parts, and it decreased the cellular viability of MCF7 and MDA-MB-231 cell lines, with one-fourth of  $IC_{50}$  values >100  $\mu\text{g/mL}$  (Table 1) [118]. The cytotoxic activity of lupeol was lower than that observed with its epoxide form, and it was related to the increase in mRNA expression levels of

apoptosis-related genes (TP53, caspase-3 and BAX) and decrease in BCL2 gene expression [118].

*Glycyrrhiza* (liquorice) contains legumes endogenous to Asia and southern Europe with reported anti-inflammatory and antiviral properties [119]. There are two *Glycyrrhiza* halophytic species with anticarcinogenic activity: *Glycyrrhiza glabra* L. and *G. uralensis* Fisch. *Glycyrrhiza glabra* is rich in phenolics, tannins, and especially glycyrrhizin, a triterpenoid saponin [120]. Glycyrrhizin reduced the proliferation of HeLa cells at 320  $\mu\text{M}$  after 24 and 48 h of incubation, via apoptosis, through mitochondrial depolarization. Moreover, nuclear condensation, cell membrane lysis, and disintegration of organelles were observed in treated cells through phase-contrast microscopy [121]. The cytotoxic potential of the roots' methanol extracts from *G. glabra*, collected in nine different areas from Italy, Turkey, Syria, Russia, Afghanistan, and Uzbekistan, was assessed against "normal" (HaCaT) and cancer cell lines (A549 and HepG2) using the MTT assay, after 24 h of incubation [122]. Results disclosed variable cytotoxicity levels depending on the samples' collection location and season, potentially related to the influence of climatic conditions on the chemical composition of the plants. Only one sample from Afghanistan was active toward HepG2 cells ( $\text{IC}_{50}$  of 248.5  $\mu\text{g}/\text{mL}$ ), while four extracts (from Italy, Afghanistan, and Syria) were cytotoxic toward A549 cells ( $\text{IC}_{50}$  between 189.1 and 238.9  $\mu\text{g}/\text{mL}$ ) [122] (Table 1). However, samples were also cytotoxic toward "normal" cells (HaCat) (Table 1). An ethanol root extract of *G. glabra* significantly reduced the proliferation of HT29 cells, at 200  $\mu\text{g}/\text{mL}$  after 24, 48, and 72 h of incubation as detected by the MTT assay [123]. Polymerase chain reaction (PCR) studies revealed a downregulation of heat-shock protein 90 (HSP90) gene expression that can be related to the reduction in cellular viability, since the HSP90 prevents tumor cells from undergoing apoptotic death; therefore, its blocking could assist active antitumor effects [124]. A reduction in HeLa cell viability above 80% was reported after application of an aqueous ethanol extract of *G. uralensis* rhizomes at the concentration of 1.84 mg/mL, but the determination of the  $\text{IC}_{50}$  value was not reported [125]. Several cytotoxic compounds were identified in that extract using UPLC-ESI-Q-TOF, including isoquercitrin, 4'-demethylpodophyllotoxin glucoside, and podophyllotoxin, all with described cytotoxic properties [125] (Table 2).

*Melilotus* contains 19 species of annual herbs widely distributed in North Africa and Eurasia [126]. *Melilotus indicus* L. All. (sweet clover), the only described halophyte within this genus, is used in traditional medicine as an analgesic, and it has reported cytotoxic activity [127,128]. A methanol extract from the aerial parts of sweet clover significantly decreased cell proliferation of HepG2 ( $\text{IC}_{50}$  = 16.60  $\mu\text{g}/\text{mL}$ ) and SNU-182 cells ( $\text{IC}_{50}$  = 13.21  $\mu\text{g}/\text{mL}$ ) after 24 h of incubation, using staurosporine as a positive control (Table 1) [127,128]. The extract was less cytotoxic toward L-02 cells (human "normal" hepatic) ( $\text{IC}_{50}$  = 90.9  $\mu\text{g}/\text{mL}$ ) (Table 1) [127,128]. The application of the extract resulted in an increase in the number of apoptotic cells and loss of mitochondrial membrane potential ( $\Delta\psi\text{m}$ ) [128] (Table 1). The chemical composition of the extract was not established by the authors.

*Prosopis* comprises 44 species, mainly small trees distributed in dry lands of America, Africa, and Asia. Several *Prosopis* species contain anti-inflammatory, antidiabetic, and anticancer compounds, namely, flavonoids, tannins, phenolics, and alkaloids [129]. *Prosopis juliflora* Sw. DC. (mesquite) is an invasive species in India with reported ethnomedicinal uses, for the treatment of eye and digestive disorders, to name a few [130]. Mesquite extracts display relevant bioactivities, including anti-inflammatory, ascribed mainly to its content in alkaloids [130]. A leaf methanol extract from *Prosopis juliflora* Sw. DC. (mesquite) had cytotoxic effects on Molt-4 cells, with  $\text{IC}_{50}$  values of 90.5, 42.5, and 20.0  $\mu\text{g}/\text{mL}$  after 24 h, 48 h, and 72 h of incubation, respectively, using mitomycin-C (6  $\mu\text{g}/\text{mL}$ ) as a positive control (Table 1). The extract was less toxic toward "normal" cells (mitogen stimulated T-lymphocyte cultures from peripheral human blood) [130]. A genotoxic assessment using a cytokinesis-block micronucleus assay reported that the number of

micronuclei showed an increasing pattern with the application of increasing concentration of the extract [130].

The genus *Sesbania* comprises 60 to 85 species of herbs, shrubs, and trees distributed mostly in the tropical and subtropical regions of Africa, Asia, Australia, and America. *Sesbania* species have relevant pharmacological properties such as anti-inflammatory and antidiabetic [131–133]. *Sesbania grandiflora* (Akatti) (sesbania or agathi), is a small perennial tree with high levels of vitamins and minerals associated with anti-inflammatory, analgesic, and antipyretic properties [134,135], and the anticancer properties of its fruit are mentioned in the Ayurvedic literature [136]. Water, ethanol, and acetone leaf extracts of *S. grandiflora* reduced the viability of IMR32 and HT29 cells with IC<sub>50</sub> values of 200 µg/mL after 24 h of incubation using doxorubicin as a control [137]. However, the chemical composition of the extract was not unraveled by the authors. Additional studies regarding the anticarcinogenic activity of other halophytes of this family are depicted in Table 1.

#### 2.11. Juncaceae Family

*Juncaceae* currently contains eight genera and 522 species [138] of herbs adapted to salty marshes or badly drained soils and accumulate different phytoconstituents such as flavonoids, triterpenes, steroids, and phenolic acid derivatives [139,140]. A phenanthrene, juncunol (1,7-dimethyl-5-vinyl-9,10-dihydrophenanthren-2-ol), was identified in a diethyl ether extract of leaves of *Juncus acutus* (L.) Torr. ex Retz. (spiny rush) and displayed selective in vitro cytotoxicity toward HepG2, HeLa, and MDA-MB-468 cell lines [141]. Juncunol had an IC<sub>50</sub> value of 18 µM in HepG2 cells after 72 h of incubation, determined using the MTT assay, and it induced an increase in the number of apoptotic cells in a concentration-dependent manner (IC<sub>50</sub> value ± 25%) accompanied by a decrease in the Δψ<sub>m</sub> [141,142]. Juncunol induced cell-cycle arrest in the G0/G1 phase, while showing no hemolytic properties. In silico studies indicate that that compound seems to bind between GC base pairs and, thus, may act as a DNA intercalator [142].

#### 2.12. Malvaceae Family

This family of flowering plants currently contains 243 genera and 5461 species [143]. A leaf decoction from *Thespesia populnea* L. Sol. ex Corrêa (portia tree) had a high content of total phenols and flavonoids and showed cytotoxic and antiproliferative properties toward HEP-2 cells, as observed in the MTT (IC<sub>50</sub> = 120.02 µg/mL after 24 h of incubation) and SRB (IC<sub>50</sub> = 77.06 µg/mL) assays [144]. Treated cells showed apoptotic characteristics including membrane blebbing, cell shrinkage, nuclear and cytoplasmic condensation, and formation of apoptotic bodies [144]. However, no attempt was made to identify the active molecules present in this sample. The chloroform-soluble fraction of a methanol bark extract of portia tree caused a reduction in cellular viability of MDA-MB-231 and MCF7 cells, with IC<sub>50</sub> values of 23.97 and 20.62 µg/mL respectively, after 24 h of incubation [145]. Chemical analysis of the extract using GC–MS revealed the presence of steroids such as cis-androsterone and fatty-acid derivatives, which display cytotoxic activity against cancer cell lines [108,145].

#### 2.13. Myrtaceae Family

*Myrtaceae* presently includes 134 genera and 6614 species [146]. Focus is given to the genus *Eucalyptus*, especially *Eucalyptus camaldulensis* Dehnh, because is a salt-tolerant species with reported antitumoral properties. *Eucalyptus* spp. are mainly cultivated for timber and paper production, but the presence of bioactive compounds, such as triterpenoids, flavonoids, and tannins, confers *E. camaldulensis* with relevant bioactivities, including antitumoral [147]. Methanol, ethyl acetate, *n*-butanol, and water extracts from leaves of *E. camaldulensis* reduced the viability of MCF7 and MDA-MB-231 cell lines, according to the MTT and SRB assays [148]. The ethyl acetate extract had the highest

cytotoxicity with IC<sub>50</sub> values of 26.7 and 7.9 µg/mL for MTT and SRB in MCF7 cells, and IC<sub>50</sub> values of 34.4 and 4.9 µg/mL for MTT and SRB in MDA-MB-231 cells after 24 h of incubation [148]. No attempt was reported by the authors to disclose the molecules responsible for the detected cytotoxic activity.

#### 2.14. Plantaginaceae Family

This family currently contains 101 genera and 2165 species [149], widely distributed worldwide but with preference for temperate zones [150]. *Bacopa monnieri* (L.) Wettst (water hyssop) is a wetland macrophyte and contains bacopaside (II) (saponine), with reported anticarcinogenic properties [151,152]. The anticarcinogenic properties of artificially digested (artificial saliva and artificial gastric juice) methanol extracts from *B. monnieri* reduced the motility (capacity of cell migration in a process of cancer development) of DU145 cells [153]. Nevertheless, the IC<sub>50</sub> value was not determined by the authors. The extracts, analyzed by HPLC, were rich in phenolic compounds such as chlorogenic, caffeic, and syringic acids and bacoside A, with reported anticarcinogenic properties [154–157].

Within the Plantaginaceae family, the *Plantago* genus is the largest with 275 annual and perennial species widespread around the world, some with traditional uses as antipyretic, anti-inflammatory, and antitumoral agents [158]. *Plantago major* L. (fleawort) is an important medicinal plant due to its richness in bioactive metabolites, such as alkaloids and flavonoids, especially luteolin-7-O-β-glucoside, with antitumoral properties [159]. Silver particles produced by combining a water extract from *P. major* seeds and silver nitrate (AgNO<sub>3</sub>, 0.1 M) significantly decreased the viability of MCF7 cells with an IC<sub>50</sub> value of approximately 12 µg/mL after 24 h of incubation [159]. However, no attempt was made by the authors to identify the bioactive compounds present in this extract. Information targeting other *Plantaginaceae* species is summarized in Table 1.

#### 2.15. Plumbaginaceae Family

*Plumbaginaceae* encompasses several species adapted to survive under saline conditions and currently includes 1138 species distributed across 26 genera [160]. The genus *Plumbago* contains 18 species (shrubs or perennial herbs) characterized by the presence of flavonoids, phenols and saponins, conferring several properties including anticarcinogenic [161,162]. The genus *Limoniastrum* Heist. ex Fabr. comprises the halophytic species *Limoniastrum monopetalum* (L.) Boiss. (≡ *Statice monopetala* L.) and *Limoniastrum guyonianum* Boiss. [163,164], thriving in coastal and saline dry areas of the Mediterranean and northern Saharan Africa [165–168]. *Limoniastrum guyonianum* is traditionally used in the treatment of gastric infections [169]. A gall aqueous extract from *L. guyonianum* reduced the viability of HeLa cells with IC<sub>50</sub> values of 170 and 140 µg/mL after 24 and 48 h of incubation, respectively [169]. This extract was rich in flavonoids and induced DNA hypomethylation and apoptosis due to its capacity to arrest cell-cycle progression in G2/M (Table 1). Due to the high levels of polyphenolic compounds detected in the extract, the antiproliferative and proapoptotic effects of such metabolites on HeLa cells were hypothesized by the researchers [169]. In another study, hexane, dichloromethane, ethanol, and methanol extracts from *L. densiflorum* (Guss.) Kuntze shoots were tested for cytotoxicity on cancer and “normal” cell lines using the resazurin reduction assay, after 48 h of treatment (Table 1) [170]. The dichloromethane extract was the most bioactive sample, with IC<sub>50</sub> values of 29 and 85 µg/mL toward A-549 and DLD-1, without significantly reducing the viability of a human skin fibroblast cell line (Table 1). The major compounds present in that extracts were identified using RP-HPLC as *trans* 3-hydroxycinnamic acid, myricetin, and isorhamnetin, which may be related to the observed cytotoxic activity [170].

#### 2.16. Poaceae Family

This grass family currently includes 11,917 species divided across 796 genera [171,172]. In this family, only two genera of halophytes are reported with anticarcinogenic properties, namely, *Cynodon* and *Echinochloa*. The most representative *Cynodon* species is *Cynodon dactylon* L. Pers (Bermuda grass), a weed with several medicinal properties, such as antidiabetic, diuretic, and purifying, as well as in vitro cytotoxic properties [173]. The in vitro antitumoral properties of petroleum ether, dichloromethane, acetone, methanol/water (3/1), and water extracts from Bermuda grass were appraised on a breast cancer cell line, and cell viability was assessed by evaluating (3H)-hypoxanthine incorporation after 48 h of incubation [174] (Table 1). The highest cytotoxicity values were observed after application of the water, acetone, and petroleum ether extract, with IC<sub>50</sub> values of 57.2, 38, and 39 µg/mL (Table 1) [174]. The acetone and ether petroleum extracts were profiled using LC-MS, allowing for the identification of several bioactive anthocyanins, namely, delphinidin, petunidin, malvidin, and cyanidin glucosides (delphinidin-3-O-acetylglucoside, petunidin-3-O-caffeoylglucoside-5-O-glucoside, petunidin-3-O-coumarylglucoside-5-O-glucoside, malvidin-3-O-monoglucoside, delphinidin-3-O-acetylglucoside-pyruvic acid, petunidin-3-O-acetylglucoside-5-O-glucoside, and cyanidin-3,5-O-diglucoside) [174] (Table 2). There is no information related to the specific anticancer properties of the detected anthocyanins; however, several others, such as delphinidin, cyanidin, malvidin, and corresponding glycosides, exhibit antitumoral properties in different cell lines [174]. An ethanol extract (70%) from grains of *Echinochloa crus-galli* (L.) P. Beauv. (cockspur grass) significantly reduced the viability of HCT-116 and HeLa cell lines, with IC<sub>50</sub> values of 11.2 and 12.0 µg/mL, respectively [175] (Table 1). The active ethanolic crude extract was fractionated using *n*-hexane, chloroform, ethyl acetate, and *n*-butanol, with the ethyl acetate fraction exhibiting the lowest IC<sub>50</sub> value (3.8 µg/mL). Fractionation of this active fraction allowed the isolation and identification of eight phenolic compounds, which exhibited significant cytotoxic activities toward the tested cells [175]. The most active molecules were 5,7-dihydroxy-3',4',5'-trimethoxy flavone and 5,7,4'-trihydroxy-3',5'-dimethoxy flavone (tricin), the latter with IC<sub>50</sub> values of 7.2, 8.6, 10.8, and 19.9 µM against HepG2, HeLa, HCT116, and MCF7 cell lines, respectively, with similar results to the commercial anticancer drug, doxorubicin [175]. Further data on the antitumoral properties of other species of the *Poaceae* family can be found in Table 1.

### 2.17. Portulacaceae Family

This family contains one genus with presently 150 species [176] distributed worldwide, including the edible *Portulaca oleracea* L. (purslane), a halophyte succulent annual plant with important nutraceutical and antioxidant properties [177,178]. The anticarcinogenic activity of this species can be ascribed to the presence of several compounds, such as portulacerebroside A, portulacanones B, and 2,2'-dihydroxy-4',6'-dimethoxychalcone [179] (Table 2). Portulacerebroside A induced apoptosis in HCCLM3 cells via activation of the p38 MAPK and JNK-triggered mitochondrial death pathway [180]. An ethanol extract from purslane seeds reduced the viability (MTT assay) of HepG2 cells in a concentration-dependent manner with IC<sub>50</sub> values of approximately 75 µg/mL after 24 h of incubation via apoptosis induction according to cellular morphology modification, i.e., a reduction in cellular size and adhesion capacity (Table 1) [181]. Oil from seeds of the same species reduced the proliferation of HepG2 and A-549 cancer cell lines, at concentrations ranging from 250 to 1000 µg/mL as observed using the MTT and neutral red uptake assays, after 24 h of incubation in both cancer cell lines [182]. However, the high concentrations of the extract used (up to 1000 µg/mL) must be highlighted; even at 100 µg/mL, no reduction in cell viability was observed (Table 1) When treated with the highest concentration, cells exhibited morphological modifications typical of apoptotic cells, such as loss of cell adhesion capacity, shrinkage, and round shape [182]. However, the chemical profile of the extract was not determined by the authors.

### 2.18. Rhizophoraceae Family

This family includes mangroves and comprises currently 148 species and 15 genera [183,184]. It has two halophytic species with reported cytotoxic properties against human cancer cells, namely, *Bruguiera gymnorhiza* (L.) Lam (black mangrove) and *Ceriops tagal* (Pers.) C. B. Rob. (oriental mangrove) [185,186]. A methanol extract from black mangrove stem bark reduced the proliferation of HeLa, RAJI, and myeloma cell lines, with IC<sub>50</sub> values of 133, 504, and 384 µg/mL after 24 h of incubation using doxorubicin as a standard control, via apoptosis induction, since DNA fragmentation was detected by fluorescence with double staining (ethidium bromide–acrydine orange) [187].

### 2.19. Solanaceae Family

*Solanaceae* contains 100 genera and 2925 species currently recognized [188], many of them with a high worldwide economic importance as cultivated crops, such as *Solanum lycopersicum* L. (tomato) [189]. The number of halophytic species with anticarcinogenic activity in this family is scarce, and reports have mainly focused on *Lycium barbarum* L. (Chinese wolfberry). Chinese wolfberry's dried fruits are widely used in traditional Chinese medicine for its anticancer properties, potentially associated with the presence of different bioactive compounds, including polysaccharides (proteoglycans), scopoletin (phytoalexin), and 2-O-β-D-glucopyranosyl-L-ascorbic acid (vitamin C analog) [190]. A polysaccharide-rich aqueous extract from Chinese wolfberry fruits reduced the proliferation of SW480 and Caco-2 cell lines, as observed by the crystal violet and MTT assays, with IC<sub>50</sub> values of 600 and 650 µg/mL respectively, after 4 days of incubation, with a high capacity to arrest cell-cycle progression in the G0/G1 phase [191]. An ethanol extract from the dried fruit residue of this species was cytotoxic toward A431 cells, with an IC<sub>50</sub> value of 873.7 µg/mL after 24 h of incubation, determined by water-soluble tetrazolium-8-[2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2Htetrazolium] monosodium salt (CCK-8) assay [192]. The same authors conducted in vivo studies in mice submitted to an intraperitoneal injection of 50 mg/kg of the crude extract, reporting a lower volume and weight of the formed tumor and a higher survival rate of tumor-bearing nude mice [192]. Several phenolic amides were isolated from an ethanol extract from the stem of *L. barbarum* and tested toward glioma cell lines, using the MTS method [180]. 4-O-Methylgrossamide and grossamide exhibited the highest capacity to decrease the viability of GSC-3# and GSC-12# cell lines, especially the latter compound, with IC<sub>50</sub> values of 6.40 and 5.85 µg/mL respectively. [193]. The precise mechanism of action was not appraised for the isolated compounds; however, there are reports that phenolic amides can inhibit tumor necrosis factor-alpha (TNF-α) and nuclear factor-kappa B (NF-κB), which may be related to the detected cytotoxic effect [194].

### 2.20. Tamaricaceae Family

*Tamaricaceae* contains five genera with currently 110 species [195], widely distributed around the world, and several are halophytic [196,197]. Tamarix plants (salt cedar) contain several bioactive metabolites such as methylferulate and syringic acid with antineoplastic properties [198]. Syringic acid was isolated from a methanol extract of *Tamarix aucheriana* (Decne.) Baum aerial parts and was cytotoxic toward SW1116 and SW837 cells in a time- (24–72 h of incubation) and dose-dependent manner, with IC<sub>50</sub> values of 0.95 and 1.2 mg/mL, respectively [199]. Cytotoxicity was related to an increase in the percentage of apoptotic cells, attributed to the antimetabolic effect of syringic acid [199]. Methylferulate was isolated from the same type of extract and significantly reduced the viability of SW1116 and SW837 cells, with IC<sub>50</sub> values of 1.7 and 1.9 mM after 24 h of incubation, respectively, being responsible for cell-cycle arrest in the S and G2/M phases and apoptosis induction, together with a reduction in NF-κB DNA-binding activity [200].

*Tamarix gallica* L. is traditionally used as an expectorant, antidiarrheal, and laxative, potentially attributed to the presence of ellagic and gallic acids, anthocyanins, tannins,

flavonones, isoflavonones, and resveratrol in its leaves [201]. A methanol extract from shoots, leaves, and flowers of *T. gallica* significantly reduced the viability of Caco-2 cells, related to a decrease in DNA biosynthesis, as observed using fluorescence microscopy after DAPI staining, while a reduction in the percentage of cells in the G0/G1 and S phases was detected using flow cytometry [202].

### 3. Conclusions and Future Perspectives

As stated in Section 1, we included in this review paper published work, which we considered relevant, related to the cytotoxic effects of halophytes. Such reports included results obtained with crude extracts, fractions, and pure compounds, using in vitro and in vivo assays, and they occasionally presented IC<sub>50</sub> values, phytochemical profile of the active samples, and identification of active molecules, along with proposed mechanisms of action. There are some drawbacks of the identified reports. For example, several authors tested considerably high concentrations of the target extracts, up to 1000 mg/mL, in some cases. Moreover, the IC<sub>50</sub> values were not reported in a high number of studies, which impairs a critical evaluation of the obtained results. Additionally, several reports failed to establish the chemical composition of the cytotoxic extracts and/or the identification of the molecules responsible for the detected activity. Lastly, information about the effect of the extracts on cell lines from nontumoral origin and, therefore, the selectivity of the extracts toward cancer cells was not always provided. According to the American National Cancer Institute (NCI), only crude extracts/fractions displaying IC<sub>50</sub> values lower than 30 µg/mL in the preliminary assay are considered promising and should be further explored as sources of antitumoral compounds. Therefore, we consider that extracts from the species and extracts included in Table 3 could be explored in detail targeting the isolation of anticancer leads, namely, *C. ambrosioides*, *S. fruticosa*, *A. millefolium*, *C. soldanella*, *C. rotundata*, *M. indicus*, *T. populnea*, *E. camaldulensis*, *P. major*, *P. lanceolata*, *E. crus-galli*, and *L. barbarum*. Moreover, isolated compounds with IC<sub>50</sub> values ≤10 µM are considered promising anticancer leads. Therefore, we highlight the compounds in Table 3, namely, 6'-*O*-(*n*-butanol) ilekudinoside B ester (*A. marina*), 5,7-dihydroxy-3',4',5'-trimethoxy flavone and tricin (*E. crus-galli*), Portulacerebroside A (*P. oleraceae*), and grossamide (*L. barbarum*).

**Table 1.** List of halophyte species, organs and extracts tested, cell lines/types of cancer assessed, and obtained half-maximal inhibitory concentrations (IC<sub>50</sub>) values.

Family	Species	Organs	Extract	Cell Line	IC <sub>50</sub>	Proposed Mechanism of Action	Ref.
<i>Acanthaceae</i>	<i>Acanthus ebracteatus</i> Vahl	Shoots	Protein hydrolysate	A431 cells (skin carcinoma)	425.9 ng/mL		[34]
	<i>Acanthus ilicifolius</i> L.	Leaves and roots	Ethyl acetate	MCF7 cells (breast carcinoma) PA1 cells (ovarian carcinoma)	24.22 µg/mL (leaf) 29.20 µg/mL (root); 15.74 µg/mL (leaf) 20.00 µg/mL (root)	nd	[33]
		Leaves	Ethanol	HepG2 cells (hepatocellular carcinoma)	100 µg/mL	Apoptosis induction (DNA damage)	[31]
		Roots	Water	HepG2 cells (hepatocellular carcinoma)	39.76 µg/mL	Apoptosis induction (DNA damage)	[32]
	<i>Avicennia alba</i> Blume	Leaves	Methanol	MCF7 and HeLa cells (breast and cervical carcinomas)	57.02 and 44.30 µg/mL	Reduction in cell size and cell detachment	[36]
		Leaves	Chloroform/methanol	WiDr cells (human colon carcinoma)	173.78 µg/mL	Apoptosis induction (cell arrest in the G0–G1 phase)	[37]
	<i>Avicennia marina</i> (Forssk.) Vierh	Leaves	Methanol	HeLa cells (cervical carcinoma)	107 µg/mL	nd	[40]
		Leaves	Methanol extract and fractions	MDA-MB-231 cells (breast carcinoma) HEK cells (human embryonic kidney)	MDA-MB-231 cells: Crude extract 250 µg/mL Active fraction (luteonin) 28 µg/mL (97 µM)	Luteonin: apoptosis induction (DNA fragmentation, decreased expression of BCL2, decreased expression of TP53).	[203]
		Fruits	Isolated compounds from ethanol/butanol fractions of ethanol extract	GSC-3# and GSC-18# cells (human glioma stem cell lines)	6'-O-( <i>n</i> -butanol) ilekudinoside B ester: 12.21 µg/mL (14 µM) for GSC-3# and 5.23 µg/mL (6 µM) for GSC-18#	nd	[204]



		Leaves and stems	Ethyl acetate	MCF-7 cells (estrogen positive breast cancer)	na	Apoptosis induction (ROS production, disruption of $\Delta\psi_m$ ), decrease in caspase-7 protein levels); autophagy	[39]
		Leaves and seeds	Water, ethanol, methanol, ethyl acetate extracts, and fractions from the latter	AU565, MDA-MB-231, and BT483 cells (breast cancer) HepG2 and Huh7 cells (liver) NIH3T3 cells (nontumoral)	Fraction 2-5: 0.75 $\mu\text{g/mL}$ on AU565 cells Fraction 3-2-9: 2.1 $\mu\text{g/mL}$ on AU565 cells	Apoptosis induction (DNA fragmentation, cell nucleus condensation and fragmentation, decreased PARP and caspase-8, increased caspase 3)	[38]
		Leaves	Ethyl acetate	Xenograft MDA-MB-231 tumor growth in nude mice	nd	Suppression of tumor growth	[38]
<i>Aizoaceae</i>	<i>Mesembryanthemum crystallinum</i> L.	Leaves	Ethanol extract, ethyl acetate and butanol fractions	HCT116 cells (colon carcinoma)	na	Ethyl acetate and butanol fractions: reduction of ROS, apoptosis induction). Butanol fraction: cell-cycle arrest at G2/M phase	[44]
	<i>Sesuvium portulacastrum</i> L.	Whole plant	Methanol, acetone, hexane, and diethyl-ether	MDA-MB-231 cells (breast carcinoma), IMR-32 cells (neuroblastoma), HCT116 cells (colon carcinoma)	Hexane extract: 942.07, 703.40 and 407.87 $\mu\text{g/mL}$ for MDA-MDB-231, IMR32 and HCT116 cells	Apoptosis induction: nuclear condensation, cell shrinkage, of apoptotic bodies	[45]
<i>Amaranthaceae</i>	<i>Arthrocnemum indicum</i> Willd. Moq.	Shoots	80% methanol	Caco-2 cells (colon carcinoma)	na	Decline of DNA synthesis, cell-cycle arrest at the G2/M phase.	[51]
	<i>Atriplex halimus</i> L.	Leaves	Ethanol	HepG2 cells (hepatocellular carcinoma) MCF-7 cells (human breast adenocarcinoma) A549 cells (lung cancer)	HepG2: 54.86 $\mu\text{g/mL}$ MCF-7: 153.6 $\mu\text{g/mL}$ A549: 101.9 $\mu\text{g/mL}$	Apoptosis induction (expression of TP53, BCL2, and BAX genes)	[54]

<i>Chenopodium album</i> L.	Branches and leaves	Petroleum ether	A549 cells (lung carcinoma)	33.31 µg/mL	Cell-cycle arrest at the G1 phase	[63]
<i>Chenopodium ambrosioides</i> L.	Whole plant	Essential oil	MCF7 cells (breast carcinoma)	18.75, 9.45 and 10.50 µg/mL after 6, 24 and 48 h of incubation	nd	[64]
<i>Chenopodium ambrosioides</i> L.	Leaves	Essential oil, ethanol extract, dichloromethane fraction	RAJI cells (lymphoblast) K562 cells (lymphoblast)	Essential oil: 1.0 g/mL; dichloromethane fraction: 34.0 g/mL (RAJI cells) Ethanol extract: 47.0 g/mL (K562 cells)	nd	[65]
<i>Chenopodium quinoa</i> Willd.	Seeds	Polysaccharide fraction	SMMC 7721 cells (liver cancer) MCF7 cells (breast carcinoma) L02 and MCF 10A cells ("normal" cell line)	121.4 µg/mL (24 h), 53.4 µg/mL (48 h) 83.5 µg/mL (24 h), 64.6 µg/mL (48 h)	nd	[66]
<i>Salicornia brachiata</i> Roxb	Shoots	Methanol	HepG2 cells (hepatocellular carcinoma)	267.84 µg/mL	Modification of cellular morphology	[205]
<i>Salicornia europaea</i> L.	Leaves	Ethyl acetate and methanol	MCF7 cells (breast carcinoma)	97.9 and 117.1 µg/mL	nd	[68]
<i>Suaeda fruticosa</i> (L.) Forssk	Leaves	Hexane	HCT116 cells (colon carcinoma) HepG2 cells (hepatocellular carcinoma), and MCF7 cells (breast carcinoma)	17.2 µg/mL 33 µg/mL 28.1 µg/mL	Apoptosis induction, cell-cycle arrest at the G0–G1 phase, chromatin condensation, membrane blebbing	[74]
<i>Suaeda fruticosa</i> (L.) Forssk	Shoots	Hexane, dichloromethane, methanol, and water	A549, DLD-1, Caco-2 and HT-29 cells (lung and colon carcinoma)	Hexane extract: 49, 10, 140 and 12 µg/mL	nd	[80]

	<i>Suaeda monoica</i> Forssk	Whole plant	Ethanol, methanol, acetone, and diethyl ether	MDA-MB-231 cells (breast carcinoma)	Ethanol: 172.38 µg/mL; methanol: 148.77 µg/mL; acetone: 185.56 µg/mL; diethyl ether: 60.18 µg/mL	nd	[206]
	<i>Suaeda palaestina</i> Eig Zohary	Shoots	Dichloromethane	A549 cells (lung carcinoma) HepG2 cells (hepatocellular carcinoma)	34.82 µg/mL 30.76 µg/mL	nd	[207]
	<i>Suaeda salsa</i> L.	Shoots	Acidic polysaccharide (molecular weight = 53.8 kDa; composition: mannose, rhamnose, glucuronic acid, galacturonic acid, galactose and xylose in a molar ratio of 0.6: 8.0: 1.0: 83.6: 5.0: 7.2).	MCF7 and MCF-10A cells (human breast carcinoma)	na	Apoptosis induction (reduction of $\Delta\psi_m$ , increase in the levels of BAX, cytochrome C, caspase-3 and caspase-9, decrease in the level of Bcl-2)	[208]
<i>Apiaceae</i>	<i>Crithmum maritimum</i> L.	Whole plant	Hexane, ethyl acetate, methanol, ethanol	Huh7 and HepG2 cells (hepatocellular carcinoma)	na	Ethyl acetate extract: cell-cycle arrest at the G0/G1 phase after 24 h of incubation and in the G2/M phase after 48 h; increase in necrotic and apoptotic cells	[85]
	<i>Eryngium maritimum</i> L.	Shoots and roots	Water	HepG2 cells (human hepatocellular carcinoma) Hep2 cells (human laryngeal epidermoid) U138-MG cells (human glioma) Vero cells (African green monkey kidney)	Shoots: 32.4 µg/mL; roots: 35.0 µg/mL Shoots: 50 µg/mL; roots: 30.3 µg/mL Shoots: 32.9 µg/mL; roots: 16.3 µg/mL	nd	[70]
<i>Asteraceae</i>	<i>Achillea millefolium</i> L.	Shoots	Methanol extract combined with bleomycin	DU145 cells (prostate carcinoma) and HFFF2	nd	nd	[88]

				cells (human non-malignant fibroblasts)			
		Aerial parts	Petroleum ether, ethyl acetate, methanol, water	K562 cells (human myelogenous leukemia), HeLa cells (human cervical carcinoma) MCF7 cells (human breast carcinoma) A549 cells (human non-small cell lung)	Ethyl acetate: 0.58 µg/mL (HeLa), 0.73 µg/mL (K562) Water: 0.87 µg/mL (MCF-7) Petroleum ether: 0.87 µg/mL (K562)		Ethyl acetate extract: pre-G1 apoptosis and cell growth arrest in G2/M (HeLa) [90]
	<i>Limbarda crithmoides</i> (L.) Dumort	Aerial parts	Methanol extract, <i>n</i> -hexane, dichloromethane, and aqueous methanol-soluble fractions, isolated compounds	OCI-AML3 cells (acute myeloid leukaemia)	nd		10-acetoxy-9Z-chloro-8,9-dehydrothymol: apoptosis induction [92]
<i>Brassicaceae</i>	<i>Cakile maritima</i> Scop.	Aerial organs	Methanol extract, and <i>n</i> -hexane, ethyl acetate, and methanol	Caco2 and HeLa cells (colon and cervical carcinoma)	Hexane fraction: 12 and 126 µg/mL		nd [96]
<i>Convolvulaceae</i>	<i>Calystegia soldanella</i> L.	Whole plant	Methylene chloride and methanol extracts, <i>n</i> -hexane, 85% methanol, <i>n</i> -butanol and water fractions	HepG2 cells (hepatocellular carcinoma)	na		Methanol fraction: cell-cycle arrest at the G0–G1 and S phases, apoptosis induction [100]
	<i>Calystegia soldanella</i> L.	Whole plant	Methanol	A549 cells (human lung cancer) Col2 cells (human colon cancer)	8.0 µg/mL 27.4 µg/mL		nd [101]
	<i>Cressa cretica</i> L.	Shoots	Hydroalcoholic	HepG2 cells (hepatocellular carcinoma)	2300 µg/mL		Increased BAX, decreased BCL2 [102]
<i>Cymodoceaceae</i>	<i>Cymodocea rotundata</i> Ehrin.	Leaves	Silver particles produced by combining water extract and silver nitrate (AgNO <sub>3</sub> , 1 M)	MG63 cells (osteosarcoma)	25.31 µg/mL		nd [105]

	<i>Cymodocea serrulata</i> (R. Br.) Aschers. & Magnus	Leaves	Silver particles produced by combining water extract and silver nitrate (AgNO <sub>3</sub> , 1 M)	A549 cells (lung carcinoma)	100 µg/mL	nd	[106]
		Shoots	Hydroethanolic	HepG2 cells (hepatocellular carcinoma)	82.92 µg/mL	nd	[107,108]
<i>Cyperaceae</i>	<i>Cyperus rotundus</i> L.	Rhizomes	Methanol, ethanol, and water.	MDA-MB-231 cells (breast carcinoma)	225 µg/mL	Apoptosis induction via upregulation of the death receptor 4 (DR4), DR5, and pro-apoptotic BAX, and downregulation of antiapoptotic BCL2	[112]
<i>Fabaceae</i>	<i>Alhagi maurorum</i> Medik	Aerial organs	Lupeol (isolated from a methanol extract)	MCF7, MDA-MB-231 and MCF 10A cells (breast carcinoma)	>100 µg/mL	Increased mRNA expression and level of TP53, caspase-3, and BAX genes, decrease in BCL2 gene expression	[118]
	<i>Glycyrrhiza glabra</i> L.	Root	Methanol	A549 cells (human lung carcinoma) HepG2 cells (human hepatocellular carcinoma) HaCaT cells (immortal human keratinocyte)	189.1–238 µg/mL 248.5 µg/mL 158.8–241.9 µg/mL	nd	[122]
		Root	Ethanol	HT-29 cells (colon carcinoma)	na	Downregulation of heat-shock protein 90 (HS90) gene expression	[123]
	<i>Glycyrrhiza uralensis</i> Fisch	Root	Ethanol/water (7:3, v/v)	HeLa cells (cervical carcinoma)	na	nd	[125]
	<i>Melilotus indicus</i> L.	Aerial parts	Methanol	HepG2 cells (human hepatocellular carcinoma) SNU-182 cells (hepatocellular carcinoma) L-02 cells (human “normal” hepatic)	16.60 µg/mL 13.21 µg/mL 90.9 µg/mL	Increase in the number of apoptotic cells, loss of mitochondrial membrane potential (Δψ <sub>m</sub> )	[128]


	<i>Prosopis cineraria</i> L. Druce	Leaves	Methanol	MCF7 cells (breast carcinoma) and HBL-100 cells (noncancerous breast)	na	nd	[209]
	<i>Prosopis juliflora</i> Sw. DC.	Leaves	Methanol	Molt-4 cells (human T-cell leukemia)	90.5, 42.5 and 20.0 µg/mL (24 h, 48 h and 72 h of incubation)	Increased number of micronuclei	[130]
	<i>Sesbania grandiflora</i> L.	Leaves	Water, ethanol, and acetone	IMR32 and HT-29 cell lines (neuroblastoma and colon carcinoma)	200 µg/mL		[137]
<i>Juncaceae</i>	<i>Juncus acutus</i> L. Torr. Ex Retz.	Shoots	Diethyl ether, chloroform, methanol, and water	HepG2 cells (hepatocellular carcinoma) S17 cells (murine non-tumoral)	Ether extract: 6.2 and 34 µg/mL Juncunol: 18 µM (HepG2 cells)	Juncunol: increased number of apoptotic cells, decrease in the $\Delta\psi_m$ , cell-cycle arrest in the G0/G1 phase, no hemolytic properties	[129,130]
<i>Malvaceae</i>	<i>Thespesia populnea</i> Sol. Ex Corrêa	Leaves	Decoction	HEP-2 cells (epidermoid carcinoma)	120.02 µg/mL	Apoptosis induction (membrane blebbing, cell shrinkage, nuclear and cytoplasmic condensation, apoptotic bodies)	[144]
		Stem bark	Methanol	MDA-MB-231 and MCF7 cells (breast carcinoma)	23.97 and 20.62 µg/mL	nd	[145]
<i>Myrtaceae</i>	<i>Eucalyptus camaldulensis</i> Dehnh	Leaves	Ethyl acetate	MDA-MB-231 and MCF7 cells (breast carcinoma)	26.7 and 34.4 µg/mL	nd	[148]
<i>Plantaginaceae</i>	<i>Bacopa monnieri</i> L. Wettst	In vitro cultures (shoots)	Methanol and artificial saliva and gastric juice extracts	DU145 cells (prostate cancer)	nd	nd	[153]
	<i>Plantago lanceolata</i> L.	Leaves	Ethanol	MCF7, AMJ13, MDAMB and CAL51 cells (breast carcinoma)	0.674, 0.726, 0.251 and 0.024 mg/mL	nd	[210]
	<i>Plantago major</i> L.	Seeds	Silver particles produced by combining water	MCF7 cells (breast carcinoma)	12 µg/mL	nd	[159]

		extract and silver nitrate (AgNO <sub>3</sub> , 0.1 M)					
Plumbaginaceae	<i>Limoniastrum densiflorum</i> (Guss.) Kuntze	Shoots	Hexane, dichloromethane, ethanol, and methanol	A-549 cells (human lung adenocarcinoma) DLD-1 cells (human colon carcinoma) WS-1 cells (human skin fibroblasts)	Dichloromethane extract: 85 µg/mL (DLD-1)-and 29 µg/mL (A-549)	nd	[170]
	<i>Limoniastrum guyonianum</i> Boiss	Gall	Water	HeLa cells (cervical carcinoma)	170 µg/mL	DNA hypomethylation and apoptosis, cell-cycle arrest at G2/M, upregulation of p16INK4A, upregulation of UHRF1 and DNMT1	[169]
Poaceae	<i>Cynodon dactylon</i> L.	Whole plant	Petroleum ether, dichloromethane, acetone, methanol/water (3/1) and water	MCF7 cells (breast carcinoma)	Water extract: 57.21 µg/mL; acetone extract: 38 µg/mL; petroleum ether extract: 39 µg/mL	nd	[174]
		Whole plant	Petroleum ether	HEP-2, HeLa, and MCF7 cells (laryngeal, cervical, and breast carcinoma)	0.20, 0.62 and 1.02 mg/mL	Apoptosis induction (DNA fragmentation)	[211]
	<i>Echinochloa crus-galli</i> L.	Grains	Ethanol extract, <i>n</i> -hexane, chloroform, ethyl acetate, and <i>n</i> -butanol fractions, isolated compounds	MCF7, HCT116, HeLa, and HepG2 cells (breast, colon, cervical, and liver carcinoma)	Crude extract: 12.0, 11.2, 18.9, and 14.2 µg/mL Ethyl acetate fraction: 3.8 µg/mL	nd	[175]
Portulacaceae	<i>Portulaca oleracea</i> L.	Seeds	Ethanol	HepG2 cells (hepatocellular carcinoma)	75 µg/mL	Round cells with reduced size and adhesion	[181]
		Seeds	Oil extraction	HepG2 and A549 cells (liver and lung carcinoma)	nd	Loss of cell adhesion capacity, shrinkage, round shape	[182]
Rhizoporaceae	<i>Bruguiera gymnorhiza</i> L. Lam	Stem bark	Methanol	HeLa cells (cervical cancer), Raji cells (lymphoma) and myeloma cells (leukemia)	133, 504, and 384 µg/mL	Apoptosis induction (DNA fragmentation)	[187]







<i>Solanaceae</i>	<i>Lycium barbarum</i> L.	Fruits	Water	SW480 and Caco-2 cells (colon carcinoma)	600 and 650 µg/mL	Cell-cycle arrest at G0/G1	[191]
		Fruits	Ethanol	A431 cells (cutaneous squamous cell carcinoma)	873.7 µg/mL	Reduced expression of Ki67 and PCNA proteins, increased expression of caspase-3, reduction of BCL2, downregulation of LC3II, reduced the phosphorylation of ERK1/2, and upregulation of JNK. Reversion of the regulation of Beclin1, LC3II, Bcl-2, and cl-caspase-3	[192]
		Fruits	Ethanol	BALB/c nude mice transplanted tumor model established by subcutaneous injection of A431 cells	nd	Increased survival rate, reduced tumor volume and weight, downregulation of Ki67 and MMP-2	
<i>Tamaricaceae</i>	<i>Tamarix aucheriana</i> (Decne.) Baum	Shoots	Syringic acid isolated from a methanol extract	SW1116 and SW837 cells (colorectal carcinoma)	0.95 and 1.2 mg/mL	Increased percentage of apoptotic cells	[199]
		Shoots	Methyl ferulate isolated from methanol extract	SW1116 and SW837 cells (colorectal carcinoma)	1.7 and 1.9 mM	Cell-cycle arrest in the S and G2/M phases, apoptosis induction, reduction in NF-κB DNA-binding activity	[200]
	<i>Tamarix gallica</i> L.	Shoots, leaves and flowers	80% methanol	Caco-2 cells (colon carcinoma)	na	Decreased DNA synthesis, cell-cycle arrest at G2/M phase; modification in the levels of cyclin B1, p38, Erk1/2, Chk1, and Chk2	[202]








nd: not determined;  $\Delta\psi$ m: mitochondrial membrane potential, na: data not available.







**Table 2.** Identified molecules in active extracts from halophyte species.






Family	Species	Chemical Compounds	Chemical Structure	Class	Ref.
<i>Acanthaceae</i>	<i>Avicennia alba</i> Blume	Catechol borane		Organoboron	[36]















		Neophytadiene		Terpenoid (diterpene)	[36]
		Hexadecanoic acid		Fatty acid (saturated)	[36]
	<i>Avicennia marina</i> (Forssk.) Vierh	Luteonin		Phenolic (flavonoid)	[203]
<i>Amaranthaceae</i>	<i>Arthrocnemum indicum</i>	Gallic acid		Phenolic (phenolic acid)	[51]
		Cyanidin		Phenolic (flavonoid)	[51]
		Chrysoeriol		Phenolic (flavonoid)	[51]







		Quercetin		Phenolic (flavonoid)	[51]
		Catechol		Phenolic (benzenediol)	[51]
		Luteolin		Phenolic (flavonoid)	[51]
	<i>Chenopodium ambrosioides L.</i>	Ascaridol		Monoterpene	[65]
<i>Asteraceae</i>	<i>Achillea millefolium</i>	Chlorogenic acid		Phenolic (phenolic acid)	[90]
		<i>p</i> -Coumaric acid		Phenolic (phenolic acid)	[90]
		Rosmarinic acid		Phenolic (polyphenol)	[90]






		Apigenin		Phenolic (flavonoid)	[90]
<i>Glycyrrhiza sp</i>		Glycyrrhizin		Saponin	[120,121]
<i>Limbarda crithmoides</i> (L.) Dumort		10-acetoxy-8,9-epoxythymol tiglate		Thymol derivative	[92]
		10-acetoxy-9Z-chloro-8,9-dehydrothymol		Thymol derivative	[92]
<i>Brassicaceae</i>	<i>Cakile maritima</i> Scop.	2-hydroxy-1,8-cineole		Monoterpenoid	[96]
		Decane		Alkane	[96]

		Limonene		Monoterpenoid	[96]
<i>Fabaceae</i>	<i>Alhagi maurorum</i> Medik	Lupeol		Terpenoid (triterpenoid)	[118]
	<i>Glycyrrhiza uralensis</i> Fisch	Isoquercitrin		(Phenolic) flavonoid	[125]
		4'-Demethylpodophyllotoxin glucoside		Lignan	[125]
		Podophyllotoxin		Lignan	[125]

<i>Juncaceae</i>	<i>Juncus acutus</i> L.	Juncunol		Phenanthrene	[141]
<i>Plantaginaceae</i>	<i>Bacopa monnieri</i> (L.) Wettst	Bacopaside (II)		Terpenoid (isoprenoid)	[153]
	<i>Plantago lanceolata</i> L.	O-Coumaric acid		Phenolic (phenolic acid)	[210]
		Rutin		Phenolic (flavonoid)	[210]
		Myricetin		Phenolic (flavonoid)	[210]
		Quercetin		Phenolic (flavonoid)	[210]


		Kaempferol		Phenolic (flavonoid)	[210]
<i>Plumbagina- ceae</i>	<i>Limonium densiflorum</i> (Guss.) Kuntze	<i>trans</i> 3-Hydroxycinnamic acid		Phenolic (phenolic acid)	[170]
		Myricetin		Flavonoid (phenolic)	[170]
		Isorhamnetin		Flavonoid (phenolic)	[170]
<i>Poaceae</i>	<i>Cynodon dactylon</i> L. Pers	Delphinidin-3-O-acetylglucoside		Anthocyanin	[174]
	<i>Echinochloa crus-galli</i>	5,7-Dihydroxy-3',4',5'-trimethoxy flavone		Phenolic (flavonoid)	[175]

		5,7,4'-Trihydroxy-3',5'-dimethoxy flavone (tricin)		Phenolic (flavonoid)	[175]
<i>Portulacaceae</i>	<i>Portulaca oleracea</i> L.	Portulacerebroside A		Cerebroside	[179,180]
		Portulacanonones B		Phenolic (flavonoid)	[179,180]
		Portulacanonones C		Phenolic (flavonoid)	[179,180]
		Portulacanonones D		Phenolic (flavonoid)	[179,180]
		2,2'-Dihydroxy-4',6'-dimethoxychalcone		Phenolic (flavonoid)	[179,180]

Solanaceae	<i>Lycium barbarum</i> L.	4-O-Methylgrossamide		Phenolic amide	[193]
		Grossamide		Phenolic amide	[193]
	<i>Lycium barbarum</i> L.	Scopoletin		Phenolic (coumarin)	[190]
		2-O-β-D-Glucopyranosyl-L-ascorbic acid		Glucoside	[190]
Tamaricaceae	<i>Tamarix aucheriana</i> (Decne.) Baum	Syringic acid		Phenolic (phenolic acid)[198,200]	



Methyl ferulate



Phenolic (phenolic acid)[198,200]

**Table 3.** Cytotoxic activity of selected halophytes species, including types of extracts, compounds, cell lines tested, obtained IC<sub>50</sub> values, and proposed mechanisms of action. The criterion of cytotoxic activity for the crude extracts, as established by the American National Cancer Institute (NCI), was considered as an IC<sub>50</sub> < 30 µg/mL upon 48 h or 72 h of incubation [24].

Family	Species	Organ	Extract/Fraction	Compound Detected in the Extract	Cell lines	IC <sub>50</sub> Values	Mechanism	Ref.
Amaranthaceae	<i>Chenopodium ambrosioides</i> L.	Whole plant	Essential oil	nd	MCF7	18.75, 9.45 and 10.50 µg/mL at 6, 24 and 48 h	DNA fragmentation	[64]
	<i>Suaeda fruticosa</i> (L.) Forssk.	Leaves	Hexane	Monoterpenes (dihydrojasnone, jasmolone, terpinene-4-ol), diterpenes (pimaric acid, steviol, and momilactone B) and phenolics (quercinol, zingerone, zingerol, neovaflan)	HCT116	17.2 µg/mL	Cell-cycle arrest at the C0–G1 phase and apoptosis induction	[74]
	<i>Suaeda fruticosa</i> (L.) Forssk.	Shoots	Dichloromethane	nd	DLD-1 HT-29	10 µg/mL 12 µg/mL	nd	[80]
Asteraceae	<i>Achillea millefolium</i> L.	Aerial parts	Ethyl acetate, water, and petroleum ether extracts	Ethyl acetate: p-coumaric acid, chlorogenic acid, and apigenin Water extract: rosmarinic acid	K562 cells HeLa cells MCF7 A549 cells	Ethyl acetate: 0.58 µg/mL (HeLa), 0.73 µg/mL (K562) Water: 0.87 µg/mL (MCF-7) Petroleum ether: 0.87 µg/mL (K562)	Ethyl acetate extract: preG1 apoptosis and cell growth arrest in G2/M (HeLa)	[90]
Brassicaceae	<i>Cakile maritima</i> Scop.	Aerial parts	Hexane	GC-MS, decane, limonene, nonaldehyde, dodecane	CaCo2	12 µg/mL	nd	[96]
Convolvulaceae	<i>Calystegia soldanella</i> (L.) R.Br. ex Roem. & Schult.	Whole plant	85% aqueous methanol fraction of crude methanol extract	nd	HepG2	< 30 µg/mL	Cell-cycle arrest at the G0–G1 and S phases, apoptosis induction	[100]
		Whole plant	Methanol	nd	A549 cells (human lung cancer) Col2 cells (human colon cancer)	8.0 µg/mL 27.4 µg/mL	nd	[101]

<i>Cymodoceae</i>	<i>Cymodocea rotundata</i> EhrenbHempr. ex Aschers.	Leaves	Water	nd	MG63	25.31 µg/mL	nd	105
<i>Fabaceae</i>	<i>Melilotus indicus</i> L. All.	Aerial parts	Methanol	nd	HepG2 SNU-182	16.6 µg/mL 13.21 µg/mL	Increase in the number of apoptotic cells, loss of Δψm.	[128]
	<i>Prosopis juliflora</i> Sw. DC.	Leaves	Methanol	nd	Molt-4	20.0 µg/mL after 72 h	Reduction in micronuclei or cell proliferation	[130]
<i>Malvaceae</i>	<i>Thespesia populnea</i> L. Sol. ex Corrêa	Bark	Chloroform fraction of methanol extract	Flavonoids, triterpenes, and tannins	MDA-MB-231 MCF7	23.97 µg/mL 20.62 µg/mL	nd	[145]
<i>Myrtaceae</i>	<i>Eucalyptus camaldulensis</i>	Leaves	Methanol, ethyl acetate, <i>n</i> -butanol, and water	nd	MCF7 MDA-MB-231	26.7 µg/mL 7.9 µg/mL 4.9 µg/mL	nd	[148]
<i>Plantaginaceae</i>	<i>Plantago major</i> L.	Seeds	Water	nd	MCF7	12 µg/mL	nd	[159]
	<i>Plantago lanceolata</i> L.	Leaves	Ethanol	Flavonoid glycosides: <i>O</i> -cumarin, rutin, myricetin, quercetin and kaempferol	CAL51	24 µg/mL	Apoptosis induction increase in the nuclei condensation	[210]
<i>Poaceae</i>	<i>Echinochloa crus-galli</i> (L.) P. Beauv.	Grains	70% ethanol	nd	MCF7 HCT116 HeLa HepG2	12.0 µg/mL 11.2 µg/mL 18.9 µg/mL 4.2 µg/mL	nd	[175]
		Grains		5,7,4-Trihydroxy-3,5-dimethoxy flavone (tricin)	MCF7 HCT116 HeLa HePG2	4.3 µg/mL 4.5 µg/mL 4.5 µg/mL 4.2 µg/mL	nd	[175]
<i>Solanaceae</i>	<i>Lycium barbarum</i> L.	Stems	Ethanol	Phenolic amides	GSC-3# GSC-12#	28 µg/mL 20 µg/mL	nd	[193]

nd: not determined; Δψm: mitochondrial membrane potential.

**Table 4.** Cytotoxic activity of selected isolated compounds from halophytes species, including cell lines tested, obtained IC<sub>50</sub> values, and proposed mechanisms of action. The criterion of cytotoxic activity for the isolated compounds was an IC<sub>50</sub> value ≤10 μM, according to the National Institutes of Health (NIH) for screening the NCI60 program [25].

Family	Species	Organ	Compound	Cell Lines	IC <sub>50</sub> Values	Mechanism	Ref.
<i>Acanthaceae</i>	<i>Avicennia marina</i> (Forssk.) Vierh	Fruits	6'-O-( <i>n</i> -butanol) ilekudinoside ester	BGSC-18# (human glioma stem cell lines)	6 μM	nd	[204]
<i>Poaceae</i>	<i>Echinochloa. crus-galli</i> (L.) P. Beauv.	Grains	5,7-dihydroxy-3',4',5'-trimethoxy flavone	HeLa	3.0 μM	nd	[175]
				HePG <sub>2</sub>	3.0 μM		
				HCT-116	10.8 μM		
			5,7,4-trihydroxy-3,5-dimethoxy flavone (tricin)	HeLa	8.6 μM	nd	
				HePG <sub>2</sub>	7.2 μM		
<i>Portulacaceae</i>	<i>Portulaca oleracea</i> L.	Aerial organs	Portulacerebroside A	HCCLM3	<3.5 μM	Apoptosis induction via activation of the p38 MAPK and JNK-triggered mitochondrial death pathway	[180]
<i>Solanaceae</i>	<i>Lycium barbarum</i> L.	Stems	Grossamide	GSC-3#	10.2 μM	nd	[193]
				GSC-12#	9.3 μM		

nd: not determined.

**Author Contributions:** P.G.-C., conceptualization, data curation, and writing—original draft preparation; P.C.-B., writing—review and editing; C.G.P., writing—review and editing; L.C., conceptualization, supervision, and writing—review and editing. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study received Portuguese national funds from FCT—Foundation for Science and Technology through projects UIDB/04326/2020, UIDP/04326/2020 and LA/P/0101/2020, and the FCT Scientific Employment Stimulus (CEECIND/00425/2017). It also received funding through the project HaloFarMs (PRIMA/0002/2019), which is part of the Partnership on Research and Innovation in the Mediterranean Area (PRIMA) Program supported by the European Union and funded by the national funding bodies of Participating States (FCT in Portugal).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Nurmik, M.; Ullmann, P.; Rodriguez, F.; Haan, S.; Letellier, E. In search of definitions: Cancer-associated fibroblasts and their markers. *Int. J. Cancer* **2020**, *146*, 895–905. <https://doi.org/10.1002/ijc.32193>.
- ACS 2020. American Cancer Society: Cancer Facts and Figures 2020. Atlanta, Ga: American Cancer Society, 2020. Available online: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2020/cancer-facts-and-figures-2020.pdf> (accessed on 03 December 2020).
- Laconi, E.; Marongiu, F.; De Gregori, J. Cancer as a disease of old age: Changing mutational and microenvironmental landscapes. *Br. J. Cancer* **2020**, *122*, 943–952. <https://doi.org/10.1038/s41416-019-0721-1>.
- GBD 2015 Risk Factors Collaborators. *Global, Regional, and National Comparative Risk Assessment of 79 Behavioural, Environmental and Occupational, and Metabolic Risks or Clusters of Risks, 1990–2015: A Systematic Analysis for the Global Burden of Disease Study 2015*; Lancet: London, UK, 2016; pp 388–1659.
- Anastasiadi, Z.; Lianos, G.D.; Ignatiadou, E.; Harissis, H.V.; Mitsis, M. Breast cancer in young women: An overview. *Updat. Surg.* **2017**, *69*, 313–317. <https://doi.org/10.1007/s13304-017-0424-1>.
- Wadowska, K.; Bil-Lula, I.; Trembecki, Ł.; Śliwińska-Mossoń, M. Genetic markers in lung cancer diagnosis: A review. *Int. J. Mol. Sci.* **2020**, *21*, 4569. <https://doi.org/10.3390/ijms21134569>.
- Arbyn, M.; Weiderpass, E.; Bruni, L.; de Sanjosé, S.; Saraiya, M.; Ferlay, J.; Bray, F. Estimates of incidence and mortality of cervical cancer in 2018: A worldwide analysis. *Lancet Glob. Health* **2020**, *8*, 191–203. [https://doi.org/10.1016/S2214-109X\(19\)30482-6](https://doi.org/10.1016/S2214-109X(19)30482-6).
- NCI (National Cancer Institute). Available online: <https://www.cancer.net/es/national-cancer-institute-nci> (accessed on 3 January 2021).
- Mattiuzzi, C.; Lippi, G. Current cancer epidemiology. *J. Epidem. Glob. Health* **2019**, *9*, 217–222. <https://doi.org/10.2991/jegeh.k.191008.001>.
- Nigam, M.; Suleria, H.A.R.; Farzaei, M.H.; Mishra, A.P. Marine anticancer drugs and their relevant targets: A treasure from the ocean. *DARU J. Pharm. Sci.* **2019**, *27*, 491–515. <https://doi.org/10.1007/s40199-019-00273-4>.
- Bracci, L.; Schiavoni, G.; Sistigu, A.; Belardelli, F. Immune-based mechanisms of cytotoxic chemotherapy: Implications for the design of novel and rationale-based combined treatments against cancer. *Cell Death Differ.* **2014**, *21*, 15–25. <https://doi.org/10.1038/cdd.2013.67>.
- Wu, Q.; Yang, Z.; Nie, Y.; Shi, Y.; Fan, D. Multi-drug resistance in cancer chemotherapeutics: Mechanisms and lab approaches. *Cancer Lett.* **2014**, *347*, 159–166. <https://doi.org/10.1016/j.canlet.2014.03.013>.
- Bhanot, A.; Sharma, R.; Noolvi, M.N. Natural sources as potential anti-cancer agents: A review. *Int. J. Phyto.* **2011**, *3*, 9–26.
- Atanasov, A.G.; Waltenberger, B.; Pferschy-Wenzig, E.M.; Linder, T.; Wawrosch, C.; Uhrin, P.; Temml, V.; Wang, L.; Schwaiger, S.; Heiss, E.H.; et al. Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotech. Adv.* **2015**, *33*, 1582–1614. <https://doi.org/10.1016/j.biotechadv.2015.08.001>.
- Willdenov, C.L. *The Principles of Botany, and of Vegetable Physiology*; University Press: Edinburgh, UK, 1805.
- Grigore, M.N. Defining halophytes: A conceptual and historical approach. In *Halophytes and Climate Change: Adaptive Mechanisms and Potential Uses*; Hasanuzzaman, M., Shabala, S., Fujita, M., Eds.; CABI: Wallingford, UK, 2019; pp. 3–18.
- Flowers, T.J.; Colmer, T.D. Salinity tolerance in halophytes. *New Phytol.* **2008**, *179*, 945–963. <https://doi.org/10.1111/j.1469-8137.2008.02531.x>.
- Ksouri, R.; Ksouri, W.M.; Jallali, I.; Debez, A.; Magné, C.; Hiroko, I.; Abdelly, C. Medicinal halophytes: Potent source of health promoting biomolecules with medical, nutraceutical and food applications. *Crit. Rev. Biotech.* **2012**, *32*, 289–326. <https://doi.org/10.3109/07388551.2011.630647>.

19. Hasanuzzaman, M.; Nahar, K.; Alam, M.; Bhowmik, P.C.; Hossain, M.; Rahman, M.M.; Prasad, M.N.V.; Ozturk, M.; Fujita, M. Potential use of halophytes to remediate saline soils. *BioMed Res. Int.* **2014**, *2014*, 589341. <https://doi.org/10.1155/2014/589341>.
20. Ksouri, R.; Megdiche, W.; Falleh, H.; Trabelsi, N.; Boulaaba, M.; Smaoui, A.; Abdelly, C. Influence of biological, environmental and technical factors on phenolic content and antioxidant activities of Tunisian halophytes. *Comptes Rendus Biol.* **2008**, *331*, 865–873. <https://doi.org/10.1016/j.crv.2008.07.024>.
21. Shahidi, F.; Ambigaipalan, P. Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects—A review. *J. Funct. Foods* **2015**, *18*, 820–897. <https://doi.org/10.1016/j.jff.2015.06.018>.
22. Arya, S.S.; Devi, S.; Ram, K.; Kumar, S.; Kumar, N.; Mann, A.; Kumar, A.; Chand, G. Halophytes: The plants of therapeutic medicine. In *Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes*; Springer: Singapore, 2019; pp. 271–287.
23. eHALOPH—Halophytes Database© Version 3.20. The University of Sussex; 2020. Available online: <https://www.sussex.ac.uk/affiliates/halophytes> (accessed on 4 November 2020).
24. Suffness, M.; Pezzuto, J.M. Assays related to cancer drug discovery. In *Methods in Plant Biochemistry: Assays for Bioactivity*; Hostettmann, K., Ed.; Academic Press: London, UK, 1990; pp. 71–133.
25. Cruz, S.; Gomes, S.E.; Borralho, P.M.; Rodrigues, C.M.P.; Gaudêncio, S.P.; Pereira, F. In Silico HCT116 Human Colon Cancer Cell-Based Models En Route to the Discovery of Lead-Like Anticancer Drugs. *Biomolecules* **2018**, *8*, 56. <https://doi.org/10.3390/biom8030056>.
26. Renna, M.; Gonnella, M. Ethnobotany, nutritional traits, and healthy properties of some halophytes used as greens in the Mediterranean basin. In *Handbook of Halophytes: From Molecules to Ecosystems towards Biosaline Agriculture*; Grigore, M.N., Eds.; Springer: Cham, Switzerland, 2020; pp. 1–19.
27. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/62332> (accessed on 18 October 2022).
28. Xu, Z.; Chang, L. Acanthaceae. In *Identification and Control of Common Weeds*; Springer: Singapore, 2017; Volume 3, pp. 329–338.
29. Khan, I.; Jan, S.A.; Shinwari, Z.K.; Ali, M.; Khan, Y.; Kumar, T. Ethnobotany and medicinal uses of folklore medicinal plants belonging to family Acanthaceae: An updated review. *MOJ Biol. Med.* **2017**, *1*, 34–38. <https://doi.org/10.15406/mojbm.2017.01.00009>.
30. Velmani, S.; Perumal, B.; Santhosh, C.; Vetrivel, C.; Maruthupandian, A. Phytochemical and traditional uses on *Acanthus ilicifolius* (L.). *J. Adv. Appl. Scient. Res.* **2016**, *1*, 43–48. <https://doi.org/10.46947/joaasr13201617>.
31. Rajamanickam, E.; Gurudeeban, S.; Satyavani, K.; Ramathan, T. In vitro anticancer effect of *Acanthus ilicifolius* on hepatocellular carcinoma cells. *Asian J. Biochem.* **2014**, *9*, 179–186.
32. Paul, T.; Ramasubbu, S. The antioxidant, anticancer and anticoagulant activities of *Acanthus ilicifolius* L. roots and *Lumnitzera racemosa* Willd. leaves, from southeast coast of India. *J. Appl. Pharm. Sci.* **2017**, *7*, 81–87. <https://doi.org/10.7324/JAPS.2017.70313>.
33. Smitha, R.B.; Madhusoodanan, P.V.; Prakashkumar, R. Anticancer activity of *Acanthus ilicifolius* linn. from Chettuva mangroves, Kerala, India. *Int. J. Bioassays* **2014**, *3*, 3452–3455.
34. Khamwut, A.; Jevapatarakul, D.; Reamtong, O. In vitro evaluation of anti-epidermoid cancer activity of *Acanthus ebracteatus* protein hydrolysate and their effects on apoptosis and cellular proteins. *Oncol. Lett.* **2019**, *18*, 3128–3136. <https://doi.org/10.3892/ol.2019.10647>.
35. Thatoi, H.; Samantaray, D.; Das, S.K. The genus *Avicennia*, a pioneer group of dominant mangrove plant species with potential medicinal values: A review. *Front. Life Sci.* **2016**, *9*, 267–291. <https://doi.org/10.1080/21553769.2016.1235619>.
36. Eswaraiah, G.; Peele, K.A.; Krupanidhi, S.; Kumar, R.B.; Venkateswarulu, T.C. Identification of bioactive compounds in leaf extract of *Avicennia alba* by GC-MS analysis and evaluation of its in-vitro anticancer potential against MCF7 and HeLa cell lines. *J. King Saud. Univ. Sci.* **2020**, *32*, 740–744. <https://doi.org/10.1016/j.jksus.2018.12.010>.
37. Illian, D.N.; Hasibuan, P.A.Z.; Sumardi, S.; Nuryawan, A.; Wati, R.; Basyuni, M. Anticancer activity of polyisoprenoids from *Avicennia alba* Blume. in WiDr cells. *Iran J. Pharm. Res.* **2019**, *18*, 1477–1487. <https://doi.org/10.22037/ijpr.2019.1100719>.
38. Huang, C.; Lu, C.K.; Tu, M.C.; Chang, J.H.; Chen, Y.J.; Tu, Y.H.; Huang, H.C. Polyphenol-rich *Avicennia marina* leaf extracts induce apoptosis in human breast and liver cancer cells and in a nude mouse xenograft model. *Oncotarget* **2016**, *7*, 35874–35893. <https://doi.org/10.18632/oncotarget.8624>.
39. Esau, L.; Sagar, S.; Bajic, V.B.; Kaur, M. Autophagy inhibition enhances the mitochondrial-mediated apoptosis induced by mangrove (*Avicennia marina*) extract in human breast cancer cells. *Eur. J. Med. Plants* **2015**, *5*, 304–317. <https://doi.org/10.9734/EJMP/2015/14181>.
40. Devi, A.S.; Joseph, J.; Bhuvaneshwari, V. Antioxidant, anticancerous and cytotoxic effects of *Avicennia marina*. *Int. J. Res. Pharm. Sci.* **2018**, *9*, 996–1001. <https://doi.org/10.26452/ijrps.v9i3.1621>.
41. Catalogue of life. Available online: <https://www.catalogueoflife.org/data/taxon/65P> (accessed on 18 October 2022).
42. Klak, C. Aizoaceae. *Bothalia* **2011**, *41*, 292–294.
43. El-Amier, Y.A.; Haroun, S.A.; El-Shehaby, O.A.; Al-hadithy, O.N. Antioxidant and antimicrobial properties of some wild Aizoaceae species growing in Egyptian desert. *J. Environ. Sci.* **2016**, *45*, 1–10.
44. Seo, J.A.; Ju, J. Antioxidant and growth inhibitory activities of *Mesembryanthemum crystallinum* L. in HCT116 human colon cancer cells. *J. Nutr. Health* **2019**, *52*, 157–167.

45. Chintalapani, S.; Swathi, M.S.; Narasu, M.L. Antiproliferative and apoptosis inducing potential of whole plant extracts of *Sesuvium portulacastrum* against different cancer cell lines. *J. Appl. Pharm. Sci.* **2019**, *9*, 38–43. <https://doi.org/10.7324/JAPS.2019.90205>.
46. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/68Z> (accessed on 18 October 2022).
47. Xu, Z.; Deng, M. *Amaranthaceae*. In *Identification and Control of Common Weeds*; Springer: Dordrecht, The Netherlands, 2017. pp. 245–302.
48. Mroczek, A. Phytochemistry and bioactivity of triterpene saponins from *Amaranthaceae* family. *Phytochem. Rev.* **2015**, *14*, 577–605. <https://doi.org/10.1007/s11101-015-9394-4>.
49. Nisar, F.; Gul, B.; Khan, M.A.; Hameed, A. Germination and recovery responses of heteromorphic seeds of two co-occurring *Arthrocnemum* species to salinity, temperature and light. *S. Afr. J. Bot.* **2019**, *121*, 143–151. <https://doi.org/10.1016/j.sajb.2018.10.035>.
50. ElNaker, N.A.; Yousef, A.F.; Yousef, L.F. A review of *Arthrocnemum (Arthrocaulon) macrostachyum* chemical content and bioactivity. *Phytochem. Rev.* **2020**, *19*, 1427–1448. <https://doi.org/10.1007/s11101-020-09686-5>.
51. Boulaaba, M.; Mkadmini, K.; Tsolmon, S.; Han, J.; Smaoui, A.; Kawada, K.; Ksouri, R.; Isoda, H.; Abdelly, C. In vitro antiproliferative effect of *Arthrocnemum indicum* extracts on Caco-2 cancer cells through cell cycle control and related phenol LC-TOF-MS identification. *Evid.-Based Compl. Altern. Med.* **2013**, *2013*, 529375. <https://doi.org/10.1155/2013/529375>.
52. Yang, Y.; Zhou, X.; Xiao, M.; Hong, Z.; Gong, Q.; Jiang, L.; Zhou, J. Discovery of chrysoeriol, a PI3K-AKT-mTOR pathway inhibitor with potent antitumor activity against human multiple myeloma cells in vitro. *J. Huazhong Univ. Sci. Technol. Med. Sci.* **2010**, *30*, 734–740. <https://doi.org/10.1007/s11596-010-0649-4>.
53. Benzarti, M.; Ben Rejeb, K.; Debez, A.; Abdelly, C. Environmental and economical opportunities for the valorisation of the genus *Atriplex*: New insights. In *Crop Improvement*; Springer: Boston, MA, USA, 2013; pp. 441–457.
54. Al-Senosi, N.K.; Abou-Eisha, A.; Ahmad, E.S. In vitro antiproliferation effect of *Atriplex halimus* L. crude extract on human cell lines by induction of apoptosis and G2/M phase arrest. *Egypt Acad. J. Biol. Sci. C Phys. Mol. Biol.* **2018**, *10*, 115–126. <https://doi.org/10.21608/eajbsc.2018.13660>.
55. Petropoulos, S.A.; Karkanis, A.; Martins, N.; Ferreira, I.C. Halophytic herbs of the Mediterranean basin: An alternative approach to health. *Food Chem. Toxicol.* **2018**, *114*, 155–169. <https://doi.org/10.1016/j.fct.2018.02.031>.
56. Al-Hazzani, A.A.; Alshatwi, A.A. Catechin hydrate inhibits proliferation and mediates apoptosis of SiHa human cervical cancer cells. *Food Chem. Toxicol.* **2011**, *49*, 3281–3286. <https://doi.org/10.1016/j.fct.2011.09.023>.
57. Di Domenico, F.; Foppoli, C.; Coccia, R.; Perluigi, M. Antioxidants in cervical cancer: Chemopreventive and chemotherapeutic effects of polyphenols. *Biochim. Biophys. Acta (BBA)—Mol. Basis Dis.* **2012**, *1822*, 737–747. <https://doi.org/10.1016/j.bbadis.2011.10.005>.
58. Roleira, F.M.; Tavares-da-Silva, E.J.; Varela, C.L.; Costa, S.C.; Silva, T.; Garrido, J.; Borges, F. Plant derived and dietary phenolic antioxidants: Anticancer properties. *Food Chem.* **2015**, *183*, 235–258. <https://doi.org/10.1016/j.foodchem.2015.03.039>.
59. Boeing, J.S.; Barizão, É.O.; Rotta, E.M.; Volpato, H.; Nakamura, C.V.; Maldaner, L.; Visentainer, J.V. Phenolic compounds from *Butia odorata* (Barb. Rodr.) Noblick fruit and its antioxidant and antitumor activities. *Food Anal Methods* **2020**, *13*, 61–68. <https://doi.org/10.1007/s12161-019-01515-6>.
60. Rha, C.S.; Jeong, H.W.; Park, S.; Lee, S.; Jung, Y.S.; Kim, D.O. Antioxidative, anti-inflammatory, and anticancer effects of purified flavonol glycosides and aglycones in green tea. *Antioxidants* **2019**, *8*, 278. <https://doi.org/10.3390/antiox8080278>.
61. Kokanova-Nedialkova, Z.; Nedialkov, P.; Nikolov, S. The genus *Chenopodium*: Phytochemistry, ethnopharmacology and pharmacology. *Pharmacogn. Rev.* **2009**, *3*, 280.
62. Choudhary, N.; Prabhu, K.S.; Prasad, S.B.; Singh, A.; Agarhari, U.C.; Suttee, A. Phytochemistry and pharmacological exploration of *Chenopodium album*: Current and future perspectives. *Res. J. Pharm. Technol.* **2020**, *13*, 3933–3940. <https://doi.org/10.5958/0974-360X.2020.00696.4>.
63. Zhao, T.; Pan, H.; Feng, Y.; Li, H.; Zhao, Y. Petroleum ether extract of *Chenopodium album* L. prevents cell growth and induces apoptosis of human lung cancer cells. *Exp. Ther. Med.* **2016**, *12*, 3301–3307. <https://doi.org/10.3892/etm.2016.3765>.
64. Jia-liang, W.; Dan-wei, M.; Ya-nan, W.; Hong, Z.; Bing, H.; Qun, L.; Zhi-yan, Z.; Jing, F. Cytotoxicity of essential oil of *Chenopodium ambrosioides* L. against human breast cancer MCF-7 cells. *Trop. J. Pharm. Res.* **2013**, *12*, 929–933. <https://doi.org/10.4314/tjpr.v12i6.10>.
65. Degenhardt, R.T.; Farias, I.V.; Grassi, L.T.; Franchi Jr, G.C.; Nowill, A.E.; Bittencourt, C.M.D.S.; Wagner, T.M.; Bella Cruz, A.; Malheiros, A. Characterization and evaluation of the cytotoxic potential of the essential oil of *Chenopodium ambrosioides*. *Rev. Bras. Farmacogn.* **2016**, *26*, 56–61. <https://doi.org/10.1016/j.bjpr.2015.08.012>.
66. Hu, Y.; Zhang, J.; Zou, L.; Fu, C.; Li, P.; Zhao, G. Chemical characterization, antioxidant, immune-regulating and anticancer activities of a novel bioactive polysaccharide from *Chenopodium quinoa* seeds. *Inter. J. Biol. Macromol.* **2017**, *99*, 622–629. <https://doi.org/10.1016/j.ijbiomac.2017.03.019>.
67. Patel, S. *Salicornia*: Evaluating the halophytic extremophile as a food and a pharmaceutical candidate. *3Biotech* **2016**, *6*, 104. <https://doi.org/10.1007/s13205-016-0418-6>.

68. Samuel, P.; Kumar, V.J.; Dhayalan, D.R.; Amirtharaj, K.; Sudarmani, D.N.P. Bioprospecting of *Salicornia europaea* L. a marine halophyte and evaluation of its biological potential with special reference to anticancer activity. *J. Pharma. Pharma. Sci.* **2017**, *2*, 138.
69. Kensil, C.R. Saponins as vaccine adjuvants. *Crit. Rev. Ther. Drug. Carr. Syst.* **1996**, *13*, 1–55.
70. Yurdakok, B.; Baydan, E. Cytotoxic effects of *Eryngium kotschyi* and *Eryngium maritimum* on Hep2, HepG2, Vero and U138 MG cell lines. *Pharma. Biol.* **2013**, *51*, 1579–1585. <https://doi.org/10.3109/13880209.2013.803208>.
71. Yildirim, I.; Kutlu, T. Anticancer agents: Saponin and tannin. *Int. J. Biol. Chem.* **2015**, *9*, 332–340.
72. Munir, U.; Perveen, A.; Qamarunnisa, S. Comparative pharmacognostic evaluation of some species of the genera *Suaeda* and *Salsola* leaf (Chenopodiaceae). *Pak. J. Pharm. Sci.* **2014**, *27*, 1309–1315.
73. Altay, V.; Ozturk, M. The Genera *Salsola* and *Suaeda* (*Amaranthaceae*) and their value as fodder. In *Handbook of Halophytes; From molecules to ecosystems towards biosaline agriculture*; Springer: Cham, Switzerland, 2020; pp. 1–12.
74. Saleh, K.A.; Albinhassan, T.H.; Al-Ghazzawi, A.M.; Mohaya, A.; Shati, A.A.; Ayoub, H.J.; Abdallah, Q.M. Anticancer property of hexane extract of *Suaeda fruticosa* plant leaves against different cancer cell lines. *Trop. J. Pharm. Res.* **2020**, *19*, 129–136. <https://doi.org/10.4314/tjpr.v19i1.20>.
75. Su, P.; Veeraraghavan, V.P.; Krishna Mohan, S.; Lu, W. A ginger derivative, zingerone-a phenolic compound-induces ROS-mediated apoptosis in colon cancer cells (HCT-116). *J. Biochem. Mol. Toxicol.* **2019**, *33*, e22403. <https://doi.org/10.1002/jbt.22403>.
76. Sobral, M.V.; Xavier, A.L.; Lima, T.C.; de Sousa, D.P. Antitumor activity of monoterpenes found in essential oils. *Sci. World. J.* **2014**, *2014*, 953451. <https://doi.org/10.1155/2014/953451>.
77. Garcia, C.; Teodósio, C.; Oliveira, C.; Oliveira, C.; Díaz-Lanza, A.; Reis, C.; Duarte, N.; Rijo, P. Naturally occurring Plectranthus-derived diterpenes with antitumoral activities. *Curr. Pharm. Des.* **2018**, *24*, 4207–4236. <https://doi.org/10.2174/1381612825666190115144241>.
78. Ahmad, S.; Ullah, F.; Zeb, A.; Ayaz, M.; Ullah, F.; Sadiq, A. Evaluation of *Rumex hastatus* D. Don for cytotoxic potential against HeLa and NIH/3T3 cell lines: Chemical characterization of chloroform fraction and identification of bioactive compounds. *BMC Compl. Altern. Med.* **2016**, *16*, 308. <https://doi.org/10.1186/s12906-016-1302-y>.
79. Gupta, E.; Kaushik, S.; Purwar, S.; Sharma, R.; Balapure, A.K.; Sundaram, S. Anticancer potential of steviol in MCF-7 human breast cancer cells. *Pharmacogn. Mag.* **2017**, *13*, 345–350. [https://doi.org/10.4103/pm.pm\\_29\\_17](https://doi.org/10.4103/pm.pm_29_17).
80. Oueslati, S.; Ksouri, R.; Falleh, H.; Pichette, A.; Abdelly, C.; Legault, J. Phenolic content, antioxidant, anti-inflammatory and anticancer activities of the edible halophyte *Suaeda fruticosa* Forssk. *Food Chem.* **2012**, *132*, 943–947. <https://doi.org/10.1016/j.foodchem.2011.11.072>.
81. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/6KC> (accessed on 18 October 2022).
82. Chen, Y.; Peng, S.; Luo, Q.; Zhang, J.; Guo, Q.; Zhang, Y.; Chai, X. Chemical and pharmacological progress on polyacetylenes isolated from the family Apiaceae. *Chem. Biodiv.* **2015**, *12*, 474–502. <https://doi.org/10.1002/cbdv.201300396>.
83. Wang, P.; Su, Z.; Yuan, W.; Deng, G.; Li, S. Phytochemical constituents and pharmacological activities of *Eryngium* L. (Apiaceae). *Pharm. Crop.* **2012**, *3*, 99–120.
84. Renna, M.; Gonnella, M.; Caretto, S.; Mita, G.; Serio, F. Sea fennel (*Crithmum maritimum* L.): From underutilized crop to new dried product for food use. *Gen. Res. Crop. Evol.* **2017**, *64*, 205–216. <https://doi.org/10.1007/s10722-016-0472-2>.
85. Gnocchi, D.; Cesari, G.; Calabrese, G.J.; Capone, R.; Sabbà, C.; Mazzocca, A. Inhibition of hepatocellular carcinoma growth by ethyl acetate extracts of apulian *Brassica oleracea* L. and *Crithmum maritimum* L. *Plant Foods Hum. Nutr.* **2020**, *75*, 33–40. <https://doi.org/10.1007/s11130-019-00781-3>.
86. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/622TP> (accessed on 18 October 2022).
87. Gao, T.; Yao, H.; Song, J.; Zhu, Y.; Liu, C.; Chen, S. Evaluating the feasibility of using candidate DNA barcodes in discriminating species of the large Asteraceae family. *BMC Evol. Biol.* **2010**, *10*, 324. <https://doi.org/10.1186/1471-2148-10-324>.
88. Shahani, S.; Hamzekanlu, N.; Zakeri, N.; Hosseinimehr, S.J. Synergistic effect of *Achillea millefolium* L. combined with bleomycin on prostate cancer cell. *Res. Mol. Med.* **2015**, *3*, 12–17. <https://doi.org/10.7508/rmm.2015.01.003>.
89. Li, Y.; Zhang, M.L.; Cong, B.; Wang, S.M.; Dong, M.; Sauriol, F.; Huo, C.-H.; Shi, Q.-W.; Gu, Y.-G.; Kiyota, H. Achillinin A, a cytotoxic guaianolide from the flower of Yarrow, *Achillea millefolium*. *Biosci. Biotechnol. Biochem.* **2011**, *75*, 1554–1556. <https://doi.org/10.1271/bbb.110234>.
90. Abou Baker, D.H. *Achillea millefolium* L. ethyl acetate fraction induces apoptosis and cell cycle arrest in human cervical cancer (HeLa) cells. *Ann. Agric. Sci.* **2020**, *65*, 42–48. <https://doi.org/10.1016/j.aos.2020.03.003>.
91. Anantharaju, P.G.; Gowda, P.C.; Vimalambike, M.G.; Madhunapantula, S.V. An overview on the role of dietary phenolics for the treatment of cancers. *Nutr. J.* **2016**, *15*, 99. <https://doi.org/10.1186/s12937-016-0217-2>.
92. Adorisio, S.; Giamperi, L.; Bucchini, A.E.A.; Delfino, D.V.; Marcotullio, M.C. Bioassay-guided isolation of antiproliferative compounds from *Limbaria crithmoides* (L.) Dumort. *Molecules* **2020**, *25*, 1893. <https://doi.org/10.3390/molecules25081893>.
93. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/7DN> (accessed on 18 October 2022).
94. Anjum, N.A.; Gill, S.S.; Ahmad, I.; Pacheco, M.; Duarte, A.C.; Umar, S.; Khan, N.A.; Pereira, M.E. The plant family Brassicaceae: An introduction. In *Plant Family Brassicaceae*; Springer: Dordrecht, The Netherlands, 2012; pp. 1–33.



95. Arbelet-Bonnin, D.; Ben-Hamed-Louati, I.; Laurenti, P.; Abdelly, C.; Ben-Hamed, K.; Bouteau, F. *Cakile maritima*, a promising model for halophyte studies and a putative cash crop for saline agriculture. In *Advances in Agronomy*; Academic Press: Cambridge, MA, USA, 2019; Volume 155, pp. 45–78. <https://doi.org/10.1016/bs.agron.2019.01.003>.
96. Omer, E.; Elshamy, A.; Taher, R.; El-Kashak, W.; Shalom, J.; White, A.; Cock, I. *Cakile maritima* Scop. extracts inhibit CaCo2 and HeLa human carcinoma cell growth: GC-MS analysis of an anti-proliferative extract. *Pharm. J.* **2019**, *11*, 258–266. <https://doi.org/10.5530/pj.2019.11.40>.
97. Hafidh, R.R.; Hussein, S.Z.; MalAllah, M.Q.; Abdulmir, A.S.; Abu Bakar, F. A high-throughput quantitative expression analysis of cancer-related genes in human HepG2 cells in response to limonene, a potential anticancer agent. *Curr. Cancer Drug Targets* **2018**, *18*, 807–815. <https://doi.org/10.2174/1568009617666171114144236>.
98. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/8J9> (accessed on 18 October 2022).
99. Sultana, R.; Rahman, A.H.M.M. Convolvulaceae: A taxonomically and medicinally important morning glory family. *Int. J. Bot. Stud.* **2016**, *1*, 47–52.
100. Lee, J.I.; Kim, I.H.; Nam, T.J. Crude extract and solvent fractions of *Calystegia soldanella* induce G1 and S phase arrest of the cell cycle in HepG2 cells. *Int. J. Oncol.* **2017**, *50*, 414–420. <https://doi.org/10.3892/ijo.2017.3836>.
101. Min, H.-Y.; Park, H.-J.; Kim, Y.; Lee, E.-J.; Hwang, H.-J.; Park, E.-J.; Lee, S.K. Cytotoxic activities of indigenous plant extracts in cultured cancer cells. *Nat. Prod. Sci.* **2002**, *8*, 170–172.
102. Pouraminaei, M.; Mirzaiey, M.R.; Khoshdel, A.; Hajizadeh, M.R.; Mahmoodi, M.; Fahmidehkar, M.A. The effect of *Cressa cretica* hydroalcoholic extract on apoptosis and the expression of Bcl2, Bax and P53 genes in hepatoma cell line HepG2. *Gene Rep.* **2020**, *20*, 100692. <https://doi.org/10.1016/j.genrep.2020.100692>.
103. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/8VQ> (accessed on 18 October 2022).
104. Petersen, G.; Seberg, O.; Short, F.T.; Fortes, M.D. Complete genomic congruence but non-monophyly of *Cymodocea* (Cymodoceaceae), a small group of seagrasses. *Taxon* **2014**, *63*, 3–8. <https://doi.org/10.12705/631.2>.
105. Abraham, J.; Saraf, S.; Mustafa, V.; Chaudhary, Y.; Sivanangam, S. Synthesis and evaluation of silver nanoparticles using *Cymodocea rotundata* against clinical pathogens and human osteosarcoma cell line. *J. Appl. Pharm. Sci.* **2017**, *7*, 55–61. <https://doi.org/10.7324/JAPS.2017.70608>.
106. Palaniappan, P.; Sathishkumar, G.; Sankar, R. Fabrication of nano-silver particles using *Cymodocea serrulata* and its cytotoxicity effect against human lung cancer A549 cells line. *Spectroch. Acta Part A Mol. Biomol. Spec.* **2015**, *138*, 885–890. <https://doi.org/10.1016/j.saa.2014.10.072>.
107. Kalpana, C.S.; Kalaivani, P.; Vanitha, V. Phytochemical screening and effect of *Cymodocea serrulata* on HepG2-human hepatocellular carcinoma cell line. *Inter. J. Res. Pharm. Sci.* **2020**, *11*, 1684–1691. <https://doi.org/10.26452/ijrps.v11i2.2058>.
108. Sofi, M.S.; Nabi, S.; Mohammed, C.; Sofi, S. The role of phytochemicals in cancer treatment: A current review. *J. Med. Plant Stud.* **2018**, *6*, 83–93.
109. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/8W5> (accessed on 18 October 2022).
110. Semmouri, I.; Bauters, K.; Léveillé-Bourret, É.; Starr, J.R.; Goetghebeur, P.; Larridon, I. Phylogeny and systematics of Cyperaceae, the evolution and importance of embryo morphology. *Bot. Rev.* **2019**, *85*, 1–39. <https://doi.org/10.1007/s12229-018-9202-0>.
111. Al-Snafi, A.E. A review on *Cyperus rotundus*: A potential medicinal plant. *IOSR J. Pharm.* **2016**, *6*, 32–48.
112. Park, S.E.; Shin, W.T.; Park, C.; Hong, S.H.; Kim, G.Y.; Kim, S.O.; Ryu, C.H.; Hong, S.H.; Choi, Y.H. Induction of apoptosis in MDA-MB-231 human breast carcinoma cells with an ethanol extract of *Cyperus rotundus* L. by activating caspases. *Oncol. Rep.* **2014**, *32*, 2461–2470. <https://doi.org/10.3892/or.2014.3507>.
113. Singh, N.; Pandey, B.R.; Verma, P.; Bhalla, M.; Gilca, M. Phyto-pharmacotherapeutics of *Cyperus rotundus* Linn. (Motha): An overview. *Ind. J. Nat. Prod. Res.* **2012**, *3*, 467–476.
114. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/623QT> (accessed on 18 October 2022).
115. Ahmad, F.; Anwar, F.; Hira, S. Review on medicinal importance of Fabaceae family. *Pharm. Online* **2016**, *3*, 151–157.
116. Muhammad, G.; Hussain, M.A.; Anwar, F.; Ashraf, M.; Gilani, A.H. Alhagi: A plant genus rich in bioactives for pharmaceuticals. *Phyto. Res.* **2015**, *29*, 1–13. <https://doi.org/10.1002/ptr.5222>.
117. Laghari, A.H.; Memon, S.; Nelofar, A.; Khan, K.M.; Yasmin, A.; Syed, M.N.; Aman, A. A new flavanone with urease-inhibition activity isolated from roots of manna plant camelthorn (*Alhagi maurorum*). *J. Mol. Struct.* **2010**, *965*, 65–67. <https://doi.org/10.1016/j.molstruc.2009.11.039>.
118. Behbahani, M. Evaluation of in vitro anticancer activity of *Ocimum basilicum*, *Alhagi maurorum*, *Calendula officinalis* and their parasite *Cuscuta campestris*. *PLoS ONE* **2014**, *9*, e116049. <https://doi.org/10.1371/journal.pone.0116049>.
119. Farag, M.A.; Porzel, A.; Wessjohann, L.A. Unequivocal glycyrrhizin isomer determination and comparative in vitro bioactivities of root extracts in four Glycyrrhiza species. *J. Adv. Res.* **2015**, *6*, 99–104. <https://doi.org/10.1016/j.jare.2014.05.001>.
120. Al-Snafi, A.E. *Glycyrrhiza glabra*: A phytochemical and pharmacological review. *IOSR J. Pharm.* **2018**, *8*, 1–17.
121. Farooqui, A.; Khan, F.; Khan, I.; Ansari, I.A. Glycyrrhizin induces reactive oxygen species-dependent apoptosis and cell cycle arrest at G0/G1 in HPV18+ human cervical cancer HeLa cell line. *Biomed Pharm.* **2018**, *97*, 752–764. <https://doi.org/10.1016/j.biopha.2017.10.147>.

122. Basar, N.; Oridupa, O.A.; Ritchie, K.J.; Nahar, L.; Osman, N.M.M.; Stafford, A.; Kushiev, H.; Kan, A.; Sarker, S.D. Comparative cytotoxicity of *Glycyrrhiza glabra* roots from different geographical origins against immortal human keratinocyte (HaCaT), lung adenocarcinoma (A549) and liver carcinoma (HepG2) cells. *Phyto. Res.* **2015**, *29*, 944–948. <https://doi.org/10.1002/ptr.5329>.
123. Nourazarian, S.M.; Nourazarian, A.; Majidinia, M.; Roshaniasl, E. Effect of root extracts of medicinal herb *Glycyrrhiza glabra* on HSP90 gene expression and apoptosis in the HT-29 colon cancer cell line. *Asian Pac. J. Cancer Prev.* **2016**, *16*, 8563–8566.
124. Gao, C.; Peng, Y.N.; Wang, H.Z.; Fang, S.L.; Zhang, M.; Zhao, Q.; Liu, J. Inhibition of heat shock protein 90 as a novel platform for the treatment of cancer. *Curr. Pharm. Des.* **2019**, *25*, 849–855. <https://doi.org/10.2174/1381612825666190503145944>.
125. Fan, H.; Xuan, J.; Zhang, K.; Jiang, J. Anticancer component identification from the extract of *Dysosma versipellis* and *Glycyrrhiza uralensis* based on support vector regression and mean impact value. *Anal. Methods* **2018**, *10*, 371–380. <https://doi.org/10.1039/C7AY02465G>.
126. Wu, F.; Ma, J.; Meng, Y.; Zhang, D.; Pascal Muvunyi, B.; Luo, K.; Di, H.; Guo, W.; Wang, Y.; Feng, B.; et al. Potential DNA barcodes for *Melilotus* species based on five single loci and their combinations. *PLoS ONE* **2017**, *12*, e0182693. <https://doi.org/10.1371/journal.pone.0182693>.
127. Ahmed, S.A.K.; Al-Refai, M. Chemical constituents and cytotoxic activities of the extracts of *Melilotus indicus*. *Eur. J. Chem.* **2014**, *5*, 503–506. <https://doi.org/10.5155/eurjchem.5.3.503-506.1070>.
128. El-Hafeez, A.A.A.; Khalifa, H.O.; Elgawish, R.A.; Shouman, S.A.; El-Twab, M.H.A.; Kawamoto, S. *Melilotus indicus* extract induces apoptosis in hepatocellular carcinoma cells via a mechanism involving mitochondria-mediated pathways. *Cytotechnology* **2018**, *70*, 831–842. <https://doi.org/10.1007/s10616-018-0195-7>.
129. Henciya, S.; Seturaman, P.; James, A.R.; Tsai, Y.H.; Nikam, R.; Wu, Y.C.; Dahms, H.-U.; Chang, F.R. Biopharmaceutical potentials of *Prosopis* spp. (Mimosaceae, Leguminosae). *J. Food Drug Anal.* **2017**, *25*, 187–196. <https://doi.org/10.1016/j.jfda.2016.11.001>.
130. Sathiyai, M.; Muthuchelian, K. Anti-tumor potential of total alkaloid extract of *Prosopis juliflora* DC. leaves against Molt-4 cells in vitro. *Afr. J. Biotech.* **2011**, *10*, 8881–8888. <https://doi.org/10.5897/AJB10.875>.
131. Bunma, S.; Balslev, H. A review of the economic botany of *Sesbania* (Leguminosae). *Bot. Rev.* **2019**, *85*, 185–251. <https://doi.org/10.1007/s12229-019-09205-y>.
132. Boddawar, G.D.; Dhawale, S.C.; Shaikh, S.S. Assessment of anti-inflammatory potential of *Sesbania bispinosa* Linn. leaf extracts and fractions by acute and chronic models. *Alex. J. Med.* **2016**, *52*, 289–293. <https://doi.org/10.1016/j.ajme.2015.10.004>.
133. Kapoor, B.B.S.; Purohit, V. Sterol contents from some fabaceous medicinal plants of Rajasthan desert. *Indian J. Pharm. Biol. Res.* **2013**, *1*, 13–15.
134. Tamboli, S.A. Anti-inflammatory activity of *Sesbania grandiflora*. *Ind. Drug.* **1996**, *33*, 504–506.
135. Thamboli, S.A.; Pal, S.C.; Kasture, S.B. Analgesic and antipyretic activity of flowers of *Sesbania grandiflora*. *Ind. Drugs-Bombay* **2000**, *37*, 95–98.
136. Laladhas, K.P.; Cheriyan, V.T.; Puliappadamba, V.T.; Bava, S.V.; Unnithan, R.G.; Vijayammal, P.L.; Anto, R.J. A novel protein fraction from *Sesbania grandiflora* shows potential anticancer and chemopreventive efficacy, in vitro and in vivo. *J. Cell Mol. Med.* **2010**, *14*, 636–646. <https://doi.org/10.1111/j.1582-4934.2008.00648.x>.
137. Ponnaniakamideen, M.; Nagalingam, M.; Vanaja, M.; Malarkodi, C.; Rajeshkumar, S. Anticancer activity of different solvent extracts of *Sesbania grandiflora* against neuroblastoma (imr-32) and colon (ht-29) cell lines. *Eur. J. Biomed. Pharm. Sci.* **2015**, *2*, 509–517.
138. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/BJG> (accessed on 18 October 2022).
139. Bús, C.; Tóth, B.; Stefkó, D.; Hohmann, J.; Vasas, A. Family Juncaceae: Promising source of biologically active natural phenanthrenes. *Phytochem. Rev.* **2018**, *17*, 833–851. <https://doi.org/10.1007/s11101-018-9561-5>.
140. El-Shamy, A.I.; Abdel-Razek, A.F.; Nassar, M.I. Phytochemical review of *Juncus* L. genus (Fam. Juncaceae). *Arab. J. Chem.* **2015**, *8*, 614–623. <https://doi.org/10.1016/j.arabjc.2012.07.007>.
141. Rodrigues, M.J.; Gangadhar, K.N.; Vizetto-Duarte, C.; Wubshet, S.G.; Nyberg, N.T.; Barreira, L.; Varela, J.; Custódio, L. Maritime halophyte species from Southern Portugal as sources of bioactive molecules. *Mar Drugs.* **2014**, *12*, 2228–2244.
142. Rodrigues, M.J.; Vizetto-Duarte, C.; Gangadhar, K.N.; Zengin, G.; Mollica, A.; Varela, J.; Custódio, L. In vitro and in silico approaches to unveil the mechanisms underlying the cytotoxic effect of juncunol on human hepatocarcinoma cells. *Pharm. Rep.* **2018**, *70*, 896–899. <https://doi.org/10.1016/j.pharep.2018.02.001>.
143. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/CDB> (accessed on 18 October 2022).
144. Lindamulage, I.K.; Soysa, P. Evaluation of anticancer properties of a decoction containing *Adenanthera pavonina* L. and *Thespesia populnea* L. *BMC Compl. Altern. Med.* **2016**, *16*, 70. <https://doi.org/10.1186/s12906-016-1053-9>.
145. Gopalakrishnan, A.; Kariyil, B.J.; John, R.; Usha, P.A. Phytochemical evaluation and cytotoxic potential of chloroform soluble fraction of methanol extract of *Thespesia populnea* in human breast cancer cell lines. *Pharm. Mag.* **2019**, *15*, 150. [https://doi.org/10.4103/pm.pm\\_329\\_18](https://doi.org/10.4103/pm.pm_329_18).
146. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/D48> (accessed on 18 October 2022).
147. Vuong, Q.V.; Chalmers, A.C.; Jyoti Bhuyan, D.; Bowyer, M.C.; Scarlett, C.J. Botanical, phytochemical, and anticancer properties of the *Eucalyptus* species. *Chem. Biodiv.* **2015**, *12*, 907–924. <https://doi.org/10.1002/cbdv.201400327>.

148. Hrubik, J.D.; Kaišarević, S.N.; Glišić, B.D.; Jovin, E.Đ.; Mimica-Dukić, N.M.; Kovačević, R.Z. *Myrtus communis* and *Eucalyptus camaldulensis* cytotoxicity on breast cancer cells. *Zb. Matice Srp. Za Prir. Nauk.* **2012**, *123*, 65–73. <https://doi.org/10.2298/ZMSPN1223065H>.
149. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/6262K> (accessed on 18 October 2022).
150. Xu, Z.; Chang, L. Plantaginaceae. In *Identification and Control of Common Weeds*; Springer: Singapore, 2017; pp. 339–374.
151. Bora, M.S.; Sarma, K.P. Phytoremediation of heavy metals/metalloids by native herbaceous macrophytes of wetlands: Current research and perspectives. In *Emerging Issues in the Water Environment during Anthropocene*; Springer: Singapore, 2020; pp. 261–284.
152. Smith, E.; Paethorpe, H.M.; Tomita, Y.; Pei, J.V.; Townsend, A.R.; Price, T.J.; Young, J.P.; Yool, A.J.; Hardingham, J.E. The purified extract from the medicinal plant *Bacopa monnieri*, bacopaside II, inhibits growth of colon cancer cells in vitro by inducing cell cycle arrest and apoptosis. *Cells* **2018**, *7*, 81. <https://doi.org/10.3390/cells7070081>.
153. Koczurkiewicz, P.; Łojewski, M.; Piska, K.; Michalik, M.; Wójcik-Pszczola, K.; Szweczyk, A.; Halaszuk, P.; Pekala, E.; Muszyńska, B. Chemopreventive and anticancer activities of *Bacopa monnieri* extracted from artificial digestive juices. *Nat. Prod. Commun.* **2017**, *12*, 337–342. <https://doi.org/10.1177/1934578X1701200306>.
154. Yamagata, K.; Izawa, Y.; Onodera, D.; Tagami, M. Chlorogenic acid regulates apoptosis and stem cell marker-related gene expression in A549 human lung cancer cells. *Mol. Cell Biochem.* **2018**, *441*, 9–19. <https://doi.org/10.1007/s11010-017-3171-1>.
155. Rezaei-Seresht, H.; Cheshomi, H.; Falanji, F.; Movahedi-Motlagh, F.; Hashemian, M.; Mireskandari, E. Cytotoxic activity of caffeic acid and gallic acid against MCF-7 human breast cancer cells: An in silico and in vitro study. *Avicenna J. Phytomed.* **2019**, *9*, 574. <https://doi.org/10.22038/AJP.2019.13475>.
156. Gheena, S.; Ezhilarasan, D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Hum. Exper. Toxicol.* **2019**, *38*, 694–702. <https://doi.org/10.1177/0960327119839173>.
157. Aithal, M.G.; Rajeswari, N. Bacoside A induced sub-G0 arrest and early apoptosis in human glioblastoma cell line U-87 MG through notch signalling pathway. *Brain Tumor Res. Treat.* **2019**, *7*, 25. <https://doi.org/10.14791/btrt.2019.7.e21>.
158. Goncalves, S.; Romano, A. The medicinal potential of plants from the genus *Plantago* (Plantaginaceae). *Ind. Crop. Prod.* **2016**, *83*, 213–226. <https://doi.org/10.1016/j.indcrop.2015.12.038>.
159. Poor, M.H.S.; Khatami, M.; Azizi, H.; Abazari, Y. Cytotoxic activity of biosynthesized Ag nanoparticles by *Plantago major* towards a human breast cancer cell line. *Rend. Lincei* **2017**, *28*, 693–699. <https://doi.org/10.1007/s12210-017-0641-z>.
160. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/627FM> (accessed on 18 October 2022).
161. Caperta, A.D.; Róis, A.S.; Teixeira, G.; Garcia-Caparros, P.; Flowers, T.J. Secretory structures in plants: Lessons from the *Plumbaginaceae* on their origin, evolution and roles in stress tolerance. *Plant Cell Environ.* **2020**, *43*, 2912–2931. <https://doi.org/10.1111/pce.13825>.
162. Singh, K.; Naidoo, Y.; Baijnath, H. A comprehensive review on the genus *Plumbago* with focus on *Plumbago auriculata* (Plumbaginaceae). *Afr. J. Trad. Compl. Altern. Med.* **2018**, *15*, 199–215. <https://doi.org/10.21010/ajtcam.v15i1.21>.
163. Crespo, M.B.; Lledó, M.D. Two new North African genera related to *Limoniastrum* (Plumbaginaceae). *Bot. J. Linn. Soc.* **2000**, *132*, 165–174. <https://doi.org/10.1111/j.1095-8339.2000.tb01212.x>.
164. Lledó, M.D.; Crespo, M.B.; Cox, A.V.; Fay, M.F.; Chase, M.W. Polyphyly of *Limoniastrum* (Plumbaginaceae): Evidence from DNA sequences of plastid *rbcL*, *trnL* intron and *trnL-F* intergene spacer. *Bot. J. Linn. Soc.* **2000**, *132*, 175–191. <https://doi.org/10.1111/j.1095-8339.2000.tb01213.x>.
165. Ozenda, P. *Flore du Sahara*, 2nd ed.; CNRS: Paris, France, 1983.
166. Dijkema, K.S.; Beeftink, W.G.; Doodly, J.P.; Géhu, J.M.; Heydemann, B.; Rivas Martinez, S. *La Végétation Halophile en Europe (Prés Salés)*; Council of Europe: Strasbourg, France, 1984.
167. Barone, R.; Scholz, S.; Mesa, R. *Limoniastrum monopetalum* (L.) Boiss. (Plumbaginaceae), adición a la flora de Fuerteventura (Islas Canarias). *Bot. Macarones.* **1995**, *21*, 59–60.
168. Ferrer-Gallego, P.P.; Iamónico, D.; Iberite, M.; Laguna, E.; Crespo, M.B. Lectotypification of two names in *Limoniastrum* (Plumbaginaceae). *Taxon* **2014**, *63*, 1342–1346. <https://doi.org/10.12705/636.4>.
169. Krifa, M.; Alhosin, M.; Muller, C.D.; Gies, J.P.; Chekir-Ghedira, L.; Ghedira, K.; Mely, I.; Bronner, C.; Mousli, M. *Limoniastrum guyonianum* aqueous gall extract induces apoptosis in human cervical cancer cells involving p16 INK4A re-expression related to UHRF1 and DNMT1 down-regulation. *J. Exp. Clin. Cancer Res.* **2013**, *32*, 30. <https://doi.org/10.1186/1756-9966-32-30>.
170. Medini, F.; Bourgou, S.; Lalancette, K.; Snoussi, M.; Mkadmini, K.; Coté, I.; Abdelly, C.; Legault, J.; Ksouri, R. Phytochemical analysis, antioxidant, anti-inflammatory, and anticancer activities of the halophyte *Limonium densiflorum* extracts on human cell lines and murine macrophages. *S. Afr. J. Bot.* **2015**, *99*, 158–164. <https://doi.org/10.1016/j.sajb.2015.04.007>.
171. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/627FW> (accessed on 18 October 2022).
172. Hodkinson, T.R. Evolution and taxonomy of the grasses (Poaceae): A model family for the study of species-rich groups. *Ann. Plant. Rev. Online* **2018**, *1*, 255–294. <https://doi.org/10.1002/9781119312994.apr0622>.
173. Ashokkumar, K.; Selvaraj, K.; Muthukrishnan, S.D. *Cynodon dactylon* (L.) Pers.: An updated review of its phytochemistry and pharmacology. *J. Med. Plants Res.* **2013**, *7*, 3477–3483. <https://doi.org/10.5897/JMPR2013.5316x>.

174. Khelifi, D.; Hayouni, E.A.; Valentin, A.; Cazaux, S.; Moukarzel, B.; Hamdi, M.; Bouajila, J. LC-MS analysis, anticancer, antioxidant and antimalarial activities of *Cynodon dactylon* L. extracts. *Ind. Crop. Prod.* **2013**, *45*, 40–247. <https://doi.org/10.1016/j.indcrop.2012.12.030>.
175. El Molla, S.G.; Motaal, A.A.; El Hefnawy, H.; El Fishawy, A. Cytotoxic activity of phenolic constituents from *Echinochloa crus-galli* against four human cancer cell lines. *Rev. Bras. Farmacogn.* **2016**, *26*, 62–67. <https://doi.org/10.1016/j.bjp.2015.07.026>.
176. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/627NK> (accessed on 18 October 2022).
177. Xu, Z.; Deng, M. Portulacaceae. In *Identification and Control of Common Weeds*; Springer, Dordrecht, The Netherlands, 2017; pp. 319–323.
178. Syed, S.; Fatima, N.; Kabeer, G. *Portulaca oleracea* L.: A mini review on phytochemistry and pharmacology. *Int. J. Biol. Biotechnol.* **2016**, *13*, 637–641.
179. Zhou, Y.X.; Xin, H.L.; Rahman, K.; Wang, S.J.; Peng, C.; Zhang, H. *Portulaca oleracea* L.: A review of phytochemistry and pharmacological effects. *BioMed. Res. Int.* **2015**, *2015*, 925631. <https://doi.org/10.1155/2015/925631>.
180. Zheng, G.Y.; Qu, L.P.; Yue, X.Q.; Gu, W.; Zhang, H.; Xin, H.L. Portulacerebroside A induces apoptosis via activation of the mitochondrial death pathway in human liver cancer HCCLM3 cells. *Phytochem. Lett.* **2014**, *7*, 77–84. <https://doi.org/10.1016/j.phytol.2013.10.005>.
181. Farshori, N.N.; Al-Sheddi, E.S.; Al-Oqail, M.M.; Musarrat, J.; Al-Khedhairi, A.A.; Siddiqui, M.A. Cytotoxicity assessments of *Portulaca oleracea* and *Petroselinum sativum* seed extracts on human hepatocellular carcinoma cells (HepG2). *Asian Pac. J. Cancer Prev.* **2014**, *15*, 6633–6638.
182. Al-Sheddi, E.S.; Farshori, N.N.; Al-Oqail, M.M.; Musarrat, J.; Al-Khedhairi, A.A.; Siddiqui, M.A. *Portulaca oleracea* seed oil exerts cytotoxic effects on human liver cancer (HepG2) and human lung cancer (A-549) cell lines. *Asian Pac. J. Cancer Prev.* **2015**, *16*, 3383–3387.
183. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/FPS> (accessed on 18 October 2022).
184. Nebula, M.; Harisankar, H.S.; Chandramohanakumar, N. Metabolites and bioactivities of Rhizophoraceae mangroves. *Nat. Prod. Bioprospect.* **2013**, *3*, 207–232. <https://doi.org/10.1007/s13659-013-0012-0>.
185. Liao, P.C.; Hwang, S.Y.; Huang, S.; Chiang, Y.C.; Wang, J.C. Contrasting demographic patterns of *Ceriops tagal* (Rhizophoraceae) populations in the South China sea. *Austr. J. Bot.* **2011**, *59*, 523–532. <https://doi.org/10.1071/BT10290>.
186. Urashi, C.; Teshima, K.M.; Minobe, S.; Koizumi, O.; Inomata, N. Inferences of evolutionary history of a widely distributed mangrove species, *Bruguiera gymnorrhiza*, in the Indo-West Pacific region. *Ecol. Evol.* **2013**, *3*, 2251–2261. <https://doi.org/10.1002/ece3.624>.
187. Warsinah, E.; Sisindari, A.; Susidarti, R.A. Antiproliferative effect and apoptosis induced in human cell lines by *Bruguiera gymnorrhiza* barks methanol extract. *Indones. J. Cancer Chemoprevent.* **2011**, *2*, 299–303. <https://doi.org/10.14499/indonesianjcanchemoprev2iss3pp299-303>.
188. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/626XM> (accessed on 18 October 2022).
189. Shah, V.V.; Shah, N.D.; Patrekar, P.V. Medicinal plants from Solanaceae family. *Res. J. Pharm. Technol.* **2013**, *6*, 143–151.
190. Tang, W.M.; Chan, E.; Kwok, C.Y.; Lee, Y.K.; Wu, J.H.; Wan, C.W.; Chan, R.Y.K.; Yu, P.H.-F.; Chan, S.W. A review of the anti-cancer and immunomodulatory effects of *Lycium barbarum* fruit. *Inflammopharmacology* **2012**, *20*, 307–314. <https://doi.org/10.1007/s10787-011-0107-3>.
191. Mao, F.; Xiao, B.; Jiang, Z.; Zhao, J.; Huang, X.; Guo, J. Anticancer effect of *Lycium barbarum* polysaccharides on colon cancer cells involves G0/G1 phase arrest. *Med. Oncol.* **2011**, *28*, 121–126. <https://doi.org/10.1007/s12032-009-9415-5>.
192. Zeng, M.; Kong, Q.; Liu, F.; Chen, J.; Sang, H. The anticancer activity of *Lycium barbarum* polysaccharide by inhibiting autophagy in human skin squamous cell carcinoma cells in vitro and in vivo. *Int. J. Polym. Sci.* **2019**, *2019*, 5065920. <https://doi.org/10.1155/2019/5065920>.
193. Zhu, P.F.; Dai, Z.; Wang, B.; Wei, X.; Yu, H.F.; Yan, Z.R.; Zhao, X.D.; Luo, X.D. The anticancer activities phenolic amides from the stem of *Lycium barbarum*. *Nat. Prod. Bioprospecting* **2017**, *7*, 421–431. <https://doi.org/10.1007/s13659-017-0134-x>.
194. Lee, Y.T.; Hsieh, Y.L.; Yeh, Y.H.; Huang, C.Y. Synthesis of phenolic amides and evaluation of their antioxidant and anti-inflammatory activity in vitro and in vivo. *RSC Adv.* **2015**, *5*, 85806–85815. <https://doi.org/10.1039/C5RA14137K>.
195. Catalogue of Life. Available online: <https://www.catalogueoflife.org/data/taxon/GWM> (accessed on 18 October 2022).
196. Sun, L.; Yang, R.; Zhang, B.; Zhang, G.; Wu, X.; Zhang, W.; Zhang, B.; Chen, T.; Liu, G. Phylogenetic relationships among species of *Tamarix* (Tamaricaceae) in China. *Biochem. Syst. Ecol.* **2016**, *69*, 213–221. <https://doi.org/10.1016/j.bse.2016.10.003>.
197. Channa, F.N.; Shinwari, Z.K.; Ali, S.I. Phylogeny of Tamaricaceae using PsbA-trnH nucleotide sequences. *Pak. J. Bot.* **2018**, *50*, 983–987.
198. Bahramsoltani, R.; Kalkhorani, M.; Zaidi, S.M.A.; Farzaei, M.H.; Rahimi, R. The genus *Tamarix*: Traditional uses, phytochemistry, and pharmacology. *J. Ethnophar.* **2020**, *246*, 112245. <https://doi.org/10.1016/j.jep.2019.112245>.
199. Abaza, M.S.; Al-Attiah, R.A.; Bhardwaj, R.; Abbadi, G.; Koyippally, M.; Afzal, M. Syringic acid from *Tamarix aucheriana* possesses antimitogenic and chemo-sensitizing activities in human colorectal cancer cells. *Pharm. Biol.* **2013**, *51*, 1110–1124. <https://doi.org/10.3109/13880209.2013.781194>.

200. Abaza, M.S.I.; Afzal, M.; Al-Attayah, R.A.J.; Guleri, R. Methylferulate from *Tamarix aucheriana* inhibits growth and enhances chemosensitivity of human colorectal cancer cells: Possible mechanism of action. *BMC Comp. Altern. Med.* **2016**, *16*, 384. <https://doi.org/10.1186/s12906-016-1358-8>.
201. KalamUrfi, M.; Mujahid, M.; Badruddeen, J.; Khalid, M.; Khan, M.I.; Usmani, A. *Tamarix gallica*: For traditional uses, phytochemical and pharmacological potentials. *J. Chem. Pharm. Res.* **2016**, *8*, 809–814.
202. Boulabaa, M.; Tsolmon, S.; Ksouri, R. Anticancer effect of *Tamarix gallica* extracts on human colon cancer cells involves Erk1/2 and p38 action on G2/M cell cycle arrest. *Cytotechnology* **2013**, *65*, 927–936. <https://doi.org/10.1007/s10616-013-9564-4>.
203. Momtazi-borojeni, A.; Behbahani, M.; Sadeghi-Aliabadi, H. Antiproliferative activity and apoptosis induction of crude extract and fractions of *Avicennia marina*. *Iran. J. Bas. Med. Sci.* **2013**, *16*, 1203.
204. Yang, X.W.; Dai, Z.; Wang, B.; Liu, Y.P.; Zhao, X.D.; Luo, X. Antitumor triterpenoid saponin from the fruits of *Avicennia marina*. *Nat. Prod. Bioprospecting* **2018**, *8*, 347–353. <https://doi.org/10.1007/s13659-018-0167-9>.
205. Santhanakrishnan, D.; Perumal, R.K.; Kanth, S.V.; Rao, J.R.; Chandrasekaran, B. Antioxidant and cytotoxic effects of methanolic extract of *Salicornia brachiata* L. in HepG2 cells. *Int. J. Res. Pharm. Sci.* **2013**, *4*, 512–517.
206. Sudarshan, S.M.; Chintalapani, S.; Narasu, M.L. Phytochemical analysis and determination of antibacterial and anticancer activities of *Suaeda monoica*. *Asian J. Pharm. Clin. Res.* **2019**, *12*, 91–96.
207. El-Attar, M.M.; Awad, A.A.; Abdel-Tawab, F.M.; Kamel, H.A.; Ahmad, S.A.; Hassan, A.I. Assessment of cytotoxic and anti-cancer activity of *Zygophyllum album* and *Suaeda palaestina* extracts on human liver cancer cell lines. *J. Agric. Sci.* **2019**, *27*, 539–544. <https://doi.org/10.21608/ajs.2019.43663>.
208. Liu, X.; Liu, F.; Zhao, S.; Guo, B.; Ling, P.; Han, G.; Cui, Z. Purification of an acidic polysaccharide from *Suaeda salsa* plant and its anti-tumor activity by activating mitochondrial pathway in MCF-7 cells. *Carbohydr. Polym.* **2019**, *215*, 99–107. <https://doi.org/10.1016/j.carbpol.2019.03.059>.
209. Sumathi, S.; Dharani, B.; Sivaprabha, J.; Raj, S.; Padma, P.R. In vitro growth inhibitory activity of *Prosopis cineraria* leaves in MCF-7 breast cancer cell line. *Int. J. Res. Pharm. Sci.* **2013**, *4*, 40–44.
210. Alsaraf, K.M.; Mohammad, M.H.; Al-Shammari, A.M.; Abbas, I.S. Selective cytotoxic effect of *Plantago lanceolata* L. against breast cancer cells. *J. Egypt Nat. Cancer Instig.* **2019**, *31*, 10. <https://doi.org/10.1186/s43046-019-0010-3>.
211. Venkateswarlu, G.; Rani, T.S.; Vani, M.; Vineela, P.A.J. In-vitro anticancer activity of petroleum ether extract of *Cynodon dactylon*. *J. Pharm. Phytochem.* **2015**, *4*, 164–168.