

The Northern Lakes and Surrounding Plains in the Nile Delta, Egypt: How Are They Now and How Will They Are in Light of Climate Changes

Khaled Abdel-Kader Ouda

Geology Department, Faculty of Science, Assiut University, Assiut, Egypt

ABSTRACT

In addition to the consensus of scientists, environmental experts, fisheries and beach research in Egypt about the pollution of the northern lakes in the Egyptian delta and the collapse of its ecosystem as a result of using these lakes as a permanent drain for agricultural, health and industrial wastewater - untreated - for the delta governorates surrounding the lakes - the continuous drying processes in the Manzala lakes Burullus, Mariout and Idku, which the government undertakes on the one hand, and individuals on the other hand, have caused a change in the morphology of the lakes and a reduction in their areas, ranging between 46% and 83% of their original areas.

Drying and filling works are still going on for these lakes, without taking into account the new climatic conditions that have hit the world, which in turn will lead to a rise in the global sea level (MGSL) during this century by an amount that may reach a maximum of one meter. This will result in the invasion of all the northern lakes of the Nile Delta, in addition to the surrounding dry low plains along the northern coast of the delta, and then the invasion of all the areas that were previously deducted and drained from the lakes with their urban, agricultural and industrial projects, as well as the remaining submerged areas of these lakes, which in turn will lose their characteristics to become part of the Mediterranean. This study aims to determine the quantitative and qualitative size of the certain risks that will face the northern lakes and the plains surrounding them in the Nile delta as a result of the global rise in sea level during this century, the coastal threat sources and the traditional defensive policies that must be followed to avoid all these risks.

How to cite this paper: Khaled Abdel-Kader Ouda "The Northern Lakes and Surrounding Plains in the Nile Delta, Egypt: How Are They Now and How Will They Are in Light of Climate Changes" Published in International Journal of Trend in Scientific Research and Development (ijtsrd), ISSN: 2456-6470, Volume-6 | Issue-6, October 2022, pp.1876-1936, URL: www.ijtsrd.com/papers/ijtsrd52183.pdf



Copyright © 2022 by author(s) and International Journal of Trend in Scientific Research and Development Journal. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0) (<http://creativecommons.org/licenses/by/4.0>)



Attribution License (CC BY 4.0) (<http://creativecommons.org/licenses/by/4.0>)

KEYWORDS: *Climate Change, Nile Delta, Lake Manzala, Lake Burullus, Lake Mariout, Lake Idku*

1. INTRODUCTION

Scientists, environmental experts, fisheries and beach researchers in Egypt have unanimously agreed on the fact of environmental pollution of all the northern lakes in the Egyptian delta, which represents the humid neutral zone that separates the sea from the dry agricultural plains, as a result of using these lakes as a permanent drain for agricultural, health and industrial wastewater - untreated - for the delta governorates surrounding the lakes on the one hand, and increasing the mixing of sea water with lakes on the other hand. This led to the concentration of cadmium, mercury, zinc, lead and copper salts not only in the lake water - but also in the plants that grow in the lakes and the fish that feed on these plants, causing the death of many fish and making the rest of them unfit for

human consumption. The mixing with sea water also led to an increase in the proportion of fresh water in the lakes, which in turn led to the spread of bushes and reeds. Also, dumping industrial and health wastes in lakes caused a lack of oxygen and a concentration of hydrogen sulfide in some of these lakes, which prompted the Ministry of Health and Population to warn against eating or trading Lake Mariout fish due to the high toxicity of fish extracted from the waters of this lake.

The encroachments and pollution led to a change in the chemical composition and the collapse of the ecosystem of all the northern lakes in the Egyptian delta, which severely affected the fish wealth in these

lakes, which represent about 60% of the total fish wealth in Egypt. In addition to this, the drying operations of the northern lakes in the delta continue, which the government is carrying out on the one hand for the purpose of urban development and the extension of roads, power lines, oil and natural gas projects, and individuals for the purpose of constructing random salinas and fish farms on the other hand. This caused the total areas of the northern lakes to be reduced and their areas to shrink between 50% and 83% of their original areas - which resulted in changing the topography of the northern coast of the Egyptian Delta - in terms of the borders of the beaches and the areas of wetlands that separate the delta from the sea (Ouda, 2010).

In a study by Mohamad (2011) on Lake Mariout, it was found that it receives 7.5 million cubic meters / day of wastewater resulting from agricultural drainage and untreated industrial and municipal wastewater, which increased the concentration of heavy metals (cadmium, lead, chromium and nickel dissolved in the lake water). According to El-Naggar and Rifaat (2017) Lake Mariout was one of the most important shallow coastal lakes north of the Nile Delta in Egypt, which produces between 50 and 70% of the total fish production of the coastal lakes, but it was widely used for the discharge of industrial waste and sewage and agricultural drainage. As a result of environmental degradation, it has changed from being the most productive fish resource of the four great brackish water lakes in Egypt to the least productive in two decades.

In 2014, Hereher explained that Lake Manzala, once the largest coastal wetland along the Mediterranean coast, is heading for its disappearance by two opposing forces, one of which is the shrinking of the water body by deposition of sediment from agricultural land and an abundance of grasses and marsh plants as well. as drying practices for agriculture, while another force includes removing the coastal sand strip separating the lake from the Mediterranean by erosion, which should eventually turn the lake into a coastal dam rather than a closed coastal lake.

In a geochemical study of the waters of Lake Manzala for both El-Badry and. Khalifa (2017), the results showed that the arsenic content in the lake ranges from 4.6 to 22 ppm (parts per million), with an average of 12 ppm, which is about 8 times the average of the Earth's crust. Selenium concentrations range from 3-5 ppm with an average of 4 ppm, which is about 80 times the average crust of the Earth. The tin in the studied lake also ranged between 25-90 ppm with an average of 46 ppm, which is about 9 times the

average of the earth's crust. The highest values of arsenic, selenium and tin extended towards the industrial zone in Port Said Governorate. To sum up what the researchers said statistically, pollution indicators reflect a high level of pollutants in the vicinity of industrial activity around Port Said and Damietta governorates, while agricultural pollution is concentrated in the banks of Al-Serw and Bahr Al-Baqar.

This was followed by Ismail and Hettiarachchi (2017), Bek et al. (2019) and Elmorsi et al. (2019) who confirmed that Lake Manzala is exposed to high inputs of pollutants from industrial, domestic and agricultural sources. According to these authors heavy metal pollution is the most important type of pollution in the lake, which receives mixed discharges from densely populated areas, and that climatic gases and large quantities of particles, nutrients, bacteria and toxic organic materials are transferred to the lake through the discharge of sewage, especially the Bahr al-Baqar drain. In an analysis of samples of fish extracted from Lake Manzala by Sallam et al. (2019) high concentrations of the elements mercury, lead and cadmium were detected in Nile tilapia, gray mullet, and African catfish, respectively. The levels of mercury and cadmium contamination showed significant differences between summer and winter in three different types of fish that were examined.

In a study by Shoman (2018) about volume changes in Lake Manzala using remote sensing and GIS, the results showed that the Manzala Lake area decreased by more than 60% during the study period from 1972 to 2017. Also, Al-Muzayen (2019), Director General of Production at the Public Authority for Fisheries Development, found that the policy of drying of Lake Manzala led to a decrease in the area of the lake to less than 25% of its original area. The draining, which is considered one of the most effective and destructive policies for fisheries, has led to a change in the topography of the lake and its collapse. According to him, the most important problems of the lake are: A- Pollution resulting from untreated sewage from the Bahr al-Baqar drain, where it pours into Lake Manzala. The quantities of untreated sewage discharged by this drain are estimated at 1.5 million cubic meters per day (of which about 1.25 million cubic meters are from Greater Cairo alone). B- Pollution as a result of sewage resulting from the remnants of housing communities whose waste is dumped on both sides of the Bahr al-Baqar drain and the shores of the lake, or directly received by sewage networks from cities and villages overlooking Lake Manzala directly, the nearest of which is Moheb and Sayala drain in Damietta, as well as the sewage

system in the Al-Heishah Al-Khayyat area in Damietta. Although there is a purification plant, it has been inoperative for two years and receives sewage waste directly without any treatment. C- Pollution resulting from untreated industrial wastewater. D- Pollution resulting from agricultural drainage laden with pesticides. The quantities of water discharged by these drains into the lake are estimated at about 6 million cubic meters / day.

As for Lake Burullus, Dumont and. El-Shabrawy (2007) explained that since the construction of the High Dam, the Nile River has stopped the flow of the annual sediment layer into the river delta, which made the decline and coastal erosion continue and now consume the sandy strip which separates the Lake from the sea. In the Annual Report on Lake Burullus (2018-2019) for the Environmental Affairs Agency, the Egyptian Lakes Environmental Monitoring Program, it was stated that the area of the lake had shrunk from 165,000 feddans to less than 70,000 feddans after nearly 60,000 feddans were deducted, and that weeds and linen cover about 25,000 thousand feddans, and that the silt rate is high due to the formation of many islands, and the pollution rate is high at rates exceeding the permissible limits in stages, as a result of dumping more than 57 billion cubic meters annually into the lake of sewage and agricultural water, and the drainage of fish farms based on the southern edges of the lake.

In a study by Mohamadin et al. (2019) the Lakes Mariout, Idku, Burullus, Manzala and Bardawil are under an increasing threat from pollution and the destruction of the surrounding wetlands. Inorganic pollutants such as heavy metals have been identified in lake sediments. Organic pollutants such as polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) have been found to strongly bind to sediments. In Lake Manzala, Hg showed the highest values and alarming toxicity levels, and is considered one of the most dangerous. Burullus, Idku and Bardawil lakes recorded the highest values for some heavy metals, while Lake Mariout got the highest values ranging for organic pollutants.

In an official report issued by the General Authority for Fisheries Development of the Ministry of Agriculture and Land Reclamation, (Almarsad Almasry, 2020) that the number of cases of infringements on state property in the northern lakes during 2015 and 2016 amounted to 4,656 cases, with a total area of infringements on Lakes Manzala, Burullus, Idku and Mariout up to 64017 feddans in the four lakes located in the governorates of

Damietta, Port Said, Daqahlia, Kafr El-Sheikh, Beheira and Alexandria, while Lake Bardawil in Sinai was not subjected to any encroachments. According to the same report, Idku Lake decreased in area from 50,000 feddans at the end of the nineteenth century to 37,000 feddans until the early fifties of this century, as a result of the siltation of large parts of it. This is in addition to the encroachments, whether by erecting bridges or fish farms, and construction and agricultural operations. An area of 4532 feddans of the Lake Idku has been encroached upon, and its area has now reached about 15,045 feddans, including a water area estimated at 14,389 feddans. Lake Idku has suffered for decades from a high level of pollution, as it was a center for receiving all kinds of pollutants from industrial and agricultural wastewater, gas companies dumping their waste into the lake, and more than 20 sewage stations pour into drains that drain directly into the lake. The water level in the lake decreased, due to the siltation process, and the insufficient dredgers to clear and deepen the lake, which led to the death of fish in the lake water.

In the Annual Report on Lake Mariout (2019-2020) for the Environmental Affairs Agency, the Egyptian Lakes Environmental Monitoring Program, it was stated that industries in the city of Alexandria receive annually more than one million cubic meters of liquid waste loaded with about 260 tons of suspended solids per day, without treatment in the sea and in a lake Mariout, south of Alexandria, in fresh water canals, and in drains and sewage sewers. The city also produces daily more than one million cubic meters of various sewage waste, mixed with industrial waste, hospital and fuel station waste. About half of this amount is received without treatment in water bodies, while the other half is dumped after preliminary treatment in Lake Mariout. The governorate also has 200,000 feddans of agricultural land, which results in agricultural drainage loaded with pesticide residues and chemical fertilizers that eventually reach Lake Mariout. Also, petroleum, cement, iron and petrochemical refineries pollute the lake with chemical residues, in addition to the spread of weeds, reeds and other aquatic plants.

On the other hand, climate change scientists have unanimously agreed that the increase in the temperature of the Earth's climate system is a fact that cannot be denied. It is derived from data, statistics and scientific observations published by the Intergovernmental Panel on Climate Change in its six reports (IPCC, 1990- IPCC, 2021), which confirm the increase in the global average temperature of the air, sea and ocean waters, the increase in ice and snow melt rates, and the rise in the global average surface level during the twentieth century. Scientists and

experts have also agreed in the IPCC reports that there is strong evidence that current policies to reduce carbon emissions will not be able to reduce these emissions over the next dozens of years. The emission of carbon dioxide gas will increase in proportion to the air by between 25-90% during the period from 2000 to 2030 and thereafter. And that the continuation of these emissions threatens to further global warming in the twenty-first century, at rates greater than what it was in the twentieth century.

All the scientific reports issued by the United Nations organizations (the Intergovernmental Authority on Climate Change “The World Bank, the United Nations Environment Program”) and scientific non-governmental organizations, universities, Egyptian and international research centers and news agencies have unanimously agreed that the Egyptian Delta is considered one of the most threatened sites by marine invasion as a result of the increase in global sea level. These risks combine with the side effects left by the construction of the High Dam to make the delta in imminent and certain danger. The situation is, thus, dangerous and requires quick treatment, and any delay means more losses. The threats arising from these risks will persist for several decades and perhaps for a few centuries, even assuming the success of the United Nations in stabilizing the concentration of greenhouse gases in the atmosphere in the near future. This is due to the fact that the Earth's climate system needs centuries to stabilize. On the other hand, the current international policies adopted to reduce the rate of emissions of these gases will not be able to reduce these emissions during the next tens of years. Added to these threats is the continuous increase in water poverty rates due to the increase in population growth rates and agricultural projects on the one hand, and the instability of fresh water resources available in Egypt on the other hand as a result of the construction of the Ethiopian Renaissance Dam and its severe impact on the amount of Nile water coming into Egypt. As well as the current environmental deterioration that afflicts the dry and wet coastal lands of the delta as a result of the river stopping pumping silt in front of the coast of the delta following the construction of the High Dam on the one hand, and unguided human activities on the other.

The present research deals with the study of the topography and geomorphology of the four northern lakes of the Nile Delta in Egypt (Al Manzala - Burullus - Mariout - Idku), the surrounding plains, and their belts separating them from the Mediterranean Sea. (Figs. 1a and b), with a comparison of the areas of these lakes now with what

they were about seventy years ago, and how these lakes will be and the plains around them if the global level of sea level increases as the most important sequential effect of the climatic changes that the world is witnessing during this century. Hence, this study aims to determine the quantitative and qualitative size of the risks that will face the northern lakes and the surrounding plains in the northern, northeastern and northwestern part of the Nile Delta as a result of the global rise in sea level during this century, the coastal threat sources and the traditional defense policies that must be followed to avoid all these risks.

The current study is based on the consideration that the maximum remainder increase in the global average sea level (GMSL) during the twenty-first century is 100 centimeters, based on the sixth report issued by the Intergovernmental Panel on Climate Change (IPCC), and the supplements published concurrently with it in December 2019-2021. As for what was mentioned in the scientific report of the World Bank directed to developing countries (Dasgupta, 2007) that the maximum sea level rise during this century could reach 300 centimeters, or what was mentioned in a study of the European Environment Agency (2021) that the rapid disintegration of the ice cover in Antarctica may lead to a sharp rise in the global sea level to 2.3 meters in 2100 - we don't work with it. Such a sharp increase in sea level is not expected by the Intergovernmental Panel to occur in this century, but rather during the next two or three centuries as a long-term effect of climate change - due to the greater ice loss expected from the Antarctic ice sheet. This is if emissions are not controlled so that the concentration of carbon dioxide in the atmosphere reaches 660-790 parts per million.

2. Methods of Study

In this study, modern international mapping programs have been used, the most important of which is the Global Mapper (versions 9-13) program to design detailed topographic and contour maps of the delta coasts and to determine the damaged and safe areas in relation to sea level based on the information and digital data for ground elevations received from the Shuttle Radar Topographic Mission (SRTM) of NASA. This shuttle has built a high-resolution World-Wide Elevation Data (3-arc- second Resolution) system for most of the Earth's land surfaces. The fourth improved version of this information released in 2008 by CGIAR-CSI was used.

The current human uses of the threatened lands in question have been determined through precise and

direct electronic projection of modern satellite images on topographical maps designed with high accuracy. Worldwide high resolution color imagery from Digital Globe, and satellite images from Landsat 7, Bing Map, Google Earth Pro, and Google Maps were used. Also, GIS programs available to determine locations were used, such as The USGS Digital Elevation Model (DEM), USGS digital Raster Graphic (DRG) data, USDA National Agriculture Imagery Program (NAIP), Digital Chart of the World, Egypt. This is in addition to field visits to some threatened neighborhoods along the northern coast of the Nile Delta and its beaches.

This study included detailed topographic maps of the northern lakes of the Nile Delta (Al-Manzala,

Burullus, Mariout, Idku) in their current condition in order to compare them with what they were in the early fifties, and then put them in the case of sea level rise; maps of short heights and colors to show the boundaries of the threatened lands in relation to the current and prospective sea level, and the sources of their threat; the satellite images corresponding to these topographic maps, represented by the lines of the new beach in the event of sea level rise, the threatened lands and their areas, and the sources of their threat; the effect of sea level rise on human uses of threatened land spaces, in order to determine the size of the gaps that permeate the coastal threat sources in order to determine the appropriate means of protection and the specifications of defense lines.




Fig 1a: Detailed topography of the Nile delta as it is the present status (after Ouda, 2010, 2011). The white lines are the boundaries of the natural delta that separate it from the desert back on both sides of the delta. The total area of the delta within these borders is 24450.0 square kilometers, extending from the Tina plain in the east to the coasts of Alexandria in the west (from Abu Qir in the north to Agamy in the south), and from the sea in the north to the west of Greater Cairo before the river branch begins directly in the south. Note that the eastern extension of the delta towards Ismailia and the Bitter Lakes falls within the natural delta range according to what was mentioned in the ancient maps before the Islamic conquest and during the Fatimid and Ayyubid eras (see the Atlas of the History of Islam - Hussein Munis 1987). The surface of the delta descends gradually from less than 20 meters above sea level at south to below zero level (below sea level) at north. The delta includes, from north to south, an area of ~ 4400 km² situated topographically below sea level, equivalent to 18% (blue area), ~7000 km² of surface level ranging from

sea level (0 meter) to 3.0 meters above sea level, equivalent to 28.6% (yellow areas), ~3600 km² of surface level ranging from more than 3.0 meters to 5.0 meters above sea level equivalent to 14.7% (light green areas), ~5700 km² of surface level ranging from more than 5.0 meters to 10.0 meters above sea level, equivalent to 23.3% (red areas), ~3660 km² surface level ranging from more than 10.0 meters to 20.0 meters above sea level equivalent to 14.9% (black areas), then sporadic 90 km² of surface level situated more than 20.0 meters above sea level equivalent to 0.4% of the total area of Delta.




Fig. 1b: A brief topography of the Nile Delta as it stands, showing the locations and areas of the northern lakes Manzala, Burullus, Idku and Mariout (with continuous white lines) as they currently appear in relation to the total area of the delta. The white dashed lines are the boundaries of the natural delta. The black areas are elevated above sea level with a maximum of 20.0 meters. The blue areas within the delta borders are the dry (2790 km²) and wet areas (1031 km²) whose level is less than sea level.

3. Results and Discussion

3.1. The current situation of the northern lakes in the Egyptian delta, compared to what it was in 1949

3.1.1. The Manzala Lake

The Lake Manzala, the largest of the Egyptian lakes, has reduced its geographical area from 1797 km² in 1949 (according to cadastral maps in 1949, Fig. 2a) to 1200 km² in 1980, after draining large areas south and east the lake. Currently, the area occupied by the lake borders is 850 km² (Fig. 2b) after separating the southern part used as a general sewage complex coming from Cairo and the neighboring governorates of the lake (Bahr al-Baqar, Bahr Hados, Faqous and other drains.). This area does not reflect the reality of the lake from the inside, as the lake was subjected to drying operations for the purpose of constructing roads and public utilities. Areas of them were also deducted and isolated as private basins by individuals for the purpose of establishing random fish farms, in addition to sand dunes, dry marsh sediments, residential villages and dry lands whose level rises above sea level.

By measuring the dry and dried areas within the current Manzala lake borders, it was found that they are about 350 km², while the boundaries of the water basins do not actually exceed 500 km², of which about 455 km² are already submerged, and the remaining 45 km² are roads, bridges and dry paths that intersect the water basins

(Figs. 3a-b). If we add the area currently submerged in the public sewage complex south of the lake, which is approximately 22 km², then the total area currently submerged in Lake Manzala is 477 km². And after the lake was overlooking five governorates, it became confined between the governorates of Port Said and Damietta. Human activity is concentrated east of Lake Manzala, along the west bank of the Suez Canal, south of the lake, and northwest of the lake near Damietta. It is worth noting that about 30% of the area of the lake has been drained to allow the Salam Canal to pass through it to deliver fresh water to the Sinai - despite the lake having been declared as a nature protectorate since 1988.

The current topography of the lake can be distributed through satellites and the application of the international mapping program as follows:

- 68 km² of dried lands and dry marsh sediments, the level of which does not exceed sea level.
- 152 km² of dry lands with a level ranging from more than zero to one meter above sea level
- 175 km² of dry land partially covered by sand dunes whose level rises more than one meter above sea level
- 455 km² is already submerged below sea level, in addition to 22 km² in the public sewage complex south of the lake (Figs. 3a and b)

The lake is isolated from the sea in the north by a belt of sand dunes that is not homogeneous topographically in terms of the latitudinal extension, the level of the sand dunes, the nature and size of the inter-depressions. This belt extends along the northern coast of the lake, east of the Nile River, in an east-west direction, with a length of 7.6 km, and then deviates sharply in a southeast direction until Port Said, with a length of about 44 km (Figs. 4a-b and 5a). The width of the belt ranges from 0.8 km to 4.75 km, with an average range from 1.7 km to 2.8 km. The area of the southeast/northwest belt of the lake is about 109 km² (Figs.5b-f), of which 33 km² (30% of the total belt area) does not exceed sea level (zero), 24 km² (22%) ranges between zero and one meter above sea level, also 24 km² (22%) is more than one meter and does not exceed two meters above sea level, and the rest of the area (28 km², or 26%) is more than two meters above sea level, with a maximum of 6 meters above sea level. The belt, south of the shore line, parallel to the coast, is interspersed with submerged depressions of varying latitudinal and longitudinal extensions that act as lakes and sub-pools of the main lake and are connected to it through channels and transverse corridors.

The western part of the southeastern belt of the lake known as Barr al-Dahra (west of al-Manasra village) constitutes the largest part of the belt area (about 75 km²). It consists of intermittent longitudinal semi-parallel chains of sand dunes ranging in level between +1 and +6 meters, and the majority of their area does not exceed 2 meters above sea level. These dunes extend longitudinally parallel to the beach line, and many submerged depressions are confined between them, representing about 30.7% of the area of this part of the belt (Figs. 5b and 5c). These depressions converge and intersect in the south at Manama Al-Sheikh Ali (south of the Coast Guard Point Al-Halq), where the migratory dune chains from the north and northwest lose their extension and decrease in density. The inter-depressions expand, which in turn forms a large sub-lake connected to the main lake (the Maidan). This lake is separated from the sea only by a dilapidated belt of scattered and intermittent sand dunes, whose level does not exceed 2 meters above sea level, and is interspersed with many gaps that serve as corridors between the sea and the lake, with a level ranging between zero and -5 meters below sea level. These gaps are spread along the coast extending from Manama Al-Sheikh to Al-Ashtum Al-Jadid (south of the coastal deflection guard point) with a length of 7.2 km. This part of the coast represents the most dangerous parts along the coast of Lake Manzala and threatens to rush sea water into the lake in case the sea level increases by an amount ranging from 50 cm to a meter.

There are also some gaps along the coast extending from Manama Sidi El-Gendy to the middle of the Al-Shenka pond northwest of Al-Manasra village, in which the beach dunes sometimes rise to about 6 meters above sea level, but these gaps (about 2.1 km long) represent a threat to the pond Al-Shenka, which lies south of the beach line, and then Lake Manzala, if the sea level rises by no less than a meter (Figs.5d and 5e). Noting that in the event of an increase in the sea level by a meter, the sea water will sweep about 59% of the coastal sandy belt that forms what is known as the Barr al-Dahra, which extends northwest from the village of Manasra to the northern border of the lake, with a length of about 23 km.

It is clear that the disintegration of the sand belt and the decrease in its level along the Barr al-Dahra is due to the continuous migration of sand dunes from the northwest to the southeast, with not being compensated by new sediments that migrate to them from the mouth of the river Damietta branch as a result of the river losing its load after the construction of the High Dam, which increased, in turn, rate of beach erosion. It is noted that there are chains of sand dunes (Bur Al-Hamar, Bur Al-Raml and Bur Sidi al-Baghdadi) that cut across the Manzala Lake

itself in several locations, extending from near the mouth of the river to the interior of the lake in a southeast direction, parallel to the coastal chains, until it intersects with the latter south of the Shenka pool (Fig.5e). As for the eastern part of the belt extending from Al-Shenka pond to Port Said, it has an area of about 34 km², and it seems more coherent and tight, despite its reduced width than it is in the Barr al-Dahra (Fig.5f), despite the fact that depressions and submerged basins occupy an area equivalent to 29% of its total area. The reason for this is that the sand dunes are a tight longitudinal chain along the shoreline with a height of 3 m to 8 m above surface level, while the depressions and submerged basins limit the dunes to the south. These depressions and basins are supported by industrial structures, as this part of the coastal strip extending east from Sidi Mohamed El Maghribi to Port Said is characterized by human, tourist, industrial and residential activity (Fig.4a). This activity has contributed to changing the morphology of this part of the sand belt and the beaches overlooking the sea.




Fig.2a: A brief topography of the northeastern coastal part of the Nile Delta, showing the boundaries of Lake Manzala (marked by yellow lines) according to the cadastral maps in 1949. The total area of the

original lake was 1797 square kilometers, including the sandy belt separating the lake from the sea, and not including the submerged areas at the bays or small separate ponds in the governorates of Dakahlia and Sharqia. The black areas are dry lands whose level rises above sea level up to 20 m, while the blue areas are below sea level. Fig. 2b: Brief topography of the northeastern coastal part of the Nile Delta, showing the current boundaries of Lake Manzala (yellow lines) after draining additional areas. The part located south of the lake is the remaining part of the sewage complex coming from Cairo and the neighboring governorates (Bahr al-Baqar, Bahr Hados, Faqous and other drains). From now and ongoing, the geographical north is directed north of the picture.




Fig.3a: Topography of the coastal strip of the current Lake Manzala - as it is in the current situation. Black areas are dry or drained lands whose level rises up to 20 meters above sea level. Blue areas are flooded, dry or drained lands below sea level. The yellow line represents the current general boundary of the lake. The red lines are the boundaries of the current water basins within the general boundaries, which represent about 59% (about 500 km²) of the general area of the current lake (850 km²). The interstitial areas that permeate the water basins are marsh deposits, sand dunes, roads and public utilities,

representing about 41% of the general area of the lake. Note that the water basins are also interspersed with 45 km of sand dunes whose level rises above sea level, Fig. 3b: Satellite image of the northeastern coastal part of the Nile River Delta, showing the development of the current Lake Manzala from 1949 until now. The white areas are the areas that have been drained from the original Manzala Lake from 1949 until 1990. The red areas are the dry natural areas inside the lake, in addition to the areas that have been drained and exploited inside and outside the lake since 1990 until now. The black ground areas are the wet areas exploited from the lake for sewage coming from the neighboring governorates of the lake.





Fig. 4a: A brief topography of the coastal strip of Lake Manzala, which extends on the Mediterranean coast between Port Said in the east and Ezbet Al-Burg in the west - if the sea level increases by one meter, showing on them the locations of the low gaps (yellow circles) that cut through the sandy belt of Lake Manzala, which in turn will serve as waterways between the sea and Lake Manzala and the dry and dried plains that surround them. Note the small gaps that cross the eastern bridge of the Nile, the Damietta

branch, between Damietta and Ezbet El Burg, which will work also as waterways assisting the flow of water from the lake to the Nile River. The black areas are dry lands whose level rises more than a meter above sea level, while the blue areas are submerged and dry or dried low lands, their level ranges from - 7.5 meters below sea level to one meter above sea level. The numbers next to the circles are the radii of the holes. Fig. 4b: A brief topography of the coastal strip extending on the Mediterranean coast between the governorates of Port Said in the east and Daqahlia in the west - if the sea level increases by one meter, explaining on it the locations of the low-lying gaps that permeate the sandy belt of Lake Manzala and the western bridge of the Suez Canal, which in turn will serve as water passages between the canal on the one hand, and Lake Manzala and the dry and dried plains that surround it on the other. The yellow line represents the borders of the lands threatened by sea invasion.



Figs. 5a & b: Detailed topography of the northern sandy belt (Fig. 5a) and the northwestern-southeastern sandy belt (Fig.5b) of Lake Manzala on the Mediterranean coast - if the sea level rises by a meter. The white line is the current beach line. Note the spread of in-between gaps (red circles) along the shoreline in both the northern and northwestern-southeastern belts, especially in the area known as Barr Al-Dahra (Fig. 5b). These depressions will serve as sea lanes between the sea and Lake Manzala in case the sea level rises. **Figs. 5c - e :** A brief topography of the northwestern-southeastern sandy belt of Lake Manzala (the northwestern part of Barr Al-Dahra, Fig. 5c, the southeastern part of Barr al-Dahra northwest of Ezbet Shalaby Al-Rawdy, Fig. 5d, and the southeastern part of the belt extending from Ezbet Shalaby Al-Rawdy in the west to Al-Manasra village in the east, Fig. 5e,) – showing the location of gaps (yellow circles) that crosses the belt and which in turn will serve as sea lanes between the sea and Lake Manzala when the sea level rises to a maximum of one meter. The black areas are dry lands whose level rises more than a meter

above sea level. The blue land areas are the submerged and drained areas of Lake Manzala and its related ponds, the level of which is below sea level (-3.0 m – -5m with a maximum of -15.0 m below sea level). The red areas are dry lands whose level rises above sea level by a maximum of one meter above sea level. The white patches are areas of unknown heights with satellites. Fig. 5 f: Detailed topography of the southeastern sandy belt of Lake Manzala extending from Al-Manasra village in the west to Port Said in the east along the coast of the Mediterranean Sea - if sea level rises by a meter. Note the existence of a continuous series of high sand dunes (from 3.0 meters to 7.0 meters and a maximum of 10.0 meters above sea level) extending directly south of the shore line and acting as a tight buffer between the sea and Lake Manzala. For legend see Figs.5a & b.

3.1.2. The Burullus Lake

Lake Burullus extends along the northern coast of the delta in a northeast/southwest direction, where its current borders start from about 12 km east of Rosetta to Baltim in the east, with a length of 47 km and a width ranging from 5 km to 12 km. The lake has undergone a morphological change from the north, west and south as a result of human activity from drying and exploitation. Its area was reduced from 869 km² in 1949 (according to the cadastral maps issued in the same year, Fig.6a) to 466 km² as a result of human activity from drying and exploitation, of which 394 km² are already submerged areas and the rest are sabkha deposits, sand dunes and islands (Fig. 6b). Bushes and reeds are widespread in the flooded areas, especially the eastern part. Drying and filling operations were concentrated along the northern, southern and western shores of the lake for the purpose of establishing fish farms, and exploiting the sand dunes on the shores to grow crops and vegetables - although the lake has been declared a nature protectorate.

The current topography of the lake is distributed as follows:

- 394 km² of submerged lands below sea level, of which about 151 km² are interspersed with sediments of sabkha, weeds and reeds (Figs. 7a and 7b).
- 57 km² of dry land and swampy sediments inside the lake whose level ranges from more than zero to one meter above sea level.
- 15 km² of dry land and sand dunes inside the lake whose level rises more than one meter above sea level.

The lake is separated from the sea by a loose belt of sand dunes that extends from Kom Mishaal in the west to the Burg in the east in a northeast-southwest direction, with a length of about 41 km, a width ranging from 700 meters to 5.5 km, and a total area of about 115.5 km². The gaps whose level is below sea level down to -5.0 meters, which permeate the belt, constitute about 44% of the belt area (Figs. 8a and 8b). The sand dunes that make up the belt have undergone a great natural morphological change due to coastal erosion processes and the continuous migration of the dunes towards the lake due to air currents, in addition to human activity. The erosion of the dunes or their migration is no longer compensated by any new sediments migrating to them from the side of the Rashid estuary due to the river losing its load since the establishment of the High Dam. Thus, the sand dunes lost their cohesion, their level decreased, and the low gaps between them whose level is less than sea level between the sand dunes whose level does not exceed 3 meters widened in the greater part of the length of the belt. This has greatly contributed to the increase in the salinity of the lake and the change in its ecological composition and fishing system as a result of the intense rush of tidal currents that penetrate the lake daily.

Sand dunes whose level does not exceed one meter above sea level occupy 23% of the belt area, and sand dunes whose level does not exceed 2.0 meters above sea level 18.6% of the total area, while the remaining 14.45% is occupied by sand dunes more than two meters above sea level. These dunes are distributed throughout the part extending between Kom Mishaal and Tell Magluba. Hence, the total area threatened by the sea invasion in case the sea level rises by a maximum of one meter is about 67% of its total area, through 12 gaps directly connected to the sea, their diameters range from 500 meters to 5000 meters, with a total length of 16.9 km, which is equivalent to 41% of the length of the belt (Figs. 8a and b). This is in addition to the huge gap that extends across the coastal sandy belt between the Green Island and Kom Mishaal, with a length of 13.5 km (Figs. 9a and 9b). All of these gaps, which are about 30 km long, will serve as wide sea lanes between the sea in the north and Lake Burullus in the south, and from there to the surrounding low plains.

Hence, it can be said that the coastal sandy belt extending between the Green Island in the west and the Burg in the east, with a length of about 55 km, is the main source of threat to the wet and dry low plains in the northern delta, where the level of this belt does not exceed sea level by 47.4%, and does not exceed one meter above sea level by 73.4%, and does not exceed two meters at a rate of 88.3% of its total area, which is about 175.5 km² (Fig. 8a). While the rest of the belt area (11.7%) contains sand dunes whose level ranges from more than 2.0

meters to 10.0 meters above sea level. As for the low gaps that permeate the belt (Figs. 10a-f), whose level ranges from -5 meters (and a maximum of -7.5 meters) below sea level to one meter above sea level, their diameters range from 500 meters to 13.5 km (the great gap that characterizes the western extension of the belt west of Kom Mishaal to the green island north of Rashid, Figs. 10b), with a total length of about 30.0 km. Hence, about 55% of the belt's length is threatened by the transformation into wide sea lanes between the sea and the wet and dry southern plains, and that 73.4 percent of the total area of the belt is threatened by direct invasion from the sea if the sea level rises by a maximum of one meter. This matter threatens to invade Lake Burullus and the dry (or dried) low plains that surround it to the east, west and south in the north of Kafr El-Sheikh Governorate, including the original lands of the lake that have previously been drained or exploited - unless the necessary defense measures are taken.

As for the sand belt that cuts between the sea in the north and the Green Island-Burg Mughaizel in the south (east of the Rashid estuary), it has an area of about 10 km², of which 6 km² does not exceed zero (sea level); some of them are already submerged (Al-Ghalyoun ponds, Al-Wastaniya, Al-Shaqaqi and Al-Mitah ponds), and others are dry or dried; and 3 km² whose level ranges between zero and one meter above sea level, and the rest of the area (1.0 km²) is covered by sand dunes whose level ranges between one and two meters above sea level. This means that the entire length of the belt (5.5 km) will be invaded by 90% of its area if the sea level rises by a meter (Fig. 10a).

Thus, the current weak areas that are below sea level in the sandy belt extending from Lisan Rashid to the Burg constitute about 47% of its total area and about 53% of its total length, which threatens to invade the sea for half of the belt and from it to Lake Burullus and the low plains located south and east the lake. In fact, the vast majority (about 62%) of the dry or drained lands extending south, east and west of Lake Burullus, north of latitude 30° 18.75' and between longitudes 30° 33.75' and 31° 3.75', with an extension southeast to Hamoul, and northeast to Baltim village, west to Rashid, and southwest to Sindion on the eastern bank of the Nile - are low lands whose level ranges between -4 meters below sea level and one meter above sea level.

In contrast to the sand belt that separates Lake Burullus from the sea, the sand belt extending from Burg (the far northeast of Lake Burullus) in the west to Gamasa in the east is considered one of the strongest natural defenses that prevent the sea from invading this coast long ago, except for the seasonal inundation of some sites. The coast extending for about 60 km in a southeast/northwest direction between Burg in the east and Gamasa in the west is distinguished from the coastal areas of the Burullus and Manzala lakes by the presence of dense fields of sand dunes (440 km²). that act as a belt parallel to the beach graduating in the transverse extension south of the beach from 1.0 km in the west (at the Burg) to 11 km east (at Gamasa), with an average of about 7 km (Fig. 11). Although there are many inter-depressions interspersing these dunes longitudinally below sea level, but there are no lakes along this coast. The disappearance of lakes from this area from the coast of the delta is due to the continuous deposition of sand dunes, which since 3000 BC covered a large area of this area (Stanley et al., 1992). The sedimentation rates of these dunes exceeded the combined rates of delta subsidence and global sea level rise during this period. This led to the accumulation of dunes that form a continuous, coherent and wide topographical ridge between the Burg in the west and Gamasa in the east, as this line acts as a natural impenetrable dam that prevents the continuation of a waterway between the sea and the inter-depressions, which prevents the sea from invading these depressions, and thus prevents the formation of permanent lakes.

As for the tourist cities overlooking the sea along the coast extending from the tower in the west to Gamasa in the east, their beaches will be exposed to direct invasion from the sea to varying degrees, as most of the beaches of these cities are artificial beaches completely cut off from the sea (Gamasa Resort) or it was expanded at the expense of the sea (Baltim Resort) instead of the natural beaches that were partially or completely eroded. The beaches of the Baltim Resort are threatened by 50% of its total area (Fig. 11), while the beaches of Gamasa are threatened with a complete invasion of 100% (Fig. 12) unless the level of these beaches is raised more than a meter by throwing sand on them. As for the cities themselves, they are safe from the invasion of the sea because their levels range from 4.0 to 8.0 meters above sea level.




Fig. 6a: A brief topography of the northern coastal part of the Nile Delta as it is in the current situation, with the boundaries of Lake Burullus indicated according to the cadastral maps in 1949. The total area of the original lake (the area marked with yellow lines) is approximately 869 square kilometers, without the sandy belt separating the lake from the sea. Black areas are dry lands whose level rises above sea level. The blue areas are wet, dry or dried lands whose level is below sea level to -3.0 meters (and a maximum of -7.5 meters). **Fig. 6b:** The same topography of Fig. 6a showing the current borders of Lake Burullus according to satellite images and information amounting to 466 square kilometers, which is equivalent to about 54% of the original area of the lake. Of this area, 394 square kilometers are submerged under sea level, and the remaining 72 square kilometers are dry marsh deposits that permeate the lake, sand dunes and lands whose level rises above sea level.




Fig. 7a: A brief topography of the northern coastal part of the Nile Delta as it is in the current situation, with the current borders of Lake Burullus (yellow lines) which has an area of about 466 square kilometers without the sandy belt separating it from the sea, including submerged areas under sea level of 394 square kilometers, and the rest of 72 square kilometers are dry marsh sediments permeating the lake, sand dunes and lands whose level rises above sea level. The white areas are the areas that have been drained since 1949 until now. Note that there are other small areas drained southeast and east of the lake, which confirms the existence of previous extensions of the lake before 1949. **Fig.7b:** Satellite image of the same northern coastal part of the Nile Delta, given its topography in Fig.7a.




Fig. 8a: A brief topography of Lake Burullus and the sandy belt separating it from the sea - when the sea level rises to a maximum of one meter, showing the locations of the low-lying gaps (red circles) that permeate this belt, which will serve as sea corridors between the sea and the dry and submerged lowlands south of the belt. The white line is the current beach line. The yellow line represents the current boundary of Lake Burullus. The blue land areas are depressions less than a meter above sea level until -5.0 meters (and a maximum of -7.5 meters near the shore line) below sea level, Black areas are lands that rise a meter or more above sea level. **Fig. 8b:** Satellite image of the same coastal belt whose topography is given in Fig. 8a, showing on it the locations of the low gaps (yellow circles) that permeate this belt.




Fig. 9a: Detailed topography of the sandy belt extending from the Green Island (north of Rashid) in the west and Kom Mishaal in the east along the Mediterranean coast - when the sea level rises to a maximum of one meter, showing the locations of the low gaps (red circles) that permeate the belt along the shoreline and which in turn will serve as sea lanes between the sea and the dry and flooded lowlands south of the belt (the blue areas). The white line is the current beach line. Note that the sand dunes that permeate the depression do not exceed an average of 2.0 meters above sea level, and therefore this coastal part of the belt is considered one of the most dangerous parts that threaten to invade the sea for all the lowlands located south of the shore line between Lake Burullus and the River Nile. **Fig. 9b:** Satellite image of the same sand belt whose topography is given in Fig. 9a.




Fig. 10a-f: Detailed topography of the sandy belt of Burullus Lake that extends from the mouth of the River Nile, the Rosetta branch at west to Al Burg at east if the sea level rises to a maximum of one meter. The white lines is the current beach line. The red circles mark the location of gaps cutting through the sandy belt and which in turn will become sea lanes between the sea in the north and low southern plains surrounding the Burullus Lake. Fig. 10a: The sang belt which separates the Green Island - Burg Mughazel (North Rashid) from the sea along the Mediterranean coast. Note the general deterioration of the sand belt north of the Green Island - Burg Mughazel, and this belt turned into large wet depressions ranging from -3,0 meters below sea level to one meter above sea level. Fig. 10b: A brief topography of the sandy belt extending from the Green Island (north of Rashid) in the west and Kom Mishaal in the east showing the collapse of the sand belt extending across the great gap (13.6 Km) which threatens to invade the sea all the low lands south of the shore line, whether submerged or dry, extending from Lake Burullus in the east to the Nile River in the west. The black areas are lands whose level rises more than above sea level, while the blue areas are low, submerged and dry lands ranging from -3.0 meters (and a maximum of -7.5 meters near the shore line) below sea level to one meter above sea level. Fig. 10c-f: Detailed topography of the sandy belt extending (north of Lake Burullus) from east of Kom Mishaal to Kom Mastrowa (Fig. 10c), from the east of Kom Mastrowa to the west of Tell El Maqluba (Fig. 10d) from Tell

Al-Maqluba in the west to Al-Burg in the east (Fig. 10e) and from the Burg in the west to Al Khashousi in the east (Fig.10f) along the coast of the Mediterranean. Note that the effect is non-existent along the shore line from Al-Burg in the west to Al-Wahhabia in the east due to the rise in the ground level from more than a meter to 8 meters or more directly above the shore.




Fig. 11: Panorama of detailed topography of the coastal strip of Baltim Resort on the Mediterranean coast between Burj in the west and Gamasa in the east when sea level rises by a maximum of one meter. The white line is the current beach line. Note that the southern depressions are isolated from the Baltim beach and summer resort by a longitudinal chain of high sand dunes (blue ground areas), and therefore they are safe from sea invasion, but the chances of being partially submerged by subsurface water increase if its level rises as a sequential effect of sea level rise.




Fig. 12: Detailed topography of the coastal strip of Gamasa Resort on the Mediterranean coast - showing the effect of sea level rise of a maximum of one meter. The white line is the current beach line. The blue beach areas delimited by black lines south of the shore line are the areas threatened by direct inundation from the sea, and they include the entire Gamasa beach. Note that the city is safe from the invasion of the sea.

3.1.3. The Idku Lake

As for Idku Lake, it is a glaring example of illegal encroachment, as the area of the lake shrank from 129 km² in 1949 (Fig. 13a-b) to 64 km² in 1982, then to 21.5 km² south of Al-Labbani until the end of 2008 (Fig. 14a), meaning that about 83% of the original lake area had been drained (Fig. 14b). As a result of the appropriation of large areas by farm owners in violation of the law, as well as due to government negligence in clearing the lake, as about 50% of the current area is made of reeds, Nile roses and weeds. Although Idku Lake is separated from the sea from the north and northeastern sides by a strong belt of sand dunes whose level ranges from one to 20 meters above sea level - the western side of the lake is not separated from the southern coast of Abu Qir Gulf by any sandy belts, except for some dunes dispersed whose level does not exceed one meter above sea level (Fig. 13a). Therefore, the sea's invasion of the southern coast of Abu Qir Gulf, when the global sea level rises by a meter, will lead to the invasion of the entire northwestern part of the Nile Delta until the Mahmoudiya Canal in the south, including what remains of Lake Idku.

While the city of Idku is located on heights ranging between 2.5 and 15.0 meters, with an average of 6.0-7.0 meters above sea level, the city is bordered by a depression in the north (the Gulf side) that extends along the beach with a level ranging from - 7.5 meters (below sea level) and one meter above sea level, which is a depression that is partially flooded and seasonally, despite the drying up of large areas of it. The city is also bordered on the west by a narrow depression perpendicular to the coast with a length of 1.5 km and a level ranging between -2.5 and +1.0 meters. It is also bordered on the east by a wide depression with a length of 2.76 km, and a ground level ranging between -2.0 and +1.0 meters. In addition, the city is surrounded on the south by low plains below sea level, part of which is submerged (remnants of Lake Idku) on the southeastern side of the city, which threatens to isolate the city from all directions and threatens the southern expansions of the city in the event of an increase in sea level by a maximum of one meter (Fig. 14b).

The southern coast of the Abu Qir Bay is not only a source of threat to Lake Idku, as this lake is no longer what remains of it after drying and exploitation as fish farms, which can be considered of great economic value - but this coast is the main source of threat to all the dry and wet lowlands (Idku and Mariout lakes) located in the south of the Bay in Northwest Delta (west of the Rashid branch, Ouda, 2022)). The sandy belt that separates the Bay from the southern plains between the Nile River in the east and the city of Alexandria in the west has been eroded not only by coastal erosion factors, but also by the population, industrial and educational activity that is concentrated on the coastal strip directly south of the Gulf between the suburb of Abu Qir and the village of Maadiyah. The topographical survey (Ouda, 2022) showed that this part of the Bay is a source of threat to all agricultural, residential and wet lands (remnants of the Idku and Mariout lakes) located in the south of the Gulf, which extends to the desert back in the south, between the Nile River - Damanhur in the east, and Alexandria in the west. The sandy coastal strip, which extends north of the Abu Qir-Maadiyah road and separates the sea from the low-lying plains to its south, which has an area of 6.1 km², is low in level, and does not exceed one meter in height above sea level at 83% of its total area. As for the gaps below sea level (down to -4 meters below sea level) that permeate the coastal strip, they occupy about 66.6% of the total area of the strip, and about 50% of the length of the coastal strip. Many of these gaps have been filled for construction purposes to a level that does not exceed sea level. It is concentrated between Abu Qir and Al Mandara, between Al Mandara and Al Tarh, and between Al Tarh and Al Maadiyah along the coast. Accordingly, an increase in sea level by a maximum of one meter would cause the sea to invade the majority of the coastal strip extending between Maadiyah and the suburb of Abu Qir (Fig. 15a-b).

Muhammad Ali's wall, which was built in 1830 in Abu Qir Bay in Alexandria, is not suitable for protecting the plains to its south. The wall is nothing but piles of stones that extend along the southern shore of the Gulf of Abu Qir that have no submersible under the soil to ensure that sea water does not seep through the subsurface soil in case the sea level rises. Also, its height is not enough to guarantee its protection from the rise in global sea level by a meter. The wall along its length is subjected to the invasion of high waves during hurricane seasons due to the high water levels during the tide, in a way that led to the flooding of the area with sea water many times, and the high waves led to collapses in many areas of the wall). The presence of leaks currently spreading on the coast directly south of the wall confirms the inability of this wall to repel severe waves if they rise for any reason of the local rise (such as severe storms and tsunami waves) as a result of the gaps and inter-holes spread in the wall ((for details see Ouda, 2022). Hence, the wear and tear of this wall, along with its inefficiency to counter the certain rise in the global sea level, will threaten the sea's invasion of all the southern plains of the Gulf of Abu Qir, including the remnants of Lake Idku and the northeastern plains of Alexandria, extending south of the neighborhoods of Maamoura, Montazah, Mandara, and Sidi Bishr al-Qibli, where their level is all under Sea level usually ranges from -2.0 m to -5.0 m, with a maximum of -7.0 m (Fig. 15a).

3.1.4. The Mariout Lake

As for Lake Mariout, opinions differed about its original area during the past centuries after the extinction of the Nile branches feeding the lake and the drying up of large parts of it, but it is certain topography that the lake was, at the least, occupying an area of not less than 570 km² south of Alexandria before 1949 (Fig. 16a), as this area is less than its current level Sea level between -3.0 m and -8.0 m below sea level although large parts of it have dried. In 1949, cadastral maps showed that the lake occupies an area of 383 km² of which 268 km² is located south of the city of Alexandria and 115 km² is the western extension of the lake to the city of Al-Hammam (Fig. 16 a-b). Since this date, the lake has been drained, especially since 1986, under the supervision of the Egyptian government and the Governorate of Alexandria until it was torn to a number of isolated lakes with a total area of 63 km² south of Alexandria (Fig. 17a), in addition to the western extension of the lake to the city of Al-Hammam, whose area does not exceed 87 km², 13 km² of which are fish farms (Fig. 17b). This means that about 76.5% of the area of the lake south of Alexandria has been deducted and drained since 1986 until now, and about 36% of the area of the western extension of the lake for the purpose of urban development and agricultural reform. In an official report issued by the Ministry of Environment, about 3.28 km² was filled in by a direct order from the Prime Minister to establish a sports city, the international park, the expansion of the international road and the sewage project in Alexandria. Within the framework of the New Alexandria project, the designs of the project included filling in about 5.67 km² in the Abu Azzam and Al-Alayma basins with the aim of establishing an urban area to which the Smart Village is attached.




Fig. 13a: Detailed topography of the northwestern coastal part of the Nile River delta - showing the boundaries of Idku Lake, according to cadastral maps in 1949, estimated at 129 square kilometers (black

borders). Note that all previous and current lake areas are below sea level without sand dunes or significant high lands interspersed. Fig. 13b: Satellite image of the same northwestern coastal part of the Nile River delta whose topography is given in Fig. 13a. - Note the areas which have been exploited as fish farms and the sabkha deposits scattered around the borders of the current lake.




Fig. 14a: Satellite image of the northwestern coastal part of the Nile Delta - showing the stages of Idku lake reduction. The white area represents the lake's borders in 1949 with a total area of 129 square kilometers. The blue areas represent the borders of the current lake with an area not exceeding 21.5 square kilometers. The reduction rate is estimated at 83%, while the remainder does not exceed 17% of the original area. **Fig. 14b:** Satellite image of the coastal strip extending between Tabiat El Gazayir in the south and Tabiat El Kalayem in the north along the eastern coast of Abu Qir Gulf - showing the effect of sea level rise of a maximum of one meter on land uses. The red areas north, east and north-east of Idku are the low plains threatened by direct marine invasion from the eastern coast of Abu Qir Bay, while the red areas south of Idku are the low plains threatened by direct marine invasion from the southern coast of Abu Qir Gulf through the wide gap (9 km) extending from El Maadiyah to the east To the suburb of Abu Qir to the west.




Fig. 15a: A brief topography of the northwestern part of the delta extending from the River Nile, the Rashid branch in the east to Alexandria in the west, and from Abu Qir Bay in the north to Mahmoudia Canal in the south - if the sea level rose by a maximum of one meter, showing the location of the main source of threat along the coast of Abu Qir Bay between the suburb of Abu Qir in the west to the village of Maadiyah in the east (large yellow circle) and the smaller low-lying gaps that permeate the thin land strip extending along the length Mahmoudiya Canal between Abu Homs in the east and Alexandria in the west (small yellow circles), which will act as auxiliary sea lanes on the rush of the Gulf waters from the northern plains to the southern plains extending between the Mahmoudiya Canal in the north and the desert backwaters in the south. The blue areas are low, dry and wet lands, with a level ranging from -7.0 meters below sea level to one meter above sea level. **Fig. 15b:** A brief topography of the northwestern part

of the Nile River delta (west of the Rosetta branch) when the sea level rises by a maximum of one meter - showing the locations of the coastal gaps (yellow circles) including the main gap on the southern coast of Abu Qir Gulf, the coastal gap extending between Wardyan and Dekheila along the Max Bay, and the small gaps that permeate the narrow land strip extending along the Mahmoudiya Canal between Abu Homs in the east and Alexandria in the west.





Fig. 16a: Detailed topography of the western coastal part of the Nile Delta, which extends south of the city of Alexandria along the Mediterranean coast, explaining on it the borders of the previous areas of Lake Mariout in 1949 (blue areas), according to the cadastral maps issued at the time, where it occupied about 383 square kilometers, of which 268 square kilometers south of Alexandria from the entrance to Abis in the east to Abu Talat in the west, and 115 square kilometers is the western extension of the lake from Abu Talat to Al- Hammam along the northern coast. Note that the lake occupied vast areas in the northwest of

the delta during ancient history when the city of Alexandria was established, but there are no documented maps that can be referenced. Fig.16 b: Satellite image of the same coastal part whose topography is given in Fig. 16a, explaining on it the previous and current borders of Lake Mariout. The white areas are the areas that have been drained and exploited from the original lake since 1949 until the present according to the cadastral maps issued at the time. The dark blue spaces represent the current remnants of the lake south of Alexandria, which is a number of stagnant ponds and isolated lakes with a total area of 63 km² (with the exception of the Nozha Airport and Airport Lake which were used before 1949), in addition to the western extension of the current lake from Abu Talat to Al- Hammam which assumes an area of no more than 87 km.



Figs. 17a and b: satellite images of the coastal strip of the city of Alexandria, extending from the entrance of Apis in the east to Al-Hammam in the west on the Mediterranean coast- showing the areas occupied by the current Lake Mariout as it is now. The dark blue areas represent the current remains of the lake, which is a number of stagnant ponds and lakes isolated from each other south of Alexandria, with a total area of 63 km² (Fig. 17a), in addition to the western extension of the lake from Abu Talat at East to Al-Hammam at West, and its current area is 87 km² (not including the fish farms that permeate it, Fig. 17b),

of which 75 square kilometers are submerged, and the rest is dry land permeating the lake. This means that about 60.0% of the area of the lake Mariout has been drained and exploited during the period from 1949 until now, and the drying operations are still going on.

The western port of Alexandria is separated from the low lands south of Alexandria, which occupies some of its area, Lake Mariout (or what is left of it), a coastal strip with a width ranging between 650 and 1600 meters, extending from Al-Qabbari station in the east to Al-Agamy in the west. This strip consists topographically of discontinuous limestone hills ranging in height from 3 m to 10 m. These hills are interspersed with depressions representing ancient drainage channels whose level ranges from -6 meters below sea level to one meter above sea level. And the elevation maps show that the middle part of this port, which extends between Wardyan and El Max, and the middle-western part that extends between El Max and Dekheila, are the weakest parts of this coastal strip, and that they represent the second source after Abu Qir Gulf to threaten all the dry and wet lowlands that lie between southern Alexandria and Damanhur until the desert back, due to the gaps between Wardyan and El-Max with a length of 2 km, and east of Dekheila with a length of 1.1 km (Figs. 15b and 18). Some of these gaps extend from the shore line to Lake Mariout, south of the port. Most of these gaps have been filled and exploited by establishing industrial facilities. However, the decrease in its current level by a meter above sea level will expose the belt to a partial invasion in the event of a rise in sea level of one meter or more, as it constitutes about 24% of the total area of the lime belt extending between the Wardyan and Dekheila (4.4 km²), and it is concentrated in the part between Wardyan and El-Max, where it constitutes about 39% of the area of this part of the belt (2.0 km²).




Fig. 18: A brief topography of the coastal strip of the western part of Alexandria, extending along the Gulf of El Max, from Qabbari in the east to Agamy in the west, including the Western Port and Dekheila Port when sea level rises by one meter. The white line is the current shore line. The yellow circle shows the location of the gap that permeate the limestone ridge that extends from Wardyan to Dekheila, especially between Wardyan and Al-Max which in turn will serve as sea passages between the sea and the southern plains. The blue areas, with their different shades, are the plains that are in danger of being invaded if the sea level rises.

Thus, it can be said that if the southern coast of the Gulf of Abu Qir is the main source of threat to all lands north of Beheira governorate, including what remains of Lake Idku, and south of Alexandria governorate due to its topographical, geological and historical conditions that pave the way for the invasion of the sea in the event of an increase in the global sea level, the middle part of the Western Port, between Wardyan and Dekheila, comes

as a second source of threat to the low plains located south of the Mahmoudiyah Canal, including what remains of Lake Mariout in case the sea level increases by not less than a meter, and not less (Ouda, 2022).

The following table summarizes the topographical scales of the northern lakes - how they were and how they became.

Total	Mariout Lake	Idku Lake	Burullus Lake	Manzala Lake	Egyptian lakes in the delta
3178 km ²	383 km ²	129 km ²	869 km ²	1797 km ²	Total area in 1949
1487.5 km ²	150 km ²	21.5 km ²	466 km ²	850 km ² Without a general sewage complex of 22 km ²	Current area
1690 km ²	233 km ²	107.5 km ²	403 km ²	947 km ²	The area that has been dehydrated or reduced
53.2 %	60.8 %	83.33 %	46.4 %	52.7 %	Area loss percentage
480 km ²	13 km ²	—	72 km ²	395 km ²	Dry plots within the existing boundaries
1007.5 km ²	137 km ²	21.5 km ²	394 km ²	455 km ² Without a general sewage complex of 22 km ²	The flooded area is already within the existing boundaries

Table 1: The northern lakes of the Nile Delta: what were they like? and How are they now

3.2. Threats to the northern lakes and the surrounding low plains in light of climate change

Based on the information and digital data received from the Shuttle Radar Topography Mission of NASA about the topography of the delta, and according to altimetry measurements using the Global Mapper program to design detailed topographic and contour maps of the Egyptian delta, it can be said that the increase in sea level by any amount greater than the current level will have dire consequences on the coasts of the Nile Delta. The delta is characterized by a continuous subsidence at its northern humid fringes, especially at the estuaries, and most of its northern parts do not exceed zero height, i.e. below sea level, with a depth inland ranging from 7.7 km to 58.5 km along the coast of the delta (Fig. 19). Therefore, the coasts of the delta are vulnerable to any rise in sea level, whether this rise is local or global. Although the local rise in sea level is limited in time and impact on the coasts of the delta, the global rise is long-term and has a great impact. A rise in this global level by a meter would sweep or submerge the dry and humid coastal lands that are lower than sea level and that are directly connected to the sea without being isolated from the sea by natural (or artificial insulators), or those separated from the sea by discontinuous chains of sand dunes with gaps between them. These lands are estimated to have an area of about 4001 km² (16.33% of the total area) at a minimum after excluding the threatened areas in the Tina Plain in the northwestern corner of the Sinai Peninsula. These lands include about 1031 km² of already submerged areas (the northern lakes of the delta currently, namely the lakes of Manzala, Burullus, Idku and Mariout, after human drying processes that affected large areas of them (Table 1), in addition to 2970 km² of dry areas extending along the northern coast of the delta (Table 2).

These lands were separated from the sea by intermittent belts of sand dunes that represented natural barriers, their height ranged between 1.5 and 14 meters above sea level, and their width ranged between 1 and 10 km along the northern arc of the delta, which extends between Port Said in the east and Abu Qir in the west. These belts arose from the accumulation of river sediments of sand and silt, which the river carried through its long journey from the headwaters to the estuaries in Damietta and Rashid, and then the sea and air currents redistributed them along the coast. But the current problem is that these belts have been severely weakened, their areas have diminished, and the gaps between them have increased as a result of their lack of the flow of river sediments from sand and silt since the start of the work of the High Dam in 1969 on the one hand, and the

continuation of the migration of the dunes due to the winds in the southeast direction on the other hand. The river's load is currently low to the point of nothing, and the transfer of river sediments to estuaries has become almost nothing. While the load of the river in 1964, before the closure of the river pass in front of the Aswan High Dam, ranged from 18 to 55 billion cubic meters annually of sediments, including the flood sediments, which amounted to 34 billion cubic meters. The river's load of suspended sediments was estimated at 111 billion kilograms annually, 93-98% of which reached estuaries during flood seasons (Ouda, 2010).

The interruption of river sediments after the construction of the High Dam contributed to the high rates of beach erosion along the northern arc of the delta. The delta outcrops inside the sea, which are called tongues, were eroded in both Rashid and Damietta. With a simple comparison between the topographical maps of the delta before the construction of the dam, and the maps designed on the basis of the information received by satellites today, it is clear that there is a wide morphological change that has occurred in both Lisan Rashid (Ahmed, 2002) Lisan Damietta (Al-Raey et al. 1999) and the sandy belts that separate the Burullus lakes from the sea, and those that separate the sea from the land in Abu Qir Bay (Ouda, 2010).

The surface of the delta falls annually at unequal rates relative to its edges as a result of its skewed position. The northeastern side of the delta is lower than the northwestern side, declining at a rate of 5 millimeters annually, while the northwestern side is declining by 3 millimeters annually. As for the middle delta, the decline rate does not exceed 4.0 mm annually (Smithsonian Institution (1985), Stanley and others, 1992). The reason for this subsidence is the continuous pressure of sediments on the subsurface rocks as well as due to some ground




Fig. 19: A brief topography of the Nile Delta if the sea level rises by a maximum of one meter - showing the borders of the plains in the northern part Nile Delta, which are threatened by direct invasion from the sea, or indirect invasion through the northern lakes. The white line is the normal delta boundary. The yellow lines are the general boundary of the threatened plains. The total affected area is 7303 km² (Table 2), representing 29.81% of the area of the delta, after excluding the Tina plain in northwestern Sinai, and the low areas not connected to surface water sources directly and could be threatened by partial inundation by subsurface water if its level rise as a sequential effect of sea level rise. The total affected area includes 5580 km² whose level does not exceed one meter above sea level, while the rest of this area, amounting to 1723², are cities, villages, dry lands and sand dunes whose level rises more than a meter above sea level but they will be besieged and isolated within the invaded areas.

movements directed by faults. This continuous decline in light of the unloaded river increases the degree of salinity of the land, and also affects the groundwater, in addition to the consequences of the exposure of the

coastal part of the delta to the invasion of the sea in the event of any local rise in sea level, even if the global sea level does not rise. If the area of 4001 km² is the minimum of wet and dry delta lands that are threatened with invasion if the global sea level rises by any amount more than the current level, then a rise of 100 centimeters in sea level, which is the maximum expected increase during the twenty-first century, will be sweeping wet and dry coastal lands with an estimated area of about 5580 km² along the northern arc of the delta from Port Said in the east to the suburb of Abu Qir in Alexandria in the west (Fig. 19). These lands do not exceed one meter above sea level. Hence, they do not include its interspersed cities, villages, roads, bridges, and sand dunes whose level is more than a meter above sea level, and which will be besieged and isolated within the invaded areas. The area of these lands is estimated at about 1723 km². Thus, the total affected areas in case of sea level rise by a maximum of one meter is 7303 km² (Table 2), representing 29.81% of the total area of the delta, after excluding the affected areas in the Tina plain in Sinai, and the low areas not connected to surface water sources directly and threatened by partial inundation by subsurface water. Its level rise as a sequential effect of sea level rise.




Fig. 20: A brief topography of the Nile Delta in case of rising sea levels by up to one meter - showing the locations of coastal gaps (yellow circles) along the northern sandy belt of the Delta from Port Said at east to Alexandria at west. The gaps are made up of 39 low-lying depressions ranging in diameters from 200 m to 13600 m. The ground level of these depressions ranges from below sea level to one meter above sea level, and thus would act as direct marine passages between the sea and the lowlands (dry and wet) lying south of the sea shore. In addition, there is a gap lying between Al Wardyan and El Dekheila along the Gulf of Max at Alexandria with a diameter of about 4000 meters, and which in turn would act as a sea corridor between the Gulf of Max and the low-lying plains south of Alexandria. Also, there are 33 depressions along the eastern and western bridges of the western branch of the Nile (Rosetta Branch, North of Fowa), ranging in diameter from 200 meters to 800 meters, and 4 depressions ranging in diameter from 400 meters to 600 meters along the eastern bridge of the eastern branch of the Nile (Damietta Branch, north of Damietta). All these depressions which assume a ground level less than one meter above sea level will turn freely as corridors between sea water in case of invading northern coasts and the River Nile. Furthermore, 21 depressions ranging in diameter from 200 meters to 800 meters occur along the western bridge of the Suez Canal between Port Said at north and El Ballah at south and which would act as marine passages between the Suez Canal and the low-lying plains situated along the western side of the Canal. Also, 12 depressions less than one meter above sea level and ranging in diameter from 200 meters to 3000 meters exist along the northern bridge of the Mahmoudiya Canal. These depressions would act as corridors freely help the rushing waters of Abu Qir Bay of the Northern Plains (North Canal Mahmoudiya) to the southern plains (South Canal Mahmoudiya).

The following is a geographical detail of the wet, dry or dried areas threatened by direct sea invasion through many coastal gaps, ranging from less than zero below sea level to one meter above sea level, permeating the sandy belt extending from Alexandria in the west to Port Said in the east (Fig. 20) These gaps will turn into sea lanes between the sea and the northern plains (including the Lakes) of the Nile Delta if the global sea level of rises by no more than one meter, The following is a geographical detail of the wet, dry or dried areas threatened by direct or indirect sea invasion through the northern lakes if the global sea level rises along the northern arc of the delta.

3.2.1. The northeastern part of the delta: Lake Manzala and the surrounding low plains

The large surface occupied by Lake Manzala and the extension of its northern coasts to the sea with a length of 52 km, as well as its eastern coasts along the Suez Canal (before drying and exploiting the eastern part of the original lake) with a length of about 53 km, with the thinness of the sandy belts that isolate it from the sea and the canal, in addition to the nature of the sandy subsurface soil of the lake - all of this makes it difficult to protect the lake from inundation as a result of seawater intrusion into the subsurface soil in case of an increase in sea level - even if it is protected from sea invasion through the sand belts surrounding it. Hence, securing its current internal borders tightly will prevent the lake waters from invading the low lands that surround it.

The area of the dry plains, the level of which does not exceed sea level, which surrounds the Manzala Lake from the east, south and west, and is connected to it, is about 389 km², including the lands that were previously deducted from the original lake, dried and exploited. This is in addition to the flooded areas used as a general sewage disposal complex south of the current lake, amounting to 22 km². Also, there is an area of about 8.0 km² of low lands below sea level extending along the Suez Canal west of Al-Ballah branch. These lands are partially submerged as a result of the continuous infiltration of the canal water through the subsurface soil. Thus, the total area of dry lands below sea level outside the current boundaries of Lake Manzala in the northeastern part of the delta is 419 km². This area is the minimum area exposed to the indirect marine invasion through the Manzala Lake if the sea invades the Manzala Lake, and then the lake level rises by any amount more than the current sea level. This is in addition to the submerged and dry areas whose level does not exceed sea level within the boundaries of Lake Manzala and threatened by direct invasion from the sea through the gaps that permeate the sandy belt of the lake. These areas are 523 km², which makes the lake lose its natural and chemical properties. Thus, the total wet and dry lands threatened by direct or indirect marine invasion through Lake Manzala, or extensive immersion due to water seeping through the subsurface soil in the northeastern part of the delta (bounded between the Suez Canal in the east and the Damietta branch in the west) is 942 km² as a minimum.

The matter becomes more dangerous if the sea level increases by a maximum of one meter, as sea water and the lake invade additional areas of dry land amounting to 686 km², of which 534 km² are dry lands outside the boundaries of the current Lake Manzala and 152 km² are dry lands that permeate the submerged areas within the boundaries of the lake. Thus, the total areas of dry and wet lands threatened by the invasion in this part of the delta, if the sea level rises by no more than one meter, will be about 1628 km² (Figs 21a-b). These lands are located east, south and west of Lake Manzala with its current borders, with a depth ranging from 1-2 km west of the lake in front of Damietta, to 14 km along the eastern bank of the Nile River from Awlad Hamam south of Damietta to Dahra, then Rawda, then Al Rahamna, and from there to Al Sarw (Farescor Center); at a depth ranging from 6.5 to 8 km south of Lake Manzala from Al-Sarw to Al-Manzala; at a depth of 47 km southeast of Al-Manzala Lake, from Al-Malaria to Ezbet Al-Rayes and then to Bar Al Sharqiah, from there to Tell Sanhour and then to Qassas Al-Sharq and Kabriteh to Al-Qantara West and then to Al-Ballah (northeast of Sharqiah Governorate, and north of Ismailia Governorate), and at a depth ranging from 9 to 10 km east of Lake Manzala along the western bank of the Suez Canal, from Al-Qabouti in the north (south of Port Said Governorate) to Al Cap in the south (north of Ismailia Governorate) (Figs. 22a-f).

The sources of threat to all of these areas are the sand belt that separates Lake Manzala from the sea, especially that part which extends between the village of Manasra in the southeast and Ezbet Al-Burg in the northwest, and the western bridge of the Suez Canal extending from south of Al-Qabouti to Tina, as well as the Manzala Lake itself, as the rise in the level of the lake surface due to the direct sweep will cause the lake to flood the surrounding land. If we take into account that the water level in the Suez Canal is higher than that of the Mediterranean, the exploited lands between the canal and Lake Manzala will face a greater threat from the eastern side than from the western side. This threat includes all the dried-up lands of Lake Manzala that are currently exploited along the western coastal strip of the canal between Port Said in the north and Al-Ballah in the south. The gaps are concentrated in the western coastal strip of the canal extending south of the Sahara to Ras Al-Ush, in addition to some sporadic gaps south of Ras Al-Ush to Al- Cap (Figs. 23a-f). These gaps range

from zero (sea level) to -2 meters below sea level. It is also noted that the size of the gaps will expand in the event of a rise in the level of the canal surface by a range of 50 cm to 100 cm to include about 4.8 km along the distance between the Sahara and Ras Al-Ush (about 12 km in length) due to the lower level of the western bridge of the canal than a meter above sea level in these distances, in addition to gaps of 2 km length in the distance between Ras Al-Ush and Al Cap. However, the last gaps are not activated unless the level of the canal surface increases by a meter.

One of the dangers of the sea water invading Lake Manzala is threatening the course of the Nile (Damietta Branch) north of Damietta City. The eastern bridge of the Nile extending from Ezbet Al-Burg in the north to Damietta in the south, its level usually does not exceed 2.0 meters above sea level and is interspersed with low gaps of less than a meter above sea level. Hence, a rise in the level of the surface of Lake Manzala by up to a meter will result in the emergence of four low gaps that permeate the eastern bank of the river between Ezbet Al-Awba and Ezbet Al-Sheikh Dargham, and between Ezbet Al-Sheikh Dergham and Ezbet Al-Ratmah, and between Ezbet Al-Ratmah and Ezbet Tabbal, and between Ezbet Tabbal and Ezbet Al-Khayat from the diameters range from 400 to 600 meters, will act as water passages between Lake Manzala and the Nile River. In the event that this level rises by an amount ranging from 1.25 meters to 1.5 meters, the connection between the Nile River and Lake Manzala will be comprehensive.




Fig. 21a: A brief topography of the northeastern part of the delta extending on the Mediterranean coast between the Suez Canal in the east and the Nile River, the Damietta branch in the west, including Lake

and the Nile River in the south, and the impact of that invasion on the Nile River facing Ezbet Sheikh Dargham. Fig. 22b:- A brief topography of - the city of Damietta and its relationship to Lake Manzala – in the case of sea level rises by a maximum of one meter. The yellow lines represent the natural lines of defense for a city Damietta, where the ground level rises more than 3 meters above sea level. The completely red areas are the areas threatened by an invasion if the sea invades the lake, and they are low, submerged and dry (or dried) areas, their level ranges from -4,0 meters below sea level to one meter above sea level. Note that the black areas confined between the yellow lines (the natural defense line) and the red area are lands often exploited in agricultural activities, and they represent a buffer belt between the lake and the city of Damietta. Figs. 22c-f: Brief topography of the western part of Lake Manzala and the low plains extending to the west along the eastern bank of the Nile River between Damietta in the north and El Rahamna in the south (Fig. 22 c), southwest of the lake to the eastern bank of the Nile River from Al-Rahmana - Kafr Al Arab in the north to Mit Salsil - Old Barnabal in the south (Fig.22d), southwest of Lake Manzala along the Nile River from Seif El-Din - Taftish Al-Saru in the north to Mit Salsil - Al-Draksa in the south (Fig.22e), and the southern part of Lake Manzala south to San Al-Hajar and southwest to Al-Draksa (Fig.22F) - when the sea level rises by a maximum of one meter, and then the invasion of Lake Manzala and its level rise.




Fig. 23a: A brief topography of the wet and dry coastal plains that lie on both sides of the Suez Canal in the northeastern part of the delta and the northwestern part of Sinai, when the sea level rises by a

maximum of one meter - showing the locations of the gaps that permeate the western bank of the Suez Canal between Al-Qabouti in the north and Al-Ballah in the south, amounting to 21 A gap ranging in diameter from 200 meters to 800 meters (yellow circles), which in turn will act as direct sea lanes between the canal and the low plains to its west if the level of the canal rises - unless due protection measures are taken. The black areas are the areas whose level rises more than one meter to 20.0 meters above sea level, while the blue areas are the dry and wet lands whose level does not exceed one meter above sea level. Figs. 23b-f: Detailed topography of the coastal strip of the (Western bank) extending between the Alexandria-Port Said road in the north (south of Al-Qabouti) and Ras El-Ush in the south (Fig. 23b) - just north of Ras El-Ush, Fig. 23c), around Ras El-Ush (Fig. 23d), between Ras El-Ush in the north and Tina in the south (Fig.23e) and between the Sahara in the north and the Al-Cap in the south (Fig. 23f) - if the sea level rises and then the Suez Canal by an amount not exceeding one meter. Note the location of numerous low gaps between the thin sandy bridge separating the canal from the western plains (red circles), which in turn will act as sea passages between the canal and the low plains to its west when the level of the canal rises. Note also that the effect on the eastern bank of the canal is very weak.

It is worth noting that the dry and wet low lands threatened by direct or indirect marine invasion through Lake Manzala in the northeast of the delta are interspersed with many cities, villages, manors, hamlets, roads, bridges, dunes, sandy hills and agricultural lands whose level rises more than a meter above sea level. These cities and villages have a total area of about 924 km². Although these heights are not threatened by invasion or inundation due to their high level, they are threatened by isolation and siege in the midst of the invaded areas. Among the cities, villages and estates threatened with isolation and total siege: Port Said, Al Qantara Al Gharbia, Al Ballah, Al Cap, Tina, Ras El Ush, Al Dahra, Al Atawi, Taftish Al-Sarw, Ezbet Al Rayes, Al Basarta, Awlad Hamam, Ezbet Al Burj, and all the tourist and residential villages within the borders of Lake Manzala. Thus, the maximum area that will be affected by the damage in the northeastern part of the delta as a result of rising sea levels by no more than one meter is about 2552 km².

There are no topographical problems threatening Port Said, although it is surrounded by the sea in the north, the Suez Canal in the east, and Lake Manzala in the south and west. The city is built on high lands ranging between 2.5 meter and 22.5 meters, with an average of 5-10 meters above sea level, interspersed with some depressions that are less than 2.5 meters above sea level. The sea level rise does not affect the city in general by a meter, with the exception of its northern beaches, and some of its southwestern beaches overlooking Lake Manzala, in which the width of the coastal part exposed to the invasion reaches about 150 meter in the northeast and 250 meters in the southwest (southwest of the international park), as well as some industrial and residential facilities built on the dried lands of the lake in Al-Qabouti, south of Port Said. The city of Damietta is also built on high lands and is isolated from the northern coast by a thick belt of sand dunes that only the port of Damietta penetrates. It is also isolated from Lake Manzala by heights of not less than 3 meters above sea level (Fig. 22b). The city is only threatened by a rise in the level of Lake Manzala by no less than a meter, as water can flood the southern outskirts of the city at Awlad Hamam and from there to Farescor on the eastern bank of the eastern Nile branch.

3.2.2. The northern part of the delta: Burullus Lake and the surrounding low plains

The vast majority (about 62%) of the dry lands extending to the south, east and west of Lake Burullus, north of latitude 30° 18.75' and between longitudes of 30° 33.75' and 31° 3.75', with an extension southeast to Hamoul, and north-east up to Baltim village, west to Rashid, and southwest to Sindion on the eastern bank of the Nile - they are low lands whose level ranges between -4.0 meters below sea level and one meter above sea level. The area of dry lands lower than sea level, which surrounds Lake Burullus from all directions, is connected to it or within it, is about 445 km², in addition to the submerged area of Lake Burullus, which is 394 km². All of these wet and dry lands (839 km²) represent the minimum areas threatened by the invasion if the sea level rises by any amount more than the current sea level. As for the total area of lands whose level does not exceed one meter above sea level and which are subject to invasion if the sea level rises by up to a meter, 1375 km², i.e. an increase of 536 km² of dry lands which 479 km² are dry lands outside the framework of Lake Burullus, and 57 km² are dry lands that permeate the submerged area inside Lake Burullus (Figs 24a-b).




Fig. 24a: A brief topography of the northern coastal part of the delta (east of the Rashid branch) extending from Al Hammad in the east to Ezbet Abu Khashaba in the west, when the sea level rises to a maximum of one meter - showing the dry and wet low areas threatened by direct invasion from the sea or indirect invasion through Lake Burullus. The yellow lines represent the general boundaries of the affected areas, amounting to 1845km², including the sand belt of Lake Burullus. The blue areas within the yellow borders are dry and wet lowlands whose level does not exceed one meter above sea level, and threatened by direct invasion from the sea, or indirect invasion through Lake Burullus, which is 1375 km², while the rest of these areas, amounting to 470 square kilometers, are cities, villages, dry lands and sand dunes whose level rises more than a meter above sea level, interspersing the depressions. Many of these villages and cities are not exposed to invasion, but all are threatened by siege and isolation as isolated islands within the areas threatened by the sea invasion. The black areas are the areas whose level rises more than

one meter to 20.0 meters above sea level. The black areas are the areas whose level rises more than one meter to 20.0 meters above sea level. Fig. 24b: Satellite image of the same northern coastal part of the delta whose topography is given in Fig.24a.

The threatened lands extend southeast of Lake Burullus with a depth of up to 38 km to the east, south of Hamoul in Markaz Biella; east of Lake Burullus; with a depth of up to 28 km, to the eastern borders of Markaz Biella, west of Stamony; south of Lake Burullus; with a depth ranging between 10 to 25 km, until the Abbasia - Ezbet Umm Sen in Markaz Kafr El-Sheikh, and Al-Qasabi - Minshat Abbas in Markaz Sidi Salem; west of Lake Burullus, at a depth of 13 km to Rashid; southwest of Lake Burullus, at a depth of 21 km, up to Sindion in Markaz Fowa; and north of Lake Burullus within the sandy belt of the lake between Tell El Magluba in the east and Kom Mishaal in the west, with a depth ranging from 1.0 km to 4.5 km (Figs. 24a-b). These areas include villages, cities, roads, bridges, lands and sand dunes whose level rises more than a meter above sea level, with a total area of 470 km², and these areas will be completely or partially besieged and will turn into separate islands within the invaded low areas. Hence, it can be said that the maximum area of land that will be damaged in the north of the delta if the sea level rises by a maximum of one meter is 1845 km² (Table 2). These lands do not include the threatened area north and east of the Rosetta estuary, which is 8.8 km², or the threatened coastal areas extending east of the tower to Ras al-Bar (see below). This portends many dangers facing the north of Kafr El-Sheikh and the north of Daqahlia Governorate - unless measures are taken to protect the beaches and isolate the lake completely from the surrounding lands.

The most important of these risks is the sea water invasion of the lands of the northern part of Markaz Fowa through the coastal sandy belt that extends between the Green Island and Kom Mishaal (Fig. 25a). As we mentioned, this belt is completely dilapidated (about 13.5 km long), and its width ranges from 1.3 km to 5.0 km, with an average of 4.0 km or a little more, and a total area of about 64 km². It is considered the most dangerous source of threat in the northern part of the delta, and in its current form, it is unable to repel any incursions of sea water from the north, and it needs complete protection along the aforementioned extension. The danger is not limited to the invasion of the agricultural plains located to the south of this belt only, but also extends to include the invasion of the course of the Nile River (Rasheed Branch) north of the city of Fowa. The sand dunes within the belt are scattered and far between and do not form insulating chains, and their height does not exceed one meter above sea level in 28% of the total area of the belt, and two meters in 12%, and more than two meters in 7% of the area of the belt. These dunes are interspersed with low, branched and intertwined gaps below sea level ranging from -2.0 m to -5.0 m (with a maximum of -7.5 or less near Kom Mishaal) constituting between them about 53% of the total area of the belt (about 64 km²).

The deterioration of the coastal sand belt extending between the Green Island and Kom Mishaal is due to the continuous migration of sand dunes towards the southeast and the failure to replace them with new dunes since the establishment of the High Dam, in addition to the human dredging operations of the coastal dunes. The area of the gaps increases to about 81% of the total area of the belt if the sea level rises by a maximum of one meter. These gaps are directly connected to the sea, where it was possible to identify about 11 small, contiguous gaps along the shore line between Green Island and Kom Mishaal, with a total diameter of about 11.5 km, or at least 82% of the length of the belt (Figs. 9a-b). These gaps will act as water passages between the sea and the southern low plains and extending to the borders of the city of Fowa - which leads us to consider this entire belt as a single gap with a diameter of about 13.5 km.




Fig. 25a: A brief topography of the northwestern part of the delta extending from Lake Burullus in the east to Abu Qir Bay in the west - in the case of a maximum sea level rise of one meter, showing the locations of the low-lying gaps that cross the eastern (or western) bank of the Nile River, the Rashid branch between Izbat Abu Khashaba in the north and Fowa in the south, and its relationship to the great gap that extends along the northern coast with a length of 13.6 km from Kom Mishaal in the east to the green island in the west. The black areas are lands whose level rises more than one meter above sea level, while the blue areas are dry (or dried) or wet low lands (Burullus Lake) whose level does not exceed one meter above sea level. The small yellow circles are low gaps numbering 33 gaps whose diameters range from 200 to 800 meters, including 30 gaps that permeate the eastern bank of the river between Ezbet Abu Khashabah in the north and Fowa in the south, in addition to 3 gaps on the West Bank between Hammad in the north and Mahmoudiya in the south. **Figs. 25b-e:** Detailed topography of the gaps (yellow circles) that permeate the eastern bank of the western branch of the Nile in front of Rashid (Fig. 23b), between Jidiyah and Breda (Fig. 25c), between Minyat al-Murshed and Ibyana (Fig. 25d), and between Motobas and Shamashira (Fig. 25e). The white lines are the current river shore lines.

On the other hand, the eastern bank of the Nile River between Abu Khashaba point (south of the Green Island) in the north and Fowa in the south, its level ranges from -1.5 meters below sea level to 2 meters above sea level, with the exception of a few limited points in front of Rashid, at Motobas and Minyat Al-Saeed and Dayrut, where this level rises more than 2 meters to 12.5 meters above sea level. The low gaps in the eastern bridge of the river abound between Abu Khashabah and Rasheed point, between Ezbet Al-Waqf and Buraidah, between

Ezbet Jazirat Al-Fars and Ezbet Al-Tijariyyin, between Minyat Al-Murshed and Ibyana, from Shamashira south of Motobas to Sindion, and Ezbet Ahmed Al-Tayeb north of Fowa (Figs. 25b-e). All of these gaps will act as waterways between sea water and river water in the event of the sea's invasion of the low plains extending between Rashid and Lake Burullus, if the sea level increases by a meter. In the event that this level rises by two meters, the sea water will sweep the vast majority of the eastern bank of the Rashid branch between Abu Khashabah in the north to Fowa in the south, and then the Rashid estuary will retreat in the south until Fowa.

3.2.3. The northwestern part of the delta: Idku and Mariout lakes and the surrounding low plains

The topography of the southern coastal strip of Abu Qir Bay portends a sea invasion of this strip, and then the flooding of vast areas of the dry and wet terrestrial plains located in the south of the Bay in case the global sea level rises, not only because of the rise in sea level, but also because of the drop in the level of these plains below sea level (Fig. 15a-b). Therefore, an increase in the sea level by a maximum of one meter would cause the sea to invade the southern exploited coastal strip of Abu Qir Bay extending from Maadiyah in the east to the suburb of Abu Qir in the west, whose level does not exceed one meter above sea level by 83% of its area. This, in turn, will lead to the invasion of the sea for about 2515 km² of the plains area extending south of Abu Qir Bay to the desert noon, approximately 88.8% of the total area of these plains (2,834 km², of which 1054 km² are north of the Mahmoudiya Canal and 1780 km² south of Mahmoudiya Cana, Ouda, 2022). The area threatened by the invasion is 2355 km² of dry land and about 160 km² of already submerged lands (the rest of the Idku and Mariout lakes), which is an increase of 344 km² over the lands whose level does not exceed zero. These areas include about 908 km² of dry (or dried) and wet lowlands extending south of Abu Qir Bay, between Rashid branch in the east and Alexandria in the west, until Mahmoudia Canal in the south (86% of the total area), whose level does not exceed one meter above sea level with an increase of 154 km² over the lands whose level is below sea level (Figs 26a-b and 27 a-b).

This is in addition to the invasion of an area of 1607 km² located between the Mahmoudiya Canal in the north and the desert valley in the south, including the submerged areas in Lake Mariout, south of Alexandria, and its western extension to Al-Hammam (137 km²). These lands range from 1 meter above sea level (190 km²) to -8 meters below sea level (1417 km²) at approximately 90% of their total area. They are separated from Abu Qir Bay only by a narrow strip of land of intermittent and scattered hills, not homogeneous in height or transverse extension, extending between Damanhour in the east and Alexandria in the west, and on which several cities and villages (such as Abu Homs, El-Karion, Kafr El-Dawwar, Kafr Selim, King Othman, Al-Bayda and Al-Khadra) are built on it (Figs.28a). It is interspersed with Mahmoudiya Canal and Damanhour-Alexandria Agricultural Road. The width of this strip ranges from 200 meters to 2000 meters.

The Mahmoudiya Canal is also interrupted, and loses its characteristics, due to the heterogeneity of the level of the course of the canal, as it takes place on a strip of land of varying level, often ranging in height between - 3 meters (below sea level) and + 5 meters above sea level. The in-between depressions that permeate this strip, the level of which is less than 3 meters below sea level, are concentrated in some locations, such as the area between Ezbet Al-Shawish, east of Al-Karion and the Al-Ghaba, the area between Al-Akrisha and Al-Karion, between King Osman and Kafr Selim, between the village of Al-Khadra and Al-Bayda, and at the upper bridge in the Al-Khadra area, the Nubar farms and the Khurshid farms - which makes the bridges of the Mahmoudiya Canal that penetrate these depressions insufficient in height to repel any marine invasion from the north in case the global sea level rises by a meter (Figs. 28a-b and 29a-f). The cities and villages located on both sides of the Mahmoudiya Canal, some of them are built on high levels ranging between 3 and 8 meters above sea level, such as Kafr Al-Dawwar, Al-Karion and Abu Homs, and others are built on lower levels ranging between - 3 meters and + 3 meters, such as Al-Bayda and Al-Bayda dyeing company, Al-Khadra, Bardala, King Osman (Figs.29a-d) and some villages whose level does not exceed zero, such as Ezab Khurshid, Ezab Nubar and Akarisha (Figs. 29e-f). In addition, all modern expansions outside the original cordon area designated for villages and cities are located in low-lying lands whose level ranges between zero and -5 meters below sea level, for example the expansions of the Al-Bayda dyeing company, the city of Al-Bayda, Kafr Selim, Al-Khadra and Kafr Al-Dawar (Figs. 29a-c).

The dry, low lands threatened by flooding include most of the agricultural basins extending between Lake Idku and the city of Alexandria, north of the Mahmoudiya Canal (Figs. 26a-b and 27a-b). These lands include the basins of the Abu Qir Qumpaniya and the former English Qumpaniya, the Minshat El- Bahariya, the south of the city of Abu Qir, the south of the neighborhoods of Montazah, Maamoura, and Sidi Bishr al-Qibli, including the new Obour city, the Montazah economic housing neighborhood, Al Tawfiqia, Ezab Dafshaw, Ezab Nubar, Ezab Khurshid, Al Khadra, and Al Bayda (Ouda, 2022). Likewise, most of the basins located south of Idku Lake are

in the Minshat Boleyn, the churches, Ghattas Lake, Nakhla El- Bahariya, Basantaway and Sekinda (Figs. 30a-b). Also, the basins located west of the Nile River, between the village of Al-Atef in the south and Jidiyah in the north (north of Mahmoudiya Center and South Markaz Rashid), northeast of Damanhour and south of Mahmoudiya Center, between Aflaqa-Zarqun in the west and Ezbet Al-Awam-Atef in the east. This is in addition to the entire agricultural area located south of the Mahmoudia Canal to the border of the desert back (Al-Nubaria Canal), which extends from the western port of Alexandria in the west to the Zawiya of Hammour-Damanhour in the east. This area includes all the agricultural basins located south of Kafr El-Dawwar, south of Abu Homs, west and southwest of Damanhour and most of the center of Abu El Matamir and south of Alexandria, including the remainder of Lake Mariout (Figs.28b and 30a-b) Hundreds of villages and cities built on high levels of more than one meter to 8 meters above sea level (a total area of 305 km²), which permeates all these agricultural basins, will turn into isolated islands such as Kafr Al-Dawwar, Al-Karion, Abu Homs, Hosh Issa, Abu Al-Matamir and others, unless protection measures are taken.




Fig. 26a: A brief topography of the coastal strip extending between Idku in the east and the village of Tarh in the west along the southern coast of Abu Qir Bay, including the remains of the current Idku Lake in the event of a sea level rise of a maximum of one meter, explaining the relationship of the coastal gap extending from Al-Maadiyah in the east to the suburb of Abu Qir in the west (yellow circle) with the dry and humid low plains south and southeast of the bay. The white line is the current beach line. The black areas defined by yellow lines are sandy hills that are more than a meter above sea level, and they form a continuous chain between Al-Maadiyah in the west and Idku in the east, where it acts as a natural buffer

between the sea in the north and the plains in the south, while the blue ground areas are the dry and wet lands (Idku Lake) which are threatened by direct invasion of sea water through the wide gap which will act as a wide sea corridor between Abu Qir Gulf and the plains extending south and southeast of the Gulf. Fig. 26b: Satellite image of the same coastal strip extending between Idku in the east and the village of Tarh in the west, along the southern coast of Abu Qir Bay, whose topography is given in Fig. 26a, including the remains of the current Idku Lake and the wide coastal gap extending from Al Maadiyah in the east to the suburb of Abu Qir in the west (the yellow circle).




Fig. 27a: A brief topography of the northwestern part of the delta extending from the Rashid branch of the Nile in the east to Alexandria in the west, and from Abu Qir Bay in the north to the Damanhour-Alexandria road in the south - if the sea level rose by a maximum of one meter, showing the boundaries of the low land area (dry and wet) north of the Damanhour-Alexandria line, which is threatened by a direct invasion from the waters of Abu Qir Gulf (the continuous red line) across the wide gap extending from the suburb of Abu Qir in the west to Al-Maadiyah in the east (the red circle). These lands are below sea level in about 72% of their total area, while 15% of their area rises above Zero to a maximum of one meter above sea level, which means that about 87% of its area is threatened by direct invasion of the Abu Qir

Bay. Fig. 27b: Satellite image of the northwestern part of the delta extending from Rashid branch of the Nile in the east to Alexandria in the west, and from Abu Qir Bay in the north to the Damanhour-Alexandria road in the south whose topography is given in Fig. 27a, showing the low, dry and wet lands, threatened by direct invasion from Abu Qir Bay (the red areas) across the wide gap extending from the suburb of Abu Qir in the west to Al-Maadiyah in the east (the yellow circle). Areas which are bounded by yellow lines (bottom right of the image) represent the areas that are threatened by partial inundation due to groundwater in case its level rises as a sequential effect of sea level rise.





Fig. 28a: A brief topography of the narrow strip of land that extends along the Mahmoudiya Canal between Abu Homs in the east and the entrance to Apis in Alexandria in the west, in the northwestern part of the delta - if the sea level rises by a maximum of one meter. The black areas are lands whose level rises more than a meter above sea level, while the blue areas are low, dry, agriculturally exploited lands,

ranging from - 8.0 meters below sea level to one meter above sea level. The yellow circles are low gaps in the narrow strip of land, numbering 12 basic gaps, and their diameters range from 200 m to 3000 m. These gaps will, in turn, act as auxiliary sea lanes on the flow of the waters of Abu Qir Gulf from the northern plains north of the Mahmoudiya Canal to the southern plains south of the Mahmoudiya Canal - unless protection measures are taken. See enlarged images of these gaps in the panel for an upcoming. Fig. 28b: A brief topography of the western part of the dry (or dried) and wet low plains (Mariout Lakes) extending south of Mahmoudia Canal along the city of Alexandria to Al-Amriya in the northwestern part of the Delta - when sea level rises by a maximum of one meter- explaining the relation of the gaps that permeate the narrow land strip extending along the Mahmoudiya Canal which, and the gap that mediates the western port in Alexandria (4 km in length) between Wardyan and Dekheila, which in turn will act as a sea corridor between the Mediterranean and the southern plains.



Figs. 29a-f: Enlarged figures showing detailed topography of the low gaps that cut through the narrow strip of land which extends along the Mahmoudiya Canal (white line) between Abu Homs in the east and the entrance to Abis in Alexandria in the west- if the sea level rises by a maximum of one meter. Fig. 29a: The big yellow circle is the gap that cuts through the Mahmoudiya Canal, between Al-Bayda and Al-Khadra, with a diameter of about 3000 meters. Fig. 29b: A moderate gap that permeate the low plains around the upper bridge and the ring road in Al Khadra district, with a diameter of 800 meters. Fig. 29c: A brief topography of the big gap whose detailed topography is given in Fig. 29a. The black areas defined by yellow lines are lands whose level rises more than a meter above sea level, while the blue areas of different colors are low, dry lands, exploited for agricultural and residential use, ranging from -7.0 meters below sea level to one meter above sea level. the sea. Fig. 29d: A big gap, about 2500 meters that permeate

the low plains between Berket Ghattas in the east and Karion in the west. The number above the circle is the radius. Fig. 29e: Detailed topography of a series of intermittent sandy hills extending on both sides of the Mahmoudiya Canal, between Karion in the east and Kafr El-Dawwar in the west - showing the locations of 6 gaps with diameters ranging from 200 to 300 meters between the canal and the hills. Fig. 29f: Detailed topography of the entrance to the agricultural city of Alexandria, Damanhour-Alexandria road, showing the locations of the gaps that permeate the Mahmoudiya Canal and the intermittent hills located on its sides. The continuous red circle is the most dangerous gap that connect the northern plains with the southern plains, while the intermittent red circles are gaps that threaten the residential neighborhoods and villages of Nubar. The black circle indicates weaknesses that must be strengthened.




Fig. 30a: A brief topography of the southern plains extending from Damanhour in the east to Alexandria in the west, and from Mahmoudia Canal in the north to Nubaria Canal in the south in the northwestern part of the delta - if the sea level rises by a maximum of one meter. The red lines are the borders of the southern low plains, dry (or dried) and wet, threatened by marine invasion through the gaps that permeate the thin land strip extending along the Mahmoudiya Canal between Abu Homs in the east and

Alexandria in the west (yellow circles), as well as the gap in the middle of the Western Port in Alexandria (4 km in length) between Wardyan and Dekheila. The blue areas are low, dry (or dried) and wet lands (Lake Mariout) with a level ranging from -8.0 meters (and a maximum of -10.0 meters) below sea level to one meter above sea level. Fig. 30b: Satellite image of the southern plains extending south of the Mahmoudiya Canal from Damanhour in the east to Alexandria in the west, whose topography is given in Fig. 30a – explaining the effect of sea level rise by a maximum of one meter. The red areas are the southern low plains, dry (or dried) and wet, threatened by the sea invasion through the gaps that permeate the thin land strip extending along the Mahmoudiya Canal between Abu Homs and Alexandria, as well as the gap in the middle of the western port in Alexandria (4 km in length) between Wardyan and Dekheila.

On the other hand, an increase in sea level by no less than a meter and its invasion of the depressions in the lime belt between Wardian and Dekheila along the middle coast of the western port of Alexandria will lead to invasion of some installations located on low levels in the belt, such as the installations of the Misr Petroleum Company, the Petroleum Cooperative Society, the Nasr Company for tanning leather, the medical stone, the Amriya nursery, and the Petroleum Pipelines Company (Ouda,2022, Fig. 18)). Also, any increase in the level of Lake Mariout - whether as a result of the rush of water from the western port of Alexandria, or as a result of the waters of Abu Qir Gulf sweeping to the southern coast and extending to the south from Abu Qir in the north to Al-Amriya in the south., it will lead to the invasion of all the southern installations located south of Dekheila, Al-Max, Al-Wardyan, Al-Qabbari, Karmouz, and Muharram Bey, and all the residential, industrial, touristic, educational and administrative installations located south of Al-Qabbari Road - Al-Nozha Airport (south of Mahmoudiya Canal). All of these installations are built on dried lands of Lake Mariout, with ground levels ranging between zero and -8 meters below sea level. This is in addition to the remaining parts of Lake Mariout, south of Alexandria, and the agricultural basins extending eastward from Alexandria to Damanhour (Figs.28b and 30a-b).

Thus, the southern coast of Abu Qir Gulf, and the middle part of the western port between Wardyan and Dekheila are the only sources of threat not only to the remaining lakes of Idku and Mariout, but also to the entire northwest of the delta during this century as a result of the certain increase in the global sea level. However, we consider the Abu Qir Gulf as the main source of threat to all the lands north of Beheira Governorate and south of Alexandria Governorate, extending to the desert back, due to its topographical, geological and historical conditions that qualify it for an invasion in the event of an increase in sea level by any amount. It is followed by the Western Port of Alexandria as a second source of threat to the low-lying areas to its south in case the sea level increases by at least a meter.

Therefore, the low, wet and dry areas threatened with invasion in the northwestern part of the delta (west of the Rosetta branch) in case the global sea level rises by any amount more than the current level with a maximum of one meter range from 2220 km² minimum to 2577 km² maximum, of which 160 km² are already submerged. (remains of Lakes Idku and Mariout), and 2,417 km² of dry lands extending south of Abu Qir Bay to the desert back, including all the areas that were drained and exploited south of Alexandria at the expense of the original Lake Mariout (Table 2). All these low lands are located within the governorates of Beheira and Alexandria. These lands are interspersed with villages, cities, high lands, dunes and sand hills whose level rises more than a meter above sea level, with an estimated area of 329 km². Although these cities, villages and lands seem safe from invasion, they are all threatened by sea blockade and isolation as isolated islands within the invaded spaces. Hence, the maximum area that will be damaged by the northwestern part of the delta due to an increase in sea level by an amount not exceeding one meter is 2906 km². This area does not include the low lands that are not connected to surface water sources and are likely to be partially submerged by groundwater south of Fowa, west and south of Rahmaniah, east of Damanhour, around Dillingat, and within the city of Alexandria (the depressions that permeate the neighborhoods of Smouha, Al Seyouf and Abu Qir, Ouda, 2022).

Total	Northeast of the Delta, Damietta, Port Said, Ismailia and Sharqia governorates	North Delta, Kafr El-Sheikh, Dakahlia and Damietta governorates	Northwest of the Delta, Beheira and Alexandria governorates	Threatened wet and dry lands
1031 km ²	477 km ²	394 km ²	160 km ²	Wet areas threatened by invasion whose level is below sea level
2970 km ²	465 km ²	445 km ²	2060 km ²	Dry areas threatened by invasion whose level is below sea level
1579 km ²	686 km ²	536 km ²	357 km ²	Dry areas threatened by invasion, whose level ranges from zero to meters above sea level
Min 4001 km² Max 5580 km²	Min 942 km² Max 1628 km²	Min 839 km² Max 1375 km²	Min 2220 km² Max 2577 km²	The total area threatened by sweeping or extensive immersion
1723 km ²	924 km ²	470 km ²	329 km ²	Cities, villages and lands that are more than 1 meter above sea level and are threatened by maritime blockade
7303 km² (29.81 %)	2552 km² (10.42 %)	1845 km² (7.53 %)	2906 km² (11.86 %)	The maximum area affected by the sea invasion and its proportion of the delta

Table 2: The dry and wet areas threatened by the marine invasion in the northern delta if the global sea level rises by a maximum of one meter

4. Conventional defense policies to be applied

The economic and human losses that may result from rising global sea levels are innumerable. They are not limited to the invasion of the northern lakes in the delta, but go beyond that to include agricultural lands, residential buildings, factories, shops, tourist sites, roads, canals, banks and installations in dry or dried up low lands. surrounding the mentioned lakes.

The effect of sea level rise extends to the increase in the salinity of groundwater. In a study on the interference of sea water in the freshwater aquifer in the Nile Delta by Sharif and Al-Rashed (2001) - it became clear that the continuous unguided withdrawal of groundwater in the delta during the past decades has caused the deterioration of groundwater in northern regions. And that sea water has overlapped in this reservoir with a width of 63 km from the shore along the bottom of the aquifer, and that climatic changes and sea level rise will affect the quality of the aquifer in two directions: First: the low areas around the shore will be flooded with sea water, which will destroy the existing reservoir under these areas; Second: The additional pressure from the sea side will cause more sea water interference in the reservoir.

The reports issued by the Beach Research Institute in Alexandria indicated that the effect of sea level rise extends to include erosion and sedimentation systems, especially in the gorges and estuaries of the two Nile branches, the ecosystems of lakes, and the drainage systems of the northern delta. World Bank experts (2007) estimate the human losses that will result from an increase in sea level by one meter at about 10% of the total population of Egypt. The residents will be forced to leave their homes due to agricultural land losses, which the bank estimates at about 12.5% of the total cultivated area in Egypt in the event of an increase in sea level by one meter. This is with the knowledge that the delta produces about 50% of the basket of crops produced by Egypt. This is in addition to the losses of industrial, commercial, tourism and fishing activities, and the change in the chemistry of groundwater and coastal lake waters - which threatens a catastrophe unless the Egyptian government begins to expedite measures to protect against the impact of climate changes and adapt to the new climatic conditions.

In a detailed statistical study of the risks arising from the rise in the global sea level on the Egyptian coasts, Ouda (2010, 2011, 2012) reported that the coastal lands of the Egyptian Delta (both dry and wet areas) are at risk of invading the Mediterranean Sea if the global sea level rises by a maximum of one meter during the current century. The threatened areas range from 4147 km (17 %) of the total area of the delta as a minimum (coastal lands whose level does not exceed sea level) to 5920 km² (24.2%) of the total area of the delta (including the

Tina plain in northwest Sinai) as a maximum (coastal lands whose level does not exceed one meter above sea level). The low-lying, dry and wet coastal lands exposed to the dangers of sea level rise include cities, villages, manors, roads, bridges, dunes, sandy hills, and dry lands whose level rises more than one meter above sea level, with a total area of 2113 km². These high areas will be subjected to complete sea blockade and isolation as isolated islands within the invaded areas; In addition to the damage that will be caused to agricultural and industrial projects and public utilities established on these lands as a result of the naval blockade. Therefore, the total affected areas in the delta as a result of the sea level rise by a maximum of one meter is estimated by Ouda (2010, 2011) as being 8033 km² at a maximum, at a rate of approximately 33% of the total area of the delta.

Therefore, building defenses must be within the framework of a comprehensive national plan to protect the northern coasts of the Egyptian Delta. The defense plans must include the following considerations.

First: general instructions

1- Completely stopping the construction of new residential or industrial facilities or tourist resorts along: a) the sandy belt extending from the Rashid estuary in the west to the village of Al-Burg in the east, b) the northwest sandy belt of Lake Manzala extending west of the Manasra village known as Barr Al-Dahra, c) the southern coast of the Abu Qir Bay, which extends between the suburb of Abu Qir in the west and Maadiyah in the east, and d) the western bank of the Suez Canal, between Al-Qabouti in the north and Al Cap in the south. As well as stopping any new construction in the agricultural plains south of the suburb of Abu Qir, south of Maamoura, south of Montazah and south of Sidi Bishr al-Qibli, until the protection measures are completed.

2- Resettlement of residents of all residential villages located on the sand belt of Lake Burullus, between Kom Mishaal in the west and Burg in the east, and the northwestern sand belt of Lake Manzala west of Manasra village known as Barr Al Dahra. Also the villages scattered within Lake Manzala and built on dry lands whose level does not exceed one meter above the sea level - for fear of the failure of the defense strategy of these belts as the first lines of defense.

3- Suspension of all expansion decisions in the cordons of cities and villages established in the coastal plains of the delta until after certainty that the level of the required expansion lands will rise above sea level by no less than 3 meters.

4- Obligating the Ministries of Environment and Irrigation to prepare long-term environmental programs to protect the coasts from marine risks, including local risks such as the sharp rise of waves arising from the intensity of storms as a direct impact of climate change, tsunami waves or global risks arising from rising sea levels. Evacuation or warning programs must also be developed to prevent the occurrence of environmental disasters, as well as programs to protect archaeological areas, in partnership with the Ministry of Culture, from climate changes, especially tourist sites located directly on the coasts. All of this will contribute to limiting the effects arising from sea level rise.

5- Working on spreading environmental culture among citizens in compliance with the decisions of the conferences of the parties in Cancun, Mexico. As there are people who remove coastal sand dunes instead of installing them in afforestation and exploit them commercially in construction work (for example: sand dunes north and east of Idku, between Idku and Al Busaili, and along the coast of the southern Bay of Abu Qir without taking into account the strategic importance of these dunes" which confirms The lack of environmental culture for the inhabitants of the Egyptian coasts of the Delta. So we recommend the criminalization of sand dune removal and the financial encouragement of all afforestation projects along the coasts.

6- Planning to build new cities along the northern coast of the Western Desert. The northern coast extending from El Alamein to El Dabaa is characterized by a sharp rise above sea level, ranging in height from 5 meters to 25 meters south of the shore, with a width ranging from 6 to 11 kilometers, which guarantees the protection of all cities built on it.

Second: Building coastal defenses

These include building concrete walls facing the beaches, coastal bridges to protect the agricultural plains south of the coast; strengthening the dilapidated dune belts separating the sea from the northern lakes and the low lands that surround them through afforestation; consolidating and raising the level of the existing bridges of the northern lakes; and strengthening the eastern low bridges in the northern part of the Nile River (Rasheed Branch) between Fowa and Rashid, and the western bridges of the northern part of the Suez Canal between Al-Qabouti and Al-Cap. This is the traditional solution followed by all coastal countries threatened by the dangers of sea encroachment. An increase of one meter in the level of the seas and oceans would sweep an estimated area of

194,000 km in 84 developing countries, which would turn about 56 million citizens in these countries into refugees. Egypt, Vietnam and the Bahamas are among the developing countries affected by rising sea levels (World Bank, 2007).

Therefore, these countries, the most important of which is Egypt, must immediately plan measures to defend the coasts. The bodies that develop strategies should set their defense priorities based on the information contained in this research and other local and international reports issued in this regard. The success of these defenses does not depend on the state's economics as much as it depends on the state's experiences in the field of environmental protection and maintenance. A country like Costa Rica is ahead of many developing coastal countries because it has given more attention to its environment in the past by protecting its agricultural lands with natural reserves, and it is also able to face future challenges such as improving evacuation operations when tornadoes occur with the intensification of storms. Kyrgyzstan is also ahead of Saudi Arabia in the field of improving water resource management and the existence of environmental programs that enable it to adapt and adapt to climate changes - despite the fact that per capita income in Saudi Arabia is 7 times the per capita income in Kyrgyzstan.

It requires a sophisticated environmental management that makes it able to take the necessary measures to create defenses against rising sea levels. These defenses do not stop at the construction of coastal bridges and sea walls, and the strengthening of dune belts, but also to be able to empty the sea water that sweeps the cities and dispose of it away from the sea. Egypt should seek the assistance of countries that have the capabilities to adapt to the increase in global warming, such as the Netherlands, when building defenses. The Netherlands has spent centuries adapting to the lands below sea level, and it has scientific talents in this regard and great experience on the impact of sea level rise on existing dams.

The following are the proposed defenses to protect the northern lakes and the surrounding low plains in the north, northeast, and northwest of the delta from the dangers of marine invasion in the event of the global rise in sea level as one of the direct consequences of climate changes that the world is witnessing this century:

1-: erecting submerged armored cement walls capable of repelling any invasion or marine intrusion into the ground in the sites where the low coastal gaps are spread. These walls are erected along shore line facing the sea in coastal sites that are not separated from the sea by strong sand belts and low sites that can be invaded, resulting in the flooding of vast areas of lands south of them. Armored walls mean concrete walls made of sand and chemically treated cement, in order to achieve the highest degree of hardness and cohesion, and to ensure complete adhesion of sand grains used in concrete facing the sea. It is required that these walls submerge below sea level at a depth ranging from 3.0 meters to 5.0 meters through soil injection operations to prevent subsurface leakage, and that they rise above sea level by no less than two meters - that is, with a total height ranging from 5.0 meters to 7.0 meters. As for the thickness of these walls in each site, it is left to the specialized engineers in proportion to the length of the gap and its extension along the shore line, and to be able to resist the pressure of the excess sea waters.


The locations, dimensions and depths of the submerged armored walls required to be erected along the northern arc of the delta were determined as follows:

- A submerged reinforced concrete wall directly in front of the shoreline facing the sea along the northern coast of Lake Manzala (Figs. 31a and 32) in the northeastern part of the delta east of Ezbet El Burj; with a length of about 8.0 km, a depth of not less than 5.0 m above sea level, and a height above sea level of not less than 2.0 m (total height of 7.0 m).
- A submerged reinforced concrete wall in front of the shoreline directly facing the sea along the southeastern coast of Lake Manzala known as Barr al-Dahra in the northeastern part of the delta (Figs.31b and 32), with a length of about 7.4 km, a depth of no less than 5.0m in sea level, and a height above sea level of not less than 2, 0 m (total height 7.0 m).
- 6 submerged armored concrete walls at spaced distances in front of the beach line directly facing the sea along the southeast coast of Lake Manzala extending from the village of Manasra in the east to Barr Al-Dahra (Figs. 31c-d and 32) ranging in length from 500 meters to 1600 meters - with a total length 5.2 km, depth below sea level of not less than 5.0 m, and height above sea level of not less than 2.0 m (total height of 7.0 m for each wall).
- A submerged reinforced concrete wall in front of the shoreline directly facing the sea along the northern coast of the delta, which extends between Kom Mishaal in the east and the Green Island (north of Ezbet Abu

Khashabah) in the west (Fig. 33), with a length of about 14.0 km and a depth of at least 5.0 meters sea level and a height Above sea level not less than 2.0 meters (with a total height 7.0 meters).

- 6 submerged reinforced concrete walls at spaced distances directly in front of the shoreline facing the White Sea along the northern coast of the delta extending between Kom Mishaal in the west and Al-Burg in the east (the sand belt of Lake Burullus, Fig. 33). Their lengths along the shore line range from 500 m to 10.5 km, with a total length of 22 km, a depth below sea level of not less than 5.0 m, and a height above sea level of not less than 2.0 m (total height of 7.0 m). They are as follows:
 1. A submerged reinforced concrete wall extends between Kom Mishaal in the west and Kom Mastrowa in the east, with a length of 5 km to fill 6 gaps ranging in diameter from 700 meters to 1000 meters, interspersing the western part of the sand belt of Lake Burullus.
 2. A submerged reinforced concrete wall at Qada'a Island, with a length of 4.0 km, to fill two gaps, their diameters range from 700 meters to 3000 meters, interspersing the middle part of the sandy belt of Lake Burullus.
 3. A submerged reinforced concrete wall at Tell Al-Maqluba with a length of 1,000 meters to fill a gap that cuts through the eastern part of the sandy belt of Lake Burullus, north of Tell Al-Maqluba. 1000 meters in diameter.
 4. A submerged reinforced concrete wall east of Tel Maqluba, with a length of 1,000 meters, to fill a gap that permeates the eastern part of the sandy belt of Lake Burullus, east of Tel Maqluba, with a diameter of 1,000 meters.
 5. A submerged armored concrete wall in the middle of the distance between Al-Burg in the east and the Maqluba hill in the west, with a length of 5,000 meters, to fill a gap that permeates the eastern part of the sandy belt of Lake Burullus, with a diameter of 5000 meters.
 6. A submerged reinforced concrete wall, directly west of the Burullus Lake, with a length of 500 meters, to fill a small gap that permeates the far eastern part of the sandy belt of Lake Burullus, with a diameter of 500 meters.
 - A submerged reinforced concrete wall in front of the shoreline directly facing the sea on the southern coast of Abu Qir Gulf, which extends in the northwest of the delta between the suburb of Abu Qir in the west and the village of Maadiyah in the east (Fig.34a), with a length of about 10 km, a depth below sea level of not less than 5 meters, and a height above sea level of not less than 2.0 meters (with a total height of 7 meters).
 - A submerged reinforced concrete wall in front of the beach line directly facing the sea on the coast of the Gulf of Max in Alexandria, between the Western Port in the east and Dekheila in the west (Fig. 34b), with a length of approximately 4.5 km, a depth of 5.0 m above sea level, and a height above sea level of not less than 2.0 m, (with a total height of 7.0 meters)
- 2- erecting cement ground bridges south of the shoreline, with a height of no less than 2.0 meters above sea level with flat surfaces and inclined sides to the north and south extending along the coast, or using existing roads after raising their level to 2.0 meters above sea level to take the place of these bridges as a second line of defense in the following locations:
- Exploitation of the international coastal road extending inside Lake Manzala between Damietta in the west and Port Said in the east as a land bridge extending south of the sandy belt of Lake Manzala, with a length of about 40 km along the shore line. This bridge is intended to strengthen the defense lines for the majority of Lake Manzala and the surrounding plains, as the road level during this path usually ranges in height between 1.0 and 2.5 meters above sea level, which makes it one of the strongest ready-made ground defense lines after raising the level of the minor gaps that cross it (Fig. 32). Hence this road can help in repelling any invasion from the sea side in the event of the partial failure of the first line of defense to confront the invasion.
 - Exploitation of the international coastal road extending from Al-Jediyah in the west to Al-Burg in the east as a land bridge extending south of the beach line within the sandy belt extending from Ezbet Abu Khashaba in the west to Al-Burg in the east along the northern coast of the delta (Fig. 33). The road level should be raised by no less than 2.0 meters above sea level. This bridge is intended to strengthen the defense lines of Kafr El-Sheikh Governorate, as the failure of the first line of defense for any reason will result in the invasion of Lake Burullus and all the dry plains surrounding it. It is possible to dispense with the heightening of the part of this road between the village of Al-Burg in the east and Tell Al-Maqluba in the west (about 14 km in length), due to the reduced thickness of the sandy belt of Lake Burullus during this distance and the approach of the first and second lines of defense to each other. It is also possible to exploit the intermittent dune chains

south of the shore line along the sandy belt of Lake Burullus extending from Kom Mishaal in the west to Al-Burg in the east as a second line of defense instead of the international coastal road, by installing the dunes by afforestation to prevent their migration while filling the gaps that permeate them.



Figs. 31a-c: Satellite image of the northern and the northwestern-southeastern sandy belts of Lake Manzala showing the proposed lines of defense. The white lines are the first lines of defense. They are represented by one submerged armored concrete wall directly in front of the shore line towards the sea, with a length of about 8.0 km along the northern sandy belt extending east of Ezbet Al-Burg (Fig.31a), one submerged armored concrete wall of 7.4 km long along the northwestern-southeastern sandy belt (the northwestern part of Barr El-Dahra) (Fig. 31b), and six submerged armored concrete walls ranging from 500 meters to 1600 meters, with a total of 5.2 km along the southeastern sand belt of Lake Manzala - extending from the village of Manasra in the east to Barr Al-Dahra in the west (Figs. 31c-d). All walls should submerged below the current sea level for depth of not less than 5 meters, and raised above the current sea level of not less than 2 meters (total height 7 meters).

- Establishment of a land bridge with a length of 10.5-11.0 km extending directly south of Al-Tabia-Rashid Road from the south of the suburb of Abu Qir in the west to Qantara El Maadiyah in the east, or using the same road as a bridge after raising its level in the low points to 2.0 meters above the sea level (Fig. 34a). This bridge is intended to strengthen the defense lines of Beheira Governorate, as the failure of the first line of defense for any reason will result in the invasion of all the plains extending south of the Gulf of Abu Qir until the desert back.
- Establishment of a land bridge extends south of the city of Alexandria along the current northern Lake Mariout shore line “from the borders of El Max in the west to East City Center and then turns around Al-Nozha Airport along the Ring Road” and from there to the northern borders of the Airport Lake, where it deviates east to Ezb Khurshid near the entrance to Apis, with a total length of 22.0 km and a height not less than 2.0 meters above sea level. This bridge is intended to protect the residential, industrial, educational and social constructions and public utilities that were erected in the south of Alexandria on drained lands of Lake Mariout from the rise in the level of the lake in the event of sea water leakage to it for any reasons. A part of the path of the bridge required to be built applies to the axis of the High Dam south of Al-Qabbari (Fig. 34b); the second part on the international road south of Moharram Bey. The third part applies to the Ring Road south of the Airport and City Center, where these roads can be exploited after raising their level to complete the bridge required to be built (Fig. 35).




Fig. 32: Satellite image of the coastal strip of Lake Manzala, extending from Port Said in the southeast to Ezbet Al-Burg in the northwest on the Mediterranean coast. Showing the second line of defense (the red intermittent line), represented by the international coastal road, with a length of about 40 km, after raising the level of parts of it to 2 meters above sea level. This is with the knowledge that the current road surface level ranges in the general average from one to 2.5 meters as shown in the attached diagram, which makes the treatment simple along the road path.

3-: Strengthening and raising the level of the existing lake bridges so that their height becomes not less than 2.0 meters above sea level in the following places:

- Strengthening and raising the level of the eastern, southern and western Manzala Lake bridges to a height of no less than 2.0 meters above sea level after separating the southern part used as a sewage complex from the lake - with a slight modification to the path of the western bridge in line with the current reality of the current lake. The length of these bridges, as the average, is about 101 km, and their current level ranges from zero to 2.0 meters (Fig. 36), meaning that the average height above the surface of the earth will not exceed one meter along the path of these bridges, with the exception of some parts of the eastern bridge near the city of Port Said, where it needs to be elevated about 2.0 meters above sea level. Raising the level of these bridges will prevent the flooding of the lake and its invasion of the surrounding plains.




Fig. 33: Satellite image of the coastal strip extending from Al- Burg in the east to Ezbet Abu Khashaba (north of Rashid) in the west on the Mediterranean coast, including the sandy belt of Lake Burullus - showing the proposed defense lines for the lake and the dry (or dried) low plains surrounding it. The continuous white lines represent the first line of defense of the lake, represented by 7 submerged armored concrete walls in front of the shore line directly facing the sea. The length of walls range from 500 meters

to 14 km with a total length of approximately 36.0 km. They should be submerged with a depth of 5.0 meters below sea level, and a height of 2.0 meters above sea level, with a total height of 7 meters. The white intermittent line is the second line of defense represented by the international coastal road extending from Al-Burg in the east to Al-Jediyah in the west with a length of 58 km after raising its level to 2 meters above the current sea level, knowing that the general average of the current road surface level is zero as shown in the attached topographical sector. Note that the second line of defense can be dispensed with during the distance from the village of Al-Burg in the east to the Maqluba hill in the west, due to the shrinkage of the thickness of the sand belt and the approach of the two defense lines to each other.

- Strengthening and raising the level of the southern and western bridges of Lake Burullus, with a length of about 70 km, to a height of no less than 2.0 meters above sea level. The average height of these bridges ranges from one meter to 2.0 meters above ground level. Raising the level of these bridges will prevent the flooding of the lake and its invasion of the southern and western plains (Fig. 37).
- Raising the level of the western bridge of the Suez Canal between Al-Qabouti in the north and Tina in the south to 2.0 meters above the current sea level: a length of 12.5 km, starting at latitude 31° 11' 15" north and ending at latitude of 31° 3' 15" south, where the bridge is cut by 16 gaps ranging in diameters from 200 meters to 800 meters, threatening the low plains that extend along the western bank of the canal. Note that the topography of the bridge along this path ranges from zero to 2.5 meters above sea level (Fig. 38a). Therefore, the height of the bridge will not exceed, on average, one meter with a length of about 12.5 km. Also, the level of the same bridge must be raised at the sites of the gaps between the Al-Salam Canal in the north and the Cape in the south, where there are two gaps with diameters ranging from 300 meters to 600 meters - and a third gap south of the Cape with a diameter of 300 meters beside a fourth with the same diameter north of the middle branch (Al-Ballah branch) (Figs 38b-c). As well as strengthening the western bridge at the intersection of the Alexandria-Port Said and Ismailia-Port Said roads, where the bridge level is less than 2.0 meters above sea level, which threatens to invade the bridge and from there to the plains west of the canal if the level of the canal surface increases by any amount more than meter.
- Strengthening and raising the level of the eastern and western bridges of the Nile River, Damietta Branch, between Ezbet Tabl in the south and Ezbet Al-Burg in the north, to a height of 2.0 meters above sea level. As for the western bridge, the path to be ascended is located in front of Ezbet El-Sheikh Dergham, with a length of 1060 meters. Knowing that the topography of the bridge along this path often ranges between zero and 0.5 meters above sea level, and then the height of the bridge will range on average from 1.5 meters to 2.0 meters with a length of about 900 meters. Hence, the height of the bridge will range on average from 1.5 meters to 2.0 meters with a length of about 900 meters. The danger of this low course lies in its being a source of threat to the Nile River from the west in the event that the waters of the Mediterranean Sea invade the low plains south of Ras El Bar, or the water seeps into these plains through the sandy subsurface soil. While the eastern bridge needs strengthening and raising the level between Ezbet Tabl and Ezbet Al-Ratmah, between Ezbet Al-Ratmah and Ezbet Al-Sheikh Dergham, and between Ezbet Al-Sheikh Dergham and Ezbet Al-Awba south of Ezbet Al-Burj with a total length of about 1000 meters and a height of 2.0 meters above sea level.
- Strengthening and raising the level of the eastern bridge of the Nile River, Rashid Branch, between Ezbet Abu Khashaba in the north and Fowa in the south, to a height of 2.0 meters above sea level, as the bridge along this path cuts through 30 gaps whose level is less than a meter above sea level (Figs. 39a-d). The diameters of these gaps range from 200 meters to 800 meters with a total length of 11.5 km from the length of the eastern bridge extending from Ezbet Abu Khashabah to Ezbet Ahmed Al-Taybani, east of Sandion. The danger of these gaps lies in the fact that they will act as auxiliary sea lanes for the rush of sea water from the north and east into the Nile River in the event that the sea invades the northern coastal part located between Lake Burullus in the east and Rasheed Branch in the west, or the sea water seeps into this part through the sandy subsurface soil. The gaps are concentrated in the following locations (Figs. 39a-d):
 1. The eastern bridge of the river extending between Ezbet Abu Khashaba and Rashid. Where there are 4 holes, their diameters range from 300 meters to 800 meters, the largest of which is located in front of the city of Rashid.
 2. The eastern bridge of the river extends from Ezbet Al-Waqf north of Al-Jediyah and Ezbet Al-Fars Island south of Buraidah, where there are 10 gaps ranging in diameters from 200 meters to 500 meters.
 3. The eastern bridge of the river extends between Izbat Jazirat Al-Farsah in the north and the ferry Mahdi (Izbat Al-Tijariyyin) in the south, where there are 6 gaps ranging in diameters from 200 meters to 400 meters.




Fig. 34a: Satellite image of the southern coast of the Gulf of Abu Qir extending from the suburb of Abu Qir in the west to the village of Maadiyah in the east - showing the proposed lines of defense (white lines, After Ouda, 2022), which are two lines, the first of them (the continuous white line), is the first line of defense. The second line (the white dashed line) is represented by a land bridge whose height is not less than 2.0 meters above the current sea level, with a length of 10.5 to 11.0 km, and extending directly south of Al-Tabia - Rashid road from south of the suburb of Abu Qir in the west to Qantara El-Maadiyah in the east. The same road that currently exists can be used to build this bridge after raising its level in low points to 2.0 meters above sea level (Ouda, 2022). **Fig. 34b:** Satellite image of the coastal strip extending on the coast of the Max Bay in, from Wardyan in the east to Dekheila in the west, and from the Mediterranean in the north to Lake Mariout in the south - showing on it the lines of defense (white lines). The continuous white line is the first line of defense. The white dashed line is the proposed second line of defense, represented by a land bridge extending along the current northern Mariout Lake shore line from the Max border in the west to the entrance of Apis in the east. Note that part of the bridge applies to the axis of the High Dam. See the extension of this bridge in the following figure (After Ouda, 2022).




Fig. 35: Google Earth satellite panorama of the coastal strip of the city of Alexandria, extending from the Sharq district in the east to El-Dekheila in the west, and from the sea in the north to Lake Mariout in the south, including the Western port and the Max Bay- showing the proposed lines of defense (white lines). The continuous white line is the proposed first line of defense, represented by a submerged reinforced concrete wall extending directly in front of the shore line towards the sea from the Western port in the east to the port of Dekheila in the west, with a total length of approximately 4.5 km and a total height of 7 meters, of which 5 meters are below sea level and 2.0 meters above sea level. The white dashed line is the proposed second line of defense, represented by a land bridge extending along the current northern Mariout Lake shore line from the Max border in the west to the City Center in the east, then extending around Nozha Airport along the ring road and from it to the northern border of the airport lake where it deviates east to Ezab Khurshid near the entrance to Apis, with a total length of about 22 km, and a height of not less than 2.0 meters above sea level. The attached diagram shows the path and terrain of this bridge, and part of this path applies to the High Dam axis road south of Qabbari, and another part applies to the new international road, south of Muharram Bey, and a third part applies to the ring road south of the airport and City Center, where these roads can be exploited after raising its level to 2.0 meters above sea level.

4. The eastern bridge of the river extending between Minyat Al-Murshed in the north and Ibiana in the south, where there are 5 gaps with diameters ranging from 200 meters to 400 meters.
5. The eastern bridge of the river extends from Shamashira 5 south of Motobas to Sindion, where there are 4 gaps ranging in diameters from 300 meters to 800 meters, the largest of which is located at Sindion in front of the city of Dayrut.
6. The eastern bridge of the river at Ezbet Ahmed Al-Tibani, east of Sindion, where there are two overlapping gaps with a diameter that can be considered as one gap with a diameter of about 500 meters.




Fig. 36: Satellite image of the coastal strip of Lake Manzala, extending from Port Said in the southeast to Ezbet Al-Burg in the northwest on the Mediterranean coast -- showing the proposed defense lines for the lake and the plains that surround it to the east, west and south. The continuous white coastal lines represent the first line of defense for the lake, and the intermittent white line is the second line of defense (see Fig. 30). The red line is the third line of defense for the dry plains exploited around the present lake, and is represented by the current bridge surrounding Lake Manzala, with the exception of some parts that we propose to modify its course north of Mataria and south of Damietta, in accordance with the reality of the current lake after isolating the part used as a drainage for Bahr al-Baqar and Bahr Hadous from the lake. The length of the bridge, as amended, is about 101 km, provided that its level is raised to two meters above the current sea level, bearing in mind that the ground level along the path of the bridge ranges, on average, from zero to 2.0 meters above sea level, as shown in the attached graph.

4-: Considering the bridges of the Mahmoudiyah Canal and the roads that border it on both sides between Damanhour and Alexandria as the second line of defense for the agricultural plains extending south of the canal until the desert back, by strengthening the bridges and roads and raising their ground level to 2 meters above sea level in the following low areas (Figs 40a-b):

- The area extending from the canal between Ezbet Al-Fakhoura, east of the city of Karion, to Al-Ghaba (west of Berket Ghattas), where there is a large gap with a diameter of 2.5 km near Ezbet Al-Shawish, west of Al-Nasiri Bank.
- The area extending from the canal between El-Akraisha (east of Kafr El-Dawwar) and El-Carryon, where there are 5 gaps, ranging in diameters from 200 meters to 300 meters, with a total of 1200 meters.
- The area extending from the canal between King Othman and Kafr Selim, where there is a 400-meter gap in front of Ezbet El Hamaleen.
- The area extending from the canal between Ezbet Al-Khadra and the city of Al-Bayda, where there is a large gap, about 3 km in diameter, that covers most of the area between Al-Bayda Dyeing Company and Al-Bayda City.




Fig. 36: Satellite image of the coastal strip extending from Al-Burg in the east to Ezbet Abu Khashaba (North of Rashid) in the west on the Mediterranean coast, including the sandy belt of Lake Burullus - showing the proposed defense lines for the lake and the dry (or dried) low plains surrounding it. The continuous white lines represent the first line of defense for the lake, and the white dashed line is the second line of defense (see Fig.31). The red line is the third line of defense, represented by the southern and western bridge of Lake Burullus, with a length of 70 km, after raising its level to at least 2.0 meters above sea level. Note that the bridges level ranges on average from -2.0 meters below sea level in the west to one meter in the east above sea level. However, by inspection, it was found that the lower areas of the bridge have been raised to more than zero, and therefore the required height above the ground usually ranges from one to 1.5 meters along the bridge path.

- The area extending from the canal around the upper bridge and the Ring Road in the Khadra district where there is a gap of 800 meters long.
- The area extending between Ezbet Al-Ubaid, near the station, east of the Khurshid Canal, and Ezab Nubar, where there are 4 holes ranging in diameter from 200 meters to 300 meters.

5-: The need to protect the city of Idku from marine isolation, as it is surrounded on all sides by wet and dry depressions. Therefore, work must be done to fill these depressions and raise their level, especially the vertical depressions on the coast that border the city from the east (about 2.75 km long) and west (about 1.5 km long), as these depressions can act as sea passages between the sea and the plains that border Idku from the south. It is also possible to reinforce the filling of the northern depression of the city, which is about 3.5 km long, by dumping sand to raise its level to become a sandy area that protects the city from coastal erosion operations in case the sea level rises by a meter and approaches the northern city limits.

6-: All these coastal defenses do not dispense with plans to empty water and drain it away from the sea. The sea water, when its level increases, can permeate the sandy subsurface soil along the coast of the delta, submerging some depressions, or mix with groundwater and increase its level, which may cause the partial inundation of the low inter-regions that lie south of the sand belt surrounding the coast of the delta. In addition, the increase in sea level will increase the height of the waves in the winter season and their ability to overcome some natural or industrial barriers along the shoreline. Therefore, it is necessary to plan for the construction of a network of pipes or channels to discharge sea water that will penetrate the coastal defenses along the coast and drain it in low desert areas far from the sea shore. The sea is not suitable as a drain for sea water.




Fig. 38a: Satellite image of the northern part of the Suez Canal, showing the locations of the gaps that cross the western bridge of the Suez Canal and their relationship to the dried-up plains extending along the western bank of the canal between Port Said in the north and Tina in the south - if the sea level rises and then the Suez Canal by a maximum of one meter. Therefore, these plains are threatened by a direct sea invasion from the waters of the Suez Canal in the east through the gaps that permeate the western bridge of the Suez Canal, which are concentrated especially between Al-Qabouti in the north and Tina in the south. The continuous white line represents the course of the western Suez Canal Bridge that is required to be raised to 2.0 meters above the current sea level, with a length of 12.5 km. **Figs. 38b-c:** Satellite image of the northern part of the Suez Canal between Ras al-Ush in the north and Qantara in the south, showing the locations of the gaps that cross the western bridge of the Suez Canal north and south of the Sahara Desert. Note that the gaps are less south of Al-Tina and their distances diverge until they are absent before Al-Ballah (see the enlarged picture of the great gap south of the Sahara, Fig. 38c). The red lines represent the paths to be raised in the western bridge of the canal to 2 meters above the current sea level.




Fig. 39a: Satellite image of the northern coastal strip of the delta extending between Lake Burullus in the east and the Nile River, Rosetta branch in the west - showing the proposed lines of defense. The continuous white lines represent the first lines of defense which are represented by submerged reinforced concrete walls extending directly north of the beach line facing the sea between Kom Mishaal to the east and the beach north of the Green Island, according to the specifications mentioned in the text. The eastern bridge of the river should be raised in its ground level to a height of 2.0 meters above sea level in the locations of the gaps. As for the intermittent white line, it is the second line of defense, represented by the international coastal road extending from Al-Burg in the east to Al-Jediyah in the west, after raising its level to 2.0 meters above sea level. **Figs. 39b-d:** Satellite images of the northern part of the Nile River, the Rashid branch, which extends between Rashid in the north to Brnbal in the south (Fig. 34b), between Minyat al-Murshed in the north and Shamashira in the south (Fig. 34c), and between Shamashira in the north and Fowa in the south (Fig. 34d)- showing the proposed lines of defense for the Nile River. The continuous white lines are the proposed lines of defense, represented by the eastern bridge of the river after raising its level to a height of 2.0 meters above sea level.




Fig. 40a: Satellite image of the ground strip that permeates the low plains extending from the Nile River, Rashid branch in the east to Alexandria in the west, in the northwestern part of the delta - showing on it the paths of the necessary defense lines (white lines) along the Mahmoudiya Canal (black line) to protect the southern plains from the sea invasion through the gaps that permeate the canal's bridges, The numbers next to the circles are the radii of the gaps. **Fig. 40b:** An enlarged satellite image of the Mahmoudiya Canal between Ghattas Pool in the east and Karion in the west The attached diagram represents the topography of the northern bridge of Mahmoudiya Canal along this gap.

5. Conclusions

The economic and human losses that may result in the north of the Egyptian delta due to the rise in global sea level are innumerable. They are not limited to the Mediterranean Sea's invasion of all the northern lakes in the delta, but go beyond that to include agricultural lands, residential buildings, factories, shops, tourist sites, roads and canals, drains and facilities in the dry or drained lowlands surrounding the northern lakes. The effect of sea level rise extends to include the high degree of salinity of groundwater, as well as to include erosion and sedimentation systems, especially in the gorges and estuaries of the two Nile branches, the ecosystems of lakes, and the drainage systems of the northern delta. The population in the governorates located north, northwest, and northeast of the delta will be forced to leave their homes because of the losses of agricultural land that will be overrun by the sea, which threatens a catastrophe unless the Egyptian government begins to expedite measures to protect against the impact of climate changes and adapt to the new climatic conditions.

The sources of threat that must be protected are diverse. The coast of the southern Abu Qir Bay, which extends between the city of Abu Qir and the village of Maadiyah, is the main source of threat to all the dry and wet low plains (what remains of the Idku and Mariout lakes) in the northwest of the delta (west of the Rashid branch), extending from the south of the Bay to the desert back between Alexandria in the west, and the Nile River to the east, due to its topographical, geological, historical and environmental conditions. The sandy buffer belt between the Gulf and the low plains to its south is subject to a marine invasion of about 66.6% to 83% of its area if the global sea level rises by a meter or less. Then comes the middle part of the western port of Alexandria, between Wardyan and Dekheila, as a second source of threat to the dry (or dried) and humid low lands extending south of Alexandria between the Western Port to the west, and the city of Damanhour to the east, in case the sea level rises by a meter or more.

In the northern delta region (between the two branches of the river), the main source of threat to it is the northern sandy belt extending from the mouth of Rashid in the west to the village of Al-Burg in the east, where the current weak areas in the belt, which are below sea level, constitute 47% of its total area and about 53 % of its total length. This means that the sea will invade half of the belt and from there to Lake Burullus and the low plains west of the lake in case the sea level increases by any amount less than a meter. In the case of an increase in sea level by a

meter, the area threatened by the invasion of the sand belt will amount to about 73.5% of its total area. The most dangerous areas of the belt are the coastal area located between the Green Island in the west and Kom Mishaal in the east, with a length of 13.5 km, as it threatens all agricultural areas south of the belt in Markaz Fowa between Lake Burullus and the Nile River until Fowa in the south. Also, the low gaps spread in the belt extending between Kom Mishaal in the west and Al-Burg in the east are the main source of threat to all the wet areas (Burullus Lake) and the surrounding dry areas. Hence, the southern and western bridges of Lake Burullus are a source of threat to the dry lowlands that extend southeast of the lake to Kom al-Hajar, north of Hamul in Markaz Biella, and east of the lake to Mastamouli in Markaz Belqas, and south of Lake to Abbasiya - Ezbet Umm Sinn in Markaz Kafr El-Sheikh, and Al-Qasabi - Mineshaft Abbas In Markaz Sidi Salem, west of Beheira until Rashid, and southwest of Beheira until Sindion in Markaz Fowa

As for the dry and wet low lands in the northeast of the delta (east of the Damietta branch), the sources of their threat are diverse, and they include the northwestern part of the sandy belt separating the sea and Lake Manzala (west of Al-Manasra village), known as Barr Al-Dahra, due to the deterioration of the belt and the spread of low gaps. The sea will cover about 59% of the area of the Barr Al-Dahra belt if the sea level rises by one meter. Also, the western bridges of the Suez Canal between Port Said in the north and Al Cap in the south, they are a major source of threat to all dry and dried lands (from the original Lake Manzala) located between Lake Manzala (with its current borders) and the Suez Canal, due to the low level of the bridges and the spread of Inter-holes that do not exceed one meter above sea level along the western coast of the canal extending south of the Sahara until Tina, and at Al- Cap as well. The western and southern bridges of Lake Manzala are also a source of threat to the dry lowlands extending to the west within Markaz Faraskour, south of Damietta, along the eastern bank of the Nile to Al-Serw, southwest to the Al-Nuzul in Markaz Dikrnis, and south to Al-Hussainiya in the northeast of Sharqia Governorate, and southeast to Western Qantara and from there to Al-Balah to the south (north of Ismailia Governorate).

Therefore, the Egyptian government must immediately plan measures to defend the northern delta coast. The bodies that develop strategies should set their defense priorities based on the information contained in this research and other local and international reports issued in this regard. The success

of these defenses does not depend on the state's economics as much as it depends on the state's experiences in the field of environmental protection and maintenance. Therefore, the defenses must be built within the framework of a comprehensive national plan to protect the northern coast of the delta. The defense plans should also include the following:

First: Construction of submerged armored cement walls facing the sea in locations where low coastal gaps are not separated from the sea by strong sandy belts, whose invasion or sea leakage through them would flood vast areas of lands south of them. These walls must be submerged below sea level by digging the subsoil at a depth ranging from 3.0 meters to 5.0 meters, and the walls should rise above sea level by not less than two meters - with a total height ranging from 5.0 meters to 7 meters. The locations, dimensions and depths of the submerged armored walls required to be erected along the northern arc of the delta are determined in this research.

Second: erecting cement ground bridges south of the shore line, with a height of no less than 2.0 meters above sea level, with flat surfaces and inclined sides to the north and south extending along the coast, or using existing roads after raising their level to 2.0 meters above sea level. to take the place of these bridges as a second line of defense in the locations whose topographical specifications and geographical locations have been determined in this research.

Third: Strengthening and raising the level of the current bridges of the northern lakes. Raising the level of these bridges will prevent the flooding of the lake and its invasion of the surrounding plains if the lake level rises. As well as strengthening and raising the lower eastern bridges in the northern part of the Nile River (Rasheed Branch) between Fowa and Rashid, the eastern and western bridges of the Nile River, the Damietta branch, between Ezbet Tabl in the south and Ezbet Al-Burg in the north, and the western bridges of the northern part of the Suez Canal between Al-Qabouti and Al-Cap, in the areas Geographically and topographically identified in this research.

Fourth: Considering the bridges of the Mahmoudiya Canal and the roads that border it on both sides between Damanhour and Alexandria as the third line of defense for the agricultural plains extending south of the canal to the desert back, by strengthening bridges and roads and raising their level in the low areas defined geographically and topographically in this research.

Fifth: The necessity of protecting the city of Idku from sea isolation.

Finally, all these coastal defenses do not dispense with plans to empty the water and drain it away from the sea in case the sea water crosses some natural or artificial barriers extending along the shore line. Therefore, it is necessary to plan for the construction of a network of pipes or channels to discharge sea water that will penetrate the coastal defenses along the coast and drain it in low desert areas far from the sea shore. The sea is not suitable as a drain for sea water.

Acknowledgement

The author wishes to express his gratitude to the Management of Assiut University for providing Lab and internet facilities necessary for the completion of this study. Deep thanks are also due to the reviewers for their insightful reviews and valuable comments on this manuscript.

Declarations Funding: No funding.

Conflicts of interest: No conflicts.

References

- [1] Ahmed M. H. (2002): Multi-temporal Conflict of the Nile Delta Coastal Changes, Egypt. Littoral 2002, The Changing Coast. Eurocoast/EUCC, Portugal: 317-323.
- [2] Almarsad Almasry (2020): Central Agency for Public Mobilization and Statistics: Egypt in Figures 2020. Heba Zain, 1, 2020. <https://marsad.ecss.com.eg/43794/>
- [3] Al-Muzayen A. A. (2019): The most important problems of Lake Manzala: The causes of low productivity. Ministry of Agriculture and Land Reclamation - General Authority for Fisheries Development <http://www.gafrd.org/information>. GAFRD@gmail.com www.GAFRD.org
- [4] Annual Report on Lake Burullus (2018-2019): . Burullus Lake: Environmental Affairs Agency, Environmental Quality Sector, for the Central Department of Water Quality: Egyptian Lakes Environmental Monitoring Program. <https://www.eeaa.gov.eg/portals/0/eeaaReports/water/reportAnnual1718/%D8%A7%D9%84%D8%A8%D8%B1%D9%84%D8%B3.pdf>
- [5] Annual Report on Lake Mariout (2019-2020): Mariout Lake: Environmental Affairs Agency, Environmental Quality Sector, for the Central Department of Water Quality: Egyptian Lakes Environmental Monitoring Program. <https://www.eeaa.gov.eg/portals/0/eeaaReports/water/reportAnnual1920/%D9%85%D8%B1%D9%8A%D9%88%D8%B7%20%D8%A7%D9%84%D8%B3%D9%86%D9%88%D9%89%202020.pdf>

- [6] Bek M. A., Lowndes I. S, Hargreaves D. M, Negm A. M. (2019): Lake Manzala Characteristics and Main Challenges. Egyptian Coastal Lakes and Wetlands: Part I, Publisher: Springer International Publishing
- [7] Dasgupta S., Laplante B., Meisner C., Wheeler d., Yan J. (2007): The impact of sea level rise on developing countries. A comparative analysis. World bank policy research Working, Paper 4136
- [8] Dumont H. J., El-Shabrawy G. M. (2007): Lake Borullus of the Nile Delta: A Short History and an Uncertain Future. *Ambio*, 36 (8): 677-682. Published By: Springer <https://www.jstor.org/stable/25547836>
- [9] El Raey M, Dewidar Kh., El Hattab, M. (1999): Adaptation to the impacts of sea level rise in Egypt. *Climate Research*, 12: 117-128.
- [10] El-Badry A. E. A., Khalifa M. M. (2017): Geochemical Assessment of Pollution at Manzala Lake, Egypt: Special Mention to Environmental and Health Effects of Arsenic, Selenium, Tin and Antimony. *Journal of Applied Sciences*, 17 (2): 72-80 DOI:10.3923/jas.2017.72.80
- [11] Elmorsi R., Abou-El-Sherbini Khaled S., Mostafa G. AH, Hamed M. A. (2019): Distribution of essential heavy metals in the aquatic ecosystem of Lake Manzala, Egypt. <https://doi.org/10.1016/j.heliyon.2019.e02276>
- [12] El-Naggar N. A., Rifaat A. E. (2017): Egyptian Coastal Lakes and Wetlands: Part I: Cite as Hydrodynamic and Water Quality Modeling of Lake Mariout (Nile Delta, Northern Egypt). Part of the Handbook of Environmental Chemistry book series, HEC, 71: 241-263
- [13] European Environmental Agency (2021): Global and European sea level rise Published in 18 Nov 2021 <https://www.eea.europa.eu/ims/global-and-european-sea-level-rise>
- [14] Hereher M. E. (2014): The Lake Manzala of Egypt: an ambiguous future. *Environmental Earth Sciences* 72: 1801–1809.
- [15] Hussein Munis, (1986). Atlas of the History of Islam (color): Publisher Dar Al-Zahra for Arab Media, 530 p
- [16] IPCC, Climate Change (1990): The Scientific Assessment. First Report of the IPCC Scientific Assessment Working Group I.
- [17] IPCC, Climate Change (1995): Synthesis of Scientific-Technical Information Relevant to Interpreting Article 2 of the UN Framework Convention on Climate Change. Contribution of Working Groups I, II and III to the Second Assessment Report of the Intergovernmental Panel on Climate Change.
- [18] IPCC, Climate Change (2001): Synthesis Report. A Contribution of Working Groups I, II, and III to the Third Assessment Report of the Intergovernmental Panel on Climate Change [Watson, R. T. and the Core Writing Team (eds.)]. Cambridge University Press, Cambridge, United Kingdom, and New York, NY, USA, 398 p.
- [19] IPCC, Climate change (2007): Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change 2007, Core Writing Team, Pachauri, R. K. and Reisinger, A. (Eds.) IPCC, Geneva, Switzerland, 104 p.
- [20] IPCC, Climate Change (2014): Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R. K. Pachauri and L. A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 p.
- [21] IPCC, Climate Change (2019): Summary for Policymakers. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H. -O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N. M. Weyer (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 3–35. <https://doi.org/10.1017/9781009157964.001>
- [22] IPCC, Climate Change (2021): The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 3–32, doi:10.1017/9781009157896.001.
- [23] Ismail A. and Hettiarachchi H. (2017): Environmental Damage Caused by Wastewater

- Discharge into the Lake Manzala in Egypt. Environmental Science. American Journal of BioScience. DOI: 10.11648/J.BIO.20170506.14Corpus ID: 55968381
- [24] Mohamad S. Z (2011): Lake Mariout: Past, Present and Future. Publisher LAP LAMBERT Academic ISBN: 978-3844325393Publishing
- [25] Mohamadin L. I., El-Sawy M. A., Bek M. A. (2019): Sediment Contaminants in Northern Egyptian Coastal Lakes. Egyptian Coastal Lakes and Wetlands: Part I. Publisher: Springer International Publishing
- [26] Ouda, Kh. A. K. (2010): Atlas of risks of climate change on the Egyptian coasts and defensive policies. Publisher: Assiut University, Assiut 71516, Egypt, 2 volumes, 955 p., 734 pl. Registration Number10847/2010. International numeration 977-17-9006-4.
- [27] Ouda, Kh. A. K. (2012): Atlas of Risks of Climate Change on The Egyptian Coasts and Defensive Policies. Humboldt kolleg. Proceedings of the Fifth International Conference of The Egyptian Society for Environmental Sciences & Suez Canal University "Climate Change. and water Resources", 7 July 2012. Published by the Egyptian Society for Environmental Sciences, pp. 26-28.
- [28] Ouda, Kh. A. K. (2011): Atlas of risks of climate change on the Egyptian coasts and defensive policies. Bulletin of the Egyptian Geographical Society, 84, pp. 185-198.
- [29] Ouda, Kh. A. K. (2022): Neighborhoods at Risk of Drowning in Alexandria in Light of Climate Change, Coastal Threat Sources and Means of Protection. International Journal of Trend in Scientific Research and Development (IJTSRD -ISSN: 2456 – 6470), 6 (5), July-August 2022 Available Online: Volume-6 | Issue-5, August 2022, pp. 390-427, URL: www.ijtsrd.com/papers/ijtsrd50484.pdf
- [30] Sallam Kh., Abd-Elghany S. M., Mohammed M. A. (2019): Heavy Metal Residues in Some Fishes from Manzala Lake, Egypt, and Their Health-Risk Assessment. J Food Sci., 84(7). doi:10.1111/1750-3841.14676.
- [31] Sherif, M., Al-Rashed, M., (2001): "Vertical and Horizontal Simulation of Seawater Intrusion in the Nile Delta Aquifer First International Conference on Saltwater Intrusion and Coastal Aquifers". Monitoring, Modeling, and Management. Essaouira, Morocco, April 23ñ25, 2001
- [32] Shoman M. M. (2018): Monitoring of some changes in Manzala Lake using remote sensing and GIS techniques. RS & GIS Unit, Soils, Water and Environment Research Institute, ARC.
- [33] Smithsonian Institution (1985): National Museum of Natural History (NMNH), Program on Nile Delta Subsidence and Sea Level Rise. www.gcrio.org/OCP/progsum/si.niledelta.htm
- [34] Stanley J. D., Warne A. G., David H. R., Bernasconi M. P., Chen Z. (1992): Nile Delta, National Geographic research & Exploration, 8 (1), pp 22-51.