

University of Groningen

Genome-wide association study in patients with posterior urethral valves

van der Zanden, Loes F.M.; Maj, Carlo; Borisov, Oleg; van Rooij, Iris A.L.M.; Quaedackers, Josine S.L.T.; Steffens, Martijn; Schierbaum, Luca; Schneider, Sophia; Waffenschmidt, Lea; Kiemeney, Lambertus A.L.M.

Published in:
Frontiers in Pediatrics

DOI:
[10.3389/fped.2022.988374](https://doi.org/10.3389/fped.2022.988374)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Zanden, L. F. M., Maj, C., Borisov, O., van Rooij, I. A. L. M., Quaedackers, J. S. L. T., Steffens, M., Schierbaum, L., Schneider, S., Waffenschmidt, L., Kiemeney, L. A. L. M., de Wall, L. L. L., Heilmann, S., Hofmann, A., Gehlen, J., Schumacher, J., Szczepanska, M., Taranta-Janusz, K., Kroll, P., Krzemien, G., ... Hilger, A. C. (2022). Genome-wide association study in patients with posterior urethral valves. *Frontiers in Pediatrics*, 10, [988374]. <https://doi.org/10.3389/fped.2022.988374>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



OPEN ACCESS

EDITED BY

Giorgio Giuseppe Orlando Selvaggio,
Children's Hospital "Vittore Buzzi", Italy

REVIEWED BY

Andrew Mallett,
Townsville University
Hospital, Australia
John Samuel Wiener,
Duke University, United States

*CORRESPONDENCE

Alina C. Hilger
alina.hilger@uk-erlangen.de

†These authors have contributed
equally to this work and share first
authorship

‡These authors share last authorship

SPECIALTY SECTION

This article was submitted to
Pediatric Urology,
a section of the journal
Frontiers in Pediatrics

RECEIVED 07 July 2022

ACCEPTED 19 August 2022

PUBLISHED 27 September 2022

CITATION

van der Zanden LFM, Maj C, Borisov O,
van Rooij IALM, Quaedackers JSLT,
Steffens M, Schierbaum L, Schneider S,
Waffenschmidt L, Kiemeny LALM, de
Wall LLL, Heilmann S, Hofmann A,
Gehlen J, Schumacher J,
Szczepanska M, Taranta-Janusz K,
Kroll P, Krzemien G, Szmigielska A,
Schreuder MF, Weber S, Zaniew M,
Roeleveld N, Reutter H, Feitz WFJ and
Hilger AC (2022) Genome-wide
association study in patients with
posterior urethral valves.
Front. Pediatr. 10:988374.
doi: 10.3389/fped.2022.988374

COPYRIGHT

© 2022 van der Zanden, Maj, Borisov,
van Rooij, Quaedackers, Steffens,
Schierbaum, Schneider,
Waffenschmidt, Kiemeny, de Wall,
Heilmann, Hofmann, Gehlen,
Schumacher, Szczepanska,
Taranta-Janusz, Kroll, Krzemien,
Szmigielska, Schreuder, Weber,
Zaniew, Roeleveld, Reutter, Feitz and
Hilger. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Genome-wide association study in patients with posterior urethral valves

Loes F. M. van der Zanden^{1†}, Carlo Maj^{2†}, Oleg Borisov²,
Iris A. L. M. van Rooij¹, Josine S. L. T. Quaedackers³,
Martijn Steffens⁴, Luca Schierbaum⁵, Sophia Schneider⁵,
Lea Waffenschmidt⁵, Lambertus A. L. M. Kiemeny¹,
Liesbeth L. L. de Wall⁶, Stefanie Heilmann⁵, Aybike Hofmann⁷,
Jan Gehlen⁸, Johannes Schumacher⁸, Maria Szczepanska⁹,
Katarzyna Taranta-Janusz¹⁰, Pawel Kroll¹¹,
Grazyna Krzemien¹², Agnieszka Szmigielska¹²,
Michiel F. Schreuder¹³, Stefanie Weber¹⁴, Marcin Zaniew¹⁵,
Nel Roeleveld¹, Heiko Reutter¹⁶, Wout F. J. Feitz^{6‡} and
Alina C. Hilger^{5,17,18**}

¹Department for Health Evidence, Radboud Institute for Health Sciences, Radboud University Medical Center, Nijmegen, Netherlands, ²Institute for Genomic Statistics and Bioinformatics, Medical Faculty, University of Bonn, Bonn, Germany, ³Department of Urology, University Medical Center Groningen, Groningen, Netherlands, ⁴Department of Urology, Isala, Zwolle, Netherlands, ⁵Institute of Human Genetics, School of Medicine and University Hospital Bonn, University of Bonn, Bonn, Germany, ⁶Division of Pediatric Urology, Department of Urology, Radboud Institute for Molecular Life Sciences, Radboudumc Amalia Children's Hospital, Nijmegen, Netherlands, ⁷Department of Pediatric Urology, Clinic St. Hedwig, University Medical Center Regensburg, Regensburg, Germany, ⁸Center for Human Genetics, University Hospital of Marburg, Marburg, Germany, ⁹Department of Pediatrics, Faculty of Medical Sciences in Zabrze, Medical University of Silesia in Katowice, Zabrze, Poland, ¹⁰Department of Pediatrics and Nephrology, Medical University of Białystok, Białystok, Poland, ¹¹Neurourology Unit, Pediatric Surgery and Urology Clinic, Poznań, Poland, ¹²Department of Pediatrics and Nephrology, Medical University of Warsaw, Warsaw, Poland, ¹³Department of Pediatric Nephrology, Radboud Institute for Molecular Life Sciences, Radboudumc Amalia Children's Hospital, Nijmegen, Netherlands, ¹⁴University Children Hospital Marburg, Philipps University Marburg, Marburg, Germany, ¹⁵Department of Pediatrics, University of Zielona Góra, Zielona Góra, Poland, ¹⁶Division of Neonatology and Pediatric Intensive Care, Department of Pediatrics and Adolescent Medicine, Friedrich-Alexander University of Erlangen-Nürnberg, Erlangen, Germany, ¹⁷Department of Pediatrics and Adolescent Medicine, Friedrich-Alexander University of Erlangen-Nürnberg, Erlangen, Germany, ¹⁸Research Center on Rare Kidney Diseases, University Hospital Erlangen, Erlangen, Germany

Congenital lower urinary tract obstructions (LUTO) are most often caused by posterior urethral valves (PUV), a male limited anatomical obstruction of the urethra affecting 1 in 4,000 male live births. Little is known about the genetic background of PUV. Here, we report the largest genome-wide association study (GWAS) for PUV in 4 cohorts of patients and controls. The final meta-analysis included 756 patients and 4,823 ethnicity matched controls and comprised 5,754,208 variants that were genotyped or imputed and passed quality control in all 4 cohorts. No genome-wide significant locus was identified, but 33 variants showed suggestive significance ($P < 1 \times 10^{-5}$). When considering only loci with multiple variants residing within <10 kB of each other showing suggestive significance and with the same effect direction in all 4 cohorts, 3 loci comprising a total of 9 variants remained. These loci resided on chromosomes 13, 16, and 20. The present GWAS and meta-analysis

is the largest genetic study on PUV performed to date. The fact that no genome-wide significant locus was identified, can be explained by lack of power or may indicate that common variants do not play a major role in the etiology of PUV. Nevertheless, future studies are warranted to replicate and validate the 3 loci that yielded suggestive associations.

KEYWORDS

genome wide association study, lower urinary tract obstruction, obstructive uropathy, posterior urethral valves, *PCDH9*, *SALL1*, *BMP7*

Introduction

Congenital lower urinary tract obstruction (LUTO, MIM # 618612) is defined by a decrease in the free passage of urine from the bladder through the urethra. LUTO can be one of the most severe diagnoses underlying obstructive uropathy and may perturb kidney development. Obstructive uropathy is the second most prevalent cause of end-stage kidney disease in children (1). LUTO is most often caused by an anatomical blockage, commonest by posterior urethral valves (PUV) (2), a male limited phenotype affecting 1 in 4,000 male live births (3). Severe forms of LUTO can be diagnosed intrauterine, presenting with megacystis, including “keyhole sign”, as well as megaureter, oligohydramnios, and often dysplastic kidneys (2). Milder LUTO forms may manifest postnatally, often with recurrent urinary tract infections (3). Little is known about the genetic background of anatomical LUTO or PUV, which occur sporadically in most cases. Familial forms, segregating through generations with mixed phenotypes (PUV and stenosis) as well as affected sib-pairs, have been described, however, suggesting a potential genetic contribution to the malformation (4, 5). Furthermore, Frese et al. (6) found higher concordance rates among monozygotic compared to dizygotic twin pairs in a classic twin study. Two genetic studies found rare copy number variants (CNVs) potentially contributing to PUV in up to 57% of cases (7) and a higher occurrence of rare CNVs among PUV patients than among controls (8). In addition, an association with a variant in the angiotensin II receptor type 2 (*AGTR2*) was found in one study (9). Variants in *basonuclin 2* (*BNC2*) as a possible first identified monogenic cause were described by Kolvenbach et al. (10). Overall, the genetic background appears heterogeneous with a multifactorial mode of inheritance likely to underlie a large proportion of patients. Here, we report the largest study using genome-wide association methods to identify genetic susceptibility loci for PUV. The final meta analyses comprised a total of 756 patients and 4,823 controls.

Materials and methods

Patients and controls and genotyping

Details of the recruitment process for patients and controls are provided in the Supplement. In short, the GWAS sample comprised 823 male PUV patients and 5,254 male controls. This sample consisted of in total 4 cohorts from The Netherlands, Germany, and Poland. Prior to inclusion, written informed consent was obtained from all subjects or from their proxies in case of legal minors. The study was approved by the institutional medical-ethics committee of the participating centers and was conducted in accordance with the principles of the Declaration of Helsinki. DNA was extracted from blood and saliva samples using standard procedures.

All Dutch samples were genotyped by deCODE Genetics (Reykjavik, Iceland). The first Dutch cohort comprised 417 patients derived from the Radboudumc AGORA data- and biobank (11, 12), which were patients born in 1981 or later who underwent a valve resection before the age of 18 years, and 2,493 controls derived from the Nijmegen Biomedical Study, a population-based study in which randomly selected inhabitants of the municipality of Nijmegen received an invitation to donate blood samples (13). This cohort was genotyped using the Infinium OmniExpress bead chips (Illumina, San Diego, CA, USA) and will further be called the Dutch Omni sample. The second Dutch cohort consisted of 62 AGORA patients and 323 controls collected in AGORA by asking 42 municipalities to provide a random sample of their inhabitants in the age range of 0–20 years and inviting these children and adolescents to provide a saliva sample. This cohort was genotyped using the Global Screening Array (GSA) from Illumina modified by deCODE (Illumina, San Diego, CA, USA), and will further be called the Dutch GSA cohort. The German cohort comprised 109 diagnosed PUV patients sampled through the CaRE for LUTO (Cause and Risk Evaluation for LUTO) study, mostly included within in the first year of life right after diagnosis of PUV and 2,144 German controls derived from a prospective

population-based cohort study, the Heinz Nixdorf Recall Study (14). The Polish sample comprised 235 diagnosed PUV patients recruited in a collaboration between the CaRE for LUTO Study and the PolTubeReg (Polish registry for tubular kidney disease) and 294 Polish controls, which were recruited as blood donors in Lodz, Poland. The German and Polish patient and control samples were genotyped using the Global Screening Array I (Illumina, San Diego, CA, USA).

Quality control and imputation

Pre-imputation quality control

Quality control was performed separately for samples from the different populations and for samples genotyped on different platforms. Variants were excluded if they had < 98% call rate, had minor allele frequencies (MAF) < 0.1%, or failed the Hardy Weinberg Equilibrium (HWE) test ($P < 1 \times 10^{-10}$ for patients or $P < 1 \times 10^{-6}$ for controls). Samples were excluded if they had discordant sex information or a call rate < 98%, or if the kinship coefficient calculated using KING (15) was >0.0442, indicating a 3rd-degree relationship or higher with another sample in the cohort. In the identification of individuals from other ethnicities, PLINK was used to compute principal components (PC) for study participants and reference samples of the 1,000 Genomes Project Phase 3 (16) and to visualize sub-structuring. Participants that clustered with a non-European population from the 1,000 Genomes project were excluded. We recomputed the PC in the remaining participants only, visualized results, and excluded participants with the first or second PC more than 6 standard deviations from the mean.

Imputation

Imputation was performed for hg19 built, separately for samples from the different populations and for samples genotyped on different platforms. Genotypes of the different samples were phased using SHAPEIT (v2) (17) and imputed using IMPUTE2 (v2) (18) using 1,000 Genomes Project Phase 3 data as a reference panel (16). After imputation, variants with an IMPUTE2 information metric (info) score < 0.6 or with a minor allele count < 20 were excluded.

Association analysis

Association analysis was performed separately for samples from the different populations and for samples genotyped on different platforms in PLINK using the logistic regression framework implemented in the glm function of PLINK 2,¹ adjusting for the first 4 principal components. Afterwards,

1 <https://www.cog-genomics.org/plink/2.0/>

results of SNPs that were genotyped or imputed and passed quality control in all cohorts were pooled in a meta-analysis using the effect size estimates and standard errors in the METAL software, SCHEME STDERR.² The qqman package in R³ was used to visualize the results by means of Manhattan and Q-Q plots. The metafor package in R⁴ was used to create forest plots for selected genetic variants of interest.

Results

The post-quality control dataset forwarded to independent GWAS comprised 402 patients and 2,213 controls for the Dutch Omni cohort, 58 patients and 314 controls for the Dutch GSA cohort, 104 patients and 2,070 controls for the German cohort, and 192 patients and 226 controls for the Polish cohort. The independent GWAS in the German cohort reached genome-wide significance for 3 genotyped variants on chromosome 16 (rs117706747, rs117689220, and rs17666927). Imputation yielded 75 additional variants in LD that also showed genome-wide significance (ORs ranging between 5.65 and 8.52 with P -values of $5E^{-8}$ to $8E^{-11}$). The results in the Polish and the Dutch GSA cohorts showed the same effect direction as in the German cohort (ORs ranging between 1.28 and 2.14 with P -values of 0.51–0.02 for the Polish cohort and OR between 1.65 and 1.70 with P -values of 0.14–0.09 for the Dutch GSA cohort), whereas the Dutch Omni cohort presented with opposite or no effect directions (ORs ranging between 0.66 and 1.18 with all P -values > 0.11) (Supplementary Table 1).

The final meta-analysis included 756 patients and 4,823 on ethnicity matched controls and comprised 5,754,208 variants that were genotyped or imputed and passed quality control in all 4 cohorts. No relevant inflation was detected ($\lambda = 1.01$) and no genome-wide significant locus was identified (Figure 1). In total, 33 variants showed suggestive significance ($P < 1 \times 10^{-5}$, see Supplementary Table 2). However, the effects were in different directions across cohorts for some of the loci (i.e., on chromosome 1, 5, and 18). When considering loci with multiple variants residing within <10 kb of each other showing suggestive significance and with the same effect direction in all 4 cohorts only, 3 loci comprising a total of 9 variants remained (Table 1). These loci reside on chromosome 13 (5 variants), 16 (2 variants), and 20 (2 variants) (see Figures 2A–C for LocusZoom plots and Figure 2D for Forest plots of leading variants at these loci). While the locus on chromosome 13 resides in an intronic region of the gene *PCDH9*, the loci on chromosome 16 and 20 reside in intergenic regions.

2 <https://genome.sph.umich.edu/wiki/METAL>

3 <https://www.r-project.org/nosvn/pandoc/qqman.html>

4 <http://www.metafor-project.org/doku.php>




TABLE 1 Association results with PUV for the variants in loci with multiple variants residing within <10 kB of each other showing suggestive significance ($P < 1 \times 10^{-5}$) and with the same effect direction in all 4 cohorts.

CHR	BP	dbSNPrs#	A1	Meta analysis results			Association results in the 4 cohorts							
							Dutch GSA		Dutch Omni		German		Polish	
				P_{het}	OR	P_{OR}	OR_a	P_a	OR_a	P_a	OR_a	P_a	OR_a	P_a
13	67498549	rs9564361	A	0.97	1.40	9×10^{-6}	1.58	0.10	1.39	4×10^{-4}	1.34	0.15	1.41	0.09
13	67498767	rs34844007	C	0.98	1.40	1×10^{-5}	1.57	0.11	1.39	4×10^{-4}	1.34	0.15	1.41	0.10
13	67499094	rs34948729	A	0.98	1.40	1×10^{-5}	1.57	0.11	1.39	4×10^{-4}	1.34	0.15	1.41	0.10
13	67502603	rs9571693	G	0.98	1.42	6×10^{-6}	1.53	0.14	1.39	5×10^{-4}	1.43	0.09	1.50	0.06
13	67503594	rs9571694	A	0.99	1.41	1×10^{-5}	1.45	0.20	1.39	7×10^{-4}	1.47	0.07	1.46	0.08
16	51567231	rs10521237	G	0.69	1.51	5×10^{-6}	1.52	0.14	1.42	2×10^{-3}	1.52	0.11	1.94	5×10^{-3}
16	51572611	rs67018781	T	0.83	1.50	6×10^{-6}	1.53	0.13	1.44	1×10^{-3}	1.43	0.16	1.82	0.01
20	55990405	rs737092	C	0.49	1.34	5×10^{-6}	1.60	0.03	1.28	2×10^{-3}	1.23	0.29	1.60	4×10^{-3}
20	55991637	rs6014993	G	0.59	1.32	9×10^{-6}	1.59	0.03	1.27	3×10^{-3}	1.20	0.34	1.50	8×10^{-3}

CHR, chromosome; BP, basepair; dbSNPrs#, rs-number as provided by dbSNP; A1, effect allele; P_{het} , P-value indicating whether statistical significant heterogeneity was present among the 4 cohorts; OR, odds ratio; P_{OR} , P-value for odds ratio; OR_a , odds ratio adjusted for the first principal component; P_a , P-value for adjusted odds ratio; Dutch GSA, Dutch cohort genotyped on the Global Screening arrays; Dutch Omni, Dutch cohort genotyped on the Human OmniExpress BeadChips; German, German cohort; Polish, Polish cohort.

Discussion

Here, we report the first GWAS aiming to identify susceptibility loci in patients with PUV. GWASs in 4 different cohorts revealed one genome-wide significant locus in the German cohort on chromosome 16. However, the strong signal could not be replicated in the other cohorts (see [Supplementary Figures 1–4](#); [Supplementary Table 1](#)). Although not genome-wide statistically significant, the cohorts genotyped on Global Screening Arrays (the German, Polish, and Dutch GSA cohorts) all show the same direction of effect, whereas the largest cohort in this study, the Dutch Omni cohort, showed the opposite effect direction. Whether this array-specific correlation might be caused by a genotyping effect or is due to the different

sample sizes remains speculative. Still, a batch effect driving the very strong German signal seems unlikely as the German and Polish patients were processed at the same time, physically sharing the same genotyping chips. Nevertheless, with one cohort showing an opposite effect direction, we consider this locus not to be a PUV-risk locus, albeit its strong positive signal in the German cohort.

In the meta-analysis of the 4 GWASs, no genome-wide significant association was identified. We would like to emphasize that we used very conservative selection criteria only including variants that passed quality control in all 4 cohorts. While the usage of multiple cohorts and thus the exclusion of many variants might be one limitation of this study, it also guaranteed that only high-quality data were used in the final




FIGURE 2

LocusZoom plots for the loci on chromosome 13, 16, and 20 with multiple variants residing within <10 kb of each other showing suggestive significance ($P < 1 \times 10^{-5}$) and with the same effect direction in all 4 cohorts. **(A)** Locus Zoom plot for chr. 13, with LD scoring referring to top SNP rs9571693. Red line and red quarter indicate rs34844007 in locus zoom plot and in position weight matrix ENCFF525TXJ targeted by POU class family genes. **(B)** Locus Zoom plot for chr. 16, with LD scoring referring to top SNP rs10521237. Heat map visualization of Hi-C interactions among parts of Chr.16q12.1 and parts of 16q12.2 from HMEC human kidney cells, as plotted by TADKB. Blue lines indicate the region plotted in the locus zoom, showing the identified locus and SALL1 to locate in the same topological associated domain. **(C)** Locus Zoom plot for chr. 20, with LD scoring referring to top SNP rs737092. **(D)** Forest plots for rs9571693, rs10521237, and rs737092, the leading SNPs of the loci on chromosome 13, 16, and 20, respectively.

analysis, excluding the possibility that batch effects or signals driven by a single cohort could lead to false positive findings in the meta-analysis. Hence, technical issues that can occur by using just a single genotyping platform could be excluded by the usage of multiple genotyping platforms. The fact that we did not observe genome-wide significant results may indicate that common variants do not play a major role in the etiology of PUV. Of note, by excluding variants with an allele count < 20 in the single cohorts, rare variants were excluded because of the low number of patients and controls, especially in the Dutch GSA ($N = 372$ individuals) and the Polish ($N = 418$ individuals) cohorts. Our lack of statistically significant results could also be explained by the relatively small number of patients and the resulting lack of power, which is a further limitation of this study and caused by the fact that PUV are a rare disease. With 756 patients and 4,823 controls in a single cohort, we would have had 80% power to detect loci with an allele frequency of 0.10 and an OR of 1.3. ORs of this magnitude are generally not observed for multifactorial disorders, although previous GWASs of similar size for other rare birth defects such as hypospadias, bladder extrophy and cleft lip and palate were able to identify loci of genome-wide significance (19–21).

We identified 3 loci with multiple variants showing suggestive significance ($P < 1 \times 10^{-5}$) and with the same effect direction in all 4 cohorts. We will discuss these loci here in the realization that our results are mainly to be regarded as suggestive evidence for association and that larger studies are needed to confirm our findings. In the first locus on chromosome 13, 5 variants reached P -values below 1×10^{-5} . The other loci, located on chromosomes 16 and 20 (Table 1), harbored 2 variants each with a P -value below 1×10^{-5} .

Chromosome 13

The signal on chromosome 13 is located within the first intron of the *PCDH9* gene (Figure 2A). Previous studies showed associations with body mass index, adult body size, and urinary potassium excretion (GWAS catalog)⁵ None of these phenotypes can be linked directly with PUV. The *PCDH9* gene encodes a member of the protocadherin family, a group of proteins that mediate cell adhesion in neural tissues in the presence of calcium (22). *PCDH9* may be involved in signaling at neuronal synaptic junctions. Hence, expression in adult tissues has been found strongest in the brain, while weak expression was found in nearly all organs including the urinary tract and prostate⁶. In the EMBL-EBI expression atlas (23), experiment # E-MTAB-6592 shows constant expression of *PCDH9* in developing human urinary bladder and genital tissues through gestational weeks 7–9. Approximately one third of genes that were found to be expressed in this experiment, however, showed higher

expression than *PCDH9*. Still, *PCDH9* might be involved in urinary tract development.

Direct functional importance of the locus is predicted in the Regulome Database⁷ (24), which gives regulatory probability scores ranging from 0 to 1, with 1 being most likely to be a regulatory variant, generated using a machine learning approach. Rs34844007, one of the 5 associated variants, has a probability score of 1 and is therefore very likely to be a regulatory variant. This variant is located within a transcription factor binding motif which is targeted by 16 members of the POU gene family (Figure 2A). Genes of this family have been shown to be crucially involved in the regulation of developmental processes in many tissues and organs (25). For example, Rieger et al. (26) showed in mice that a missense mutation in *Pou3f*, one of the genes predicted to target the transcription factor binding motif, reduces nephron numbers and impairs development of the thick ascending limb of the Loop of Henle. Nevertheless, the chromatin state in all investigated embryonic cell lines in the Regulome DB is indicative of low or quiescent transcription. Thus, association of the specific transcription factor binding motif with the development of the lower urinary tract awaits further evaluation.

Chromosome 16

The 2 variants with suggestive significance located on chromosome 16 are located within an intergenic region (Figure 2B). Located 40 kb upstream from our top variants, the GWAS catalog lists a genome-wide significant association with the use of diuretics and the use of agents acting on the renin-angiotensin system (top variant rs62039768, p -value 1×10^{-8} , location chr16:51526850) (27), but no functional annotation was given in this publication. Both medications are frequently used in patients with chronic kidney disease, which is often seen in PUV patients. The closest protein coding gene, located 14kb downstream of our top locus, is *HNRNPA1P48*. This gene is a pseudogene to *HNRNPA1*, variants of which have been implicated in amyotrophic lateral sclerosis, a phenotype not associated with LUTO (28). The pseudogene is likely to be expressed in the nucleus (www.proteinatlas.org) and little is known about its functional implications. The locus is flanked by a second gene, *SALL1*, located 387 kb upstream the lead SNP at this locus. According to existing data of topological associated domains (TAD) within several human adult and embryonic cell types (HMEC, NHEK, IMR-90, HUVEC, data source)⁸, our top locus locates in one topologically associated domain with *SALL1* (Figure 2B). This is particularly interesting as autosomal dominantly inherited variants in *SALL1* are known to cause Townes-Brocks syndrome (MIM# 107480), with genitourinary malformations being one of the syndromes features (29).

⁵ <https://www.ebi.ac.uk/gwas/>

⁶ <http://www.proteinatlas.org> version 20.1.

⁷ <https://regulomedb.org>

⁸ <http://dna.cs.miami.edu/TADKB/>

Especially as PUV have been reported in single cases of Townes-Brocks syndrome (30, 31). Functional studies could shed more light on this correlation.

Chromosome 20

The signal on chromosome 20 is also located within an intergenic region (Figure 2C). Both variants reaching suggestive significance in this locus have been described in several GWASs and a meta-analysis focusing on erythrocyte traits and erythropoiesis (32). In these studies, no direct functional annotation was provided.

Several genes lie in close proximity to this locus, with *RBM38* being the closest. In mice this gene is described to be weakly expressed during development of genitourinary system, but KO-mice do not show any genitourinary phenotype.⁹ Interestingly, *BMP7*, which locates within 150 kb distance, is described to be expressed in the developing murine urinary tract. Loss of *Bmp7* has been shown to result in arrest of cloacal septation and in severe abnormalities of the morphogenesis of the genital urethra and mesenchyme (33). Furthermore, heterozygous deletion of this region in human, including the here identified risk locus of suggestive significance, was described in a patient with anatomical LUTO and in one patient from Decipher with hydronephrosis (7). As these are only single case descriptions, larger CNV studies on LUTO patients need to prove that heterozygous loss of the region can be causal for the phenotype.

Conclusion

The present GWAS and meta-analysis is the largest genetic study on PUV performed to date. The fact that no genome-wide significant loci could be identified, may indicate that common variants do not play a major role in the etiology of PUV or could be explained by a lack of power. Nevertheless, 3 loci yielded suggestive associations. Future studies are warranted to support these loci. Ideally, by confirming them either, in a replication study of a single very large cohort or by adding more cohorts to the performed meta-analysis to reach genomewide significance. Nevertheless, considering PUV being a rare disease sampling such a cohort will take some time. Nevertheless, in the meantime, loci could be further explored by performing sequencing of the complete locus regions in cases and controls to identify variants in exonic, intronic, or extragenic regions that are overrepresented in cases compared to controls, suggesting them to be implicated in the phenotype.

Data availability statement

The data presented in the study are deposited in the GWAS Catalog (<https://www.ebi.ac.uk/gwas/>), accession number GCP000376.

⁹ <http://www.informatics.jax.org/> version MGI 6.21.

Ethics statement

The Arnhem-Nijmegen Regional Committee on Research Involving Human Subjects approved the AGORA project, the Nijmegen Biomedical Study, and the current study. The board of directors of the Isala clinics and UMCG approved implementation of AGORA in these centers. The German and Polish studies involving human participants were reviewed and approved by Ethikkommission der Friedrich-Alexander-Universität Erlangen-Nürnberg, Krankenhausstraße 12, 91054 Erlangen. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

Author contributions

CM and OB performed GWAS analysis. LS, SS, LdW, SH, JG, JS, and ACH performed genotyping of different patient and control cohorts. LvZ, IvR, JQ, MSt, LK, LdW, AH, JS, MSz, KT-J, PK, GK, AS, MFS, SW, MZ, NR, HR, WF, and ACH recruited families for the study. LvZ and ACH designed and oversaw the entire study and wrote the manuscript together with CM and HR. All authors approved and commented the manuscript.

Funding

This work was supported by a Kolff junior postdoc grant from the Dutch Kidney Foundation (13OKJ36) and a ZonMW-VENI grant from the Netherlands Organization for Scientific Research (91618036) to LvZ, by the German Research Foundation (DFG, RE 1723/4-1) and by the Else-Kröner-Fresenius-Stiftung (EKFS, 2014_A14) to HR and by BONFOR grant O-149.0123, and the Else Kröner-Fresenius-Stiftung and the Eva Luise und Horst Köhler Stiftung—Project No: 2019_KollegSE.04 to ACH. We acknowledge financial support by Deutsche Forschungsgemeinschaft and Friedrich-Alexander-Universität Erlangen-Nürnberg within the funding program “Open Access Publication Funding.”

Acknowledgments

We would like to thank everyone involved in data collection, all patients and parents for their cooperation in this study, and deCODE Genetics (Reykjavik, Iceland) for performing the microarray experiments of the Dutch cohorts. Part of the collaborations were supported by the European Reference Networks eUROGEN and ERKNet and the German society of pediatric nephrology (GPN).

Conflict of interest

Author SH was employed by Life&Brain GmbH.

The remaining authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or

claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher. Provide the complete details for the following references.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fped.2022.988374/full#supplementary-material>

References

- Chapter 4 *Demography of the UK Paediatric Renal Replacement Therapy Population in 2016*. - PubMed. Available online at: <https://pubmed.ncbi.nlm.nih.gov/29991000/> (accessed April 13, 2022).
- Haeri S. Fetal lower urinary tract obstruction (LUTO): a practical review for providers. *Matern Health Neonatol Perinatol.* (2015) 1:26. doi: 10.1186/s40748-015-0026-1
- Brownlee E, Wragg R, Robb A, Chandran H, Knight M, McCarthy L, et al. Current epidemiology and antenatal presentation of posterior urethral valves: outcome of BAPS CASS National Audit. *J Pediatr Surg Februar.* (2019) 54:318–21. doi: 10.1016/j.jpedsurg.2018.10.091
- Hanlon-Lundberg KM, Verp MS, Loy G. Posterior urethral valves in successive generations. *Am J Perinatol Januar.* (1994) 11:37–9. doi: 10.1055/s-2007-994532
- Schreuder MF, van der Horst HJR, Bökenkamp A, Beckers GMA, van Wijk JAE. Posterior urethral valves in three siblings: a case report and review of the literature. *Birt Defects Res Clin Mol Teratol April.* (2008) 82:232–5. doi: 10.1002/bdra.20439
- Frese S, Weigert A, Hoppe B, Feldkötter M, Ludwig M, Weber S, et al. A classic twin study of lower urinary tract obstruction: report of 3 cases and literature review. *Low Urin Tract Symptoms April.* (2019) 11:O85–8. doi: 10.1111/luts.12222
- Boghossian NS, Sicko RJ, Kay DM, Rigler SL, Caggana M, Tsai MY, et al. Rare copy number variants implicated in posterior urethral valves. *Am J Med Genet A März.* (2016) 170:622–33. doi: 10.1002/ajmg.a.37493
- Caruana G, Wong MN, Walker A, Heloury Y, Webb N, Johnstone L, et al. Copy-number variation associated with congenital anomalies of the kidney and urinary tract. *Pediatr Nephrol Berl Ger März.* (2015) 30:487–95. doi: 10.1007/s00467-014-2962-9
- Lakshmi NK, Narasimhan KL, Khullar M, Madhu K, Kaur B, Balpinder K, et al. Association of angiotensin converting enzyme and angiotensin type 2 receptor gene polymorphisms with renal damage in posterior urethral valves. *J Pediatr Urol Dezember.* (2010) 6:560–6. doi: 10.1016/j.jpuro.2010.01.002
- Kolvenbach CM, Dworschak GC, Frese S, Japp AS, Schuster P, Wenzlitschke N, et al. Rare variants in BNC2 are implicated in autosomal-dominant congenital lower urinary-tract obstruction. *Am J Hum Genet 2 Mai.* (2019) 104:994–1006. doi: 10.1016/j.ajhg.2019.03.023
- van Rooij IALM, van der Zanden LFM, Bongers EMHF, Renkema KY, Wijers CHW, Thonissen M, et al. AGORA, a data- and biobank for birth defects and childhood cancer. *Birt Defects Res A Clin Mol Teratol.* (2016) 106:675–84. doi: 10.1002/bdra.23512
- Manders P, Lutomski JE, Smit C, Swinkels DW, Zielhuis GA. The radboud biobank: a central facility for disease-based biobanks to optimise use and distribution of biomaterial for scientific research in the Radboud University Medical Center, Nijmegen. *Open J Biosour.* (2018) 5:2. doi: 10.5334/ojb.36
- Galesloot TE, Vermeulen SH, Swinkels DW, de Vegt F, Franke B, den Heijer M, et al. Cohort profile: the Nijmegen biomedical study (NBS). *Int J Epidemiol.* (2017) 46:1099–1100j. doi: 10.1093/ije/dyw268
- Schermund A, Möhlenkamp S, Stang A, Grönemeyer D, Seibel R, Hirche H, et al. Assessment of clinically silent atherosclerotic disease and established and novel risk factors for predicting myocardial infarction and cardiac death in healthy middle-aged subjects: rationale and design of the Heinz Nixdorf RECALL Study Risk Factors, Evaluation of Coronary Calcium and Lifestyle. *Am Heart J.* (2002) 144:212–8. doi: 10.1067/mhj.2002.123579
- Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. Robust relationship inference in genome-wide association studies. *Bioinforma Oxf Engl.* (2010) 26:2867–73. doi: 10.1093/bioinformatics/btq559
- Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A global reference for human genetic variation. *Nature.* (2015) 526:68–74. doi: 10.1038/nature15393
- Delaneau O, Howie B, Cox AJ, Zagury JF, Marchini J. Haplotype estimation using sequencing reads. *Am J Hum Genet.* (2013) 93:687–96. doi: 10.1016/j.ajhg.2013.09.002
- Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet.* (2009) 5:e1000529. doi: 10.1371/journal.pgen.1000529
- van der Zanden LFM, van Rooij IALM, Feitz WFJ, Knight J, Donders ART, Renkema KY, et al. Common variants in DGKK are strongly associated with risk of hypospadias. *Nat Genet.* (2011) 43:48–50. doi: 10.1038/ng.721
- Zhang R, Knapp M, Suzuki K, Kajioka D, Schmidt JM, Winkler J, et al. ISL1 is a major susceptibility gene for classic bladder exstrophy and a regulator of urinary tract development. *Sci Rep.* (2017) 7:42170. doi: 10.1038/srep42170
- van Rooij IA, Ludwig KU, Welzenbach J, Ishorst N, Thonissen M, Galesloot TE, et al. Non-syndromic cleft lip with or without cleft palate: genome-wide association study in Europeans identifies a suggestive risk locus at 16p121 and supports SH3PXD2A as a clefting susceptibility gene. *Genes.* (2019) 10:E1023. doi: 10.3390/genes10121023
- Strehl S, Glatt K, Liu QM, Glatt H, Lalande M. Characterization of two novel protocadherins (PCDH8 and PCDH9) localized on human chromosome 13 and mouse chromosome 14. *Genomics.* (1998) 53:81–9. doi: 10.1006/geno.1998.5467
- Athar A, Füllgrabe A, George N, Iqbal H, Huerta L, Ali A, et al. ArrayExpress update - from bulk to single-cell expression data. *Nucleic Acids Res.* (2019) 47:D711–5. doi: 10.1093/nar/gky964
- Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, et al. Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res.* (2012) 22:1790–7. doi: 10.1101/gr.137323.112
- Ryan AK, Rosenfeld MG. POU domain family values: flexibility, partnerships, and developmental codes. *Genes Dev.* (1997) 11:1207–25. doi: 10.1101/gad.11.10.1207
- Rieger A, Kemter E, Kumar S, Popper B, Aigner B, Wolf E, et al. Missense mutation of POU domain class 3 transcription factor 3 in Pou3f3L423P mice causes reduced nephron number and impaired development of the thick ascending limb of the loop of Henle. *PLoS ONE.* (2016) 11:e0158977. doi: 10.1371/journal.pone.0158977
- Wu Y, Byrne EM, Zheng Z, Kemper KE, Yengo L, Mallett AJ, et al. Genome-wide association study of medication-use and associated disease in the UK Biobank. *Nat Commun.* (2019) 10:1891. doi: 10.1038/s41467-019-09572-5
- Kim HJ, Kim NC, Wang YD, Scarborough EA, Moore J, Diaz Z, et al. Mutations in prion-like domains in hnRNPA2B1 and hnRNPA1 cause multisystem

proteinopathy and ALS. *Nature*. (2013) 495:467–73. doi: 10.1038/nature11922

29. Miller EM, Hopkin R, Bao L, Ware SM. Implications for genotype-phenotype predictions in Townes-Brocks syndrome: case report of a novel SALL1 deletion and review of the literature. *Am J Med Genet A*. (2012) 158A:533–40. doi: 10.1002/ajmg.a.34426

30. Kurnit DM, Steele MW, Pinsky L, Dibbins A. Autosomal dominant transmission of a syndrome of anal, ear, renal, and radial congenital malformations. *J Pediatr*. (1978) 93:270–3. doi: 10.1016/S0022-3476(78)80518-6

31. Salerno A, Kohlhase J, Kaplan BS. Townes-Brocks syndrome and renal dysplasia: a novel mutation in the SALL1 gene. *Pediatr Nephrol Berl Ger*. (2000) 14:25–8. doi: 10.1007/s004670050006

32. Vuckovic D, Bao EL, Akbari P, Lareau CA, Mousas A, Jiang T, et al. The polygenic and monogenic basis of blood traits and diseases. *Cell*. (2020) 182:1214–31.e11. doi: 10.1016/j.cell.2020.08.008

33. Wu X, Ferrara C, Shapiro E, Grishina I. Bmp7 expression and null phenotype in the urogenital system suggest a role in re-organization of the urethral epithelium. *Gene Expr Patterns GEP*. (2009) 9:224–30. doi: 10.1016/j.gep.2008.12.005