

甲状腺TSH受容体の精製，分布高圧構造とその臨床的意義

著者	水上 勇治
著者別表示	Mizukami Yuji
雑誌名	昭和62(1987)年度 科学研究費補助金 一般研究(C) 研究成果報告書概要
巻	1986 1987
ページ	2p.
発行年	1989-03-29
URL	http://doi.org/10.24517/00067915



1987 Fiscal Year Final Research Report Summary

Purification, distribution and clinical significance of TSH-receptor in human thyroid.

Research Project

Project/Area Number

61570166

Research Category

Grant-in-Aid for General Scientific Research (C)

Allocation Type

Single-year Grants

Research Field

Experimental pathology

Research Institution

Kanazawa University

Principal Investigator

MIZUKAMI Yuji Kanazawa University Hospital, Pathology Section, 医学部附属病院・病理部, 講師 (60110540)

Co-Investigator(Kenkyū-buntansha)

HASHIMOTO Takuma Kanazawa University Hospital, Central Clinical Laboratory, 医学部附属病院・検査部, 助教授 (10124710)

Project Period (FY)

1986 - 1987

Keywords

TSH-receptor / Ras-oncogene product / EGF / EGF-receptor

Research Abstract


We have attempted to produce an antibody for human TSH-receptor. Using this antibody, distribution and changes in distribution of TSH-receptor was examined. First, we attempted to purify the TSH-receptor antibody using the serum of patients with high titer of TSH-receptor antibody. The serum containing high titer of TSH-receptor antibody and porcine thyroid membrane fraction were incubated. The binding IgG was released by acidic buffer and released IgG was further purified with IgG affinity chromatography. Using this IgG the localization of TSH-receptor was examined in human thyroid follicular cells. The positive staining was observed at the apical plasma membrane. The only one band was found at Mol. wt 130,000 using the immunoblotting. We also examined the ultrastructural localization of ras p21 product, which participates on GTP regulation of adenylate cyclase. Similar finding that ras p21 product was localized at apical cell membrane was obtained. Epidermal growth factor (EGF) and EGF-receptor were also examined immunohistochemically in


human thyroid. These combined evidences strongly suggested that TSH-receptor is restricted at apical plasma membrane of thyroid follicular cells. The changes in distribution of TSH-receptro in various thyroid diseases have been studied at present.

Research Products (2 results)

All Other

All Publications (2 results)

[Publications] Mizukami Y;Nonomura A;Hashimoto T;Terahata S;Matsubara F;Michigishi T;Noguchi M: CANCER. 61. (1988) 

[Publications] Mizukami Y, Nonomura A, Hashimoto T, Terahata S, Matsubara F, Michigishi T, Noguchi M.: "Immunohistochemical demonstration of ras p21 oncogene product in normal, benign, and malignant human thyroid tissues." Cancer. 61. (1988) 

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-61570166/615701661987kenkyu_seika_hokoku_

Published: 1989-03-29