

抗精神病薬の局所脳血流と脳代謝に及ぼす影響について-臨床的・実験的研究

著者	倉知 正佳
著者別表示	Kurachi Masayoshi
雑誌名	昭和62(1987)年度 科学研究費補助金 一般研究(C) 研究成果報告書概要
巻	1985 1987
ページ	2p.
発行年	1989-03-29
URL	http://doi.org/10.24517/00067941



1987 Fiscal Year Final Research Report Summary

The effect of antipsychotics on regional cerebral blood flow and metabolim - clinical and experimetal study.

Research Project

Project/Area Number

60570496

Research Category

Grant-in-Aid for General Scientific Research (C)

Allocation Type

Single-year Grants

Research Field

Psychiatric science

Research Institution

Toyama medical and pharmaceutical University (1986-1987)
Kanazawa University (1985)

Principal Investigator

KURACHI Masayoshi Toyama medical and pharmaceutical Univ., 医学部, 教授 (80019603)

Co-Investigator(Kenkyū-buntansha)

MORI Hirohumi Kanazawa Univ. Radioisotope Center, アイソトープ総合センター, 助教授 (90019604)

Project Period (FY)

1985 - 1987

Keywords

¹³³Xe inhalation / regional cerebral blood flow / haloperidol / ¹²⁵I-IMP / オートラジオグラフ

Research Abstract

1. Clinical study

To study the effect of antipsychotics on cerebral blood flow, we examined regional cerebral blood flow (rCBF) using ¹³³Xe inhalation technique in eleven patients (9 schizophrenics and 2 schizophreniform disorder; mean age, 22.9 years) before and about two weeks after medication. The measurements were also made with a two-week interval for ten normal volunteers without medication (mean age, 31.2 years). RCBF measurements showed good reproducibility in the normal controls. their mean hemispheric blood flow being between 49.8-51.4 ml/100g/min. RCBF in the patients after two-week's medication (haloperidol 2.25-6.0 mg/day or sulpiride 300mg/day) showed a statistically significant decrease in the left frontal and right temporo-parietal region

compared with that before treatment, but their mean hemispheric blood flow only showed a tendency to decrease. On the Brief Psychiatric Rating Scale, the score of suspiciousness has significantly decreased.

2. Experimental Study

The acute effect of antipsychotics on rCBF in 37 discrete regions of the rat brain was studied by the quantitative autoradiographic N-isopropyl-p-[125-I]-Iodoamphetamine technique. There was no significant difference in mean hemispheric blood flow between controls (172.1ml/ 100g/min) and haloperidol 0.1 or 1.0 mg/kg i.v. administered groups (180.8; 164.5). For the regional distribution value of the blood flow, bilateral habenular nucleus showed a significant increase, whereas, medial prefrontal cortex and nucleus accumbens showed a tendency to decrease. The effect of chronic haloperidol administration is not yet completed, but these clinical and experimental studies would contribute to the understanding of the antipsychotic effects of neuroleptics.

Research Products (2 results)

All Other

All Publications (2 results)

[Publications] Kurachi M, et al. edited by Takahashi et al: "Regional cerebral blood flow in patients with schizophrenic disorders. In: Cerebral dynamics, laterality and psychopathology" Elsevier, Amsterdam, 493-501 (1987) ▼

[Publications] Kurachi et al.;: Regional cerebral blood flow in patients with schizophrenic disorders : In : Cerebral dynamics, laterality and psychopathology (edited by Takahashi, et al). Elsevier, Amsterdam, pp9 493-501 (1987) ▼

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-60570496/605704961987kenkyu_seika_hokoku_

Published: 1989-03-29